THE UNIVERSITY OF RHODE ISLAND

University of Rhode Island DigitalCommons@URI

Biomedical and Pharmaceutical Sciences Faculty Publications

Biomedical and Pharmaceutical Sciences

9-2012

Pharmacy Research at URI: Mining Red Maple (Acer rubrum) Trees for Novel Therapeutics to Manage Diabetes

Navindra P. Seeram University of Rhode Island, nseeram@uri.edu

Jialin Xu

Liya Li

Angela Slitt University of Rhode Island, angela_slitt@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/bps_facpubs

Citation/Publisher Attribution

Seeram, N., Xu, J., Li, L., & Slitt, A. (2012). Pharmacy Research at URI: Mining Red Maple (*Acer rubrum*) Trees for Novel Therapeutics to Manage Diabetes. *Medicine & Health Rhode Island*, *95*(9), 283-284. Retrieved from http://www.rimed.org/medhealthri/2012-09/2012-09-283.pdf. Available at: http://www.rimed.org/medhealthri/2012-09/2012-09-283.pdf

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Biomedical and Pharmaceutical Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Pharmacy Research at URI: Mining Red Maple (Acer rubrum) Trees for Novel Therapeutics to Manage Diabetes

Terms of Use All rights reserved under copyright.

This article is available at DigitalCommons@URI: https://digitalcommons.uri.edu/bps_facpubs/8

Pharmacy Research at URI: Mining Red Maple (Acer rubrum) Trees for Novel Therapeutics to Manage Diabetes

Navindra P. Seeram, PhD, Jialin Xu, PhD, Liya Li, PhD, and Angela Slitt, PhD

Type 2 DIABETES MELLITUS (T2DM) accounts for about 90% of all diagnosed cases of diabetes in adults. Over 200 million people suffer from this disease worldwide. In the United States alone, in 2007, 10% of American adults had diabetes and the cost of management was \$174 billion and this figure is expected to skyrocket (Centers for Disease Control and Prevention, 2010). Plants and their derived products have been used for centuries by various cultures as traditional medicines for the management of diabetes. Plants contain secondary metabolites (known as phytochemicals; 'phyto' means plant), which are implicated in the prevention and treatment of several chronic human diseases, including diabetes. Among these natural products, polyphenols and phenolic glycosides, have attracted significant interests for their anti-diabetic properties.

Plant polyphenols as α -glucosidase inhibitors. Dietary carbohydrates are hydrolyzed by pancreatic α -amylase with absorption aided by α -glucosidases and thus, inhibition of the activities of these enzymes is a promising approach for managing T2DM. In fact, the clinical a-glucosidase inhibitor, acarbose, has been shown to effectively reduce glycated hemoglobin levels when given as monotherapy or as an add-on to other antidiabetic drug treatment.1 Additionally, a fixed dose combination of acarbose and metformin was reported to be superior to metformin alone in controlling HbA1c, fasting blood glucose, and post prandial blood glucose levels in T2DM patients,² as well as, enhancing the blood glucose control.3 Multiple plant-derived polyphenols have been investigated as *a*-glucosidase inhibitors. Extracts from grape seeds and green tea been demonstrated to exert inhibition on α-glucosidase activity.4 We recently have shown a similar activity for phenolic-enriched extracts from the bark of Acer rubrum, commonly known as red maple.

Study of maple for novel therapeutics. The sugar maple (*Acer saccharum*) and red maple plant species are native to eastern North America and highly regarded for their sap which is used to produce the natural sweetener, maple syrup. Our laboratory has recently identified a number of phenolic compounds native in maple plant parts that are also present in maple syrup.⁵

Here we report a case study in which the ability of red maple bark extract (MBE) to decrease elevation of blood glucose levels was investigated. Male adult C57BL/6 mice were administered MBE in combination with a bolus sucrose challenge to evaluate the effect of MBE or acarbose on α -glucosidase activity in vivo according to described methods.6 As expected, mice receiving a bolus sucrose challenge increased blood glucose concentration 30, 60, and 90 minutes after administration compared to initial blood glucose concentration. (Figure 1) Both MBE and acarbose significantly lowered blood glucose concentrations at these time intervals after the sucrose challenge. Compared to acarbose, MBE was less effective, but did impart a significant inhibitory effect. We now hope to further purify and evaluate individual phytochemicals of MBE to evaluate for activity against sucrose-induced glucose elevation.

We have demonstrated that phenolic glycosides are present in various maple constituents. The isolated compounds and extracts inhibit α -glucosidase activity in vitro.7,8 Recently, multiple gallotannins, named maplexins A-E, were isolated from red maple stems and bark.7,8 These maplexin compounds were shown in vitro to possess more activity than acarbose for inhibition of α -glucosidase activity (IC₅₀ = 8 vs. 160 µM; maplexin E vs. acarbose, respectively). These interesting maplexin compounds, that are as effective as acarbose in vitro, will be further evaluated and developed to determine whether they impart beneficial properties in preventing blood glucose elevations in vivo. Overall, we determined that MBE contains bioactive constituents that can potentially aid



Figure 1. Adult male C57BL/6 mice were administered sucrose (3g/kg BW), sucrose and acarbose (3g/kg BW + 3 mg/kg BW), or 3) sucrose and maple bark extract (3g/kg BW+150mg/kg BW) by oral gavage. Blood glucose levels were determined by tail bleed at 0, 30, 60, 90, and 120 minutes post sucrose administration. An asterisk (*) denotes statistical significance (p<0.05) from the sucrose treatment group.

283

in the regulation of blood glucose control after a carbohydrate challenge.

We will continue to explore how bioactive extracts and molecules isolated from plants can impart beneficial effects on metabolic syndrome. As metabolic syndrome encompasses T2DM, dyslipidemia, systemic inflammation, and nonalcoholic fatty liver disease, the potential therapeutics isolated will be tested in various cell-based and in vivo models. We are currently exploring anti-inflammatory effects of maple-derived extracts and compounds, as well as, the potential for plant-derived extracts to augment weight loss and T2DM therapies. Ultimately, we hope that identification of beneficial components in foods commonly consumed can arm the consumer with information to make better food choices that can be synergistic with drug therapies used to treat T2DM. We also will continue to mine botanicals for potential therapies to treat T2DM or enhance activity of commonly prescribed anti-diabetes therapies.

Funding Sources

This project was supported by funding provided by Agriculture and Agri-Food Canada through the **Developing Innovative Agri Products (DIAP)** initiative and, in part, by The National Institute of Health (ES-016042-03 and ES-016042-2S2). The authors would like to thank Mr. J. Peter Morgan for assistance with plant authentication.

REFERENCES

- Derosa G, Maffioli P. Efficacy and safety profile evaluation of acarbose alone and in association with other antidiabetic drugs: a systematic review. *Clin Ther.* Jun 2012;34(6):1221–36.
- Jayaram S, Hariharan RS, Madhavan R, Periyandavar I, Samra SS. A prospective, parallel group, open-labeled, comparative, multi-centric, active controlled study to evaluate the safety, tolerability and benefits of fixed dose combination of acarbose and metformin versus metformin alone in type 2 diabetes. J Assoc Physicians India. Nov 2010;58:679–82, 687.
- Schnell O, Mertes G, Standl E. Acarbose and metabolic control in patients with type 2 diabetes with newly initiated insulin therapy. *Diabetes Obes Metab.* Nov 2007;9(6):853–8.
- Yilmazer-Musa M, Griffith A, Michels AJ, Schneider E, Frei B. Grape seed and tea extracts and catechin 3-gallates are potent inhibitors of alpha-amylase and alpha-glucosidase activity. J Agric Food Chem. Jun 15 2012.
- Li L, Seeram NP. Further investigation into maple syrup yields 3 new lignans, a new phenylpropanoid, and 26 other phytochemicals. J Agric Food Chem. Jul 27 2011;59(14):77–16.
- Honma A, Koyama T, Yazawa K. Anti-hyperglycaemic effects of the Japanese red maple Acer pycnanthum and its constituents the ginnalins B and C. J Enzyme Inhib Med Chem. Apr 2011;26(2):176–80.
- Wan C, Yuan T, Li L, et al. Maplexins, new alpha-glucosidase inhibitors from red maple (Acer rubrum) stems. *Bioorg Med Chem Lett.* Jan 1 2012;22(1):597–600.
- Yuan T, Wan C, Liu K, Seeram NP. New maplexins F-I and phenolic glycosides from red maple (Acer rubrum) bark. *Tetrahedron*. 2012;68:959–64.

Navindra P. Seeram, PhD, is an Assistant Professor in the Department of Biomedical and Pharmaceutical Sciences at the University of Rhode Island.

Jialin Xu, PhD, is a Postdoctoral Fellow, Slitt, in the Department of Biomedical and Pharmaceutical Sciences at the University of Rhode Island.

Liya Li, PhD, is a Postdoctoral Fellow, Slitt, in the Department of Biomedical and Pharmaceutical Sciences at the University of Rhode Island.

Angela Slitt, PhD, is an Assistant Professor in the Department of Biomedical and Pharmaceutical Sciences at the University of Rhode Island.

Disclosure of Financial Interests

The authors and/or their spouses/ significant others have no financial interests to disclose.

CORRESPONDENCE

Angela Slitt, PhD Department of Biomedical and Pharmaceutical Sciences College of Pharmacy University of Rhode Island 7 Greenhouse Road Kingston, RI 02881 phone: (401) 874-5020 e-mail: aslitt@uri.edu



Copyright of Medicine & Health Rhode Island is the property of Rhode Island Medical Society and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.