

2015

# A plasmid-based reporter system for live cell imaging of dengue infected cells

Carey L. Medin

University of Rhode Island, carey\_medin@uri.edu

Sierra Valois

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.uri.edu/immunology\\_facpubs](https://digitalcommons.uri.edu/immunology_facpubs)

**The University of Rhode Island Faculty have made this article openly available.  
Please let us know how Open Access to this research benefits you.**

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

## Citation/Publisher Attribution

Carey L. Medin, Sierra Valois, Chinmay G. Patkar, Alan L. Rothman, A plasmid-based reporter system for live cell imaging of dengue virus infected cells, Journal of Virological Methods, Volume 211, January 2015, Pages 55-62, ISSN 0166-0934, <http://dx.doi.org/10.1016/j.jviromet.2014.10.010>. (<http://www.sciencedirect.com/science/article/pii/S0166093414004005>)

This Article is brought to you for free and open access by the Institute for Immunology and Informatics (iCubed) at DigitalCommons@URI. It has been accepted for inclusion in Institute for Immunology and Informatics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

---

**Authors**

Carey L. Medin, Sierra Valois, Chinmay G. Patkar, and Alan Rothman

1 A plasmid-based reporter system for live cell imaging of dengue infected cells.

2 Running title: live cell imaging of dengue virus infection

3

4 Carey L. Medin<sup>1#</sup>, Sierra Valois<sup>1</sup>, Chinmay G. Patkar<sup>2</sup>, and Alan L. Rothman<sup>1,2\*</sup>

5

6 <sup>1</sup>Institute for Immunology and Informatics, Department of Cell and Molecular Biology,

7 University of Rhode Island, Providence, RI 02903

8

9 <sup>2</sup>Division of Infectious Diseases and Immunology, Department of Medicine, University of

10 Massachusetts Medical School, Worcester, MA 01655

11 \* Corresponding author:

12 Carey L. Medin, Ph.D.

13 Institute for Immunology and Informatics

14 80 Washington St., Room 334

15 Providence, RI 02903

16 Tel: (401) 277-5252

17 FAX: (401) 277-5154

18 e-mail: cmedin.uri@gmail.com

19

20 Abstract: 109 words

21 Text: 3560 words

22

23

1 Abstract

2 Cell culture models are used widely to study the effects of dengue virus (DENV)  
3 on host cell function. Current methods of identification of cells infected with an  
4 unmodified DENV requires fixation and permeablization of cells to allow DENV-specific  
5 antibody staining. This method does not permit imaging of viable cells over time. In this  
6 report, a plasmid-based reporter was developed to allow non-destructive identification of  
7 DENV-infected cells. The plasmid-based reporter was demonstrated to be broadly  
8 applicable to the four DENV serotypes, including low-passaged strains, and was  
9 specifically cleaved by the viral protease with minimal interference on viral production.  
10 This study reveals the potential for this novel reporter system to advance the studies of  
11 virus-host interactions during DENV infection.

12 Keywords

13 Dengue virus; viral reporter; protease; fluorescence microscopy

## 1 1. Introduction

2           Dengue virus (DENV) is a mosquito-borne human pathogen of global medical  
3 importance (Bhatt et al., 2013; Simmons et al., 2012). DENV causes an acute febrile  
4 illness that, in some patients, is associated with a life-threatening plasma leakage  
5 syndrome, dengue hemorrhagic fever (DHF) (Anonymous, 2009). While there is ongoing  
6 debate regarding the contribution of different mechanisms in dengue illness, there is  
7 substantial evidence supporting both viral and host factors in disease pathogenesis.  
8 Most models propose a cascade involving systemic infection of cells of the myeloid cell  
9 lineage and cytokine production through both direct (infected cells) and indirect  
10 (bystander and immune cells) pathways (Halstead et al., 2010; Rothman, 2010;  
11 Whitehorn and Farrar, 2010). There is ongoing scientific need and interest in examining  
12 the direct effects of DENV infection on cellular metabolic processes and gene  
13 expression, which will give insight into the initial events that may impact disease  
14 severity.

15           Given the limitations of clinical studies and existing animal models, cell culture  
16 models remain an important approach to studying DENV infection and host responses  
17 (Becerra et al., 2009; Chase et al., 2011; Heaton and Randall, 2010; Hibberd et al.,  
18 2006; Pena and Harris, 2012; Ramirez-Ortiz et al., 2006). These studies have involved  
19 both continuous cell lines and primary cells from human and arthropod hosts. The  
20 analysis of host cell responses to infection has been accomplished through imaging,  
21 immunoassays, and RT-PCR. These methods are subject to at least two technical  
22 limitations- difficulty in discriminating infected and uninfected cells in the same culture,  
23 and collection of static data from individual time points. Detection of individual DENV-  
24 infected cells generally relies on immunostaining. Because most DENV proteins are not  
25 expressed on the cell surface, sensitive detection of infected cells requires fixation and  
26 permeabilization of the cells, an approach that is incompatible with ongoing observation

1 of living cells and potentially introduces artifactual changes in cellular ultrastructure. One  
2 approach to investigate early events during DENV infection in living cells used virus  
3 labeled with a lipophilic fluorescent probe that fluoresces upon viral fusion with  
4 membranes of acidified vesicles. This approach is limited to analysis of events prior to  
5 viral fusion (van der Schaar et al., 2008; van der Schaar et al., 2007). Several groups  
6 have reported recently the construction of DENV replicons or infectious viruses that  
7 encode reporter molecules such as luciferase or GFP (Leardkamolkarn and Sirigulpanit,  
8 2012; Leardkamolkarn et al., 2012; Mattia et al., 2011; Schoggins et al., 2012; Yang et  
9 al., 2013; Zou et al., 2011). These constructs can effectively label cells infected with the  
10 engineered virus or replicon. However, recombinant modified DENV genomes are  
11 invariably attenuated in comparison to wild-type DENV, which is likely to complicate the  
12 interpretation of virus-host interactions. Furthermore, these strategies will require further  
13 extensive manipulation to permit the study of other DENV variants, limiting the  
14 generalizability of the findings.

15 To facilitate studies of live cells infected with DENV, a novel plasmid-based  
16 reporter, p4B5-EGFP, was created that utilize the viral protease cleavage site resulting  
17 in nuclear localization of GFP in DENV-infected cells. The 4B5-EGFP reporter was  
18 shown to be effective in detecting infection of cells by all four DENV serotypes, including  
19 low-passaged viral strains. The results demonstrate the capability of the 4B5-EGFP  
20 reporter to identify DENV-infected cells without destruction of the cell.

## 21 2. Materials and Methods

### 22 2.1. Cell lines and virus

23 Vero cells were obtained from the ATCC and maintained in Dulbecco's  
24 Modification of Eagle's Medium (DMEM) supplemented with 10% fetal bovine serum,  
25 100 U/ml penicillin and 100 U/ml streptomycin, with cell media also containing

1 nonessential amino acid (1X, HyClone). Cells were incubated in a humidified chamber at  
2 37 °C and 5% CO<sub>2</sub>.

3         Prototype DENV strains DENV-1 Hawaii, DENV-2 16681, DENV-2 New Guinea  
4 C (NGC), DENV-3 CH53489, and DENV-4 814669 were originally obtained from the  
5 American Type Culture Collection and the Walter Reed Army Institute of Research, and  
6 were passaged multiple times in C6/36 cells (ATCC, USA). Low passaged DENV-2  
7 strains C0102/96 and C0112/96 were isolated from plasma collected in 1996 as part of a  
8 prospective study (Vaughn et al., 2000); these viruses were initially isolated by  
9 intrathoracic injection of *Toxorhynchites splendens* mosquitoes and then passaged up to  
10 4 times in C6/36 cells. Virus titers were determined by immunostained plaque assay on  
11 Vero cells based on the method of Liu et al with minor modifications (Liu et al., 2012).  
12 Briefly, Vero cells (1×10<sup>5</sup> cells in 50 µl/well) were added to replicate wells of 96-well  
13 white-bottom plates with 50 µl of serial 0.5 log dilutions of virus. Plates were incubated  
14 for 2 h and then 100 µl of overlay containing 1% carboxymethylcellulose was added.  
15 Plates were stained after 3 d incubation using anti-DENV antibody MAB8705 (EMD  
16 Millipore, Billerica, MA, 1:1000), horseradish peroxidase-conjugated anti-mouse Ig  
17 (Southern Biotech, 1:2000), and TMB substrate (Mabtech, Cincinnati, OH). Stained  
18 regions were read using an ELISpot plate reader to give focus-forming units per ml  
19 (ffu/ml). The ffu/ml was log transformed and graphed using Graph Pad Prism 6.0  
20 software.

## 21 **2.2. Construction of the DENV reporter plasmid**

22         The DENV reporter plasmid, p4B5-EGFP, was constructed to encode the full-  
23 length DENV-2 NS4B protein (without sequences encoding the 2k peptide) and the first  
24 10 amino acids of the DENV-2 NS5 protein fused to the SV40 nuclear localization signal  
25 sequence (NLS, PKKKRKVG (Cressman et al., 2001)) and the enhanced GFP (EGFP)

1 protein in the pcDNA3.1 vector (Life Technologies, Grand Island, NY). The primers used  
2 for PCR synthesis are shown in Table 1. The DENV sequences were originally amplified  
3 from a DENV-2 NGC infectious clone, which was kindly provided by Dr. Barry Falgout  
4 (Polo et al., 1997). A plasmid generated in our lab containing DENV-2 sequences from  
5 nucleotides 6757 to 7599, which includes NS4B and the first 30 nucleotides of NS5, was  
6 used to insert the SV40 NLS and GFP sequences downstream of the NS4B-5 cleavage  
7 site. Briefly, to generate a fragment containing the SV40 NLS upstream of GFP, a  
8 forward primer 'NLSGFP-EcoRI' that incorporated a 5' EcoRI restriction site and the  
9 SV40 NLS sequence and the reverse primer 'GFP XhoI' that contained a 3'XhoI  
10 restriction site were used to amplify from the pTRE-eGFP plasmid (Clontech) by PCR.  
11 The PCR fragment was digested with EcoRI and XhoI, gel purified, and ligated into the  
12 vector downstream of nucleotide 7599. To generate the p4B5-EGFP, the 'NS4B HindIII'  
13 forward primer and the 'GFP XhoI' reverse primer was used to amplify the reporter  
14 sequence by PCR. The product of the PCR reaction and pcDNA 3.1 (Life Technologies,  
15 Grand Island, NY) were then digested with HindIII and XhoI, gel purified and ligated  
16 together. The identities of the clones were confirmed by DNA sequencing.

17 The plasmid pNS2B3 expressing the DENV-2 NS2B3 protease was constructed  
18 using DENV-2 NGC RNA as a template. Sense and antisense primers (Table 1) were  
19 designed to generate a cDNA fragment encompassing nucleotides 4132 to 6375 of  
20 DENV-2 NGC using SuperScript® One-Step RT-PCR for long templates (Life  
21 Technologies, Grand Island, NY). The PCR fragment and the pcDNA3.1 V5-His vector  
22 (Life Technologies, Grand Island, NY) were digested with HindIII and XbaI, gel purified  
23 and ligated together. The identities of the clones were confirmed by DNA sequencing.

### 24 **2.3. Transfection and DENV infection**

25 Vero cells were transfected using GeneJuice® Transfection Reagent (EMD  
26 Millipore, Billerica, MA) following the manufacturer's instructions. Briefly, cells were



1 seeded in an 8-chambered Nunc Lab-Tek slide (Thermo Fisher Scientific, Rockford, IL)  
2 with a glass coverslip bottom at  $2 \times 10^4$  cells per well 24 hrs prior to transfection. For  
3 transfection, 1.2  $\mu$ l of GeneJuice® Transfection Reagent was diluted in 15  $\mu$ l serum-free  
4 media and incubated at room temperature for 5 minutes, and then 0.55  $\mu$ g of plasmid  
5 were added to the diluted GeneJuice® Transfection Reagent and incubated for 15  
6 minutes at room temperature. The complex was then added to the cells. Vero cells were  
7 infected with DENV at a multiplicity of infection of 1 as previously described (Medin and  
8 Rothman, 2006).

9 For cotransfection with p4B5-EGFP and pNS2B3, Vero cells were transfected  
10 with 22.5  $\mu$ g of each plasmid.

#### 11 **2.4. Western Blot**

12 Whole cell extracts were prepared using lysis buffer (10% glycerol, 20 mM Tris  
13 (pH 7), 150 mM NaCl, 0.5 mM EDTA, 1% Nonidet P-40) freshly supplemented with a  
14 protease inhibitor cocktail (Sigma-Aldrich, St. Louis, MO) and 25 U of the Pierce  
15 Universal Nuclease (Thermo Fisher Scientific, Rockford, IL). Cells were placed on ice for  
16 15 min and centrifuged for 15 min at 14,000 rpm. Protein concentrations were  
17 determined using Bradford reagent in the Envision plate reader (Perkin Elmer, Waltham,  
18 Massachusetts) using bovine serum albumin as a standard. 15  $\mu$ g of protein were  
19 separated using a 12% SDS-PAGE precast gel (Biorad, Hercules, CA) and then  
20 transferred onto nitrocellulose membranes using the Trans-Blot® Turbo™ Transfer  
21 System (Biorad, Hercules, CA). Blots were blocked with 5% skim milk and subjected to  
22 immunoblot analysis using anti-GFP (Thermo Fisher Scientific, Rockford, IL, cat. 22169)  
23 diluted 1:3000 and anti- $\beta$ -actin (Abcam, Cambridge, MA, cat. 8226) diluted 1:5000  
24 antibodies as indicated in the figure legends.

## 1 **2.5. Staining and immunofluorescence microscopy**

2 In experiments analyzing nuclear localization of GFP, nuclear DNA was stained  
3 immediately prior to imaging with NucBlue™ Live Cell Stain or NucBlue™ Fixed Cell  
4 Stain following the manufacturer's protocol (Life Technologies, Grand Island, NY). For  
5 detection of ER membranes, cells were washed with 1X DPBS and stained with 500nM  
6 ER-Tracker Blue-White DPX solution (Life Techonologies, Grand Island, NY) for 30  
7 minutes following the manufacturer's protocol. Staining solution was replaced with cell  
8 culture media and immediately imaged by fluorescence microscopy using a DAPI filter.

9 Fluorescence images were obtained using an EVOS® fl digital inverted  
10 fluorescence microscope (Advanced Microscopy Group, USA). To obtain a time series of  
11 images of nuclear localization of GFP after DENV infection, Vero cells transfected with  
12 p4B5-EGFP were infected with DENV-2 16681. The wells were washed after two hours  
13 and fresh media was added to each well. The plate was placed into Cytation3 Cell  
14 Imager (Biotek, Winooski, VT) that contains a temperature and CO<sub>2</sub> controlled chamber.  
15 The images were acquired every 45 minutes starting at 4 h after adding virus to the cell  
16 culture using a 20x objective.

17 Individual cells were analyzed for total cellular and nuclear GFP fluorescence  
18 using imageJ's Analyze and Measure function. Nuclear and cytoplasmic GFP  
19 fluorescence was defined by setting parameters based on NucBlue staining of the  
20 nucleus. Background fluorescence was measured from three regions outside of the cell.  
21 The product of the average background and area of the cell was subtracted from the  
22 integrated density (sum of the intensity of the pixels) for both the total fluorescence and  
23 nuclear fluorescence giving a corrected fluorescence as described previously (Gavet  
24 and Pines, 2010). The ratios of nuclear to total fluorescence were calculated.

25 Total fluorescence data were not distributed normally necessitating use of  
26 nonparametric methods. Wilcoxon-Mann-Whitney rank sum test was used to quantify

1 differences between groups. Data presented includes analyses of individual cells with n  
2 = 24-36 cells. All statistical analyses were performed utilizing Graph Pad Prism 6.0  
3 software.

4 To confirm DENV infection, cells were stained for DENV antigen at 24 and 48 hrs  
5 post-infection. Cells were washed once with PBS, fixed with 4% paraformaldehyde for  
6 30 min at room temperature, and permeabilized with Cytoperm (BD Biosciences,  
7 Franklin Lakes, NJ) for 15 min at 4°C. DENV antigen was detected by indirect  
8 immunofluorescence staining using DENV complex-specific monoclonal antibody (EMD  
9 Millipore, Billerica, MA, cat. MAB8705) diluted 1:300 as the primary antibody and  
10 allophycocyanin (APC)-conjugated goat anti-mouse immunoglobulin G antibody (BD  
11 Biosciences, Franklin Lakes, NJ, cat. 558026) diluted 1:100 as the secondary antibody.

12 To confirm NS2B3 expression, cells were stained for NS3 at 48hrs post  
13 transfection using an anti-NS3 primary antibody (Genetex, Irvine, CA ) and APC-  
14 conjugated goat anti-mouse IgG as the secondary antibody.

### 15 3. Results

#### 16 **3.1. Construction and characterization of DENV reporter plasmid**

17 The DENV reporter, p4B5-EGFP, was designed to take advantage of two events  
18 that occur during DENV infection- localization of the DENV NS4B protein to the  
19 endoplasmic reticulum (ER) and proteolytic processing of the cleavage site between  
20 NS4B and NS5 by the NS2B3 protease (Bera et al., 2007; Miller et al., 2006) (Figure  
21 1a). The processing of the cleavage site by NS2B3 is expected to release the GFP  
22 component (with the small fragment of NS5) to relocate from the cytoplasm to the  
23 nucleus.

24 The expression and cellular distribution of the reporter construct in living cells  
25 was first evaluated by transfecting cells with the reporter plasmid and staining with an  
26 ER membrane stain. In the absence of infection with DENV (Figure 1b), transfected cells

1 showed cytoplasmic green fluorescence that colocalized with the ER membrane stain,  
2 indicating ER retention of the reporter.

3 To determine whether infection with DENV would induce proteolytic cleavage  
4 between NS4B and NS5(10) and mobilize GFP to the nucleus, cells were transfected  
5 with p4B5-EGFP and infected with DENV-2 16681. Figure 1c shows a DENV-infected  
6 Vero cell culture expressing 4B5-EGFP and stained with the nuclear dye NucBlue for  
7 live-cell labeling (Invitrogen). The two cells show different phenotypes of GFP  
8 localization at 24hrs after infection, with one cell showing cytoplasmic GFP (arrowhead)  
9 and the other cell showing nuclear GFP (arrow). Analysis of nuclear to total cellular  
10 fluorescence for individual cells at 24 h showed at least a 2.4 fold increase in infected  
11 cells when compared to uninfected cells and were significantly different ( $p=0.0001$ )  
12 (Figure 1d). These results confirmed that GFP localized to the nucleus in infected cells.  
13 Since these images were taken on a standard fluorescence microscope, the small  
14 number of cells in the uninfected 24h condition showing an increase in fluorescence  
15 intensity was due to the fluorescence bleed from cytoplasmic region into the nuclear  
16 region.

17 To verify cleavage of the 4B5-EGFP, Western blot analysis was performed on  
18 cell lysates using an antibody against GFP. As seen in Figure 1e, uninfected cells  
19 transfected with p4B5-EGFP showed a single band localizing to approximately 60kD,  
20 which is the expected size of the expressed fusion protein. In cells infected with DENV-2  
21 at an m.o.i. of 1 for 24 hours, the 60kD band was reduced in intensity and a lower 30kD  
22 band appeared. The 30kD band corresponds to the expected size of the NS5 fragment  
23 fused to eGFP. These results confirm that the 4B5-EGFP is cleaved during DENV  
24 infection.

### 25 **3.2. Kinetics of nuclear localization of GFP after DENV infection**

1 To determine the earliest time point that the 4B5-EGFP reporter could detect  
2 DENV infection, transfected cells infected with DENV were analyzed by fluorescence  
3 microscopy. Relocation of green fluorescence into the nucleus was seen as early as 8 h  
4 post-infection (data not shown). Time-lapse images of Vero cells in a DENV-infected  
5 culture were taken to further determine the time required to visually identify GFP  
6 movement from the cytoplasm to the nucleus (Figure 2). Movement of cytoplasmic GFP  
7 to the nucleus was visible in four of the five cells in Figure 2 by 7 h after addition of virus.  
8 One cell remained uninfected during this analysis (Figure 2, arrowhead). Calculation of  
9 the ratio of nuclear to total fluorescence at each time point for a representative cell  
10 (Figure 2, arrow) showed successive increase in nuclear fluorescence over time.

### 11 **3.3. Nuclear GFP correlates with DENV antigen staining**

12 To verify that the cells displaying nuclear localization of GFP were infected with  
13 DENV, cells were fixed, permeabilized and stained using a DENV-specific monoclonal  
14 antibody. Figure 3a shows a pair of cells in a DENV-infected cell culture. The arrow  
15 shows a cell with nuclear GFP, indicating viral infection, whereas the arrowhead shows a  
16 cell with only cytoplasmic GFP, indicating no viral infection. As seen in the third and  
17 fourth panels of Figure 3a, the bottom left cell (with nuclear GFP) stained with the anti-  
18 DENV antibody whereas the upper cell (with only cytoplasmic GFP) did not.

### 19 **3.4. Expression of the DENV NS2B-3 protease is sufficient for cleavage of 4B5- 20 EGFP**

21 Previous reports have shown that both NS2B and NS3 are required for efficient  
22 proteolytic cleavage of the DENV polyprotein (Brinkworth et al., 1999; Falgout et al.,  
23 1991). A construct that expresses DENV-2 NS2B3 was created to determine if the  
24 NS2B3 protease alone could induce nuclear localization of GFP. Cells were transfected  
25 with the p4B5-EGFP alone or cotransfected with the NS2B3 plasmid. As seen in Figure  
26 3b, cells transfected with p4B5-EGFP showed cytoplasmic GFP expression only. In

1 contrast, cells expressing NS2B3 and 4B5-EGFP showed nuclear localization of GFP  
2 (Figure 3c). In these experiments, nuclear localization was detected 48 and 72 hs post-  
3 transfection. This delay in protease activity may reflect a lower level of protease  
4 expression from the plasmid compared to DENV-infected cells. Alternatively, NS2B3  
5 may require additional viral factors to cleave efficiently the NS4B-5 junction.

### 6 **3.5. DENV serotype and strain specificity of the 4B5-EGFP reporter**

7 The viral gene segment in p4B5-EGFP was derived from DENV-2 NGC, a  
8 laboratory-adapted virus originally isolated in 1944 and was tested using DENV-2 16681,  
9 another laboratory-adapted virus that was isolated in 1964 (Vaughn et al., 1996). Each  
10 DENV serotype can be subdivided into specific genotypes, which can have up to 6%  
11 genetic divergence (Rico-Hesse, 1990; Twiddy et al., 2002). To assess whether the  
12 construct would be processed by other DENV-2 strains, low passaged recent clinical  
13 isolates were used to infect cells. Transfected cells were infected with two viral isolates,  
14 C0102/96 and C0112/96, which were originally isolated from the plasma of patients with  
15 dengue in 1996. Nuclear localization of GFP and DENV antigen staining was analyzed  
16 at 24 hours post-infection. As seen with DENV-2 16681, DENV antigen staining  
17 correlated with nuclear GFP for both C0102/96 (data not shown) and C0112/96 (Figure  
18 4b).

19 The four serotypes of DENV have up to 70% sequence homology across their  
20 genomes (Blok, 1985). Previous reports have shown that the DENV protease is less  
21 selective than the proteases of other flaviviruses and can tolerate amino acid changes  
22 within the cleavage site (Shiryaev et al., 2007). Additionally, the NS2B3 protease  
23 cleaves at least five sites with little sequence homology within the same DENV  
24 polyprotein (Bera et al., 2007). Therefore, to analyze whether this reporter plasmid could  
25 be used to detect infection with other DENV serotypes, cells transfected with the reporter  
26 plasmid were infected with each of the four DENV serotypes. For all four serotypes,

1 nuclear localization of GFP was detected at 24 hours post-infection (Figure 4). The  
2 specificity of nuclear GFP for DENV-infected cells was verified by co-staining for DENV  
3 antigen (red).

#### 4 **3.6. Effects of 4B5-EGFP on DENV replication**

5 Full length NS4B contains a cleavable 2K signal peptide at the N terminus, and  
6 has been shown to be required for IFN antagonism (Munoz-Jordan et al., 2005). The  
7 NS4B in the pNS4B5-EGFP was designed to be expressed without the 2K region and  
8 therefore, should have no effect on inhibiting the antiviral response. To determine  
9 whether transfection of the p4B5-EGFP had an effect on viral production compared to  
10 untransfected cells, viral titers in supernatants of DENV-infected Vero cells transfected  
11 with p4B5-EGFP or a control plasmid, pTracer, expressing GFP alone were assessed by  
12 plaque assay. Viral titers in virally infected cells without transfection were compared to  
13 cells that were transfected. Since Vero cells do not have an intact innate immune system  
14 (Mosca and Pitha, 1986), parallel experiments with CV-1 cells, which do elicit an innate  
15 immune response to dsRNA was also performed (Thacore et al., 1990). Cells  
16 transfected with pNS4B5-EGFP and pTracer showed a 3 to 7-fold reduction in viral titers  
17 when compared to infected cells that were untransfected depending on the serotype  
18 used for infection (Figure 5). There were minimal differences in viral titers between cells  
19 transfected with either the control plasmid or cells transfected with p4B5-EGFP. These  
20 results indicate that transfection alone reduced viral titers. Additionally, cells transfected  
21 with p4B5-EGFP and pTracer were comparable at both 24 and 48h post-infection in both  
22 Vero and CV-1 cells indicating that the reporter did not affect viral replication (data not  
23 shown).

#### 24 4. Discussion

25 In the present study, a DENV-specific plasmid-based reporter system, 4B5-  
26 EGFP, was characterized for detection of infected cells in live cell culture for all four

1 DENV serotypes. Although systems for analyzing DENV replication in culture have  
2 expanded in recent years, detection methods applicable to imaging and detection of  
3 cells infected with native viral isolates in unfixed cells have not been available. The 4B5-  
4 EGFP reporter does not require genetic modification of the viral genome.

5 The 4B5-EGFP reporter also showed detection of cellular infection with each of  
6 the DENV serotypes. Previous reports have analyzed sequence variation within the  
7 NS4B-NS5 junction between various flaviviruses (Bera et al., 2007; Lin et al., 1993;  
8 Shiryayev et al., 2007). The amino acids in the P1, P2 and P1' positions are similar for all  
9 four DENV serotypes as well as Yellow Fever virus, however, amino acids occupying P2'  
10 and P3' are different (Shiryayev et al., 2007). Those experiments were done with  
11 synthetically produced peptides; the role of surrounding amino acids and membrane  
12 tethering of the proteins may be an additional factor in the selectivity of trans cleavage  
13 by flaviviral proteases (Clum et al., 1997). Further analysis using other flaviviruses would  
14 be useful to establish the specificity of this reporter system.

15 The advantage of using the 4B5-EGFP reporter the detection of GFP can be  
16 carried out with fluorescence microscopy. Additionally, the ability to monitor infection by  
17 fluorescence relocalization in living cells also suggests the possibility of continuous  
18 visualization of DENV infection. The 4B5-EGFP reporter was proteolytically processed  
19 as early as 8 hrs post-infection, which correlates with synthesis of new viral protein  
20 (Shrivastava et al., 2011). The ability to detect viral infection during initial translation of  
21 the viral genome will extend analysis of cellular events to early time points.

22 Numerous studies have demonstrated that in vitro infection with DENV induces  
23 significant changes in cellular gene expression and metabolic processes (Becerra et al.,  
24 2009; Chase et al., 2011; Heaton and Randall, 2010; Hibberd et al., 2006; Pena and  
25 Harris, 2012; Ramirez-Ortiz et al., 2006). The experiments conducted in those studies  
26 processed unsorted cultures of cells exposed to DENV. Therefore, the effects measured



1 in those studies reflect both direct effects of viral infection on the infected cell as well as  
2 secondary effects on both infected and uninfected cells in the same culture (Nightingale  
3 et al., 2008; Palmer et al., 2005). The 4B5-EGFP reporter in this study has the potential  
4 to identify effects on individually infected and uninfected cells within the same culture.  
5 Live-cell analysis of virus-infected cells by fluorescence microscopy represents a  
6 promising approach to investigate virus-cell interactions (Coller et al., 2012; Jones et al.,  
7 2010; Lee-Huang et al., 2011; Paloheimo et al., 2011; Rand et al., 2012; Sivaraman et  
8 al., 2011; van der Schaar et al., 2008; van der Schaar et al., 2007). This approach has  
9 been used for other viruses to study virion entry and assembly and the dynamics of  
10 movement of viral proteins and cell structures within the infected cell. For several  
11 viruses, live cell imaging has revealed new mechanisms critical to viral replication or cell-  
12 to-cell spread and has been used for antiviral drug screening.

13         A theoretical concern with this approach was that the DENV protein segment  
14 expressed by the plasmid might itself affect cellular function. For example, DENV NS4B  
15 has been reported to inhibit type I IFN signaling (Munoz-Jordan et al., 2003). Munoz-  
16 Jordan et al reported, however, that deletion of the N-terminal 2K segment eliminated  
17 the inhibition of IFN without affecting ER localization (Munoz-Jordan et al., 2005). The  
18 4B5-EGFP reporter plasmid did not contain the 2K segment, and, consistent with the  
19 published results, was not different from a control plasmid in its effects on DENV  
20 replication. Viral replication was lower in transfected cells than in untransfected cells,  
21 likely due at least in part to induction of type I IFN (Park et al., 2003). A cell line stably  
22 transfected with the 4B5-EGFP reporter would be expected to overcome this limitation.

23         The reagents developed in this project have some limitations for other types of  
24 analyses. To this effect the 4B5-EGFP transfected cells fluoresce green regardless of  
25 infection status, therefore the cells are not amenable to cell sorting for downstream  
26 analyses. Since eGFP is a commonly used fluorescence molecule, exchanging eGFP for

1 a less commonly used color such as BFP or YFP will permit multicolor fluorescence  
2 target analyses. Additionally, there are potential applications of the 4B5-EGFP reporter  
3 in drug screening and the development of novel animal or cell culture models of DENV  
4 infection.

5 The 4B5-EGFP reporter described is a promising strategy for identifying live  
6 DENV-infected cells by fluorescence microscopy. Potential applications of the 4B5-  
7 EGFP reporter include identifying infectious virus directly from patient samples. To this  
8 effect, nuclear GFP was seen in transfected cell cultures infected with low-passaged  
9 isolates.

#### 10 Acknowledgements

11 We thank Drs. Kate Fitzgerald and Shanaka Rodrigo for helpful discussions,  
12 Diane Lang, Susannah Colt and Jurand Janus for technical assistance, Dr. Barry  
13 Falgout for providing the DENV-2 infectious clone and Dr. Jennifer Friedman for  
14 statistical analysis. This work was supported by NIH grants P01 AI034533, P20  
15 GM103430-12 and U01 AI070484.

1 References

- 2 Anonymous, 2009. Dengue: guidelines for diagnosis, treatment, prevention and control -  
3 new edition. World Health Organization, Geneva.
- 4 Becerra, A., Warke, R.V., Martin, K., Xhaja, K., de Bosch, N., Rothman, A.L., Bosch, I.,  
5 2009. Gene expression profiling of dengue infected human primary cells identifies  
6 secreted mediators in vivo. *Journal of medical virology* 81, 1403-1411.
- 7 Bera, A.K., Kuhn, R.J., Smith, J.L., 2007. Functional characterization of cis and trans  
8 activity of the Flavivirus NS2B-NS3 protease. *The Journal of biological chemistry* 282,  
9 12883-12892.
- 10 Bhatt, S., Gething, P.W., Brady, O.J., Messina, J.P., Farlow, A.W., Moyes, C.L., Drake,  
11 J.M., Brownstein, J.S., Hoen, A.G., Sankoh, O., Myers, M.F., George, D.B., Jaenisch, T.,  
12 Wint, G.R., Simmons, C.P., Scott, T.W., Farrar, J.J., Hay, S.I., 2013. The global  
13 distribution and burden of dengue. *Nature* 496, 504-507.
- 14 Blok, J., 1985. Genetic relationships of the dengue virus serotypes. *The Journal of*  
15 *general virology* 66 ( Pt 6), 1323-1325.
- 16 Brinkworth, R.I., Fairlie, D.P., Leung, D., Young, P.R., 1999. Homology model of the  
17 dengue 2 virus NS3 protease: putative interactions with both substrate and NS2B  
18 cofactor. *The Journal of general virology* 80 ( Pt 5), 1167-1177.
- 19 Chase, A.J., Medina, F.A., Munoz-Jordan, J.L., 2011. Impairment of CD4+ T cell  
20 polarization by dengue virus-infected dendritic cells. *J Infect Dis* 203, 1763-1774.
- 21 Clum, S., Ebner, K.E., Padmanabhan, R., 1997. Cotranslational membrane insertion of  
22 the serine proteinase precursor NS2B-NS3(Pro) of dengue virus type 2 is required for  
23 efficient in vitro processing and is mediated through the hydrophobic regions of NS2B.  
24 *The Journal of biological chemistry* 272, 30715-30723.

1 Coller, K.E., Heaton, N.S., Berger, K.L., Cooper, J.D., Saunders, J.L., Randall, G., 2012.  
2 Molecular determinants and dynamics of hepatitis C virus secretion. *PLoS Pathog* 8,  
3 e1002466.

4 Cressman, D.E., O'Connor, W.J., Greer, S.F., Zhu, X.S., Ting, J.P., 2001. Mechanisms  
5 of nuclear import and export that control the subcellular localization of class II  
6 transactivator. *Journal of immunology* 167, 3626-3634.

7 Falgout, B., Pethel, M., Zhang, Y.M., Lai, C.J., 1991. Both nonstructural proteins NS2B  
8 and NS3 are required for the proteolytic processing of dengue virus nonstructural  
9 proteins. *Journal of virology* 65, 2467-2475.

10 Gavet, O., Pines, J., 2010. Progressive activation of CyclinB1-Cdk1 coordinates entry to  
11 mitosis. *Developmental cell* 18, 533-543.

12 Halstead, S.B., Mahalingam, S., Marovich, M.A., Ubol, S., Mosser, D.M., 2010. Intrinsic  
13 antibody-dependent enhancement of microbial infection in macrophages: disease  
14 regulation by immune complexes. *Lancet Infect Dis* 10, 712-722.

15 Heaton, N.S., Randall, G., 2010. Dengue virus-induced autophagy regulates lipid  
16 metabolism. *Cell Host Microbe* 8, 422-432.

17 Hibberd, M.L., Ling, L., Tolfvenstam, T., Mitchell, W., Wong, C., Kuznetsov, V.A.,  
18 George, J., Ong, S.H., Ruan, Y., Wei, C.L., Gu, F., Fink, J., Yip, A., Liu, W., Schreiber,  
19 M., Vasudevan, S.G., 2006. A genomics approach to understanding host response  
20 during dengue infection. *Novartis Found Symp* 277, 206-214; discussion 214-207, 251-  
21 203.

22 Jones, C.T., Catanese, M.T., Law, L.M., Khetani, S.R., Syder, A.J., Ploss, A., Oh, T.S.,  
23 Schoggins, J.W., MacDonald, M.R., Bhatia, S.N., Rice, C.M., 2010. Real-time imaging of  
24 hepatitis C virus infection using a fluorescent cell-based reporter system. *Nat Biotechnol*  
25 28, 167-171.

1 Leardkamolkarn, V., Sirigulpanit, W., 2012. Establishment of a stable cell line  
2 coexpressing dengue virus-2 and green fluorescent protein for screening of antiviral  
3 compounds. *Journal of biomolecular screening* 17, 283-292.

4 Leardkamolkarn, V., Sirigulpanit, W., Chotiwan, N., Kumkate, S., Huang, C.Y., 2012.  
5 Development of Dengue type-2 virus replicons expressing GFP reporter gene in study of  
6 viral RNA replication. *Virus research* 163, 552-562.

7 Lee-Huang, S., Lin Huang, P., Lee Huang, P., 2011. Live-cell real-time imaging reveals  
8 role of mitochondria in cell-to-cell transmission of HIV-1. *Biochem Biophys Res Commun*  
9 415, 384-389.

10 Lin, C., Chambers, T.J., Rice, C.M., 1993. Mutagenesis of conserved residues at the  
11 yellow fever virus 3/4A and 4B/5 dibasic cleavage sites: effects on cleavage efficiency  
12 and polyprotein processing. *Virology* 192, 596-604.

13 Liu, L., Wen, K., Li, J., Hu, D., Huang, Y., Qiu, L., Cai, J., Che, X., 2012. Comparison of  
14 plaque- and enzyme-linked immunospot-based assays to measure the neutralizing  
15 activities of monoclonal antibodies specific to domain III of dengue virus envelope  
16 protein. *Clin Vaccine Immunol* 19, 73-78.

17 Mattia, K., Puffer, B.A., Williams, K.L., Gonzalez, R., Murray, M., Sluzas, E., Pagano, D.,  
18 Ajith, S., Bower, M., Berdough, E., Harris, E., Doranz, B.J., 2011. Dengue reporter virus  
19 particles for measuring neutralizing antibodies against each of the four dengue  
20 serotypes. *PloS one* 6, e27252.

21 Medin, C.L., Rothman, A.L., 2006. Cell type-specific mechanisms of interleukin-8  
22 induction by dengue virus and differential response to drug treatment. *The Journal of*  
23 *infectious diseases* 193, 1070-1077.

24 Miller, S., Sparacio, S., Bartenschlager, R., 2006. Subcellular localization and membrane  
25 topology of the Dengue virus type 2 Non-structural protein 4B. *The Journal of biological*  
26 *chemistry* 281, 8854-8863.

1 Mosca, J.D., Pitha, P.M., 1986. Transcriptional and posttranscriptional regulation of  
2 exogenous human beta interferon gene in simian cells defective in interferon synthesis.  
3 *Molecular and cellular biology* 6, 2279-2283.

4 Munoz-Jordan, J.L., Laurent-Rolle, M., Ashour, J., Martinez-Sobrido, L., Ashok, M.,  
5 Lipkin, W.I., Garcia-Sastre, A., 2005. Inhibition of alpha/beta interferon signaling by the  
6 NS4B protein of flaviviruses. *Journal of virology* 79, 8004-8013.

7 Munoz-Jordan, J.L., Sanchez-Burgos, G.G., Laurent-Rolle, M., Garcia-Sastre, A., 2003.  
8 Inhibition of interferon signaling by dengue virus. *Proceedings of the National Academy*  
9 *of Sciences of the United States of America* 100, 14333-14338.

10 Nightingale, Z.D., Patkar, C., Rothman, A.L., 2008. Viral replication and paracrine effects  
11 result in distinct, functional responses of dendritic cells following infection with dengue 2  
12 virus. *Journal of leukocyte biology* 84, 1028-1038.

13 Palmer, D.R., Sun, P., Celluzzi, C., Bisbing, J., Pang, S., Sun, W., Marovich, M.A.,  
14 Burgess, T., 2005. Differential effects of dengue virus on infected and bystander  
15 dendritic cells. *J Virol* 79, 2432-2439.

16 Paloheimo, O., Ihalainen, T.O., Tauriainen, S., Valilehto, O., Kirjavainen, S., Niskanen,  
17 E.A., Laakkonen, J.P., Hyoty, H., Vihinen-Ranta, M., 2011. Coxsackievirus B3-induced  
18 cellular protrusions: structural characteristics and functional competence. *J Virol* 85,  
19 6714-6724.

20 Park, M.S., Shaw, M.L., Munoz-Jordan, J., Cros, J.F., Nakaya, T., Bouvier, N., Palese,  
21 P., Garcia-Sastre, A., Basler, C.F., 2003. Newcastle disease virus (NDV)-based assay  
22 demonstrates interferon-antagonist activity for the NDV V protein and the Nipah virus V,  
23 W, and C proteins. *Journal of virology* 77, 1501-1511.

24 Pena, J., Harris, E., 2012. Early dengue virus protein synthesis induces extensive  
25 rearrangement of the endoplasmic reticulum independent of the UPR and SREBP-2  
26 pathway. *PLoS one* 7, e38202.

1 Polo, S., Ketner, G., Levis, R., Falgout, B., 1997. Infectious RNA transcripts from full-  
2 length dengue virus type 2 cDNA clones made in yeast. *Journal of virology* 71, 5366-  
3 5374.

4 Ramirez-Ortiz, Z.G., Warke, R.V., Pacheco, L., Xhaja, K., Sarkar, D., Fisher, P.B., Shaw,  
5 S.K., Martin, K.J., Bosch, I., 2006. Discovering innate immunity genes using differential  
6 display: a story of RNA helicases. *Journal of cellular physiology* 209, 636-644.

7 Rand, U., Rinas, M., Schwerk, J., Nohren, G., Linnes, M., Kroger, A., Flossdorf, M.,  
8 Kaly-Kullai, K., Hauser, H., Hofer, T., Koster, M., 2012. Multi-layered stochasticity and  
9 paracrine signal propagation shape the type-I interferon response. *Mol Syst Biol* 8, 584.

10 Rico-Hesse, R., 1990. Molecular evolution and distribution of dengue viruses type 1 and  
11 2 in nature. *Virology* 174, 479-493.

12 Rothman, A.L., 2010. Cellular immunology of sequential dengue virus infection and its  
13 role in disease pathogenesis. *Current topics in microbiology and immunology* 338, 83-  
14 98.

15 Schoggins, J.W., Dorner, M., Feulner, M., Imanaka, N., Murphy, M.Y., Ploss, A., Rice,  
16 C.M., 2012. Dengue reporter viruses reveal viral dynamics in interferon receptor-  
17 deficient mice and sensitivity to interferon effectors in vitro. *Proc Natl Acad Sci U S A*  
18 109, 14610-14615.

19 Shiryayev, S.A., Kozlov, I.A., Ratnikov, B.I., Smith, J.W., Lebl, M., Strongin, A.Y., 2007.  
20 Cleavage preference distinguishes the two-component NS2B-NS3 serine proteinases of  
21 Dengue and West Nile viruses. *The Biochemical journal* 401, 743-752.

22 Shrivastava, N., Sripada, S., Kaur, J., Shah, P.S., Cecilia, D., 2011. Insights into the  
23 internalization and retrograde trafficking of Dengue 2 virus in BHK-21 cells. *PloS one* 6,  
24 e25229.

25 Simmons, C.P., Farrar, J.J., Nguyen v, V., Wills, B., 2012. Dengue. *The New England*  
26 *journal of medicine* 366, 1423-1432.

1 Sivaraman, D., Biswas, P., Cella, L.N., Yates, M.V., Chen, W., 2011. Detecting RNA  
2 viruses in living mammalian cells by fluorescence microscopy. *Trends Biotechnol* 29,  
3 307-313.

4 Thacore, H.R., Lin, H.Y., Davis, P.J., Schoenl, M., 1990. Effect of protein kinase C  
5 inhibitors on interferon-beta production by viral and non-viral inducers. *The Journal of*  
6 *general virology* 71 ( Pt 12), 2833-2839.

7 Twiddy, S.S., Farrar, J.J., Vinh Chau, N., Wills, B., Gould, E.A., Gritsun, T., Lloyd, G.,  
8 Holmes, E.C., 2002. Phylogenetic relationships and differential selection pressures  
9 among genotypes of dengue-2 virus. *Virology* 298, 63-72.

10 van der Schaar, H.M., Rust, M.J., Chen, C., van der Ende-Metselaar, H., Wilschut, J.,  
11 Zhuang, X., Smit, J.M., 2008. Dissecting the cell entry pathway of dengue virus by  
12 single-particle tracking in living cells. *PLoS Pathog* 4, e1000244.

13 van der Schaar, H.M., Rust, M.J., Waarts, B.L., van der Ende-Metselaar, H., Kuhn, R.J.,  
14 Wilschut, J., Zhuang, X., Smit, J.M., 2007. Characterization of the early events in  
15 dengue virus cell entry by biochemical assays and single-virus tracking. *Journal of*  
16 *virology* 81, 12019-12028.

17 Vaughn, D.W., Green, S., Kalayanarooj, S., Innis, B.L., Nimmannitya, S., Suntayakorn,  
18 S., Endy, T.P., Raengsakulrach, B., Rothman, A.L., Ennis, F.A., Nisalak, A., 2000.  
19 Dengue viremia titer, antibody response pattern, and virus serotype correlate with  
20 disease severity. *The Journal of infectious diseases* 181, 2-9.

21 Vaughn, D.W., Hoke, C.H., Jr., Yoksan, S., LaChance, R., Innis, B.L., Rice, R.M.,  
22 Bhamarapravati, N., 1996. Testing of a dengue 2 live-attenuated vaccine (strain 16681  
23 PDK 53) in ten American volunteers. *Vaccine* 14, 329-336.

24 Whitehorn, J., Farrar, J., 2010. Dengue. *Br Med Bull* 95, 161-173.

25 Yang, C.C., Tsai, M.H., Hu, H.S., Pu, S.Y., Wu, R.H., Wu, S.H., Lin, H.M., Song, J.S.,  
26 Chao, Y.S., Yueh, A., 2013. Characterization of an efficient dengue virus replicon for



1 development of assays of discovery of small molecules against dengue virus. Antiviral  
2 research.

3 Zou, G., Xu, H.Y., Qing, M., Wang, Q.Y., Shi, P.Y., 2011. Development and  
4 characterization of a stable luciferase dengue virus for high-throughput screening.  
5 Antiviral Res 91, 11-19.

6

7

8

1 TABLE 1. Oligonucleotide primers used for PCR amplification.

Oligonucleotide	Sequence <sup>a</sup>
NS4B HindIII F	5'- CATTGGCAA <b>AAGCTT</b> <u>GCCACCATG</u> GCGAACGAGATGGGTTTCCTAGAAAAACGAAG- 3'
NS5(10aa) EcoRI R	5'-CATTCTC <b>GAATTC</b> TCCAAGCGTCTCTCCTATGTTGCCAGTCCCCTTC-3'
SV40NLS-eGFP EcoRI F	5'- CGCG <b>GAATTC</b> <u>GCCACCATG</u> CCGAAGAAAAAGCGGAAGGTTGGCGTGAGCAAGGGC GAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGT-3'
eGFP XhoI R	5'-CGCGCTGC <b>CTCGAG</b> TACTTGTACAGCTCGTCCATGCCGAGAGTGATC-3'
NS2B3 HindIII F	5'-CAAGAAAAGGA <b>AAGCTT</b> <u>GCCACCATG</u> AGCTGGCCATTAATGAGGCTATCATG-3'
NS2B3 XbaI R	5'-GGTCAGAGAT <b>CTAGACT</b> TTTCTCCGGCTGCAAATTC-3'

<sup>a</sup> underlined text = the Kozak sequence, bold text = restriction endonuclease, italics = SV40 NLS

2

### 3 **Figure Legends**

4 Figure 1. Construction and characterization of p4B5-EGFP.

5 (a) Schematic of the p4B5-EGFP construct containing NS4B and the first 30

6 nucleotides of NS5 of the DENV-2 genome tagged with the SV40 NLS and

7 eGFP. The arrow indicates the cleavage site utilized by the DENV NS2B3

8 protease. The vertical lines represent restriction endonuclease sites.

9 (b) Unfixed Vero cells transfected with p4B5-EGFP were stained with ER-Tracker

10 Blue-White DPX dye (Cyan) 24hrs post-transfection and immediately imaged to

11 detect colocalization of NS4B5-EGFP (reporter, green) with ER membranes

12 (blue, magnification, x100). Inset reflects ER staining of cells alone. (c) Unfixed

13 Vero cells transfected with p4B5-EGFP were infected with DENV-2 16681. Nuclei

14 were counterstained with NucBlue for live cells (Invitrogen) (blue) to detect

15 nuclear localization of GFP (green, magnification, x100). Cytoplasmic expression

16 of GFP is indicated by the arrowhead and nuclear expression of GFP is indicated

17 by an arrow (magnification, x100). (d) Analysis of nuclear to total fluorescence

18 intensity ratios of GFP (nuclear:total fl) in uninfected and DENV infected cells at

1 24 h was performed using ImageJ software. Each symbol represents analysis of  
2 a single cell. Statistical analysis was performed using nonparametric Wilcoxon-  
3 Mann-Whitney rank sum test. Straight line represents the median. (e) Cellular  
4 lysates were prepared 24 h post infection and subjected to Western blotting for  
5 GFP or  $\beta$ -actin. The 5-EGFP fragment represents the cleavage product resulting  
6 from cleavage between NS4B and NS5-EGFP during viral infection.

7

8 Figure 2. Kinetics of nuclear localization of GFP after DENV infection. Vero cells  
9 transfected with p4B5-EGFP were infected with DENV-2 16681 and sequential  
10 images were acquired to assess the time for GFP to localize into the nucleus.  
11 Each set of images shows the expression of GFP in p4B5-EGFP transfected  
12 cells (top row), nuclear stain using NucBlue (middle row) and transmitted images  
13 (bottom row) at each time point. Time after addition of virus to the culture is  
14 located at the lower left of each image (magnification, x20). Nuclear/total  
15 fluorescence (N/T) was calculated at each time point for a representative cell  
16 (arrow). The arrowhead shows a cell that remains uninfected over time. Data are  
17 representative of at least five experiments.

18

19 Figure 3. Nuclear localization of GFP correlates with DENV antigen and co-  
20 expression of the NS2B3 protease. (a) Vero cells were transfected with p4B5-  
21 EGFP (green) and infected with DENV-2 16681 at an m.o.i of 1. 24 hours post-  
22 infection, cells were fixed, permeabilized and stained with antibody against  
23 DENV complex. The NS4B5-EGFP panel shows cytoplasmic expression of GFP

1 (right, arrow neighboring cytoplasmic and nuclear expression of GFP (left,  
2 arrowhead) (magnification, x100). DENV Ag panel shows DENV antigen staining  
3 in cells infected with DENV. Overlay of GFP and antigen staining (red) shows that  
4 nuclear localization of GFP correlates with DENV antigen staining (magnification,  
5 x100). (b and c) Vero cells transfected with the p4B5-EGFP alone (b) or  
6 cotransfected with pNS2B3 (c) were analyzed for nuclear localization of GFP at  
7 48hrs post-transfection (magnification, x100). NS4B5-EGFP panel shows  
8 location of GFP within the cells (green). The nucleus panel shows the nucleus  
9 stained with NucBlue (blue, Life Technologies). The NS2B3 panel shows indirect  
10 antibody staining for NS3 (red). The overlay shows that nuclear GFP expression  
11 correlates with NS2B3 expression. Data are representative of at least six (a) and  
12 two (b) experiments.

13

14 Figure 4. All four DENV serotypes induce cleavage of p4B5-EGFP to localize  
15 GFP to the nucleus. Vero cells transfected with p4B5-EGFP (green) were  
16 infected with each of the four DENV serotypes at an MOI of 1. Cells were fixed,  
17 permeabilized and stained for DENV antigen (red) and nuclear DNA (cyan). Cells  
18 were analyzed at 24 hours post-infection by fluorescence microscopy. Each row  
19 is a representative image of DENV infected cells in bright field (trans) and  
20 fluorescence images showing nuclear stain (cyan), DENV antigen stain (red),  
21 4B5-EGFP expression (green). The overlay is a composite of the nucleus, DENV  
22 and NS4B-EGFP images. (a) DENV-1 Hawaii, (b) DENV-2 C0112/96, (c) DENV-3

1 CH53489 and (d) DENV-4 814669. Data are representative of at least four  
2 experiments each.

3

4 Figure 5. Expression of 4B5-EGFP reporter has minimal effects on DENV  
5 replication.

6 Vero cells were transfected with pTracer or p4B5-EGFP or untransfected and  
7 infected at an m.o.i. of 1 with DENV 24h later. After 24 h supernatants were  
8 collected for viral titration. Each time point represents the geometric mean and  
9 95% confidence interval (CI) of six experiments.

10