

University of Rhode Island

DigitalCommons@URI

---

Institute for Immunology and Informatics  
Faculty Publications

Institute for Immunology and Informatics  
(iCubed)

---

12-5-2014

## Tuberculosis interferon-gamma responses in the breast milk of human immunodeficiency virus infected mothers

Lisa Cranmer

Mercy Kanyugo

Barbara Lohman-Payne  
*University of Rhode Island*, b.lohman.payne@gmail.com

Ken Tapia

Grace John-Stewart

Follow this and additional works at: [https://digitalcommons.uri.edu/immunology\\_facpubs](https://digitalcommons.uri.edu/immunology_facpubs)

The University of Rhode Island Faculty have made this article openly available.  
Please let us know how Open Access to this research benefits you.

### Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

---

### Citation/Publisher Attribution

Cranmer, L.M. et al. "Tuberculosis interferon-gamma responses in the breast milk of human immunodeficiency virus infected mothers." *The International Journal of Tuberculosis and Lung Disease*. December 5, 2014. Available: <http://dx.doi.org/10.5588/ijtld.14.0384>

This Article is brought to you for free and open access by the Institute for Immunology and Informatics (iCubed) at DigitalCommons@URI. It has been accepted for inclusion in Institute for Immunology and Informatics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons-group@uri.edu](mailto:digitalcommons-group@uri.edu).

---

## Tuberculosis interferon-gamma responses in the breast milk of human immunodeficiency virus infected mothers

The University of Rhode Island Faculty have made this article openly available.  
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

### Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).



## Tuberculosis IFN- $\gamma$ Responses in Breastmilk of HIV-infected Mothers

Journal:	<i>The International Journal of Tuberculosis and Lung Disease</i>
Manuscript ID:	IJTLD-05-14-0384.R1
Manuscript Type:	Short Communication
Date Submitted by the Author:	n/a
Complete List of Authors:	Cranmer, Lisa; University of Washington, Pediatrics; University of Nairobi, Paediatrics Kanyugo, Mercy; University of Nairobi, Paediatrics Lohman-Payne, Barbara; University of Nairobi, Paediatrics Tapia, Ken; University of Washington, Global Health John-Stewart, Grace; University of Washington, Medicine; University of Washington, Epidemiology; University of Washington, Global Health
Key Words:	TB, IGRA, HIV, maternal, infant

**Title:** Tuberculosis IFN- $\gamma$  Responses in Breastmilk of HIV-infected Mothers

**Authors and affiliations:** Lisa M. Cranmer<sup>1,2,7</sup>, Mercy Kanyugo<sup>2</sup>, Barbara Lohman-Payne<sup>2,3</sup>, Ken Tapia<sup>4,5</sup>, Grace C. John-Stewart<sup>5-8</sup>

<sup>1</sup>Department of Pediatrics, Emory University School of Medicine and Children's Healthcare of Atlanta, Atlanta, Georgia; <sup>2</sup>Department of Paediatrics and Child Health, University of Nairobi, Nairobi, Kenya; <sup>3</sup>Department of Cell and Molecular Biology, University of Rhode Island, Providence, Rhode Island, Departments of <sup>4</sup>Biostatistics, <sup>5</sup>Global Health, <sup>6</sup>Medicine, <sup>7</sup>Pediatrics, and <sup>8</sup>Epidemiology, University of Washington, Seattle, Washington, USA

**Author Contributions:** LC contributed to study design, data analysis, and drafted the manuscript. MK performed laboratory assays and assisted with revision of manuscript. BL-P assisted with study design, oversight of laboratory procedures, and revision of manuscript. KT assisted with data analysis and manuscript revision. GJ-S designed the study and revised the manuscript.

**Funding and Conflicts of Interest:** Supported by the US National Institutes Health NIH HD058477-01. LC was a Fellow of the Pediatric Scientist Development Program (PSDP) supported by an American Pediatric Society/American Academy of Pediatrics grant and NICHD K12-HD000850. BL-P was a scholar in the International AIDS Training and Research Program, NIH Research Grant D43 TW000007, funded by the Fogarty International Center and

the Office of Research on Women's Health. GJ-S is supported by NIH K24 HD054314-04. This publication was made possible with help from the University of Washington Center for AIDS Research (CFAR), an NIH funded program (P30 AI027757), which is supported by the following NIH Institutes and Centers (NIAID, NCI, NIMH, NIDA, NICHD, NHLBI, NIA). The authors have no other funding or conflicts of interest to disclose.

**Corresponding author:**

Lisa M. Cranmer, MD, MPH, Division of Pediatric Infectious Diseases, Emory University School of Medicine, 2015 Uppergate Drive NE, Suite 500, Atlanta, Georgia 30322. Telephone: (206) 987-2073, Fax: (206) 987-3890. Email: lisa.cranmer@emory.edu.

**Running Head:** TB IFN- $\gamma$  Responses in Breastmilk**Key words:** TB, IGRA, HIV, maternal, infant**Summary word count:** 72**Text word count:** 996**Number of tables:** 1**Number of figures:** 1**Number of references:** 7

Tuberculosis (TB) cellular immune responses were examined in breastmilk of HIV-infected mothers using the T-SPOT.TB interferon gamma release assay (IGRA). Positive TB IFN- $\gamma$  responses were detected in 6 of 8 (75%) valid breast milk assays. Among 7 mothers with paired breastmilk and blood assays, TB IFN- $\gamma$  responses were higher in breastmilk compared to blood ( $p=.02$ ). The magnitude of TB IFN- $\gamma$  responses in maternal breastmilk and blood were correlated. Elucidating the influence of breastmilk TB immune responses on infant TB susceptibility and immunity may inform future maternal TB vaccine strategies.

For Review Only

## 1 INTRODUCTION

2 Cell-mediated immunity in breastmilk has been associated with protection from  
3 infectious pathogens and modulation of infant immune development.<sup>1,2</sup> Tuberculosis  
4 (TB)-specific cellular immunity in breastmilk has not been well-characterized and its role  
5 in infant TB immunity is undefined. Small historical studies suggest that tuberculin  
6 sensitivity can be transferred by breastmilk to infants.<sup>3,4</sup> Breastfed infants born to  
7 tuberculin positive mothers had evidence of tuberculin sensitivity, while infants of non-  
8 breastfed tuberculin positive or tuberculin negative mothers did not.<sup>4</sup> We adapted a  
9 standardized TB interferon-gamma (IFN- $\gamma$ ) release assay to measure breastmilk IFN- $\gamma$   
10 responses in HIV-infected mothers and compared these to TB IFN- $\gamma$  responses in  
11 peripheral blood of mothers and their infants.

12

## 13 STUDY POPULATION AND METHODS

14 In a historical perinatal HIV cohort, TB IFN- $\gamma$  responses were measured in  
15 cryopreserved maternal breastmilk cells (BMCs) and peripheral blood mononuclear  
16 cells (PBMCs) using the T-SPOT.TB ELISPOT assay (Oxford Immunotec, Oxfordshire,  
17 UK). This study was approved by the University of Washington Institutional Review  
18 Board and the Kenyatta National Hospital Ethics and Research Committee. HIV-  
19 infected women in Nairobi, Kenya were enrolled during pregnancy in 2002 and mother-  
20 infant pairs followed monthly for one year postpartum.<sup>5</sup> Breastmilk and peripheral blood  
21 were collected at 1 month postpartum, centrifuged, and cryopreserved in 10% dimethyl  
22 sulfoxide-90% fetal calf serum (FCS) (Sigma, St. Louis, Missouri, USA). In 2011,  
23 cryopreserved PBMCs and BMCs were thawed and incubated for 4 hours in RPMI 1640  
24 supplemented with 10% FCS (Sigma). Lymphocytes were counted manually in the  
25 presence of Trypan Blue and viable cells were plated at mean concentration  $1.5 \times 10^5$   
26 BMCs and  $2.5 \times 10^5$  PBMCs per well to perform the T-SPOT.TB assay as previously  
27 described.<sup>1,6</sup> Infant T-SPOT.TB results from 6 months of age were available from a  
28 prior study.<sup>6</sup> Total BMC or PBMC spot-forming cells (SFCs) were normalized to the  
29 number of spots per  $2.5 \times 10^5$  cells, and an assay was considered valid if there were  
30  $\geq 20$  SFCs in the positive control well. A test was positive if  $\geq 6$  spots above the nil  
31 control for either antigen ESAT-6 or CFP-10 when the nil control was  $< 10$  SFCs. If the

32 nil control was >10 SFCs, a test was considered positive if the SFCs for either ESAT-6  
33 or CFP-10 were at least twice the SFCs in the nil control.

34

35 Statistical analyses were performed using STATA software, version 11.2 (StataCorp,  
36 College Station, Texas, USA). After exclusion of invalid assays, TB IFN- $\gamma$  responses  
37 were described as dichotomous (positive/negative) and continuous (SFCs above nil)  
38 measures. Median IFN- $\gamma$  SFCs in response to ESAT-6 and CFP-10 above nil in BMCs  
39 and PBMCs were compared using Wilcoxon signed-rank tests. Data were log-  
40 transformed and Spearman's correlation of BMC and PBMC IFN- $\gamma$  responses to ESAT-6  
41 and CFP-10 were assessed.

42

### 43 RESULTS

44 T-SPOT.TB assays were conducted in paired BMC and PBMC specimens from 11 HIV-  
45 infected mothers at one month postpartum. Prior T-SPOT.TB results were available for  
46 10 of their infants at 6 months of life.<sup>6</sup> Median maternal CD4 count was 676 cells/mm<sup>3</sup>.  
47 In the first 6 months postpartum, mothers exclusively breastfed for a median of 6  
48 months (IQR 3-6); one mother delayed breastfeeding initiation until 1 month postpartum.  
49 No mothers or infants were diagnosed with active TB disease during follow-up. All  
50 infants were HIV negative by HIV DNA PCR at 6 months of age.

51

52 Of 11 paired maternal BMC and PBMC assays, 8 BMC and 10 maternal PBMC assays  
53 were valid. Of valid assays, TB IFN- $\gamma$  responses to either ESAT-6 or CFP-10 were  
54 present in 6 (75%) BMCs and 4 (40%) maternal PBMCs. For BMC assays, median  
55 response to ESAT-6 was 27 SFCs (IQR 2-221) and CFP-10 was 18 SFCs (IQR 4-240).  
56 Among maternal PBMC assays, median IFN- $\gamma$  response to ESAT-6 was 3 SFCs (IQR 1-  
57 38) and CFP-10 was 3 SFCs (IQR 0-21) (Table 1). Median IFN- $\gamma$  response to ESAT-6  
58 and CFP-10 were higher in BMCs than PBMCs ( $p=0.02$  for both antigens) (Figure 1a).  
59 Maternal BMC and PBMC TB IFN- $\gamma$  SFCs were highly correlated (ESAT-6  $r=.91$ ,  $p=.005$   
60 and CFP-10  $r=.86$ ,  $p=.01$ ) (Figure 1b).

61



62 We previously conducted T-SPOT.TB assays on the PBMCs of 10 infants of these  
63 mothers.<sup>6</sup> Three of 8 infants (37%) with a valid assay had a positive T-SPOT.TB result.  
64 Median infant IFN- $\gamma$  response to ESAT-6 was 0 SFCs (IQR 0-4) and CFP-10 was 0  
65 SFCs (IQR 0-2). Magnitude of infant PBMC TB IFN- $\gamma$  response to ESAT-6 and CFP-10  
66 did not correlate with maternal BMCs ( $r=.53$ ,  $p=.36$  and  $r=0$ ,  $p>.9$ , respectively). Of 5  
67 mother-infant pairs with valid assays, 3 were positive in both maternal BMC and infant  
68 PBMC, 1 was negative in both maternal BMC and infant PBMC, and 1 pair was  
69 discordant (maternal BMC positive, infant negative in pair with delayed breastfeeding).

70

## 71 DISCUSSION

72 In this proof of concept study, we measured TB-specific cellular immunity in breastmilk  
73 using the T-SPOT.TB assay. We found TB IFN- $\gamma$  responses in breastmilk cells of most  
74 HIV-infected mothers examined. Breastmilk IFN- $\gamma$  responses to ESAT-6 and CFP-10  
75 were of higher magnitude and strongly correlated with maternal PBMC IFN- $\gamma$  responses.

76

77 Measurement of TB IFN- $\gamma$  in breastmilk may be useful in future prospective studies to  
78 determine if breastmilk cell-mediated immunity may alter infant TB immunity or TB  
79 susceptibility. TB-specific immune cells in breastmilk could provide passive infant TB  
80 protection in the oral or respiratory mucosa or enhance development of infant TB  
81 immune responses. Maternal immunity may shape antigen sensitization, as observed  
82 in children aged 12 to 36 months who were tuberculin positive after breastfeeding from  
83 tuberculin positive mothers.<sup>3</sup> We did not find a significant association between maternal  
84 breastmilk and longitudinal infant TB IFN- $\gamma$  production. However our finding that 4 of 5  
85 mother-infant pairs had concordant TB IFN- $\gamma$  responses in maternal breastmilk and  
86 infant blood at 6 months of age is intriguing.

87

88 Use of cryopreserved samples may have decreased T-SPOT.TB assay sensitivity<sup>7</sup>, and  
89 small sample size limits the generalizability of our findings. Future studies to determine  
90 whether and how breastmilk TB cellular immunity influences infant immunity and  
91 susceptibility to TB will be important, and may inform complementary maternal vaccine  
92 strategies that potentially enhance protection of infants via breastmilk.

R1 Acknowledgements

7

We thank the participants of the perinatal cohort and the clinical and laboratory staff at the Paediatric Research Laboratory, Kenyatta National Hospital. We thank Dr. Cheryl Day for graphical assistance.

For Review Only

## R1 References

8

1. Lohman-Payne B, Slyker JA, Moore S, et al. Breast milk cellular HIV-specific interferon gamma responses are associated with protection from peripartum HIV transmission. *AIDS* 2012;26:2007-16.
2. Garofalo R. Cytokines in human milk. *J Pediatr* 2010;156:S36-40.
3. Mohr JA. The possible induction and-or acquisition of cellular hypersensitivity associated with ingestion of colostrum. *J Pediatr* 1973;82:1062-4.
4. Schlesinger JJ, Covelli HD. Evidence for transmission of lymphocyte responses to tuberculin by breast-feeding. *Lancet* 1977;2:529-32.
5. John-Stewart GC, Mbori-Ngacha D, Payne BL, et al. HIV-1-specific cytotoxic T lymphocytes and breast milk HIV-1 transmission. *J Infect Dis* 2009;199:889-98.
6. Cranmer LM, Kanyugo M, Jonnalagadda SR, et al. High prevalence of tuberculosis infection in HIV-1 exposed Kenyan infants. *Pediatr Infect Dis J* 2014;33:401-6.
7. Meier T, Eulenbruch HP, Wrighton-Smith P, Enders G, Regnath T. Sensitivity of a new commercial enzyme-linked immunospot assay (T SPOT-TB) for diagnosis of tuberculosis in clinical practice. *Eur J Clin Microbiol Infect Dis* 2005;24:529-36.

## Tables

Table 1: IFN- $\gamma$  SFCs per  $2.5 \times 10^5$  cells in response to *M. tuberculosis* in BMCs and maternal and infant PBMCs

ID	Cells	PHA	ESAT-6	CFP-10	Nil	Result	Breastfed* (months)
189	Maternal BMC	2432	7.5	15	2.5	Positive	
	Maternal PBMC	888	0	0	0	Negative	
	Infant PBMC	211	169	2	0	Positive	6
181	Maternal BMC	48	5	9	0	Positive	
	Maternal PBMC	665	3	3	0	Negative	
	Infant PBMC	200	20	44	16	Positive	6
143	Maternal BMC	1661	0	0	0	Negative	
	Maternal PBMC	348	0	0	0	Negative	
	Infant PBMC	0	4	4	4	Invalid	6
134	Maternal BMC	1207	373	297	17	Positive	
	Maternal PBMC	474	83	204	0	Positive	
	Infant PBMC	nd	nd	nd	nd	nd	5
194	Maternal BMC	168	0	0	0	Negative	
	Maternal PBMC	1	0	0	0	Invalid	
	Infant PBMC	288	0	1	0	Negative	2
173	Maternal BMC	0	0	0	0	Invalid	
	Maternal PBMC	921	1	1	0	Negative	
	Infant PBMC	708	24	52	77	Negative	6
141	Maternal BMC	1	0	0	0	Invalid	
	Maternal PBMC	961	59	31	0	Positive	
	Infant PBMC	0	0	0	0	Invalid	6
118	Maternal BMC	1087	744	488	115	Positive	
	Maternal PBMC	625	38	21	0	Positive	
	Infant PBMC	93	235	28	4	Positive	1
113	Maternal BMC	458	86	24	0	Positive	
	Maternal PBMC	576	39	6	1	Positive	
	Infant PBMC	69	0	0	0	Negative	3 ¶
140	Maternal BMC	800	50	200	0	Positive	
	Maternal PBMC	973	4	4	0	Negative	
	Infant PBMC	6	0	0	0	Invalid	3
107	Maternal BMC	0	0	0	0	Invalid	
	Maternal PBMC	861	1	0	0	Negative	
	Infant PBMC	82	0	0	0	Negative	6

\*Months mother reported exclusive breastfeeding during the first 6 months postpartum;

¶ Mother delayed initiation of breastfeeding until 1 month postpartum; nd = not done

R1 Figure




Figure 1: Magnitude and correlation of IFN- $\gamma$  response to ESAT-6 and CFP-10 in Maternal BMCs and PBMCs. A. The T-SPOT.TB assay was performed on maternal PBMCs and BMCs. SFCs per  $2.5 \times 10^5$  cells in response to antigens ESAT-6 (closed circles) and CFP-10 (open circles) are shown after subtraction of background in the nil control. Horizontal lines represent the median. Differences were determined using the Wilcoxon signed-rank test. B. Data were log-transformed and Spearman's correlation of BMC and PBMC IFN- $\gamma$  responses to ESAT-6 (closed circles) and CFP-10 (open circles) were assessed.