

University of Rhode Island

DigitalCommons@URI

Past Departments Faculty Publications (CELS) College of the Environment and Life Sciences

1988

Serologic analyses of *Peromyscus leucopus*, a rodent reservoir for *Borrelia burgdorferi*, in northeastern United States

Louis A. Magnarelli

John F. Anderson

Kerwin E. Hyland

University of Rhode Island

Durland Fish

Jay B. McAninch

Follow this and additional works at: https://digitalcommons.uri.edu/cels_past_depts_facpubs

Citation/Publisher Attribution

Magnarelli, L. A., Anderson, J. F., Hyland, K. E., Fish, D., & McAninch, J. B. (1988). Serologic analyses of *Peromyscus leucopus*, a rodent reservoir for *Borrelia burgdorferi*, in northeastern United States. *J. Clin. Microbiol.*, 26(6), 1138-1141. Retrieved from <http://jcm.asm.org/content/26/6/1138>.

Available at: <http://jcm.asm.org/content/26/6/1138>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Past Departments Faculty Publications (CELS) by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Serologic analyses of *Peromyscus leucopus*, a rodent reservoir for *Borrelia burgdorferi*, in northeastern United States

Terms of Use

All rights reserved under copyright.

Serologic Analyses of *Peromyscus leucopus*, a Rodent Reservoir for *Borrelia burgdorferi*, in Northeastern United States

LOUIS A. MAGNARELLI,^{1*} JOHN F. ANDERSON,¹ KERWIN E. HYLAND,² DURLAND FISH,³
AND JAY B. MCANINCH⁴

Department of Entomology, The Connecticut Agricultural Experiment Station, New Haven, Connecticut 06504¹;
Department of Zoology, University of Rhode Island, Kingston, Rhode Island 02881²;
Medical Entomology Laboratory,
New York Medical College, Armonk, New York 10504³;
Institute of Ecosystem Studies,
The New York Botanical Garden, Millbrook, New York 12545⁴

Received 19 January 1988/Accepted 9 March 1988

An enzyme-linked immunosorbent assay (ELISA) and indirect fluorescent-antibody test were used to detect antibodies to *Borrelia burgdorferi*, the causative agent of Lyme disease, in *Peromyscus leucopus* (white-footed mouse). Of the 661 mice captured in Connecticut, Rhode Island, and New York during 1980 and 1983 to 1987, 166 (25.1%) had antibodies to *B. burgdorferi* by ELISA. Comparative analyses of 210 serum specimens, collected in areas where Lyme disease is endemic, revealed a threefold difference in sensitivity between the ELISA (38.1% positive) and the indirect fluorescent-antibody method (12.4%). Although prevalence of seropositive *P. leucopus* was highest during June, elevated amounts of antibody (1:1,280 to 1:2,560) were detected in mice that harbored spirochetes during all seasons. Being reservoirs for *B. burgdorferi*, these rodents are suitable for monitoring spirochete infections at foci and should be included in field evaluations of control programs aimed at suppressing Lyme disease.

The etiologic agent of Lyme disease, *Borrelia burgdorferi*, is transmitted by ticks of the *Ixodes ricinus* complex to mammals and birds. In northeastern United States, Wisconsin, and Minnesota, *Ixodes dammini* is the primary vector (2, 3, 5, 7, 12, 15, 18). Although in its immature stages this hard-bodied tick is known to feed on a variety of vertebrate hosts (6, 13, 20, 25), including rodents, humans, domestic animals, and passerine birds, *Peromyscus leucopus* (the white-footed mouse) is considered a chief reservoir for *B. burgdorferi* in or near woodlands (1, 2, 5, 10, 11, 14, 17). The detection of antibody to this bacterium (15, 20, 21), frequent isolation of *Borrelia* spirochetes in areas where Lyme disease is endemic (1, 2, 4, 5), and the ability of *P. leucopus* to infect immature *I. dammini* (14, 17) reinforce the epidemiological significance of this mammal.

The association of *I. dammini* larvae and nymphs with *P. leucopus* is well documented (6, 13, 15, 20, 25). It is, therefore, appropriate to select this mammal for surveillance programs. Indirect fluorescent-antibody (IFA) methods have been used to detect antibody to *B. burgdorferi* (15, 20), but among the mammals studied, the prevalence of seropositive white-footed mice has been relatively low. It is unclear whether these rodents have weak immunologic responses to *B. burgdorferi* or whether the results are due to low test sensitivity. Accordingly, this investigation was conducted to further evaluate an enzyme-linked immunosorbent assay (ELISA) developed earlier (24), to compare results with those obtained by IFA, and to determine whether ELISA is more suitable for monitoring immunologic responses to *B. burgdorferi* in *P. leucopus* populations.

MATERIALS AND METHODS

Sampling. White-footed mice were captured in Sherman box traps from woodlands in Connecticut, Rhode Island, and New York during 1980 and 1983 to 1987. Although rodents were collected during all seasons, the majority were ob-

tained from May through August. Sites in Connecticut (East Haddam, Norwich, and Salem) are located in the south-central or southeastern part of the state, a region where Lyme disease is prevalent in human populations (21, 28). Mice were marked, released, and recaptured at East Haddam as previously described (20). In Rhode Island, *P. leucopus* was collected from inland areas, Block Island, Prudence Island, and Jamestown Island, whereas in New York, animals were captured in Westchester and Dutchess Counties. All study sites are within the geographical range of *I. dammini*. For controls, 60 additional mice were obtained from Newtown and Woodbridge, Conn., towns where Lyme disease and *I. dammini* are rare.

Serologic tests. Details on the use of IFA and ELISA methods and on quantitating total immunoglobulins to *B. burgdorferi* have been reported previously (20, 24). To improve the sensitivity of the ELISA, the concentration of washed, whole-cell antigen used to coat the solid phase was increased from 7 µg of protein per ml to 15 µg of protein per ml. In addition, the dilution of unconjugated rabbit anti-*P. leucopus* immunoglobulins was increased from 1:200 to 1:400, and the concentration of horseradish peroxidase-labeled goat anti-rabbit immunoglobulin G antibodies in phosphate buffer solutions was changed from 1:800 to 1:2,500. With these modifications, test sera were screened at dilutions \geq 1:160; net absorbance values (i.e., net optical densities) greater than 0.39, 0.34, and 0.29 were graded as positive for the respective serum dilutions of 1:160, 1:320, and \geq 1:640. These critical regions were determined by statistical analyses (3 standard deviations + mean) of net absorbance values for the respective serum dilutions of 60 normal specimens. Positive controls included serum samples from *P. leucopus* that had been found harboring *B. burgdorferi*. Procedures for blocking nonspecific binding sites and for the preparation of diluents, washes, and substrate remained unchanged. When new lots of enzyme-labeled antibodies or other reagents were purchased, tests were reevaluated with positive and negative serum controls and

* Corresponding author.

TABLE 1. Comparison of IFA and ELISA results for *P. leucopus* sera

| Site | Yr | Total no. of sera tested | IFA | | ELISA | |
|----------------|-----------|--------------------------|------------------|-------------------------------------|------------------|-------------------------------------|
| | | | No. (%) positive | Antibody titer ^a (range) | No. (%) positive | Antibody titer ^a (range) |
| Norwich, Conn. | 1986 | 50 | 0 | | 6 (12) | 320-2,560 |
| Salem, Conn. | 1986 | 135 | 23 (17) | 32-128 | 69 (51.1) | 320-2,560 |
| Rhode Island | 1983-1985 | 25 | 3 (12) | 32-512 | 5 (20) | 320-1,280 |

^a Reciprocal antibody titers ≥ 32 (IFA positive) and ≥ 320 (ELISA positive).

procedures were standardized accordingly. Uniformity of assay sensitivity was monitored daily by including the same positive and negative control sera on each plate. Controls were also included for antigens, conjugates, buffer solutions, and diluents.

To assess the specificity of the IFA method, 8 serum samples with antibodies to *B. burgdorferi* and 12 specimens that were nonreactive to this spirochete were screened against *Borrelia hermsii* (a relapsing fever spirochete), *Treponema pallidum* (the causative agent of syphilis in humans), and the following serovars of *Leptospira interrogans*: canicola, icterohaemorrhagiae, and pomona. The sources and preparation of these antigens have been reported previously (19).

Isolations. The blood, spleen tissues, or kidney tissues of white-footed mice were aseptically removed and processed for spirochete isolations as described previously (1, 5, 16). Briefly, 20 μ l of blood or 1:10 dilutions of spleen and kidney tissues, triturated in 7 ml of Barbour-Stoenner-Kelly medium, were placed into duplicate tubes of this medium containing 0.1% agarose. Cultures were kept at 31°C for 3 to 6 weeks and examined for spirochetes by dark-field microscopy. Isolates were subcultured and identified as *B. burgdorferi* by fixing washed, whole cells to glass microscope slides and by applying murine monoclonal antibody (H5332) and appropriate conjugates in IFA tests (8, 9). This monoclonal antibody was directed against outer surface protein A, a polypeptide with an approximate molecular mass of 31 kilodaltons. This protein is common to North American isolates of *B. burgdorferi*. Results verifying the presence of *B. burgdorferi* in rodents and arthropods from sites in Connecticut, Rhode Island, and New York have been published before (1, 2, 5, 7, 12, 19).

Statistical analyses. To determine significant differences in percentages of positive sera, values were analyzed by a Student *t* test after arcsin transformation (27).

RESULTS

Comparative analyses of 210 *P. leucopus* sera revealed threefold greater sensitivity by modified ELISA (Table 1). In areas where Lyme disease is prevalent in human populations, the number of positive serum samples by ELISA ($n = 80$) greatly exceeded that determined by the IFA method ($n = 26$). Antibody titers were also correspondingly higher in the ELISA than in the IFA test. In analyses of an additional 24 serum samples, representing *P. leucopus* specimens from areas where *I. dammini* and Lyme disease are rare, results were negative in both assays. To assess reproducibility of the ELISA, 24 positive sera were retested. There were no differences in antibody titers for 10 samples, whereas twofold ($n = 12$) or fourfold ($n = 2$) variations were recorded for the remaining sera in the second trial.

White-footed mice collected in Connecticut, Rhode Island, and New York (Westchester County) contained antibodies to *B. burgdorferi*. However, prevalence of seropositive animals by ELISA was highly variable (Table 2). For example, in 1984 and 1985 the number of *P. leucopus* specimens with antibodies in East Haddam, Conn., differed by nearly threefold. During 1986, more than half of the 135 *P. leucopus* specimens captured in Salem, Conn., had been exposed to Lyme disease spirochetes. Antibody titers ranged from 1:320 to 1:2,560, and among these, 1:640 was recorded most frequently (40.7% of 172 positive sera) followed by 1:320 (23.3%), 1:1,280 (21.5%), and 1:2,560 (14.5%).

Although antibodies to *B. burgdorferi* were present in *P. leucopus* during all months of the year in East Haddam, Conn. (Table 3), there were significantly more positive sera collected during June through August (22%) than during March through May (12%) ($t = 2.1, P < 0.03$). Antibody titers varied during all seasons, but titration endpoints peaked (1:2,560) during May, July, and August.

TABLE 2. Prevalence of *P. leucopus* specimens with antibodies to *B. burgdorferi*

| Site and yr | No. of serum samples tested | No. (%) positive ^a | No. of specimens with reciprocal titer of: | | | |
|--------------------------------|-----------------------------|-------------------------------|--|-----|-------|-------|
| | | | 320 | 640 | 1,280 | 2,560 |
| East Haddam, Conn. | | | | | | |
| 1980 | 147 | 25 (17) | 8 | 12 | 3 | 2 |
| 1983 | 63 | 11 (17.5) | 2 | 7 | 1 | 1 |
| 1984 | 81 | 9 (11.1) | 2 | 6 | 1 | 0 |
| 1985 | 61 | 18 (29.5) | 3 | 6 | 7 | 2 |
| Salem, Conn. | | | | | | |
| 1986 | 135 | 69 (51.1) | 19 | 25 | 13 | 12 |
| 1987 | 25 | 6 (24) | 2 | 2 | 0 | 2 |
| Norwich, Conn., 1986 | 50 | 6 (12) | 2 | 2 | 1 | 1 |
| Rhode Island, 1983-1985, 1987 | 57 | 10 (17.5) | 1 | 3 | 5 | 1 |
| Westchester County, N.Y., 1986 | 47 | 18 (38) | 1 | 7 | 6 | 4 |
| Dutchess County, N.Y., 1986 | 20 | 0 | | | | |
| Total | 686 | 172 (25.1) | 40 | 70 | 37 | 25 |

^a Analyzed by ELISA.

TABLE 3. Prevalence of white-footed mice with antibodies to *B. burgdorferi* in East Haddam during 1980 and 1983 to 1985

| Mo | Total sera tested | No. (%) ELISA positive ^a | Reciprocal antibody titer | |
|-----------|-------------------|-------------------------------------|---------------------------|---------------------|
| | | | Range | Geometric \bar{x} |
| January | 12 | 1 (8.3) | 640 | |
| February | 10 | 2 (20) | 640-1,280 | 905 |
| March | 19 | 3 (15.8) | 320-640 | 508 |
| April | 41 | 3 (7.3) | 320-1,280 | 640 |
| May | 40 | 6 (15) | 320-2,560 | 718 |
| June | 58 | 18 (31) | 320-1,280 | 616 |
| July | 54 | 9 (16.7) | 320-2,560 | 941 |
| August | 47 | 8 (12.8) | 320-2,560 | 761 |
| September | 39 | 7 (18) | 320-1,280 | 640 |
| October | 14 | 2 (14.3) | 320-640 | 453 |
| November | 16 | 3 (18.8) | 640-1,280 | 806 |
| December | 2 | 1 (50) | 640 | |
| Total | 352 | 63 (17.9) | 320-2,560 | 630 |

^a Prevalences based on sample sizes <19 not included in statistical comparisons.

Multiple serum samples were obtained from 82 *P. leucopus* specimens in East Haddam during 1980, 1983, and 1984. Seroconversions (i.e., a negative-to-positive change in test results) were recorded for 6 mice from May through September (Table 4). Rising antibody titers were detected in 4 animals by ELISA, but IFA test results for these sera remained negative. The second or subsequent serum specimens from the remaining 76 mice were negative in either or both assays.

Infected white-footed mice had antibodies to spirochetes during all seasons. Of the 206 animals tested for *B. burgdorferi*, 72 (35%) harbored this bacterium in their blood, kidneys, or spleen (Table 5). The majority of these (57.8% of 72) had detectable amounts of antibody to *B. burgdorferi* with titers ranging from 1:320 to 1:2,560 in ELISA. Although spirochetes were not isolated from the remaining 134 *P. leucopus* specimens, antibodies were detected in similar amounts in 10 mice.

Serum antibodies from 8 mice collected in Salem, an area where Lyme disease is highly endemic, cross-reacted in IFA tests against *B. hermsii* and *T. pallidum*. Similar tests with serovars of *L. interrogans* were negative. Antibody titers to *B. hermsii* differed twofold or less from homologous reactions to *B. burgdorferi* in all serum samples, whereas titration endpoints to *T. pallidum* were at least fourfold lower than those to the Lyme disease agent. All 12 serum specimens without antibodies to *B. burgdorferi* were likewise negative when tested against all other spirochetal antigens.

TABLE 5. Prevalence of *Borrelia*-infected *P. leucopus* specimens with antibodies to spirochetes

| Site | Yr | Total no. infected ^a | No. (%) of infected mice ^b | |
|--------------------|------------------------|---------------------------------|---------------------------------------|--------------------|
| | | | With antibodies | Without antibodies |
| East Haddam, Conn. | 1984 ^c | 6 | 4 (67) | 2 (33) |
| East Haddam, Conn. | 1985 ^c | 33 | 14 (42.4) | 19 (57.6) |
| Rhode Island | 1984-1986 ^d | 6 | 4 (67) | 2 (33) |
| New York | 1986 ^c | 27 | 16 (59.3) | 11 (40.7) |
| Total | | 72 | 38 (57.8) | 34 (47.2) |

^a Some isolates reported earlier (1, 2, 5).

^b Tested by ELISA.

^c Increase in isolations due to improved growth medium and the testing of spleen and kidney tissues.

^d Mice from Prudence Island.

^e Mice from Westchester County.

DISCUSSION

Antibodies to *B. burgdorferi* can be detected by either an IFA test or a modified ELISA, but the latter had greater sensitivity. The relatively low number of seropositive *P. leucopus* specimens reported earlier (20) was a conservative estimate of *B. burgdorferi* presence in endemic areas. Therefore, for surveillance purposes, ELISA is preferred because of greater sensitivity and ease of automation and standardization and because test results are more objective.

White-footed mice can harbor *B. burgdorferi* during all seasons. Anderson et al. (1) reported infection rates of $\geq 66\%$ during the summer and $\leq 33\%$ during the winter. In addition, the number of mice with antibodies to the Lyme disease agent varied seasonally and geographically. The highest titers of antibody and number of seroconversions occurred during the summer, a time when nymphs of *I. dammini* were active. Therefore, the warmer months appear to be most suitable for monitoring serologic responses to *B. burgdorferi* in these rodents.

Antibodies to *B. burgdorferi* were present in mice that had spirochete infections. Compared with those for other mammals, such as humans or dogs, antibody titers for these rodents were relatively low, regardless of the assay method used. Since some infected mice had no detectable antibody, immunologic response in *P. leucopus* may be delayed during initial and possibly late stages of infection. Even when immunoglobulins are produced, we suspect that they may be relatively ineffective in quickly neutralizing the spirochetes. Reservoir competence of *P. leucopus* has been confirmed (14, 17), and although spirochetemia periods are unknown, these and possibly other vertebrate hosts (2, 3, 6) are

TABLE 4. Antibody titers for *P. leucopus* specimens recaptured in East Haddam, Conn., during 1980 and 1983

| Mouse no. ^a | Date of capture | Reciprocal antibody titer at capture | | Date of recapture | Reciprocal antibody titer at recapture | |
|------------------------|------------------|--------------------------------------|----------------|-------------------|--|-------|
| | | IFA | ELISA | | IFA | ELISA |
| 1 | 14 May 1980 | Neg ^b | Neg | 23 July 1980 | Neg | 640 |
| 2 | 7 August 1980 | Neg | Neg | 20 August 1980 | Neg | 640 |
| 3 ^c | 25 April 1983 | Neg | — ^d | 10 May 1983 | 1:32 | — |
| 4 | 19 May 1983 | Neg | — | 9 June 1983 | Neg | 640 |
| 5 | 9 June 1983 | Neg | Neg | 24 June 1983 | Neg | 1,280 |
| 6 | 9 September 1983 | Neg | Neg | 29 September 1983 | — | 320 |

^a Isolation of spirochetes not attempted.

^b Neg, Negative.

^c Additional serum specimens collected on 19 April (IFA, negative) and 19 May (ELISA, 1:640).

^d —, Not tested because of insufficient amounts of serum.

important sources of infection. Transovarial transmission of *B. burgdorferi* occurs at a low rate in *I. dammini* (22, 26). Therefore, uninfected larval or nymphal ticks acquire *B. burgdorferi* by feeding on spirochetemic hosts and transmit the spirochetes when they refeed in later life stages.

White-footed mice are abundant in woodlands, have relatively small home ranges, are easily caught, and often have numerous larvae and nymphs of *Dermacentor variabilis* and *I. dammini* attached (2, 6, 13, 15, 20, 25). Antibodies produced against *B. burgdorferi* are now readily detected by ELISA, and methods are available for isolating this pathogen. Therefore, *P. leucopus* is ideal for studies designed to provide more precise information on the distribution and prevalence of *B. burgdorferi* infections in nature. Serologic analyses for the Lyme disease agent should be conducted along with isolation procedures. Furthermore, with growing interest in finding ways to decrease *I. dammini* populations and to reduce the risk of human infection, white-footed mice should be included in field evaluations of control efforts.

Tests on the specificity of our IFA method revealed cross-reactivity among the *Borrelia* and *Treponema* species. This has been demonstrated before (19, 23) for antibodies in sera from humans and white-tailed deer (*Odocoileus virginianus*). However, since tick-borne relapsing fever is not known to occur in eastern United States and since *T. pallidum* does not infect *P. leucopus*, cross-reactivity with *B. burgdorferi* is probably insignificant. However, other unclassified spirochetes have been isolated from *P. leucopus* specimens and short-tailed shrews (*Blarina brevicauda*) in Connecticut and Minnesota (4). These organisms appear to be serologically distinct from species of *Borrelia*, *Treponema*, *Leptospira*, and *Spirochaeta*. Nonetheless, there might be other unknown spirochetes in nature that could stimulate immune responses in *P. leucopus* and lessen the specificity of these assays. Therefore, serologic evidence of *B. burgdorferi* infections should be supported by isolation and characterization of spirochetes in study areas.

ACKNOWLEDGMENTS

We thank George Hansen, Clifford Snow III, Patricia Trzcinski, and Carol Lemmon for their technical assistance.

LITERATURE CITED

- Anderson, J. F., R. C. Johnson, and L. A. Magnarelli. 1987. Seasonal prevalence of *Borrelia burgdorferi* in natural populations of white-footed mice, *Peromyscus leucopus*. *J. Clin. Microbiol.* **25**:1564-1566.
- Anderson, J. F., R. C. Johnson, L. A. Magnarelli, and F. W. Hyde. 1985. Identification of endemic foci of Lyme disease: isolation of *Borrelia burgdorferi* from feral rodents and ticks (*Dermacentor variabilis*). *J. Clin. Microbiol.* **22**:36-38.
- Anderson, J. F., R. C. Johnson, L. A. Magnarelli, and F. W. Hyde. 1986. Involvement of birds in the epidemiology of the Lyme disease agent *Borrelia burgdorferi*. *Infect. Immun.* **51**:394-396.
- Anderson, J. F., R. C. Johnson, L. A. Magnarelli, F. W. Hyde, and T. G. Andreadis. 1987. New infectious spirochete isolated from short-tailed shrews and white-footed mice. *J. Clin. Microbiol.* **25**:1490-1494.
- Anderson, J. F., R. C. Johnson, L. A. Magnarelli, F. W. Hyde, and J. E. Myers. 1987. Prevalence of *Borrelia burgdorferi* and *Babesia microti* in mice on islands inhabited by white-tailed deer. *Appl. Environ. Microbiol.* **53**:892-894.
- Anderson, J. F., and L. A. Magnarelli. 1984. Avian and mammalian hosts for spirochete-infected ticks and insects in a Lyme disease focus in Connecticut. *Yale J. Biol. Med.* **57**:627-641.
- Anderson, J. F., L. A. Magnarelli, and J. B. McAnich. 1987. *Ixodes dammini* and *Borrelia burgdorferi* in northern New England and upstate New York. *J. Parasitol.* **73**:419-421.
- Barbour, A. G., R. A. Heiland, and T. R. Howe. 1985. Heterogeneity of major proteins in Lyme disease borreliae: a molecular analysis of North American and European isolates. *J. Infect. Dis.* **152**:478-484.
- Barbour, A. G., S. L. Tessier, and W. J. Todd. 1983. Lyme disease spirochetes and ixodid tick spirochetes share a common surface antigenic determinant defined by a monoclonal antibody. *Infect. Immun.* **41**:795-804.
- Bosler, E. M., B. G. Ormiston, J. L. Coleman, J. P. Hanrahan, and J. L. Benach. 1984. Prevalence of the Lyme disease spirochete in populations of white-tailed deer and white-footed mice. *Yale J. Biol. Med.* **57**:651-659.
- Bosler, E. M., and T. L. Schulze. 1986. The prevalence and significance of *Borrelia burgdorferi* in the urine of feral reservoir hosts. *Zentralbl. Bakteriol. Mikrobiol. Hyg. Ser. A* **263**:40-44.
- Burgdorfer, W., A. G. Barbour, S. F. Hayes, J. L. Benach, E. Grunwaldt, and J. P. Davis. 1982. Lyme disease—a tick-borne spirochetosis? *Science* **216**:1317-1319.
- Carey, A. B., W. L. Krinsky, and A. J. Main. 1980. *Ixodes dammini* (Acari:Ixodidae) and associated ixodid ticks in south-central Connecticut, USA. *J. Med. Entomol.* **17**:89-99.
- Donahue, J. G., J. Piesman, and A. Spielman. 1987. Reservoir competence of white-footed mice for Lyme disease spirochetes. *Am. J. Trop. Med. Hyg.* **36**:92-96.
- Godsey, M. S., T. E. Amundson, E. C. Burgess, W. Schell, J. P. Davis, R. Kaslow, and R. Edelman. 1987. Lyme disease ecology in Wisconsin: distribution and host preferences of *Ixodes dammini*, and the prevalence of antibody to *Borrelia burgdorferi* in small mammals. *Am. J. Trop. Med. Hyg.* **37**:180-187.
- Johnson, R. C., N. Marek, and C. Kodner. 1984. Infection of Syrian hamsters with Lyme disease spirochetes. *J. Clin. Microbiol.* **20**:1099-1101.
- Levine, J. F., M. L. Wilson, and A. Spielman. 1985. Mice as reservoirs of the Lyme disease spirochete. *Am. J. Trop. Med. Hyg.* **34**:355-360.
- Loken, K. I., C. Wu, R. C. Johnson, and R. F. Bey. 1985. Isolation of the Lyme disease spirochete from mammals in Minnesota. *Proc. Soc. Exp. Biol. Med.* **179**:300-302.
- Magnarelli, L. A., J. F. Anderson, C. S. Apperson, D. Fish, R. C. Johnson, and W. A. Chappell. 1986. Spirochetes in ticks and antibodies to *Borrelia burgdorferi* in white-tailed deer from Connecticut, New York State, and North Carolina. *J. Wildl. Dis.* **22**:178-188.
- Magnarelli, L. A., J. F. Anderson, W. Burgdorfer, and W. A. Chappell. 1984. Parasitism by *Ixodes dammini* (Acari:Ixodidae) and antibodies to spirochetes in mammals at Lyme disease foci in Connecticut, USA. *J. Med. Entomol.* **21**:52-57.
- Magnarelli, L. A., J. F. Anderson, and W. A. Chappell. 1984. Geographic distribution of humans, raccoons, and white-footed mice with antibodies to Lyme disease spirochetes in Connecticut. *Yale J. Biol. Med.* **57**:619-626.
- Magnarelli, L. A., J. F. Anderson, and D. Fish. 1987. Transovarial transmission of *Borrelia burgdorferi* in *Ixodes dammini* (Acari:Ixodidae). *J. Infect. Dis.* **156**:234-236.
- Magnarelli, L. A., J. F. Anderson, and R. C. Johnson. 1987. Cross-reactivity in serological tests for Lyme disease and other spirochetal infections. *J. Infect. Dis.* **156**:183-188.
- Magnarelli, L. A., J. M. Meegan, J. F. Anderson, and W. A. Chappell. 1984. Comparison of an indirect fluorescent-antibody test with an enzyme-linked immunosorbent assay for serological studies of Lyme disease. *J. Clin. Microbiol.* **20**:181-184.
- Main, A. J., A. B. Carey, M. G. Carey, and R. H. Goodwin. 1982. Immature *Ixodes dammini* (Acari:Ixodidae) on small animals in Connecticut, U.S.A. *J. Med. Entomol.* **19**:655-664.
- Piesman, J., J. G. Donahue, T. N. Mather, and A. Spielman. 1986. Transovarially acquired Lyme disease spirochetes (*Borrelia burgdorferi*) in field-collected larval *Ixodes dammini* (Acari:Ixodidae). *J. Med. Entomol.* **23**:291.
- Snedecor, G. W., and W. G. Cochran. 1967. *Statistical methods*. Iowa State University Press, Ames.
- Steere, A. C., and S. E. Malawista. 1979. Cases of Lyme disease in the United States: locations correlated with distribution of *Ixodes dammini*. *Ann. Intern. Med.* **91**:730-733.