

9-3-2018

Financial Impact of Incentive Spirometry

Adam E. M. Eltorai

Grayson L. Baird

Joshua Pangborn

Ashley Szabo Eltorai

Valentin Antoci

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/nursing_facpubs

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial 4.0 License](https://creativecommons.org/licenses/by-nc/4.0/)

Citation/Publisher Attribution

M., A. E., Baird, G. L., Pangborn, J., Eltorai, A. S., Antoci, V., Paquette, K., ... Daniels, A. H. (2018). Financial Impact of Incentive Spirometry. *INQUIRY: The Journal of Health Care Organization, Provision, and Financing*. <https://doi.org/10.1177/0046958018794993>

Available at: <https://doi.org/10.1177/0046958018794993>

This Article is brought to you for free and open access by the College of Nursing at DigitalCommons@URI. It has been accepted for inclusion in College of Nursing Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.


Authors

Adam E. M. Eltorai, Grayson L. Baird, Joshua Pangborn, Ashley Szabo Eltorai, Valentin Antoci, Katherine Paquette, Kevin Connors, Jacqueline Barbaria, Kimberly J. Smeals, Barbara Riley, Shyam A. Patel, Saurabh Agarwal, Terrance T. Healey, Corey E. Ventetuolo, Frank W. Sellke, and Alan H. Daniels

Financial Impact of Incentive Spirometry

INQUIRY: The Journal of Health Care
Organization, Provision, and Financing
Volume 55: 1–8
© The Author(s) 2018
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/0046958018794993
journals.sagepub.com/home/inq



Adam E. M. Eltorai, PhD¹ , Grayson L. Baird, PhD^{1,2},
Joshua Pangborn, BS¹, Ashley Szabo Eltorai, MD³,
Valentin Antoci Jr, MD, PhD¹, Katherine Paquette, PhD, RN⁴,
Kevin Connors, BA, RRT, CPFT², Jacqueline Barbaria, RRT²,
Kimberly J. Smeals, RRT², Barbara Riley, DNP, RN²,
Shyam A. Patel, MD¹, Saurabh Agarwal, MD¹, Terrance T. Healey, MD¹,
Corey E. Ventetuolo, MD, MS^{1,2}, Frank W. Sellke, MD¹,
and Alan H. Daniels, MD¹

Abstract

Despite largely unproven clinical effectiveness, incentive spirometry (IS) is widely used in an effort to reduce postoperative pulmonary complications. The objective of the study is to evaluate the financial impact of implementing IS. The amount of time nurses and RTs spend each day doing IS-related activities was assessed utilizing an online survey distributed to the relevant national nursing and respiratory therapists (RT) societies along with questionnaire that was prospectively collected every day for 4 weeks at a single 10-bed cardiothoracic surgery step-down unit. Cost of RT time to teach IS use to patients and cost of nurse time spent reeducating and reminding patients to use IS were used to calculate IS implementation cost estimates per patient. Per-patient cost of IS implementation ranged from \$65.30 to \$240.96 for a mean 9-day step-down stay. For the 566 patients who stayed in the 10-bed step-down in 2016, the total estimated cost of implementing IS ranged from \$36 959.80 to \$136 383.36. Using national survey workload data, per-patient cost of IS implementation costed \$107.36 (95% confidence interval [CI], \$97.88–\$116.98) for a hospital stay of 4.5 days. For the 9.7 million inpatient surgeries performed annually in the United States, the total annual cost of implementing postoperative IS is estimated to be \$1.04 billion (95% CI, \$949.4 million–\$1.13 billion). The cost of implementing IS is substantial. Further efficacy studies are necessary to determine whether the cost is justifiable.

Keywords

incentive spirometry, health care costs, nursing, respiratory therapy, length of stay, inpatients, workload, motivation, postoperative period, treatment outcome, respiratory function tests, surveys and questionnaires, spirometry

What do we already know about this topic?

Despite largely unproven clinical effectiveness, incentive spirometry (IS) is widely used in an effort to reduce postoperative pulmonary complications.

How does your research contribute to the field?

The current cost of implementing IS is substantial.

What are your research's implications toward theory, practice, or policy?

Further clinical effectiveness studies are necessary to determine whether the cost of IS is justifiable.

¹The Warren Alpert Medical School of Brown University, Providence, RI, USA

²Rhode Island Hospital, Providence, USA

³Yale University, New Haven, CT, USA

⁴The University of Rhode Island, Providence, USA

Received 22 November 2017; revised 24 July 2018; revised manuscript accepted 26 July 2018

Corresponding Author:

Adam E. M. Eltorai, The Warren Alpert Medical School of Brown University, 222 Richmond St, Providence, RI 02903, USA.

Email: adam_eltorai@brown.edu



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Introduction

Implementation of incentive spirometry (IS) involves a team of providers including physicians, respiratory therapists (RT), and nurses.¹ Clinical practice guidelines recommend IS usage in acute care inpatient, skilled nursing facility, extended care, and home care settings to optimize pulmonary function.² IS has been reported to be used in 95% of US hospitals after surgery.³ Most commonly, RT lead initial patient education on IS use, followed by nurse reeducation and reminding. Provider time and costs of IS implementation have not previously been reported.

Despite widespread use of IS, there is a paucity of data demonstrating therapeutic value.⁴⁻²² One systematic review²³ stated, "Presently, the evidence does not support the use of IS for decreasing the incidence of postoperative pulmonary complications." Another combined analysis¹⁶ concluded, "routine use of respiratory physiotherapy after abdominal surgery does not seem to be justified." The most recent meta-analyses found "no evidence" of benefit from IS.²⁴⁻²⁶ Last, the American Association For Respiratory Care (AARC) Clinical Practice Guideline: Incentive Spirometry: 2011² states that "evidence strongly suggests that IS alone may be inappropriate to prevent or treat postoperative complications."

In an increasingly cost-conscious health care environment,²⁷⁻³³ the goal of this study was to assess the costs of implementing IS.

Methods

Installation Costs

To assess the amount of time spent by RTs on IS-related activities, an IRB (Institutional Review Board)-approved, secure online survey was created and distributed via social media and newsletters by the AARC, from September to December 2016. As a complementary analysis, the survey was also distributed to RTs at a single academic medical center (AMC). Survey results were de-identified.

RTs national hourly wage data were gathered from the U.S. Department of Labor (2015). Using the latest Consumer Price Index data from the U.S. Department of Labor Bureau of Labor Statistic, wages were inflation-adjusted to 2017 dollars.³⁴ RTs' per-minute wage was calculated by dividing hourly wage by 60 minutes.

Individual IS unit (4000 mL Coach 2 [Smiths Medical, Saint Paul, Minnesota]) cost was determined based on the wholesale case price paid to the hospital supplier by AMC.³⁵ This commonly used IS model represents the IS device type recommended by the AARC.²

Maintenance Costs

To assess the amount of time spent per patient by nurses on IS-related activities, an IRB-approved, secure online survey, as described, was distributed via social media and newsletters by the following national nursing societies from September to December 2016: Academy of Medical-Surgical Nurses (AMSN), American Association of Critical-Care Nurses (AACN), and American Society of Peri-Anesthesia Nurses (ASPAN). The survey was also distributed to the nurses at the AMC. Survey results were de-identified.

Questionnaires on IS workload were prospectively collected every day for 4 weeks in December 2016 from nurses on a 10-bed cardiothoracic surgery step-down unit at the AMC. On that unit, nursing orders for IS are "Routine, Now then every 4 hours." IS orders are bundled as part of the admission order set, specifically requiring opt-out for exclusion. Surveys were collected at 6:45 PM from nurses who worked 7:00 AM to 7:00 PM. Nurses were asked how much time (minutes) during their shift they spent educating or reminding patients to use their ISs. Nurses completed one survey for each of their patients.

Registered nurses' national hourly wage data were gathered from the Bureau of Labor Statistics of the U.S. Department of Labor (2011) and 2017 inflation-adjusted numbers.³⁴ Nurses' per-minute wage is calculated by dividing hourly wage by 60 minutes.

Calculations

The following equation was used to calculate IS implementation costs per patient (Table 1):

The data from the AMC study were used to estimate the annual cost of IS implementation on the 10-bed cardiothoracic surgery step-down unit. Annual costs for implementing IS in the United States were estimated using national mean hospital length of stay (4.5 days),³⁷ number of inpatient surgeries per year (9.7 million),³⁸ and provider time spent doing IS-related activities from our national surveys. Point and interval estimates of time were calculated using a generalized linear model assuming a negative binomial distribution using SAS/GLIMMIX 9.4 (SAS Inc, Cary, North Carolina).

Table 1. Estimating Equation Cost Per Patient.

$IS + (\$_{RT} \times t_{RT}) + (\$_{RN} \times t_{RN} \times 16 \text{ hours}^3/\text{day} \times \text{LOS})$
IS = Mean IS unit cost (\$)
$\$_{RT}$ = RTs' mean per-minute wage (\$)
t_{RT} = RTs' spent initially educating a patient to use their IS
$\$_{RN}$ = Nurses' mean per-minute wage (\$)
t_{RN} = Minutes per hour nurses spend doing IS-related activities per patient
LOS = Mean hospital length of stay (days)
^a Number of hours per day patients are not asleep to perform IS: 24 hours-8 hours (recommended number of hours of sleep per night for adults) ³⁶

Note. IS = incentive spirometry; RTs = respiratory therapists.

Table 2. RT Survey Respondent Characteristics.

	Answer options	AARC % (n)	AMC % (n)
Degree (highest)	AS	48.9 (174/356)	61.1 (11/18)
	BS	51.1 (182/356)	38.9 (7/18)
Years in practice, mean \pm SD (n)	—	21.4 \pm 13.5 (368)	23.6 \pm 14.0 (19)
Primary practice location	PACU	0.3 (1/350)	0 (0/19)
	ICU	42.3 (148/350)	57.9 (11/19)
	Step-down unit	5.1 (18/350)	0 (0/19)
	Medical/surgical wards	35.4 (124/350)	21.1 (4/19)
	Rehabilitation	5.7 (20/350)	0 (0/19)
	In-home	1.1 (4/350)	0 (0/19)
	Other	10.0 (350/350)	21.1 (4/19)
In an average 8-hour shift, typically how much time do you spend educating or reminding a patient to use their IS? (minutes), mean (95% CI)		16.4 (14.7-18.4)	21.6 (1.1-42.2)
	Minutes per hour (above/8 hours), mean (95% CI)	2.1 (1.8-2.3)	2.7 (0.1-5.3)

Note. RT = respiratory therapists; AARC = American Association for Respiratory Care; AMC = academic medical center; PACU = post anesthesia care unit; ICU = intensive care unit; IS = incentive spirometry; CI = confidence interval; AS = Associate of Science; BS = Bachelor of Science.

Microsoft Excel (Microsoft Corporation, Redmond, Washington) was used for standard data analysis.

Results

Installation Costs

There were a total of 416 unique respondents from the AARC and 19 from the AMC with various educational backgrounds, years of experience, and primary practice locations (Table 2). Survey completion rates were 80.3% for AARC and 78.9% for AMC. Given the distribution methodology, response rate cannot be determined due to the inability to identify the total number of individuals the survey may have reached. AARC respondents reported spending an average of 16.4 (95% confidence interval [CI], 14.7-18.4) minutes per 8-hour shift educating or reminding patients to use IS, equating to 2.1 (95% CI, 1.8-2.3) minutes per hour. AMC RT respondents reported an average of 21.6 (95% CI, 1.1-42.2) minutes per 8-hour shift, equating to 2.7 (95% CI, 0.1-5.3) minutes per hour. There was no significant difference between AMC and AARC times spent educating or reminding patients ($P = .5349$).

RTs' national 2017-adjusted mean per-minute wage is \$0.49 (\$29.37/60 min). Individual IS unit cost was \$12.92.

Maintenance Costs

There were a total of 1265 unique respondents from the ASPAN, AMSN, and AACN and 126 from the AMC. Survey completion rates were 84.3% for ASPAN, 84.8% for AMSN, 90.1% for AACN, and 84.9% for AMC. The respondents included nurses with various educational backgrounds, years of experience, and primary practice locations (Table 3). Given the distribution methodology, response rate cannot be

determined due to the inability to identify the total number of individuals the survey may have reached. Combined national society (ASPAN, AMSN, AACN) respondents reported spending an average of 15.8 (95% CI, 14.5-17.2) minutes per 8-hour shift educating or reminding patients to use their ISs, equating to 2.0 (95% CI, 1.8-2.2) minutes per hour. AMC nurse respondents reported an average of 15.1 (95% CI, 11.4-18.9) minutes per 8-hour shift, equating to 1.9 (95% CI, 1.4-2.4) minutes per hour. There was no significant difference between AMC and aggregated nursing society times spent educating or reminding patients ($P = .7465$).

A total of 217 questionnaires from 23 the AMC step-down unit nurses were prospectively collected, representing a 100% response rate. Nurses reported spending on average 8.8 (95% CI, 7.8-9.9) minutes per 12-hours shift, equating to 0.7 (95% CI, 0.6-0.8) minutes per hour educating or reminding patients to use their ISs. In 2016, there were 566 patients, who each spent an average of 9 days on the AMC step-down unit.

Registered nurses' national 2017-adjusted mean per-minute wage is \$0.60 (\$35.87/60 min).

Implementation Cost Estimates

Incorporating mean data into the estimating equation, the new estimating equation is simplified (Table 4).

Using AMC mean data, per-patient cost of IS implementation ranged from \$65.30 to \$240.96 (Table 5). For the 566 patients who stayed in the AMC's 10-bed cardiothoracic step-down unit in 2016, total annual costs of implementing IS ranged from \$36 959.80 to \$136 383.36.

Using national survey mean data, per-patient cost of IS implementation costs \$107.36 (95% CI, \$97.88-\$116.98) (Table 6). For the 9.7 million inpatient surgeries performed annually in the United States (Project HCaU, 2015), total annual cost of implementing IS is \$1.04 billion (95% CI, \$949.4 million-\$1.13 billion).

Table 3. Nurse Survey Respondent Characteristics.

	Answer options	ASPN % (n)	AMSN % (n)	AACN % (n)	Aggregated % (n)	AMC % (n)
Degree (highest)	Nurse: Diploma	7.1 (56/788)	2.9 (8/272)	2.9 (3/102)	5.8 (67/1162)	0 (0/124)
	Nurse: ADN	14.5 (114/788)	16.9 (46/272)	6.9 (7/102)	14.4% (167/1162)	17.7 (22/124)
	Nurse: LPN	0.1 (1/788)	0.0 (0/272)	0.0 (0/102)	0.1 (1/1162)	0 (0/124)
	Nurse: BSN	59.5 (469/788)	52.9 (144/272)	61.8 (63/102)	58.2 (676/1162)	75.0 (93/124)
	Nurse: MSN	14.5 (114/788)	21.0 (57/272)	19.6 (20/102)	16.4 (191/1162)	3.2 (4/124)
	Nurse: DNP	0.4 (3/788)	0.4 (1/272)	3.9 (4/102)	0.7 (8/1162)	0.8 (1/124)
	Nurse: PhD	0.3 (2/788)	2.2 (6/272)	3.9 (4/102)	1.0 (12/1162)	0 (0/124)
	Nurse: Other	3.7 (29/788)	3.7 (10/272)	1.0 (1/102)	3.4 (40/1162)	3.2 (4/124)
Years in practice, mean \pm SD (n)	—	26.8 \pm 11.1 (793)	17.6 \pm 12.8 (279)	15.7 \pm 12.6 (104)	23.7 \pm 12.5 (1176)	12.4 \pm 10.1 (121)
Primary practice location	PACU	85.6 (670/783)	0.4 (1/276)	2.9 (3/102)	58.1 (674/1161)	9.0 (11/122)
	ICU	0.9 (7/783)	1.8 (5/276)	62.7 (64/102)	6.5 (76/1161)	19.7 (24/122)
	Step-down unit	0.4 (3/783)	7.6 (21/276)	16.7 (17/102)	3.5 (41/1161)	13.9 (17/122)
	Medical/surgical wards	1.8 (14/783)	83.3 (230/276)	14.7 (15/102)	22.3 (259/1161)	52.5 (64/122)
	Rehabilitation	0.1 (1/783)	0.7 (2/276)	1.0 (1/102)	0.3 (4/1161)	1.6 (2/122)
	In-home	0.0 (0/783)	0.7 (2/276)	0 (0/102)	0.2 (2/1161)	0 (0/122)
	Other	11.2 (88/783)	5.4 (15/276)	2.0 (2/102)	9.0 (105/1161)	3.3 (4/122)
In an average 8-hour shift, typically how much time do you spend educating or reminding a patient to use their IS? (minutes), mean (95% CI)	Minutes per hour (above/8 hours), mean (95% CI)	16.0 (14.7-17.3)	16.2 (14.3-18.4)	14.0 (11.3-17.4)	15.8 (14.5-17.2)	15.1 (11.4-18.9)
		2.0 (1.8-2.2)	2.0 (1.8-2.3)	1.8 (1.4-2.2)	2.0 (1.8-2.2)	1.9 (1.4-2.4)

Note. ASPAN = American Society of Peri-Anesthesia Nurses; AMSN = Academy of Medical-Surgical Nurses; AACN = American Association of Critical-Care Nurses; AMC = academic medical center; PACU = post anesthesia care unit; ICU = intensive care unit; IS = incentive spirometry; CI = confidence interval; AND = Associate's degree in Nursing; LPN = licensed practical nurse; BSN = bachelor of science in nursing; MSN = Master's of Science degree in Nursing; DNP = Doctor of Nursing Practice.

Table 4. Simplified Cost Estimating Equation of Per Patient.

$$IS + (\$_{RT} \times t_{RT}) + (\$_{RN} \times t_{RN} \times 16 \text{ hours}^a/\text{day} \times LOS) = \$12.92 + (\$0.49/\text{minute} \times t_{RT}) + (\$0.60/\text{minute} \times t_{RN} \times 16 \text{ hours}^a/\text{day} \times LOS)$$

IS = Mean IS unit cost (\$)
 \$_{RT} = RTs' mean per-minute wage (\$)
 t_{RT} = RTs' spent initially educating a patient to use their IS
 \$_{RN} = Nurses' mean per-minute wage (\$)
 t_{RN} = Minutes per hour nurses spend doing IS-related activities per patient
 LOS = Mean hospital length of stay (days)
^aNumber of hours per day patients are not asleep to perform IS: 24 hours-8 hours (recommended number of hours of sleep per night for adults)

Note. IS = incentive spirometry; RTs = respiratory therapists.

Table 5. Per-Patient IS Implementation Cost Estimates Using AMC Data.

IS (\$)	+	\$ _{RT} (\$/min)	×	t _{RT} (min) (95% CI)	+	\$ _{RN} (\$/min)	×	t _{RN} (min/h) (95% CI)	×	16 h/d	×	LOS (days)	Cost (\$) (95% CI)
12.92		0.49		21.6 (1.1-42.2)		0.60		0.7 (0.6-0.8) ^a 1.9 (1.4-2.4) ^b				9	83.98 (65.30-102.72) 187.66 (134.42-240.96)

Note. IS = incentive spirometry; AMC = academic medical center; \$_{RT} = RT's mean per-minute wage (\$); RT = respiratory therapist; t_{RT} = RTs' spent initially educating a patient to use their IS; CI = confidence interval; \$_{RN} = nurses' mean per-minute wage (\$); t_{RN} = minutes per hour nurses spend doing IS-related activities per patient; LOS = mean hospital length of stay (days).
^aProspectively collected AMC step-down unit questionnaire data.
^bAMC online nurse survey responses.

Table 6. Per-Patient IS Implementation Cost Estimates Using National Survey Data.

IS (\$)	+	\$ _{RT} (\$/min)	×	t _{RT} (minutes) (95% CI)	+	\$ _{RN} (\$/min)	×	t _{RN} (min/h) (95% CI)	×	16 h/d	×	LOS (days)	Cost (\$) (95% CI)
12.92		0.49		16.4 (14.7-18.4)		0.60		2.0 (1.8-2.2)				4.5	107.36 (97.88-116.98)

Note. IS = incentive spirometry; \$_{RT} = RT's mean per-minute wage (\$); RT = respiratory therapist; t_{RT} = RTs' spent initially educating a patient to use their IS; CI = confidence interval; \$_{RN} = nurses' mean per-minute wage (\$); t_{RN} = minutes per hour nurses spend doing IS-related activities per patient; LOS = mean hospital length of stay (days).

Discussion

This study estimated the cost of implementing IS. IS is used in more than 95% of hospitals across the United States, even though no clear clinical or cost benefit has been documented. The current study attempted to define the cost of using IS in an academic cardiothoracic surgery step-down unit and correlate those data with the national average costs. To accomplish that, a survey was conducted at AMC and nationally by targeting relevant nursing and respiratory therapy organizations.

Nurses and RTs report spending a substantial portion of their time performing IS-related activities. This is the first study to report IS implementation cost per patient (≥\$65.30), annual cost per a 10-bed cardiothoracic step-down unit at AMC (≥\$36 959.80), and annual expenditures in the United States (≥\$949.4 million). Provider time dedicated to IS was not significantly different between the AMC and national survey responses, validating the accuracy of the cost analysis.

Despite a dearth of evidence in reducing postoperative pulmonary complications, IS began to be prescribed in the 1970s after intermittent positive pressure breathing was proven therapeutically ineffective.³⁶ In addition to IS's benign safety profile² and physicians' lack of knowledge regarding costs,³⁹⁻⁴² the relatively low per-patient financial impact of implementation may explain why IS continues to be so widely prescribed today relative to the lack of evidence establishing its benefit. Until now, IS-prescribing physicians had no data on the time and effort required of nurses and RTs to implement IS. When compared directly with costs of other tests or interventions, IS may appear relatively low. However, when implementation is considered at the step-down unit, hospital, or nationwide levels, the estimated \$1.04 billion cost spent on IS in the United States is substantial, especially given the lack of efficacy data.

As health care costs continue to rise in an unsustainable fashion, so does the need to cut out wasteful spending.⁴³ Per Berwick et al, major sources of waste in medicine include

(1) failures of care delivery, (2) failures of care coordination, (3) overtreatment, (4) administrative complexity, (5) pricing failures, and (6) fraud and abuse.⁴³ Until now, IS-prescribing physicians had no data on the time and effort required of nurses and RTs to implement IS. In 2008, health care expenditure totaled \$2.4 trillion comprising 16% of the US gross domestic product.⁴⁴ However, when implementation is considered at the step-down unit, hospital, or nationwide levels, the estimated \$1.04 billion cost spent on IS in the United States is substantial, especially given the lack of efficacy data. IS falls into the category, as delineated by Berwick et al, of “Overtreatment”—when patients are subjected to care not necessarily grounded in literature or science.⁴³ Other examples of overtreatment include excessive use of antibiotics, excessive and even unwanted care toward the end of life, and the use of expensive, more lucrative, treatment modalities which conservative treatment is sufficient.⁴³ The category of “Overtreatment” as a whole is thought to comprise between 158 and 226 billion dollars of wasteful spending.

This study has several potential limitations. National survey respondents were members of professional societies, which may have created a sampling bias of individuals who tend to have certain perspectives. Surveys ideally would be distributed nationwide to all RTs and nurses. Whether respondents had an Master of Science in Respiratory Care degree was not surveyed. Time reported by providers is subjectively recollected and may be influenced by recall bias. The ideal observation method would be quantified provider time measured via video recording; however, implementation of such procedures is not realistic. Last, the local survey was performed in a cardiothoracic step-down unit, and estimates may not be generalizable to all inpatient bed or other types of step-down units. More data are needed regarding specific uses, costs, efficacy, and compliance. Without adequate clinical efficacy data, whether or not IS should continue to be prescribed warrants consideration. As overall costs of health care continue to rise,²⁹ allocation of resources to early mobilization^{45,46} and optimizing pain control,⁴⁷⁻⁵⁶ as well as other evidence-based therapies, may be more appropriate.

IS makes clinical sense but has limited support in the literature and is associated with very high costs. The current data highlight a broader need for greater critical inquiry concerning the evidence supporting and costs associated with common medical practices in an effort to avoid unnecessary spending and optimize quality care. Further studies are needed to determine clinical efficacy and cost-effectiveness of IS to determine whether costs are justifiable.

Acknowledgments

For distributing the survey, special thanks to the national societies mentioned.


Declaration of Conflicting Interests

The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: Several hypotheses are being explored from the data from the survey to the national societies (American Society of Peri-Anesthesia Nurses [ASPAN], American Association For Respiratory Care [AARC], Academy of Medical-Surgical Nurses [AMSN], American Association of Critical-Care Nurses [AACN]), with articles under consideration elsewhere related to providers' perspectives on incentive spirometry (IS), IS use protocols, and IS use compliance. Although the respondents are the same, the hypotheses and most variables under investigation are distinct necessitating separation of discussion. Dr. Daniels reports relationships with DePuy, Globus Medical, Orthofix, Springer, and Stryker. Dr. Eltorai reports relationships with Springer, Quitbit, and Lippincott and being listed as an inventor on the patent application related to the reminder. Dr. Sellke reports a relationship with Stryker and Boehringer Ingelheim. The other authors report no conflicts.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Adam E. M. Eltorai  <https://orcid.org/0000-0001-5749-6631>

References

- Cassidy MR, Rosenkranz P, McCabe K, Rosen JE, McAneny D. I COUGH: reducing postoperative pulmonary complications with a multidisciplinary patient care program. *JAMA Surg*. 2013;148(8):740-745.
- Restrepo RD, Wettstein R, Wittnebel L, Tracy M. Incentive spirometry: 2011. *Respir Care*. 2011;56(10):1600-1604. doi:10.4187/respcare.01471.
- O'Donohue WJ. National survey of the usage of lung expansion modalities for the prevention and treatment of postoperative atelectasis following abdominal and thoracic surgery. *Chest*. 1985;87(1):76-80.
- Agostini P, Singh S. Incentive spirometry following thoracic surgery: what should we be doing? *Physiotherapy*. 2009;95(2):76-82.
- Ballantyne JC, Carr DB, deFerranti S, et al. The comparative effects of postoperative analgesic therapies on pulmonary outcome: cumulative meta-analyses of randomized, controlled trials. *Anesth Analg*. 1998;86(3):598-612.
- Bapojé SR, Whitaker JF, Schulz T, Chu ES, Albert RK. Preoperative evaluation of the patient with pulmonary disease. *Chest*. 2007;132(5):1637-1645.
- Crowe JM, Bradley CA. The effectiveness of incentive spirometry with physical therapy for high-risk patients after coronary artery bypass surgery. *Phys Ther*. 1997;77(3):260-268.
- Desai PM. Pain management and pulmonary dysfunction. *Crit Care Clin*. 1999;15(1):151-166, vii.
- Dias CM, Vieira Rde O, Oliveira JF, Lopes AJ, Menezes SL, Guimaraes FS. Three physiotherapy protocols: effects on pulmonary volumes after cardiac surgery. *J Bras Pneumol*. 2011;37(1):54-60.

10. Gosselink R, Schrever K, Cops P, et al. Incentive spirometry does not enhance recovery after thoracic surgery. *Crit Care Med.* 2000;28(3):679-683.
11. Guimaraes MM, El Dib R, Smith AF, Matos D. Incentive spirometry for prevention of postoperative pulmonary complications in upper abdominal surgery. *Cochrane Database Syst Rev.* 2009(3):CD006058.
12. Hall JC, Tarala R, Harris J, Tapper J, Christiansen K. Incentive spirometry versus routine chest physiotherapy for prevention of pulmonary complications after abdominal surgery. *Lancet.* 1991;337(8747):953-956.
13. Hall JC, Tarala RA, Tapper J, Hall JL. Prevention of respiratory complications after abdominal surgery: a randomised clinical trial. *BMJ.* 1996;312(7024):148-152; discussion 152-153.
14. Mang H, Kacmarek RM. Prevention of pulmonary complications after abdominal surgery. *Lancet.* 1991;338(8762):312-313.
15. Pasquina P, Tramer MR, Walder B. Prophylactic respiratory physiotherapy after cardiac surgery: systematic review. *BMJ.* 2003;327(7428):1379.
16. Pasquina P, Tramer MR, Granier JM, Walder B. Respiratory physiotherapy to prevent pulmonary complications after abdominal surgery: a systematic review. *Chest.* 2006;130(6):1887-1899.
17. Schwieger I, Gamulin Z, Forster A, Meyer P, Gemperle M, Suter PM. Absence of benefit of incentive spirometry in low-risk patients undergoing elective cholecystectomy. *Chest.* 1986;89(5):652-656.
18. Thomas JA, McIntosh JM. Are incentive spirometry, intermittent positive pressure breathing, and deep breathing exercises effective in the prevention of postoperative pulmonary complications after upper abdominal surgery? a systematic overview and meta-analysis. *Phys Ther.* 1994;74(1):3-10; discussion 10-16.
19. Varela G, Ballesteros E, Jimenez MF, Novoa N, Aranda JL. Cost-effectiveness analysis of prophylactic respiratory physiotherapy in pulmonary lobectomy. *Eur J Cardiothorac Surg.* 2006;29(2):216-220.
20. Vats N. Effect of deep breathing exercises and incentive spirometry in the prevention of postoperative pulmonary complications in the patients of cancer esophagus undergoing esophagectomy. *Ind J Physiother Occup Ther.* 2009;3(3):321-325.
21. Weiner P, Man A, Weiner M, et al. The effect of incentive spirometry and inspiratory muscle training on pulmonary function after lung resection. *J Thorac Cardiovasc Surg.* 1997;113(3):552-557.
22. Winters BA. Older adults with traumatic rib fractures: an evidence-based approach to their care. *J Trauma Nurs.* 2009;16(2):93-97.
23. Overend TJAC, Lucy SD, Bhatia C, Jonsson BI, Timmermans C. The effect of incentive spirometry on postoperative pulmonary complications. *Chest.* 2001;120(3):971-978.
24. Carvalho CR, Paisani DM, Lunardi AC. Incentive spirometry in major surgeries: a systematic review. *Rev Bras Fisioter.* 2011;15(5):343-350.
25. do Nascimento JP, MÓdolo NS, Andrade S, Guimarães MM, Braz LG, El Dib R. Incentive spirometry for prevention of postoperative pulmonary complications in upper abdominal surgery. *Cochrane Database Syst Rev.* 2014(2):CD006058.
26. Freitas ER, Soares BG, Cardoso JR, Atallah AN. Incentive spirometry for preventing pulmonary complications after coronary artery bypass graft. *Cochrane Database Syst Rev.* 2012(9):CD004466.
27. Berwick DM, Nolan TW, Whittington J. The triple aim: care, health, and cost. *Health Aff (Millwood).* 2008;27(3):759-769.
28. Cohen ER, Feinglass J, Barsuk JH, et al. Cost savings from reduced catheter-related bloodstream infection after simulation-based education for residents in a medical intensive care unit. *Simul Healthc.* 2010;5(2):98-102.
29. Cutler DM, Rosen AB, Vijan S. The value of medical spending in the United States, 1960-2000. *N Engl J Med.* 2006;355(9):920-927.
30. Emanuel EJ. Where are the health care cost savings? *JAMA.* 2012;307(1):39-40.
31. James BC, Savitz LA. How Intermountain trimmed health care costs through robust quality improvement efforts. *Health Aff (Millwood).* 2011;30(6):1185-1191.
32. Lawson EH, Hall BL, Louie R, et al. Association between occurrence of a postoperative complication and readmission: implications for quality improvement and cost savings. *Ann Surg.* 2013;258(1):10-18.
33. Thorpe KE. The rise in health care spending and what to do about it. *Health Aff (Millwood).* 2005;24(6):1436-1445.
34. Coinnews Media Group. *US Inflation Calculator.* San Antonio, TX: CoinNews Media Group LLC. 2015.
35. Tri-anim. *Coach 2 Incentive Spirometers: SMITHS MEDICAL ASD INC: Incentive Spirometer.* Coach 2, 4000 ml Capacity, No 1-Way Valve, Mouthpiece Holder: 26-22-4001EA. <https://www.tri-anim.com/coach-2-incentive-spirometers-group-26838-3827.aspx?search=26-22-4001EA>. Published 2015. Accessed August 10, 2018.
36. Handelsman H. Intermittent positive pressure breathing (IPPB) therapy. *Health Technol Assess Rep.* 1991;(1):1-9. <https://www.ncbi.nlm.nih.gov/pubmed/1810351>
37. Weiss AJ. *Overview of Hospital Stays in the United States, 2012.* Statistical Brief #180. Rockville, MD: Healthcare Cost and Utilization Project, Agency for Healthcare Research and Quality; 2014.
38. Healthcare Cost and Utilization Project. *Inpatient vs. Outpatient Surgeries in U.S. Hospitals.* Rockville, MD: Agency for Healthcare Research and Quality; 2015.
39. Allan GM, Lexchin J, Wiebe N. Physician awareness of drug cost: a systematic review. *PLoS Med.* 2007;4(9):e283.
40. Allan GM, Lexchin J. Physician awareness of diagnostic and nondrug therapeutic costs: a systematic review. *Int J Technol Assess Health Care.* 2008;24(2):158-165.
41. Ryskina KL, Smith CD, Weissman A, et al. U.S. internal medicine residents' knowledge and practice of high-value care: a national survey. *Acad Med.* 2015;90(10):1373-1379.
42. Tek Sehgal R, Gorman P. Internal medicine physicians' knowledge of health care charges. *J Grad Med Educ.* 2011;3(2):182-187.
43. Hackbarth AD, Hackbarth AD. Eliminating waste in US health care. *JAMA.* 2012;307(14):1513-1516. doi:10.1001/jama.2012.362.
44. Fred HL. Cutting the cost of health care: the physician's role. *Texas Hear Inst J.* 2016;43(1):4-6. doi:10.14503/THIJ-15-5646.
45. Haines KJ, Skinner EH, Berney S, Austin Health POST Study Investigators. Association of postoperative pulmonary complications with delayed mobilisation following major abdominal surgery: an observational cohort study. *Physiotherapy.* 2013;99(2):119-125.

46. van der Leeden M, Huijsmans R, Geleijn E, et al. Early enforced mobilisation following surgery for gastrointestinal cancer: feasibility and outcomes. *Physiotherapy*. 2016;102(1):103-110.
47. Berrizbeitia LD, Tessler S, Jacobowitz IJ, Kaplan P, Budzilowicz L, Cunningham JN. Effect of sternotomy and coronary bypass surgery on postoperative pulmonary mechanics. Comparison of internal mammary and saphenous vein bypass grafts. *Chest*. 1989;96(4):873-876.
48. Daganou M, Dimopoulou I, Michalopoulos N, et al. Respiratory complications after coronary artery bypass surgery with unilateral or bilateral internal mammary artery grafting. *Chest*. 1998;113(5):1285-1289.
49. Ferguson MK. Preoperative assessment of pulmonary risk. *Chest*. 1999;115(5 suppl):58S-63S.
50. Groeneveld AB, Verheij J, van den Berg FG, Wisselink W, Rauwerda JA. Increased pulmonary capillary permeability and extravascular lung water after major vascular surgery: effect on radiography and ventilatory variables. *Eur J Anaesthesiol*. 2006;23(1):36-41.
51. Groeneveld AB, Jansen EK, Verheij J. Mechanisms of pulmonary dysfunction after on-pump and off-pump cardiac surgery: a prospective cohort study. *J Cardiothorac Surg*. 2007;2:11.
52. Hallbook TLB, Lindroth B, Wolff T. Prophylaxis against pulmonary complications in patients undergoing gall-bladder surgery. A comparison between early mobilization, physiotherapy with and without bronchodilatation. *Ann Chir Gynaecol*. 1984;73(2):55-58.
53. Hulzebos EH, Van Meeteren NL, De Bie RA, Dagnelie PC, Helder PJ. Prediction of postoperative pulmonary complications on the basis of preoperative risk factors in patients who had undergone coronary artery bypass graft surgery. *Phys Ther*. 2003;113(5):994-1002.
54. Kips J. Preoperative pulmonary evaluation. *Acta Clinica Belgica*. 1997;52(5):301-305.
55. Kulkarni SR, Fletcher E, McConnell AK, Poskitt KR, Whyman MR. Pre-operative inspiratory muscle training preserves post-operative inspiratory muscle strength following major abdominal surgery—a randomised pilot study. *Ann R Coll Surg Engl*. 2010;92(8):700-707.
56. Mohr DN, Lavender RC. Preoperative pulmonary evaluation. Identifying patients at increased risk for complications. *Postgrad Med*. 1996;100(5):241-256.