

11-19-2015

22. Relativity

Gerhard Müller
University of Rhode Island, gmuller@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/elementary_physics_2

Abstract

Lecture slides 22 for Elementary Physics II (PHY 204), taught by Gerhard Müller at the University of Rhode Island.

Some of the slides contain figures from the textbook, Paul A. Tipler and Gene Mosca. *Physics for Scientists and Engineers*, 5th/6th editions. The copyright to these figures is owned by W.H. Freeman. We acknowledge permission from W.H. Freeman to use them on this course web page. The textbook figures are not to be used or copied for any purpose outside this class without direct permission from W.H. Freeman.

Recommended Citation


Müller, Gerhard, "22. Relativity" (2015). *PHY 204: Elementary Physics II (2015)*. Paper 6.
https://digitalcommons.uri.edu/elementary_physics_2/6https://digitalcommons.uri.edu/elementary_physics_2/6

This Course Material is brought to you for free and open access by the Physics Open Educational Resources at DigitalCommons@URI. It has been accepted for inclusion in PHY 204: Elementary Physics II (2015) by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Is There Absolute Motion?



Forces between two long, parallel, charged rods




- $\frac{F_E}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1 \lambda_2}{d}$ (left), $\frac{F_E^*}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1^* \lambda_2^*}{d}$, $\frac{F_B}{L} = \frac{\mu_0}{2\pi} \frac{I_1 I_2}{d}$, (right)
- $\frac{F_E^* - F_B}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1^* \lambda_2^*}{d} \left(1 - \frac{v^2}{c^2}\right) = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1 \lambda_2}{d}$
- $c = \frac{1}{\sqrt{\epsilon_0 \mu_0}} = 2.998 \times 10^8 \text{ ms}^{-1}$ (speed of light)
- $\lambda_1^* = \frac{\lambda_1}{\sqrt{1 - v^2/c^2}}$, $\lambda_2^* = \frac{\lambda_2}{\sqrt{1 - v^2/c^2}}$ (due to length contraction)

Catching Up with a Photon? (1)




Forces between two long, parallel, charged rods in relative motion.



- Galilean kinematics predicts $u = v_1 + v_2$.
- Relativistic kinematics requires $v_1 < c$, $v_2 < c$, $u < c$.
- Relativistic dynamics requires $F_E^* - F_B = F_E$.
- Length-contracted charge densities: $\lambda(v) = \frac{\lambda(0)}{\sqrt{1 - v^2/c^2}}$.
- Electric currents: $I(v) = \lambda(v)v$.

Catching Up with a Photon? (2)



- $\frac{F_E}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda(0)\lambda(v_1)}{d}, \quad \frac{F_E^*}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda(v_2)\lambda(u)}{d}.$
- $\frac{F_B}{L} = \frac{\mu_0}{2\pi} \frac{[\lambda(v_2)v_2][\lambda(u)u]}{d} = \frac{1}{2\pi\epsilon_0} \frac{\lambda(v_2)\lambda(u)}{d} \frac{v_2u}{c^2}.$
- $\frac{F_E^*}{L} - F_B = \frac{F_E}{L} \Rightarrow \frac{1}{2\pi\epsilon_0} \frac{\lambda(v_2)\lambda(u)}{d} \left(1 - \frac{v_2u}{c^2}\right) = \frac{1}{2\pi\epsilon_0} \frac{\lambda(0)\lambda(v_1)}{d}$
- $\Rightarrow \frac{1}{\sqrt{1 - v_2^2/c^2}} \frac{1}{\sqrt{1 - u^2/c^2}} \left(1 - \frac{v_2u}{c^2}\right) = \frac{1}{\sqrt{1 - v_1^2/c^2}}$ to be solved for u .
- Relativistic kinematic predicts $u = \frac{v_1 + v_2}{1 + v_1v_2/c^2} < c.$