

2018

Draft Genome Sequence of *Bowmanella denitrificans* JL63, a Bacterium Isolated from Whiteleg Shrimp (*Litopenaeus vannamei*) That Can Inhibit the Growth of *Vibrio Parahaemolyticus*

Jason P. LaPorte
University of Rhode Island

Edward J. Spinard
University of Rhode Island

See next page for additional authors

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Follow this and additional works at: https://digitalcommons.uri.edu/favs_facpubs

Citation/Publisher Attribution

LaPorte J. P., Spinard E. J., Gomez-Chiarri, M., Rowley D. C., Mekalanos, J. J., & Nelson, R. D. (2018). Draft genome sequence of *Bowmanella denitrificans* JL63, a bacterium isolated from whiteleg shrimp (*Litopenaeus vannamei*) that can inhibit the growth of *Vibrio parahaemolyticus*. *Genome Announcements* 6:e00215-18. <https://doi.org/10.1128/genomeA.00215-18>
Available at: <https://doi.org/10.1128/genomeA.00215-18>

This Article is brought to you for free and open access by the Fisheries, Animal and Veterinary Sciences at DigitalCommons@URI. It has been accepted for inclusion in Fisheries, Animal and Veterinary Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors

Jason P. LaPorte, Edward J. Spinard, Marta Gomez-Chiarri, David C. Rowley, John J. Mekalanos, and David R. Nelson



Draft Genome Sequence of *Bowmanella denitrificans* JL63, a Bacterium Isolated from Whiteleg Shrimp (*Litopenaeus vannamei*) That Can Inhibit the Growth of *Vibrio parahaemolyticus*

Jason P. LaPorte,^a Edward J. Spinard,^a Marta Gomez-Chiarri,^b David C. Rowley,^c John J. Mekalanos,^d  David R. Nelson^a

^aDepartment of Cell and Molecular Biology, University of Rhode Island, Kingston, Rhode Island, USA

^bDepartment of Fisheries, Animal and Veterinary Sciences, University of Rhode Island, Kingston, Rhode Island, USA

^cDepartment of Biomedical and Pharmaceutical Sciences, University of Rhode Island, Kingston, Rhode Island, USA

^dDepartment of Microbiology and Immunobiology, Harvard Medical School, Boston, Massachusetts, USA

ABSTRACT *Bowmanella denitrificans* strain JL63 was isolated from a whiteleg shrimp (*Litopenaeus vannamei*) and was determined to have antibacterial activity against an acute hepatopancreatic necrosis disease (AHPND) strain of *Vibrio parahaemolyticus*. Here, we report the draft genome sequence of this strain and identify genes that are potentially involved in its antibacterial activity.

Bowmanella, a genus in the family Alteromonadaceae within the Gammaproteobacteria, was first identified in 2006 (1). Currently, only three species belonging to this genus have been described, *Bowmanella denitrificans*, *Bowmanella pacifica*, and *Bowmanella dokdonensis*. *B. denitrificans* is a chemoorganotrophic bacterium capable of respiratory, but not fermentative, metabolism (1). *B. denitrificans* strain BD1^T, the first strain of this species to be identified, is capable of anaerobic growth by carrying out denitrification, whereas *B. denitrificans* strain S088 has been shown to produce a potent heat-stable algicidal compound (1, 2). *B. denitrificans* JL63 was isolated from a whiteleg shrimp (*Litopenaeus vannamei*). A zone-of-inhibition assay (3) determined that *B. denitrificans* JL63 can inhibit the growth of an acute hepatopancreatic necrosis disease (AHPND) strain of *Vibrio parahaemolyticus* on an agar surface. The *B. denitrificans* JL63 genome sequence reported here is the first draft genome sequence of a *Bowmanella* species.

B. denitrificans JL63 was grown overnight in yeast-tryptone broth supplemented with 3% artificial sea salts (LB30IOS) at 27°C. Genomic DNA was isolated using the Promega Wizard genomic DNA purification kit, and DNA was resuspended in 2 mM Tris-HCl buffer (Bio Basic). Sequencing was performed at the Rhode Island Genomics and Sequencing Center using an Illumina MiSeq sequencer. Sequence trimming was performed using CLC Genomics Workbench (version 9.5.3), resulting in 2,641,396 paired-end reads averaging 180 bp in size. Contigs with a coverage of $\geq 34\times$ were assembled using the SPAdes genomic assembler (version 3.1.1) (4). The resulting contigs were processed using the CLC Microbial Genome Finishing module. The draft genome consists of 39 contigs, with a total sequence length of 5,479,914 bp and a G+C content of 50.4%. Gene annotation was performed using the Rapid Annotations using Subsystems Technology (RAST) server and resulted in 4,980 open reading frames (5–7). The 16S rRNA gene of *B. denitrificans* JL63 is 99.8% similar to that of *B. denitrificans* BD1^T, 99.0% similar to that of *B. pacifica* W3-3A, and 95.0% similar to that of *B. dokdonensis* UDC354. The *B. denitrificans* JL63 *gyrB* and *rpoD* genes are 98.6% and 98.3% similar to

Received 20 February 2018 Accepted 16 March 2018 Published 5 April 2018

Citation LaPorte JP, Spinard EJ, Gomez-Chiarri M, Rowley DC, Mekalanos JJ, Nelson RD. 2018. Draft genome sequence of *Bowmanella denitrificans* JL63, a bacterium isolated from whiteleg shrimp (*Litopenaeus vannamei*) that can inhibit the growth of *Vibrio parahaemolyticus*. Genome Announc 6:e00215-18. <https://doi.org/10.1128/genomeA.00215-18>.

Copyright © 2018 LaPorte et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to David R. Nelson, dnelson@uri.edu.

those of *B. denitrificans* BD1^T and 81.2% and 80.8% similar to those of *B. pacifica* W3-3A, respectively.

The genome of *B. denitrificans* JL63 encodes several gene clusters that are potentially involved in the production of the following antibacterial compounds: lanthionine, colicin V, a secreted hemolysin-type calcium-binding bacteriocin, and the broad-spectrum antibacterial protein marinocin (encoded by the *lodAB* operon). The genome also encodes an antibiotic biosynthesis monooxygenase, a type VI secretion system (T6SS), and two iron acquisition systems (hemin and TonB), including the full complement of proteins responsible for the formation of the TonB-ExbB-ExbD complex. Gene clusters for a type IV pilus and a mannose-sensitive hemagglutinin type IV pilus system are also present. The genome contains gene clusters for denitrification and for nitrate/nitrite ammonification. Three secondary metabolic gene clusters in the JL63 genome were predicted using antiSMASH (8). Gene clusters predicted to synthesize a nonribosomal peptide and two bacteriocins, including lantipeptide, were identified.

Accession number(s). This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. [PEBU00000000](https://doi.org/10.1093/jgs.0.64306-0). The version described in this paper is the first version, PEBU01000000.

ACKNOWLEDGMENTS

This work was supported by an award from CP Foods to D. R. Nelson, M. Gomez-Chiarri, D. C. Rowley, and J. J. Mekalanos. The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

We thank William Robins at the Mekalanos Lab, Harvard Medical School, for providing us with the whiteleg shrimp (*L. vannamei*) specimens from which we isolated *B. denitrificans* JL63.

REFERENCES

- Jean WD, Chen JS, Lin YT, Shieh WY. 2006. *Bowmanella denitrificans* gen. nov., sp. nov., a denitrifying bacterium isolated from seawater from An-Ping Harbour, Taiwan. *Int J Syst Evol Microbiol* 56:2463–2467. <https://doi.org/10.1099/ijs.0.64306-0>.
- Jiang X, Ren C, Hu C, Zhao Z. 2014. Isolation and algicidal characterization of *Bowmanella denitrificans* S088 against *Chlorella vulgaris*. *World J Microbiol Biotechnol* 30:621–629. <https://doi.org/10.1007/s11274-013-1478-y>.
- Zhao W, Dao C, Karim M, Gomez-Chiarri M, Rowley D, Nelson DR. 2016. Contributions of tropodithietic acid and biofilm formation to the probiotic activity of *Phaeobacter inhibens*. *BMC Microbiol* 16:1. <https://doi.org/10.1186/s12866-015-0617-z>.
- Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol* 20:714–737. <https://doi.org/10.1089/cmb.2013.0084>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, III, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.