

2016

# The Game Saturation Number of a Graph

James M. Carraher

William B. Kinnersley

University of Rhode Island, billk@uri.edu

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.uri.edu/math\\_facpubs](https://digitalcommons.uri.edu/math_facpubs)

**The University of Rhode Island Faculty have made this article openly available.  
Please let us know how Open Access to this research benefits you.**

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

## Citation/Publisher Attribution

Carraher, J. M., Kinnersley, W. B., Reiniger, B., & West, D. B. (2016). The Game Saturation Number of a Graph. *Journal of Graph Theory*, 85(2), 481-495. doi:10.1002/jgt.22074  
Available at: <https://doi.org/10.1002/jgt.22074>

This Article is brought to you for free and open access by the Mathematics at DigitalCommons@URI. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

---

**Authors**

James M. Carraher, William B. Kinnersley, Benjamin Reiniger, and Douglas B. West

# The Game Saturation Number of a Graph

James M. Carraher\*, William B. Kinnersley†,  
Benjamin Reiniger‡, Douglas B. West§

May 10, 2016

## Abstract

Given a family  $\mathcal{F}$  and a host graph  $H$ , a graph  $G \subseteq H$  is  $\mathcal{F}$ -saturated relative to  $H$  if no subgraph of  $G$  lies in  $\mathcal{F}$  but adding any edge from  $E(H) - E(G)$  to  $G$  creates such a subgraph. In the  $\mathcal{F}$ -saturation game on  $H$ , players *Max* and *Min* alternately add edges of  $H$  to  $G$ , avoiding subgraphs in  $\mathcal{F}$ , until  $G$  becomes  $\mathcal{F}$ -saturated relative to  $H$ . They aim to maximize or minimize the length of the game, respectively;  $\text{sat}_g(\mathcal{F}; H)$  denotes the length under optimal play (when *Max* starts).

Let  $\mathcal{O}$  denote the family of odd cycles and  $\mathcal{T}_n$  the family of  $n$ -vertex trees, and write  $F$  for  $\mathcal{F}$  when  $\mathcal{F} = \{F\}$ . Our results include  $\text{sat}_g(\mathcal{O}; K_n) = \lfloor \frac{n}{2} \rfloor \lceil \frac{n}{2} \rceil$ ,  $\text{sat}_g(\mathcal{T}_n; K_n) = \binom{n-2}{2} + 1$  for  $n \geq 6$ ,  $\text{sat}_g(K_{1,3}; K_n) = 2 \lfloor \frac{n}{2} \rfloor$  for  $n \geq 8$ , and  $\text{sat}_g(P_4; K_n) \in \{\lfloor \frac{4n}{5} \rfloor, \lceil \frac{4n}{5} \rceil\}$  for  $n \geq 5$ . We also determine  $\text{sat}_g(P_4; K_{m,n})$ ; with  $m \geq n$ , it is  $n$  when  $n$  is even,  $m$  when  $n$  is odd and  $m$  is even, and  $m + \lfloor n/2 \rfloor$  when  $mn$  is odd. Finally, we prove the lower bound  $\text{sat}_g(C_4; K_{n,n}) \geq \frac{1}{21}n^{13/12} - O(n^{35/36})$ . The results are very similar when *Min* plays first, except for the  $P_4$ -saturation game on  $K_{m,n}$ .

## 1 Introduction

The archetypal question in extremal graph theory asks for the maximum number of edges in an  $n$ -vertex graph that does not contain a specified graph  $F$  as a subgraph. The answer is called the *extremal number* of  $F$ , denoted  $\text{ex}(F; n)$ . The celebrated theorem of Turán [22] gives the answer when  $F$  is the complete graph  $K_r$  and determines the largest  $n$ -vertex

---

\*Department of Mathematics and Statistics, University of Nebraska, Kearney, NE: carraherjm2@unk.edu. Research supported by NSF grant DMS 09-14815

†Department of Mathematics, University of Rhode Island, Kingston, RI, USA, 02881: billk@uri.edu

‡Mathematics Department, Ryerson University, Toronto ON: reiniger@ryerson.ca

§Mathematics Departments, Zhejiang Normal University (Jinhua, Zhejiang, China) and University of Illinois (Urbana, IL, U.S.A): west@math.uiuc.edu. Research supported by Recruitment Program of Foreign Experts, 1000 Talent Plan, State Administration of Foreign Experts Affairs, China.

graphs not containing  $K_r$  (the *size* of a graph is its number of edges, and by *largest* we mean “of maximum size”).

We consider maximal graphs not containing  $F$ . The concept extends to a family  $\mathcal{F}$  of graphs. A graph  $G$  is  $\mathcal{F}$ -saturated if no subgraph of  $G$  belongs to  $\mathcal{F}$  but  $G + e$  contains a member of  $\mathcal{F}$  whenever  $e \in E(\overline{G})$ . The extremal number  $\text{ex}(\mathcal{F}; n)$  is the maximum size of an  $\mathcal{F}$ -saturated  $n$ -vertex graph. (In all notation involving families of graphs, we write  $\mathcal{F}$  as  $F$  when  $\mathcal{F}$  consists of a single graph  $F$ .)

One may also ask for the minimum size of an  $\mathcal{F}$ -saturated  $n$ -vertex graph; this is the *saturation number* of  $\mathcal{F}$ , denoted  $\text{sat}(\mathcal{F}; n)$ . Erdős, Hajnal, and Moon [6] initiated the study of graph saturation by determining  $\text{sat}(K_r; n)$ .

Generalizing further, a subgraph  $G$  of a host graph  $H$  is  $\mathcal{F}$ -saturated relative to  $H$  if no subgraph of  $G$  lies in  $\mathcal{F}$  but adding any edge of  $E(H) - E(G)$  to  $G$  completes a subgraph belonging to  $\mathcal{F}$ . Although  $\text{ex}(\mathcal{F}; n)$  and  $\text{sat}(\mathcal{F}; n)$  concern saturation relative to  $K_n$ , saturation has also been studied relative to other graphs. For example, Zarankiewicz’s Problem, which asks for the largest subgraph of  $K_{n,n}$  not containing  $K_{t,t}$ , involves saturation relative to  $K_{n,n}$ . When two agents have opposing interests in creating a large or a small  $\mathcal{F}$ -saturated graph, we obtain the “saturation game”.

**Definition 1.1.** The  $\mathcal{F}$ -saturation game on a host graph  $H$  has players *Max* and *Min*. The players jointly construct a subgraph  $G$  of  $H$  by iteratively adding one edge of  $H$ , constrained by  $G$  having no subgraph that lies in  $\mathcal{F}$ . The game ends when  $G$  becomes  $\mathcal{F}$ -saturated relative to  $H$ . Max aims to maximize the length of the game; Min aims to minimize it. When both players play optimally, the length of the game is the *game  $\mathcal{F}$ -saturation number* of  $H$ , denoted  $\text{sat}_g(\mathcal{F}; H)$  when Max starts the game and by  $\text{sat}'_g(\mathcal{F}; H)$  when Min starts it. For clarity and for consistency with the extremal and saturation numbers, we write the values as  $\text{sat}_g(\mathcal{F}; n)$  and  $\text{sat}'_g(\mathcal{F}; n)$  when playing on  $K_n$ .

The saturation game generalizes to any hereditary family of sets. Let  $D$  be a family of subsets of a set  $X$  such that every subset of a member of  $D$  also belongs to  $D$ . The saturated subsets are the maximal elements of  $D$ . Max and Min alternately add elements of  $X$  to a set that always lies in  $D$ . The game ends when a saturated set is reached, with Max and Min having the same goals as before. We denote the length of the game under optimal play by  $\text{sat}_g(D; X)$  when Max starts and by  $\text{sat}'_g(D; X)$  when Min starts. In the  $\mathcal{F}$ -saturation game on  $H$ , we have  $X = E(H)$ , and avoiding subgraphs in  $\mathcal{F}$  defines the hereditary family  $D$ .

Patkós and Vizer [21] introduced this general model and studied the case where  $X$  is the family of  $k$ -element subsets of  $\{1, \dots, n\}$  and  $D$  is the set of intersecting families of  $k$ -sets. View  $X$  as the  $n$ -vertex complete  $k$ -uniform hypergraph  $K_n^{(k)}$ . Let  $M$  denote the forbidden subgraph consisting of two disjoint edges. The Erdős–Ko–Rado Theorem [7] then states  $\text{ex}(M; K_n^{(k)}) = \binom{n-1}{k-1} \sim \frac{1}{(k-1)!} n^{k-1}$ . Füredi [12] proved that  $\text{sat}(M; K_n^{(k)}) \leq \frac{3}{4} k^2$  when

a projective plane of order  $k/2$  exists. For  $k \geq 2$ , Patkós and Vizer [21] proved  $\Omega(n^{\lfloor k/3 \rfloor - 5}) \leq \text{sat}_g(M; K_n^{(k)}) \leq O(n^{k - \sqrt{k}/2})$ .

The saturation game is also related to other well-studied graph games. In a *Maker-Breaker* game, the players Maker and Breaker take turns choosing edges of a host graph  $H$ , typically  $K_n$ . Maker wins by claiming all of the edges in a subgraph of  $H$  having some specified property  $\mathcal{P}$ , and Breaker wins by preventing this. For example, Hefetz, Krivelevich, Stojaković, and Szabó [16] studied Maker-Breaker games played on  $K_n$  in which Maker seeks to build non-planar graphs, non- $k$ -colorable graphs, or  $K_t$ -minors. Several papers have considered the minimum number of turns needed for Maker to win (see [10, 17]). In this context, Breaker behaves like Max in the saturation game, making the game last as long as possible. In the saturation game both players contribute edges, but here Maker cannot use the edges taken by Breaker.

The  $\mathcal{F}$ -saturation game on  $H$  is also related to the process in which edges are added at random, which in the literature has been called *the  $\mathcal{F}$ -free process*; the length of the process is the number of moves to reach a graph that is  $\mathcal{F}$ -saturated relative to  $H$ . Usually  $H = K_n$  (see [3, 4, 8, 20]), but [1] is more general. When  $\mathcal{F} = \{C_4\}$  and  $H = K_{n,n}$ , the lower bound of [1] specializes to  $\Omega(n^{4/3} \log^{1/3} n)$ .

The saturation game on graphs was introduced by Füredi, Reimer, and Seress [14]; they studied  $\text{sat}_g(K_3; n)$ , calling it “a variant of Hajnal’s triangle-free game”. In Hajnal’s original “triangle-free game”, the players aim only to avoid creating triangles, and the loser is the player first forced to create one (Ferrara, Jacobson, and Harris [11] considered the generalization of Hajnal’s loser criterion to arbitrary  $\mathcal{F}$  and  $G$ ). Since the  $F$ -saturation game always ends with an  $F$ -saturated graph,  $n - 1 = \text{sat}(K_3; n) \leq \text{sat}_g(K_3; n) \leq \text{ex}(K_3; n) = \lfloor n^2/4 \rfloor$ ; hence  $\text{sat}_g(K_3; n) \in \Omega(n) \cap O(n^2)$ . Füredi et al. [14] proved  $\text{sat}_g(K_3; n) \in \Omega(n \log n)$ . Erdős claimed  $\text{sat}_g(K_3; n) \leq n^2/5$ , but his proof has been lost; the strongest bound appearing in print is the recent result of Biró, Horn, and Wildstrom [2], who proved  $\text{sat}_g(K_3; n) \leq \frac{26}{121}n^2 + o(n^2)$ . The correct order of growth remains unknown.

The  $P_3$ -saturation game was studied by Cranston, Kinnersley, O, and West [5]; here  $P_k$  denotes the  $k$ -vertex path. The subgraphs of  $H$  that are  $P_3$ -saturated relative to  $H$  are precisely the maximal matchings in  $H$ . Thus the game  $P_3$ -saturation number is just the *game matching number*, with  $\alpha'_g(G)$  and  $\hat{\alpha}'_g(G)$  denoting the values of the Max-start and Min-start games since  $\alpha'(G)$  denotes the maximum size of a matching in  $G$ . They proved  $\alpha'_g(G) \geq \frac{2}{3}\alpha'(G)$  for every graph  $G$  (with equality for some split graphs) and  $\alpha'_g(G) \geq \frac{3}{4}\alpha'(G)$  when  $G$  is a forest (with equality for some trees). They also showed that the minimum of  $\alpha'_g(G)$  over  $n$ -vertex 3-regular graphs is between  $n/3$  and  $7n/18$ .

We have mentioned bounds on  $\alpha'_g$  but not  $\hat{\alpha}'_g$  because the two parameters never differ by more than 1 (see [5]). This does not hold for  $\mathcal{F}$ -saturation in general. For example, when the host graph is obtained from a star with  $m$  edges by subdividing one edge, the

Max-start  $2K_2$ -saturation number is  $m$ , but the Min-start  $2K_2$ -saturation number is 2. As a less artificial example, we will show that  $|\text{sat}_g(P_4; K_{m,n}) - \text{sat}'_g(P_4; K_{m,n})|$  can be large, where  $K_{m,n}$  is the complete bipartite graph with part-sizes  $m$  and  $n$ . In most instances that we study, the choice of the starting player matters little.

In Section 2, we study the  $\mathcal{F}$ -saturation games on  $K_n$  for  $\mathcal{F} \in \{\mathcal{O}, \mathcal{T}_n, \{K_{1,3}\}, \{P_4\}\}$ , where  $\mathcal{O}$  is the family of all odd cycles and  $\mathcal{T}_n$  is the family of all  $n$ -vertex trees. We first prove  $\text{sat}_g(\mathcal{O}; n) = \text{sat}'_g(\mathcal{O}; n) = \lfloor \frac{n}{2} \rfloor \lceil \frac{n}{2} \rceil$ , achieving the trivial upper bound  $\text{ex}(\mathcal{O}; n)$ . For  $n \geq 3$ , we prove  $\text{sat}_g(\mathcal{T}_n; n) = \text{sat}'_g(\mathcal{T}_n; n) = \binom{n-2}{2} + 1$ , except  $\text{sat}_g(\mathcal{T}_5; 5) = 6$  and  $\text{sat}'_g(\mathcal{T}_4; 4) = 3$ ; note that  $\text{ex}(\mathcal{T}_n; n) = \binom{n-1}{2}$ . Hefetz et al. [15] have since studied more general versions of both of these problems. They studied  $\text{sat}_g(\mathcal{C}_k; n)$  and  $\text{sat}_g(\mathcal{X}_k; n)$  where  $\mathcal{C}_k$  is the family of  $k$ -connected graphs with  $n$  vertices and  $\mathcal{X}_k$  is the family of non- $k$ -colorable graphs. In both cases, the value is close to the extremal number. Lee and Riet [19] have generalized the tree problem in a different direction, studying  $\text{sat}_g(\mathcal{T}_k; n)$  for  $k \leq n$ .

Since  $\text{sat}(K_{1,3}; n) = n - 1$  and  $\text{ex}(K_{1,3}; n) = n$ , both  $\text{sat}_g(K_{1,3}; n)$  and  $\text{sat}'_g(K_{1,3}; n)$  always lie in  $\{n, n - 1\}$ . The two values differ when  $n \notin \{2, 3, 4, 7\}$ , with  $\text{sat}_g(K_{1,3}; n)$  being even and  $\text{sat}'_g(K_{1,3}; n)$  being odd. That is,  $\text{sat}_g(K_{1,3}; n) = 2 \lfloor n/2 \rfloor$  and  $\text{sat}'_g(K_{1,3}; n) = 2 \lceil n/2 \rceil - 1$  when  $n \geq 8$ . For  $n > r > 2$ , we have checked by computer that  $\text{sat}_g(K_{1,r+1}; n) = \lfloor \frac{rn-1}{2} \rfloor$  when  $n \leq 8$ . We ask whether this holds for larger  $n$ ; note that  $\text{ex}(K_{1,r+1}; n) = \lfloor \frac{rn}{2} \rfloor$ . Kászonyi and Tuza [18] proved  $\text{sat}(K_{1,r+1}) = \lceil \frac{rn}{2} - \frac{(r+1)^2}{8} \rceil$  for  $n \geq 3r/2$ . Lee and Riet [19] proved  $\text{sat}_g(K_{1,r+1}; n) \geq \frac{rn}{2} - r + 1$ .

For the  $P_4$ -saturation game on  $K_n$ , the value is not asymptotic to the extremal number. We prove that  $\text{sat}_g(P_4; n)$  and  $\text{sat}'_g(P_4; n)$  lie in  $\{\lfloor \frac{4n}{5} \rfloor, \lceil \frac{4n}{5} \rceil\}$  when  $n \geq 5$ , while  $\text{ex}(P_4; n) \in \{n, n - 1\}$ . Lee and Riet [19] proved  $n - 1 \leq \text{sat}_g(P_5; n) \leq n + 2$ .

In Section 3, we study the  $P_4$ -saturation game on  $K_{m,n}$ , for  $m \geq n \geq 2$ . The choice of who starts the game can matter a lot, as do the parities of  $m$  and  $n$ . The value of  $\text{sat}_g(P_4; K_{m,n})$  is  $n$  when  $n$  is even,  $m$  when  $m$  is even and  $n$  is odd, and  $m + \lfloor n/2 \rfloor$  when  $mn$  is odd. The value of  $\text{sat}'_g(P_4; K_{m,n})$  is  $m$  when  $n = 2$  and  $m + \lfloor n/2 \rfloor - \epsilon$  when  $n > 2$ , where  $\epsilon = 0$  when  $mn$  is even and  $\epsilon = 1$  when  $mn$  is odd.

Note that the difference is  $m - 2$  when  $n = 2$ ; for larger  $n$  the difference is  $m - n/2$  when  $n$  is even. In fact, when  $n = o(m)$ , we have  $\text{sat}_g(P_4; K_{m,n}) = o(\text{sat}'_g(P_4; K_{m,n}))$  when  $n$  is even, but  $\text{sat}_g(P_4; K_{m,n}) = (1 - o(1))(\text{sat}'_g(P_4; K_{m,n}))$  when  $n$  is odd and  $m$  is even. Note also that  $\text{sat}(P_4; K_{m,n}) = n$ , so when  $n$  is even we obtain an example where  $\text{sat}_g(\mathcal{F}; H) = \text{sat}(\mathcal{F}; H)$ . We ask whether there are other interesting examples where  $\text{sat}_g(\mathcal{F}; H)$  or  $\text{sat}'_g(\mathcal{F}; H)$  equals  $\text{sat}(\mathcal{F}; H)$ ; [9] provides a survey of saturation numbers as of 2011.

In Section 4, we study the  $C_4$ -saturation game on  $K_{n,n}$ . This game is the natural bipartite analogue of the triangle-saturation game on  $K_n$  studied by Füredi, Reimer, and Seress [14]. A subgraph that is  $C_4$ -saturated relative to  $K_{n,n}$  must be connected, and the spanning tree having adjacent vertices of degree  $n$  is  $C_4$ -saturated, so  $\text{sat}(C_4; K_{n,n}) = 2n - 1$ . On the other

hand, Füredi [13] proved  $\text{ex}(C_4; K_{n,n}) = n^{3/2} + O(n^{4/3})$ , so  $2n - 1 \leq \text{sat}_g(C_4; K_{n,n}) \leq O(n^{3/2})$ . Our main result increases the exponent from the trivial lower bound:  $\text{sat}_g(C_4; K_{n,n}) \geq \frac{1}{21}n^{13/12} - O(n^{35/36})$ .

Our results leave many open questions. The most interesting specific question is the order of growth of  $\text{sat}_g(C_4; K_{n,n})$ . One would also like to understand the conditions under which  $\text{sat}_g(\mathcal{F}; n)$ ,  $\text{sat}_g(\mathcal{F}; K_{m,n})$ , or  $\text{sat}_g(\mathcal{F}; H)$  does not differ much from the value of the corresponding Min-start game.

We mention some general terminology. A graph is *nontrivial* if it has at least one edge. The disjoint union of graphs  $G$  and  $H$  is denoted  $G + H$ , and the graph  $kG$  is the disjoint union of  $k$  copies of  $G$ . When we say that a player *creates*  $F$ , *makes*  $F$ , or *forms*  $F$ , we mean that the player selects an edge whose addition to the graph of selected edges produces a copy of  $F$  (that is, a subgraph isomorphic to  $F$ ).

## 2 Saturation games on complete graphs

We begin with saturation games on the complete graph  $K_n$ .

**Theorem 2.1.**  $\text{sat}_g(\mathcal{O}; n) = \text{sat}'_g(\mathcal{O}; n) = \lfloor n/2 \rfloor \cdot \lceil n/2 \rceil = \text{ex}(\mathcal{O}; n)$ .

*Proof.* A bipartite graph is *balanced* if the parts of the bipartition have equal size; *nearly balanced* if the sizes differ by 1. Every  $\mathcal{O}$ -saturated graph is a complete bipartite graph. The largest such  $n$ -vertex graph is  $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$ . It therefore suffices to give Max a strategy that ensures the following condition after each move by Max:

- (\*) Each nontrivial component of the selected graph is balanced, except when  $n$  is odd and no isolated vertices remain, in which case all components are balanced except for one nearly balanced component.

Condition (\*) ensures that when only one component remains, it is balanced or nearly balanced. When  $n \geq 4$ , Max can ensure (\*) after his first move by choosing an isolated edge, no matter who moves first, and the forced copy of  $P_3$  when  $n = 3$  and Min moves first also satisfies (\*).

If (\*) holds after a move by Max, then Min cannot produce a component that fails to be balanced or nearly balanced, and Min can increase the number of nontrivial nearly balanced components only by absorbing an isolated vertex into a balanced component. If Min does that, then there were still isolated vertices in the graph, all components were balanced before Min's move, and Max can restore (\*) by absorbing another isolated vertex into the nearly balanced component, unless  $n$  is odd and no isolated vertices remain.

In the remaining case, (\*) holds after the move by Min. If Max plays within a component that is not complete bipartite, or connects two nontrivial components, or connects two isolated vertices, then (\*) again holds. If no such move is available, then there is only one

nontrivial component, it is complete bipartite, and there is at most one isolated vertex. If there is an isolated vertex, then the nontrivial component is balanced, and Max connects the isolated vertex to it. Otherwise, the game is over.  $\square$

**Theorem 2.2.** *If  $n \geq 3$ , then  $\text{sat}_g(\mathcal{T}_n; n) = \text{sat}'_g(\mathcal{T}_n; n) = \binom{n-2}{2} + 1$ , except that  $\text{sat}_g(\mathcal{T}_5; 5) = 6 = \text{ex}(\mathcal{T}_5; 5)$  and  $\text{sat}'_g(\mathcal{T}_4; 4) = 3 = \text{ex}(\mathcal{T}_4; 4)$ .*

*Proof.* The value is 1 when  $n = 3$ , so we may assume  $n \geq 4$ . Every  $\mathcal{T}_n$ -saturated subgraph of  $K_n$  is  $K_r + K_{n-r}$  for some  $r$ . The largest is  $K_{n-1} + K_1$ , with  $\binom{n-1}{2}$  edges, and next is  $K_{n-2} + K_2$ , with  $\binom{n-2}{2} + 1$  edges. We show that the final subgraph under optimal play is  $K_{n-2} + K_2$  except when  $n = 5$  and Max starts or when  $n = 4$  and Min starts. Let  $s$  and  $t$  denote the current numbers of nontrivial components and isolated vertices, respectively.

Max can force the smaller component to have at most two vertices by always leaving  $s = 1$  after his move while  $t \geq 1$ . Min can create a second nontrivial component only by joining two isolated vertices, and Max can then absorb it into the first component. Only when Min connects the last two isolated vertices can a second nontrivial component survive.

Max does better in two cases. When  $n = 4$  and Min starts, Max forms  $P_3 + K_1$  on his first move. When  $n = 5$  and Max starts, the second move by Max forms  $P_4 + K_1$ . In both cases, the final graph is  $K_{n-1} + K_1$ , and we may henceforth exclude these cases.

In the remaining cases, it suffices to show that Min can ensure  $s = 2$  at the end. Suppose first that some move by Max leaves  $s \geq 2$ . If  $t \geq 2$ , then Min connects two isolated vertices to obtain  $s \geq 3$ , and the move by Max again leaves  $s \geq 2$ . If  $t = 1$ , then Min plays an edge incident to the isolated vertex, leaving  $s \geq 2$ , and guarantees  $s = 2$  at the end of the game.

Therefore, to have a chance of reaching  $K_{n-1} + K_1$ , Max must always leave  $s = 1$  after his move. If Min increases  $s$ , then Max must connect the new isolated edge to the existing nontrivial component. If at some point  $t$  is even and Min is to move, then Min can enforce  $s = 2$  at the end by repeatedly making isolated edges, which Max must absorb, until Max cannot absorb the last edge. It thus suffices to show that Min can achieve even  $t$  before one of her moves when the first move by Max (with  $s = 1$ ) leaves  $t$  odd.

If Max starts, then having  $t$  odd requires  $n$  odd. Min forms  $P_3$ . Max now must enlarge this component to  $P_4$  or  $K_{1,3}$  to keep  $t$  odd and  $s = 1$ . There remain three unplayed edges among these four vertices, and Min plays next. Eventually, Max will be forced to absorb an isolated vertex or make an isolated edge, reaching  $t$  odd or  $s = 2$ , respectively.

If Min starts, then Max must form  $P_3$  to keep  $s = 1$ , so having  $t$  odd requires  $n$  even. Min next forms  $K_3$ , and again Max must leave  $t$  odd or  $s = 2$  after his next move.  $\square$



**Theorem 2.3.** For  $n \in \mathbb{N}$ ,

$$\begin{aligned} \text{sat}_g(K_{1,3}; n) &= \begin{cases} n & \text{when } n \in \{3, 7\} \cup 2\mathbb{N} - \{2\} \\ n - 1 & \text{otherwise} \end{cases} \\ \text{sat}'_g(K_{1,3}; n) &= \begin{cases} n - 1 & \text{when } n \in \{1\} \cup 2\mathbb{N} - \{4\} \\ n & \text{otherwise} \end{cases} \end{aligned}$$

*Proof.* All  $K_{1,3}$ -saturated graphs are disjoint unions of cycles plus possibly one isolated vertex or one isolated edge (not both). Hence the only possible outcomes are  $n$  (call this *Max wins*) or  $n - 1$  (call this *Min wins*). Let  $X(n)$  and  $Y(n)$  denote the Max-start and Min-start  $K_{1,3}$ -saturation games on  $K_n$ , respectively.

For  $n \geq 5$ , our claim is that the first player wins when  $n$  is even and the second player wins when  $n$  is odd, except that Max wins  $X(7)$ . After giving specific strategies for  $n \leq 8$ , we provide general strategies for  $n \geq 9$  that reduce the problem to the case  $n \in \{5, 6, 8\}$ .

When  $n \leq 3$ , no copy of  $K_{1,3}$  is possible; Min wins when  $n \leq 2$  and Max wins when  $n = 3$ , no matter who starts.

Max wins  $X(4)$  by forming  $P_4$  (second round) and  $Y(4)$  by forming  $2K_2$  (first round).

Min wins  $X(5)$  by forming  $2K_2$  first and then  $C_3$  or  $C_4$ .

Max wins  $Y(5)$  by forming  $2K_2$  first and then  $P_5$ , forcing  $C_5$  as the final graph.

Max wins  $X(6)$ . If Min first makes  $P_3$ , then Max completes the triangle, reducing to  $Y(3)$ , which Max wins. If Min first makes  $2K_2$ , then Max forms  $3K_2$  and next  $P_6$  to win.

Min wins  $Y(6)$  by making  $P_4$  on the second move and then  $C_4$  or  $C_5$ .

Max wins  $X(7)$ . If Min first makes  $P_3$ , then Max completes the triangle and wins  $Y(4)$ . If Min first makes  $2K_2$ , then Max makes  $3K_2$ . After Min's move, Max next completes  $C_3$  or  $C_4$ , and the remainder of the game completes a cycle on the other vertices.

Max wins  $Y(7)$ . Max first makes  $P_3$ . If Min completes the cycle, then Max wins the remaining  $X(4)$ . If Min forms  $P_4$ , then Max completes the cycle and wins  $Y(3)$ . If Min forms  $P_3 + P_2$ , then Max can form  $P_3 + P_3$  and force the game to end with  $C_3 + C_4$  or  $C_7$ .

Max wins  $X(8)$ . Max makes  $P_4$  on his second move. If Min closes the cycle or plays an isolated edge, then Max forms  $C_4 + P_2$ , and the remaining game is equivalent to  $X(4)$ , which Max wins. If Min extends the path to  $P_5$ , then Max closes the cycle and wins  $Y(3)$ .

Min wins  $Y(8)$ . If after four edges the selected graph contains a triangle, then the remaining game is  $X(5)$  in which the first move has been played, which is won by Min. To avoid this, Max must make  $2K_2$ , and after Min makes  $P_3 + P_2$  Max must make  $P_3 + 2P_2$ , extend one of the paths, or join the paths to form  $P_5$ . If  $P_3 + 2P_2$ , then again Min closes the triangle, and the remainder is a copy of  $X(5)$  in which Min has played a winning  $2K_2$ . In the other cases, Min forms  $P_6$ ; the final graph is  $C_6 + P_2$  or  $C_7 + P_1$ , with Min winning.

Now assume  $n \geq 9$ . We denote by Player 1 and Player 2 the player who moves first and the player who moves second, respectively. We begin by showing that Player 1 wins when  $n$

is even. Player 1 always leaves an even number of isolated vertices, some number of cycles, and one nontrivial path, until the number of isolated vertices is 6. This is true after the initial move. If Player 2 extends the path or makes an isolated edge, then Player 1 further extends the path or combines it with the new isolated edge to restore the condition. If Player 2 completes the path to a cycle, then Player 1 starts a new path. In each case, there is still one nontrivial path component, and the number of isolated vertices has decreased by 2.

When six isolated vertices remain, Player 2 is ready to move. If the nontrivial path is a single edge, then the remaining game is a copy of the game on eight vertices in which Player 1 has made the arbitrary first move, and we have proved that Player 1 wins this game. Hence we may assume that the nontrivial path is longer. If Player 2 completes it to a cycle (and Player 1 chooses an isolated edge) or Player 2 chooses an isolated edge and lets Player 1 close the cycle, then the remaining game is equivalent to the game with  $n = 6$  in which Player 1 has made the first move; we have shown that Player 1 wins this game. If Player 2 extends the path, then Player 1 closes the cycle to leave a game with  $n = 5$  started by Player 2. Since 5-vertex games are won by the player who makes the second move, again Player 1 wins.

Suppose now that  $n$  is odd; we claim that Player 2 wins. On his first move, Player 2 makes  $P_3$ . The graph now consists of one nontrivial path and some isolated vertices. Moreover, the number of isolated vertices is even and at least 6. Player 2 now wins by following the strategy outlined above for Player 1.  $\square$

In the  $K_{1,3}$ -saturation game on  $K_n$ , each player prefers to move first when  $n$  is even and second when  $n$  is odd. Hefetz, Krivelevich, Naor, and Stojaković [15] showed that the  $kP_2$ -saturation game on  $K_n$  behaves similarly (where  $k \leq n/2$ ). In this game, each player prefers to move first if and only if  $k$  is even. For the  $kP_2$ -saturation game, the identity of the first player makes a dramatic difference: Hefetz et al. showed that when  $k$  is even,  $\text{sat}_g(kP_2; n) \geq n-1$  and  $\text{sat}'_g(kP_2; n) \leq \binom{2k-1}{2}$ . Likewise, when  $k$  is odd,  $\text{sat}_g(kP_2; n) \leq \binom{2k-1}{2}$  and  $\text{sat}'_g(kP_2; n) \geq n-1$ .

Because there are only two possible (consecutive) lengths of the  $K_{1,3}$ -saturation game on  $K_n$ , given the identity of the first player, the outcome is determined by who plays last. Ferrara, Jacobson, and Harris [11] studied that question explicitly; in their game, the player who moves last wins. Although their analysis is similar to ours due to the structure of  $K_{1,3}$ -saturated graphs, their result is different: in their game, for  $n \geq 5$ , the first player wins if and only if  $n$  is even, except  $n = 7$ . In particular, under their criterion for winning, the number of moves played will always be  $n - 1$  (except  $n = 7$ ).

Our final game on  $K_n$  is the  $P_4$ -saturation game. In a graph not containing  $P_4$ , all components must be stars or triangles. Since Max seeks a large ratio of number of edges to number of vertices, triangles and large stars are beneficial to Max, while small stars are beneficial to Min. However, stars with two edges are dangerous for Min, since they may

become triangles. This intuition motivates the strategies for the players. It turns out that optimal play by both players produces components isomorphic to  $K_{1,4}$ , which explains the form of the answer.

**Theorem 2.4.** *For  $n \geq 5$ , both  $\text{sat}_g(P_4; n)$  and  $\text{sat}'_g(P_4; n)$  lie in  $\{\lfloor 4n/5 \rfloor, \lceil 4n/5 \rceil\}$ .*

*Proof.* Every  $P_4$ -saturated subgraph of  $K_5$  contains exactly four edges, so we may suppose  $n \geq 6$ . As noted above, all components of the graph being constructed must be stars or triangles. Let the *value*  $\rho$  of the current position count a contribution for each component: 0 for an isolated vertex or triangle,  $\frac{1}{2}$  for  $P_2$  or  $P_3$ , and 1 for a larger star. The value can decrease only by a move that turns a copy of  $P_3$  into a triangle. In this proof every mention of a subgraph refers to a component of the current selected graph. As in the proof of Theorem 2.2, let  $t$  denote the current number of isolated vertices during the game. Let  $m$  be the final number of edges.

*Upper bound: Min strategy.* While  $t \geq 1$ , if a copy of  $P_3$  exists, then Min converts it to  $K_{1,3}$ . If there is no  $P_3$  and  $t \geq 2$ , then Min makes  $P_2$ . Otherwise,  $t = 1$  and there is no copy of  $P_3$ , in which case Min adds the isolated vertex to a largest star, or  $t = 0$ , in which case the only possible move completes a copy of  $P_3$  to a triangle.

With this strategy, each move by Min increases  $\rho$  by  $\frac{1}{2}$ , except when  $t \leq 1$ . Min never forms a 3-vertex component, except when  $t = 1$  and all other components are isolated edges, yielding  $K_3 + \frac{n-3}{2}P_2$  with  $\frac{n+3}{2}$  edges. Any earlier copy of  $P_3$  formed by Max is converted immediately to  $K_{1,3}$ . If Max makes  $P_3$  with the last isolated vertex, then Min must complete the triangle. Hence except for one possible triangle, all components are stars. Regardless of whether a triangle appears at the end, the number of stars is  $n - m$ .

Since the strategy also prevents Max from decreasing  $\rho$  (with the exception of the final triangle when Max stupidly makes isolated edges as mentioned above), the final value of  $\rho$  is at least  $\frac{1}{2} \frac{m-2}{2} - \frac{1}{2}$ , which equals  $\frac{m-4}{4}$ . (When Min absorbs the last isolated vertex,  $\rho$  reaches at least  $\frac{m}{4}$ .) Also, at the end  $\rho$  is at most the number of components that are stars. We obtain  $\frac{m-4}{4} \leq n - m$ , which simplifies to  $m \leq \frac{4n+4}{5}$ . The same computation yields  $m \leq \frac{4n+3}{5}$  in the Min-start game, so in both cases  $m \leq \lceil 4n/5 \rceil$ .

*Lower bound: Max strategy.* Max never chooses an isolated edge, except on the first turn of the Max-start game or when all nontrivial components are triangles. If there is an isolated edge and  $t \geq 1$ , then Max turns it into  $P_3$ . Otherwise, if some component is  $P_3$ , then Max completes the triangle. If none of these moves is available (and the game is not over), then Max enlarges a star with at least three edges.

Note that Max increases  $\rho$  only when all nontrivial components are triangles. Except for the first move in the Max-start game, each move by Max that increases  $\rho$  is thus preceded by a move by Min that decreases  $\rho$  by completing a triangle. With each Min move increasing  $\rho$  by at most  $\frac{1}{2}$ , the upper bound on  $\rho$  is  $\frac{m+2}{4}$  (or  $\frac{m+1}{4}$  in the Min-start game). Moreover, if

Min ever completes a triangle, then  $\rho$  is at most  $\frac{m-2}{4}$  in the Max-start game or  $\frac{m-3}{4}$  in the Max-start game.

Any isolated edge formed by Min is immediately converted to  $P_3$  by Max (provided  $t \geq 1$ ). Max himself forms an isolated edge only when none are already present and all nontrivial components are triangles (including on the first turn of the game). When  $t = 2$ , Min may create a second copy of  $P_2$  that Max cannot convert to  $P_3$ . Thus, the final graph contains at most two copies of  $P_2$ . If Min always makes  $P_2$ , then Max turns these components into  $P_3$  and they become triangles at the end, yielding at least  $n - 2$  edges, which is good for Max since  $n - 2 \geq \lfloor \frac{4n}{5} \rfloor$  when  $n \geq 6$ . If at some point Min does not make  $P_2$ , then Max turns any remaining isolated edge into  $P_3$ . Max later creates another isolated edge only if every nontrivial component of the graph is a triangle; in this case, Min's preceding move must have completed a triangle. Hence we may assume that either the final graph has at most one isolated edge, or it has two but at some point Min completed a triangle.

Components that are triangles have no net effect on  $\rho$  or on  $n - m$ . Also,  $n - m$  is the number of stars. When the game ends with at most one isolated edge, we thus have  $\rho \geq n - m - \frac{1}{2}$  at the end, no matter who starts. Using the weaker upper bound on  $\rho$  from the Max-start game, we obtain  $\frac{m+2}{4} \geq n - m - \frac{1}{2}$ , which simplifies to  $m \geq \frac{4n-4}{5}$ . When the game ends with two isolated edges and Min having completed a triangle, we have the weaker lower bound  $\rho \geq n - m - 1$  but a stronger upper bound on  $\rho$ . This leads to  $\frac{m-2}{4} \geq n - m - 1$ , hence  $m \geq \frac{4n-2}{5}$ . In either case,  $m \geq \lfloor 4n/5 \rfloor$ .  $\square$

A referee pointed out that a more delicate analysis shows that for  $n \geq 8$  the answer is always  $\lfloor 4n/5 \rfloor$ , except for the Min-start game when  $n$  is congruent to 1 or 3 modulo 5, where the game lasts one additional move.

### 3 The $P_4$ -saturation game on $K_{m,n}$

Now we study the  $P_4$ -saturation game on the complete bipartite graph  $K_{m,n}$ . Since  $K_{m,n}$  contains no triangles, during the game all components are stars. Throughout this section,  $X$  and  $Y$  are the partite sets of  $K_{m,n}$ , with  $|X| = m \geq n = |Y|$ . Let an  $X$ -star or a  $Y$ -star be a star having at least two leaves in  $X$  or in  $Y$ , respectively. We use  $\alpha'(G)$  for the maximum size of a matching in  $G$ .

**Lemma 3.1.** *A graph  $G$  that is  $P_4$ -saturated relative to  $K_{m,n}$  has at most  $m + n - \alpha'(G)$  edges. If it contains both an  $X$ -star and a  $Y$ -star (or an isolated edge), then equality holds.*

*Proof.* Any even cycle contains  $P_4$ , so  $G$  is a forest. To avoid  $P_4$ , edges of a matching must lie in distinct components, so  $\alpha'(G)$  is the number of nontrivial components. Since  $G$  is a forest,  $|E(G)|$  is the number of vertices minus the number of components, so  $|E(G)| \leq m + n - \alpha'(G)$ .

A saturated subgraph containing both an  $X$ -star and a  $Y$ -star (or an isolated edge) cannot have isolated vertices. All components are then nontrivial stars, so equality holds in the computation above.  $\square$

Call a  $P_4$ -saturated subgraph that contains both an  $X$ -star and a  $Y$ -star a *full subgraph*. A  $P_4$ -saturated subgraph that is not full has stars of only one of these types (plus isolated edges, possibly) and thus has only  $m$  or  $n$  edges. Hence Max wants to make a full subgraph. When  $m$  or  $n$  is even, Min can prevent this in the Max-start game; in particular, when  $n$  is even, we obtain  $\text{sat}_g(P_4; K_{m,n}) = n = \text{sat}(P_4; K_{m,n})$ . When Max can make a full subgraph, Lemma 3.1 encourages Min to create a large matching.

**Theorem 3.2.** *For  $m \geq n \geq 2$ , the  $P_4$ -saturation numbers of  $K_{m,n}$  are given by*

$$\text{sat}_g(P_4; K_{m,n}) = \begin{cases} n & \text{when } n \text{ is even,} \\ m & \text{when } n \text{ is odd and } m \text{ is even,} \\ m + \lfloor \frac{n}{2} \rfloor & \text{when } mn \text{ is odd.} \end{cases}$$

and

$$\text{sat}'_g(P_4; K_{m,n}) = \begin{cases} m & \text{when } n \leq 2, \\ m + \lfloor \frac{n}{2} \rfloor & \text{when } n > 2 \text{ and } mn \text{ is even,} \\ m + \lfloor \frac{n}{2} \rfloor - 1 & \text{when } n > 2 \text{ and } mn \text{ is odd.} \end{cases}$$

*Proof.* We will consider cases based on who moves first and the parity of  $m$  and  $n$ . Let  $G$  denote the  $P_4$ -saturated subgraph built during the game. Again “making” a subgraph means producing it as a component of the current graph.

*Upper bounds.* We give strategies for Min. If Max moves first and  $n$  is even, then Min ensures that only  $Y$ -stars are created, by immediately extending isolated edges made by Max to such stars and otherwise enlarging such stars. If  $m$  is even and  $n$  is odd, then Min similarly creates  $X$ -stars. The final number of edges is then  $|Y|$  or  $|X|$  in these two cases.

In the other cases, Min just ensures a large matching. If Max moves first and  $mn$  is odd, or Min moves first and  $mn$  is even, then Min makes isolated edges until a matching of size  $\lfloor \frac{n}{2} \rfloor$  is built, later playing any legal move. By Lemma 3.1, at most  $m + \lfloor \frac{n}{2} \rfloor$  moves are played.

If Min moves first and  $mn$  is odd, then Min can do slightly better. If Max responds to the first move by making an  $X$ -star or a  $Y$ -star, then the parity allows Min to ensure that only  $X$ -stars or  $Y$ -stars, respectively, will be played, yielding an outcome of  $|X|$  or  $|Y|$ . Hence Max must immediately make another isolated edge. The moves by Min still yield a matching of size  $\lfloor \frac{n}{2} \rfloor$ , and with the extra edge made by Max the bound improves by 1.

*Lower bounds.* We give strategies for Max. Since the game cannot end with an isolated vertex in each part, at least  $n$  moves are played. If an  $X$ -star is made, then no isolated vertex can be left in  $X$ , and at least  $m$  moves are made. In the Max-start game with  $n$  odd and  $m$  even, Min can prevent an  $X$ -star only by ensuring that, after each of her moves, every

nontrivial component is a  $Y$ -star. After  $n - 1$  moves, Max makes an isolated edge using the last isolated vertex of  $Y$ , and then Min is forced to make an  $X$ -star. In the Min-start game with  $n \leq 2$ , Max makes an  $X$ -star immediately.

In the other cases, we may assume  $n \geq 3$ . Max wants to force a full subgraph and keep  $\alpha'(G)$  small. Since  $\alpha'(G)$  equals the number of components in  $G$ , Max avoids making many isolated edges.

First consider the Min-start game with  $n$  even. Max responds to the first move by making a  $Y$ -star. While at least two isolated vertices remain in  $Y$ , Max enlarges the  $Y$ -star. If no isolated vertices remain in  $Y$ , then Max plays any legal move. Suppose that exactly one isolated vertex remains in  $Y$ . If the graph does not contain an  $X$ -star or an isolated edge, then Max plays an isolated edge; otherwise, Max again adds to the  $Y$ -star. After Max's move, either the graph is already full or it contains an isolated edge. In the latter case, note that every move in the game has reduced the number of isolated vertices in  $Y$ , but not all moves have reduced the number of isolated vertices in  $X$ . Consequently, although  $Y$  no longer contains isolated vertices,  $X$  does. Thus, Min is forced to transform an isolated edge into an  $X$ -star on her next move, making the graph full. Finally, note that if it is Min's turn when exactly one isolated vertex remains in  $Y$ , then by parity of  $Y$ , at least one of Min's moves did not reduce the number of isolated vertices in  $Y$ ; that is, Min created an  $X$ -star. In any case, the graph becomes full.

Throughout the game, Max makes at most one isolated edge. If in fact Max makes no isolated edges, then Min can make at most  $n/2$ , since on each turn the number of isolated vertices in  $Y$  decreases. Moreover, if Min makes an isolated edge on each of her turns, then Max is not forced to make one himself. Consequently, if Max does make an isolated edge, then Min makes fewer than  $n/2$ . In either case, the game ends with at most  $n/2$  components. Thus  $\alpha'(G) \leq n/2$  so, by Lemma 3.1, the number of edges in  $G$  is  $m + n - n/2$ , which equals  $m + \lfloor n/2 \rfloor$ .

Now consider the Min-start game with  $n$  odd and  $m$  even. Max responds to Min's initial move by creating an  $X$ -star. After this initial move, Min may either enlarge the  $X$ -star or play an isolated edge. If Min always enlarges the  $X$ -star, then Max does the same, until exactly one isolated vertex remains in  $X$ . Since  $m$  is even, it is now Max's turn, and he plays an isolated edge. Since  $n \geq 3$ , at least one isolated vertex remains in  $Y$ , while no isolated vertices remain in  $X$ ; consequently, Min must transform the isolated edge into a  $Y$ -star. The graph is full, there are only two components, and no new components can be added. Hence  $\alpha'(G) = 2$ , so the number of edges in  $G$  is  $m + n - 2$ , which is at least  $m + \lfloor n/2 \rfloor$ .

Suppose instead that Min does eventually play an isolated edge. Since  $n \geq 3$ , at least one isolated vertex remains in  $Y$ . Max transforms the isolated edge into a  $Y$ -star, making the graph full. Henceforth, as long as it remains possible, Max always enlarges this  $Y$ -star. When the  $Y$ -star is first created, the graph has two components,  $n - 3$  isolated vertices remain in  $Y$ , and it is Min's turn. While  $Y$  still has isolated vertices, each of Max's moves

decreases the number by one, as does each move by Min that creates a new component. Thus at most  $(n-3)/2$  more components are created throughout the game, so  $\alpha'(G) \leq 2 + (n-3)/2 = (n+1)/2$ . Now by Lemma 3.1, the number of edges in  $G$  is  $m+n - (n+1)/2$ , which equals  $m + \lfloor n/2 \rfloor$ .

For the Min-start game with  $mn$  odd, Max cannot do quite as well. As noted when discussing upper bounds, if Max makes an  $X$ -star or  $Y$ -star on move 2, then Min can limit the final number of edges to  $m$  or  $n$ , respectively. If  $n=3$ , then the claimed upper bound reduces to  $m$ , so Max makes an  $X$ -star. Otherwise, Max makes an isolated edge on move 2. If Min makes  $K_{1,2}$ , then Max makes the other type of star. If Min makes an isolated edge, then Max makes an  $X$ -star and can make a  $Y$ -star on the next round.

Hence the graph becomes full. Max subsequently enlarges  $Y$ -stars until  $Y$  has no more isolated vertices. After Max's second turn, the graph has at most three components and  $Y$  has exactly  $n-3$  isolated vertices. As before, at most  $(n-3)/2$  more components are created throughout the game. Now  $\alpha'(G) \leq 3 + (n-3)/2 = (n+3)/2$ , so  $|E(G)| \geq m+n - (n+3)/2 = m + \lfloor n/2 \rfloor - 1$ .  $\square$

## 4 The $C_4$ -saturation game on $K_{n,n}$

In this section, we study the  $C_4$ -saturation game on  $K_{n,n}$ , the natural bipartite analogue of the Füredi-Reimer-Seress problem. As we have noted, the trivial lower bound and the result of [13] yield  $2n-1 \leq \text{sat}_g(C_4, K_{n,n}) \leq O(n^{3/2})$ .

Our main result is a polynomial improvement of the lower bound:  $\text{sat}_g(C_4, K_{n,n}) = \Omega(n^{13/12})$ . We first prove a technical lemma giving a lower bound on the size of a restricted type of graph that is also  $C_4$ -saturated graph relative to  $K_{n,n}$ . Here our interest is the exponent on  $n$ ; we make no attempt to optimize lower-order terms or the leading coefficient.

**Lemma 4.1.** *Let  $G$  be a graph that is  $C_4$ -saturated relative to  $K_{n,n}$ , and let  $c$  and  $d$  be positive constants. If there exists  $S \subseteq V(G)$  with at least  $cn$  vertices in each partite set such that  $|N(v) \cap S| \leq d\sqrt{n}$  for all  $v \in V(G)$ , then  $|E(G)| \geq an^{13/12} - O(n^{35/36})$ , where  $a = \min\{\frac{1}{2}(\frac{c^2}{2d^2})^{2/3}, \frac{c^2}{2d}\}$ .*

*Proof.* Let  $S_X$  and  $S_Y$  be the subsets of  $S$  in the two partite sets. Consider  $x \in S_X$  and  $y \in S_Y$  such that  $xy \notin E(G)$ . Since  $G$  is  $C_4$ -saturated relative to  $K_{n,n}$ , it contains a copy of  $P_4$  with endpoints  $x$  and  $y$ . Each vertex in  $S_X$  has at most  $d\sqrt{n}$  neighbors in  $S_Y$  and hence at least  $cn - d\sqrt{n}$  nonneighbors in  $S_Y$ . Thus  $G$  contains at least  $c^2n^2 - cdn^{3/2}$  copies of  $P_4$  with endpoints in  $S_X$  and  $S_Y$ ; call such paths *essential paths*. Since each essential path has endpoints in  $S_X$  and  $S_Y$ , and since no vertex has more than  $d\sqrt{n}$  neighbors in  $S$ , no edge is the central edge of more than  $d^2n$  essential paths.

Let  $T$  be the set of vertices of  $G$  with degree at least  $n^{5/12}$ , and let  $b = (\frac{c^2}{2d^2})^{2/3}$ . If  $|T| \geq bn^{2/3}$ , then  $\sum_{v \in T} d(v) \geq bn^{13/12}$ , which yields  $|E(G)| \geq \frac{b}{2}n^{13/12}$ . Otherwise, let

$H$  be the subgraph of  $G$  induced by  $T$ . Since  $C_4 \not\subseteq H$ , the result of Füredi [13] yields  $|E(H)| \leq (bn^{2/3})^{3/2} + O((bn^{2/3})^{4/3})$ , which simplifies to  $|E(H)| \leq \frac{c^2}{2d^2}n + O(n^{8/9})$ . Multiplying by  $d^2n$ , we conclude that at most  $\frac{c^2}{2}n^2 + O(n^{17/9})$  essential paths have central edges in  $H$ .

Thus at least  $\frac{c^2}{2}n^2 - O(n^{17/9})$  essential paths have central edges incident to a vertex with degree less than  $n^{5/12}$ . Each such edge is the central edge of at most  $dn^{11/12}$  essential paths; hence  $G$  has at least  $\frac{c^2}{2d}n^{13/12} - O(n^{35/36})$  such edges.  $\square$

Though the hypotheses of Lemma 4.1 seem technical, they apply whenever  $\Delta(G) \leq d\sqrt{n}$ . Hence we obtain a corollary for ordinary saturation (using  $c = 1$ ).

**Corollary 4.2.** *If  $G$  is  $C_4$ -saturated relative to  $K_{n,n}$  and  $\Delta(G) \leq d\sqrt{n}$ , then  $|E(G)| \geq an^{13/12} - O(n^{35/36})$ , where  $a = \min\{\frac{1}{2}(\frac{1}{2d^2})^{2/3}, \frac{1}{2d}\}$ . (If  $d \leq \frac{1}{4}$ , then  $a = \frac{1}{2d}$ ).*

Our main result for the  $C_4$ -saturation game on  $K_{n,n}$  follows easily from Lemma 4.1.

**Theorem 4.3.**  $\text{sat}_g(C_4, K_{n,n}) \geq \frac{1}{21}n^{13/12} - O(n^{35/36})$ , and similarly for  $\text{sat}'_g(C_4; K_{n,n})$ .

*Proof.* We provide a strategy for Max that forces the final subgraph of  $K_{n,n}$  to satisfy the hypotheses of Lemma 4.1. This strategy governs almost the first  $2n/3$  moves for Max, after which Max plays arbitrarily.

Let  $k = \lfloor \sqrt{n/3} \rfloor - 1$ . Max arranges to give degree  $k$  to  $k$  specified vertices in each partite set. Each move by Max makes an isolated vertex adjacent to one of the specified vertices; hence it cannot complete a 4-cycle. Fewer than  $n/3$  vertices are needed by Max in each part, so Min cannot exhaust the isolated vertices in either part with fewer than  $2n/3$  moves. After this phase, Max may play any legal move.

At the end of the first phase, let  $S$  be the set of leaves attached to the  $2k$  specified stars. By construction, these stars are disjoint, so  $S$  has about  $n/3 - 2\sqrt{n/3}$  vertices in each part.

In the final subgraph  $G$ , no vertex has more than  $2\sqrt{n/3}$  neighbors in  $S$ , since each vertex other than the center of a star is adjacent to at most one leaf of the star. (However, in addition to the leaves of its own star, the center of a star may be adjacent to one leaf in each of the other stars whose center is in the same partite set.) Thus  $G$  satisfies the hypotheses of Lemma 4.1 with  $c$  being any constant less than  $1/3$  and  $d = 2\sqrt{1/3}$ , from which the claim follows.  $\square$

While Theorem 4.3 does establish a nontrivial asymptotic lower bound for  $\text{sat}_g(C_4; K_{n,n})$ , the correct order of growth remains undetermined. Theorem 4.3 can perhaps be strengthened by improving the bounds in Lemma 4.1. This suggests the following question: What is the minimum number of edges in a graph with maximum degree  $D$  that is  $C_4$ -saturated relative to  $K_{n,n}$ ?

### Acknowledgment

The authors thank the referees for extremely careful reading that led to improvements of several proofs and results.



## References

- [1] P. Bennett and T. Bohman, *A note on the random greedy independent set algorithm*, arXiv:1308.3732.
- [2] C. Biró, P. Horn, and J. Wildstrom, *An upper bound on the extremal version of Hajnal's triangle-free game*, Discrete Applied Math. (2015), <http://dx.doi.org/10.1016/j.dam.2015.06.031>.
- [3] T. Bohman and P. Keevash, *The early evolution of the  $H$ -free process*, Invent. Math. **181** (2010), no. 2, 291–336.
- [4] B. Bollobás and O. Riordan, *Constrained graph processes*, Electron. J. Comb. **7**, R18 (2000).
- [5] D. W. Cranston, W. B. Kinnersley, S. O, and D. B. West, *Game matching number of graphs*, Discrete Applied Math. **161** (2013), 1828–1836.
- [6] P. Erdős, A. Hajnal, and J. W. Moon, *A problem in graph theory*, Amer. Math. Monthly **71** (1964), 1107–1110.
- [7] P. Erdős, C. Ko, and R. Rado, *Intersection theorems for systems of finite sets*, Quart. J. Math. Oxford Ser. (2) **12** (1961), 313–320.
- [8] P. Erdős, S. Suen, and P. Winkler, *On the size of a random maximal graph*, Random Struct. Algorithms **6**, (1995), 309–318.
- [9] J.R. Faudree, R.J. Faudree, and J.R. Schmitt, *A survey of minimum saturated graphs*, Elec. J. Combin. **18** (2011), Dynamic Survey 19, 36 pages.
- [10] O. N. Feldheim and M. Krivelevich, *Winning fast in sparse graph construction games*, Combin. Probab. Comput. **17** (2008), no. 6, 781–791.
- [11] M. Ferrara, M. S. Jacobson, and A. Harris, *The game of  $\mathcal{F}$ -saturation*, Discrete Appl. Math. **158** (2010), no. 3, 189–197.
- [12] Z. Füredi, *On maximal intersecting families of finite sets*, J. Combin. Theory Ser. A **28** (1980), no. 3, 282–289.
- [13] Z. Füredi, *New asymptotics for bipartite Turán numbers*, J. Combin. Theory Ser. A **75** (1996), no. 1, 141–144.
- [14] Z. Füredi, D. Reimer, and Á. Seress, *Hajnal's triangle-free game and extremal graph problems*, Proceedings of the Twenty-second Southeastern Conference on Combinatorics, Graph Theory, and Computing (Baton Rouge, LA, 1991), vol. 82, 1991, pp. 123–128.
- [15] D. Hefetz, M. Krivelevich, A. Naor, and M. Stojaković, *On saturation games*, arXiv:1406.2111.
- [16] D. Hefetz, M. Krivelevich, M. Stojaković, and T. Szabó, *Planarity, colorability, and minor games*, SIAM J. Discrete Math. **22** (2008), no. 1, 194–212.
- [17] D. Hefetz, M. Krivelevich, M. Stojaković, and T. Szabó, *Fast winning strategies in Maker-Breaker games*, J. Combin. Theory Ser. B **99** (2009), no. 1, 39–47.

- [18] L. Kászonyi and Zs. Tuza, *Saturated graphs with minimal number of edges*, J. Graph Theory **10** (1986), no. 2, 203–210.
- [19] J. D. Lee and A.-E. Riet,  *$\mathcal{F}$ -Saturation Games*, <http://arxiv.org/abs/1406.1500>
- [20] D. Osthus and A. Taraz, *Maximal  $H$ -free graphs*, Random Struct. Algorithms **18** (2001), 61–82.
- [21] B. Patkós and M. Vizer, *Game saturation of intersecting families*, Cent. Eur. J. Math. **12** (2014), no. 9, 1382–1389.
- [22] P. Turán, *Eine Extremalaufgabe aus der Graphentheorie*, Mat. Fiz. Lapok **48** (1941), 436–452.