10-2013

Feasibility of using low density polyethylene sheets to detect atmospheric organochlorine pesticides in Alexandria, Egypt

Mohammed A. Khairy

Rainer Lohmann

University of Rhode Island, rlohmann@uri.edu

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

The University of Rhode Island Faculty have made this article openly available.
[Please let us know](#) how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Mohammed A. Khairy, Rainer Lohmann, Feasibility of using low density polyethylene sheets to detect atmospheric organochlorine pesticides in Alexandria, Egypt, *Environmental Pollution*, Volume 181, October 2013, Pages 151-158.

Available at: <http://dx.doi.org/10.1016/j.envpol.2013.06.031>.

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30

4 Mohammed A. Khairy^{1,2*}, Rainer Lohmann¹

6 ¹Graduate School of Oceanography, University of Rhode Island, 215 South Ferry Road, Narragansett, Rhode Island
7 02882, United States

8 ²Department of Environmental Sciences, Faculty of Science, Alexandria University, 21511 Moharam Bek,
9 Alexandria, Egypt

*Corresponding author. E-mail address: mohabdel@mail.uri.edu

Phone: 401-874-6765; Fax 401-874-6811

14 Abstract

Egypt is a major agricultural country in Africa with a known past of organochlorine pesticides (OCPs) application, yet data on atmospheric levels of OCPs in Egypt is sparse. Low density polyethylene (LDPE) passive samplers were therefore deployed for 3 weeks each at 11 locations in July, 2010 and January, 2011 in Alexandria to screen for gas-phase OCPs. Performance reference compounds were used to investigate the uptake kinetics. Field-derived sampler-air partitioning coefficients (K_{PE-A}) for OCPs were significantly correlated against the compounds' subcooled liquid vapour pressure ($\log P_L$): [$\log K_{PE-A} = -0.77 \pm 0.07 * \log P_L + 6.35 \pm 0.13$ ($R^2 = 0.90$; $n = 17$; $SE = 0.19$; $p < 0.001$)]. Estimated and measured OCP concentrations in Alexandria agreed well (factor difference ≤ 2) indicating the feasibility of monitoring OCPs using LDPEs. OCP concentrations ranged from $<LOD$ to 168 pg/m^3 . Calculated isomeric ratios indicated recent usage of chlordanes and endosulfans.

28 Keywords

29 Passive sampling; partitioning constants; Egypt; organochlorine pesticides; Sources

31 **Capsule Abstract**

32 LDPE passive air samplers spiked with PRCs can be used as a practical low cost technique for
33 monitoring gas phase OCPs in Egypt.

35 **1. Introduction**

37 Organochlorine pesticides (OCPs) are a group of organic pollutants with high
38 bioaccumulation (Nakata et al., 2002), toxicity (Lemaire et al., 2004; Souza et al., 2004; Tiido et
39 al., 2005; Murvoll et al., 2006), persistence and long range transportation in the environment
40 (Bailey et al. 2000; Li et al. 2007) causing them to be readily detected even in remote regions
41 and pristine areas such as the Arctic (Hung et al., 2010). Despite being banned for decades in
42 Canada, U.S. and European countries, DDTs, hexachlorocyclohexanes (HCHs), chlordanes and
43 dieldrin continue being detected in the ambient air on regional and continental scales (Van
44 Drooge et al., 2002; Jaward et al., 2004; Shen et al., 2004, 2005; Messing et al., 2013).

45 In some developing countries, however, several OCPs are still in use (Baek et al., 2013).
46 Pesticides were introduced in Egypt in 1952 and about one million metric tons of commercial
47 pesticides were used until 2003 (Mansour, 2004). In addition, illegal pesticides application
48 cannot be ignored due to the poor enforcement of environmental laws. In Egypt, 70 % of the
49 pesticides are applied on cotton crops, while the rest **are** applied on corn, rice, sugarcane
50 plantations, vegetables and fruits (Mansour, 2004). DDT, lindane and endrin were among the
51 pesticides used to control cotton pests. Three decades ago, the use of DDT in agriculture was
52 officially banned. Following the restriction on DDT, the use of other organochlorine pesticides
53 (e.g., aldrin, dieldrin, chlordane, heptachlor, lindane) was gradually restricted in Egypt (Abou-
54 Arab et al., 1995; Mansour, 2004).

55 Despite the fact that Egypt was a signatory of Stockholm Convention held on May, 2002,
56 and the use of OCPs **was** banned in Egypt, these toxic compounds are still detected in various

57 physical and biological environmental compartments in the country (EL Nemr and Abd Allah,
58 2004; Sallam and Morshedy, 2008; Barakat et al. 2012 a,b; Khairy et al., 2012; Barakat et al.,
59 2013). To our knowledge, limited studies have been carried out to assess the atmospheric levels
60 of OCPs in Egypt, possibly due to the difficulties associated with the conventional active
61 sampling techniques. According to the Stockholm Convention, parties to the Convention should
62 develop a national inventory of POPs. No information exists about the levels, sources and health
63 effects associated with the exposure to atmospheric OCPs in Egypt. Yet frequent measurements
64 of air concentrations in different locations as well as monitoring studies on various levels from
65 local point sources up to the continental scale are a matter of a great importance. Accordingly,
66 cheap techniques that accurately measure atmospheric concentrations of POPs are highly
67 desirable.

68 Passive air samplers have been proven to be a powerful monitoring technique for POPs,
69 as they are cheaper and require less labour compared to active samplers (Lohmann et al., 2001).
70 Polyurethane foam (PUF) disks (Jaward et al., 2004, 2005; Zhang et al., 2008; Devi et al., 2011)
71 and XAD-2 resin (Choi et al., 2008; Wang et al., 2010; Baek et al., 2013) are the most widely
72 used passive air samplers for monitoring OCPs worldwide. Low density polyethylene (LDPE)
73 has been successfully used as passive air samplers for PAHs (Bartkow et al., 2004; Kennedy et
74 al., 2007; Khairy and Lohmann, 2012).

75 The purpose of the current study was to investigate the possibility of using LDPE as
76 passive air samplers for OCPs in the atmospheric environment of Alexandria, Egypt.
77 Accordingly, we undertook two major 21-days sampling campaigns of 15 deployments during
78 summer (July, 2010) and winter (January, 2011) across 11 different sites. By deploying samplers

in different locations and in different seasons, we were able to assess the pollution levels, temporal variations and the possible sources of OCPs in Alexandria using isomeric ratios.

2. Materials and methods

2.1. Site Description

Alexandria is the second-largest city of Egypt, with a total surface area of 2300 km² and a population of 4.1 million, extending about 90 km along the coast of the Mediterranean Sea on the northwestern side of the Nile Delta (Figure SI 1). The city's climate shows Mediterranean characteristics, namely mild, variably rainy winters and hot summers that, at times, can be very humid. More details about the study area can be found in Khairy and Lohmann (2012). The land use map of Alexandria (Figure SI 1) indicated the existence of different land uses including the agriculture, which represented ~ 27% of the governorate area and occurred mainly in the south and southeastern parts (Switch, 2011).

2.2. Preparation and Deployment of LDPEs

LDPE sheets were cut from commercial sheeting (Carlisle Plastics, Inc., Minneapolis, MN) with a thickness of 51 µm, yielding a 10 x 30 cm strip of ~1-2 g each. Samplers were cleaned with DCM and n-hexane and spiked with four performance reference compounds (PRCs) according to the method developed by Booij et al. (2002), namely 2,5-dibromobiphenyl (PBB 9), 2,2',5,5'-tetrabromobiphenyl (PBB 52), 2,2',4,5',6-pentabromobiphenyl (PBB 103) and octachloronaphthalene (see supplementary data, text SI 1 for more details). Once spiked, PEs were strung on stainless steel wires, placed in precleaned aluminum foil packets, numbered, and frozen in plastic bags until the time of deployment.

LDPE sheets were deployed ~ 2 meters above the ground surface in 11 different locations across Alexandria city (Figure SI 1) each inside two inverted bowls providing protection from direct radiation and rainfall. At each location, LDPEs were deployed for 21 days in July, 2010 (summer season) and January 2011 (winter season) (site and deployment details are given in Table SI 1).

2.3. Active Air Sampling

Active sampling was performed to validate LDPE as passive air samplers for OCPs (see text SI 1 for more details about the preparation of the active sampling media). Samples were collected using a high volume sampler at the 11 sites in July, 2010 and January 2011 (for meteorological details and sampling volumes, see text SI 2 and Table SI 2). Air was first drawn through a glass fiber filter (GFF) to collect the particulate bound compounds followed by a PUF to retain the vapor phase compounds. Particulate bound data are not presented in the current study.

2.4. Extraction of LDPEs and PUFs

LDPEs were cold extracted twice in DCM for 24 hours after spiking with 10 µL of a surrogate standard mixture composed of labeled OCPs (¹³C₆-hexachlorobenzene and ¹³C₁₂-p,p'-DDT; 5 ng/µL in nonane). Extracts were concentrated to ~1 mL on a rotary evaporator, solvent exchanged to hexane, and concentrated to ~50 µL. Ten µL of 2,4,6-tribromobiphenyl (5 ng/µL), was added as an injection standard before analysis.

PUF samples were extracted using a Dionex ASE 350 (Dionex Corporation, Sunnyvale, CA 94088) accelerated solvent extraction device after spiking with the surrogate standard

125 mixture. Extracts were concentrated to a final volume of ~1 mL using a rotary evaporator (after
126 solvent exchange into hexane) and passed on a glass chromatographic cleanup column packed
127 with silica gel/alumina (2:1 wt/wt) in order to remove the interfering compounds (Khairy and
128 Lohmann, 2012). The collected fraction was concentrated to a final volume of ~ 50 μ L. Finally,
129 2,4,6-tribromobiphenyl was added as the injection standard before analysis.

130

131 **2.5. Instrumental Analysis and Quality Control**

132 All samples were analyzed for OCPs with an Agilent 6890 Series GC System connected
133 to an Agilent 5973 Network Mass Selective Detector. DDTs (p,p'-DDE, o,p'-DDD, p,p'-DDD,
134 o,p'-DDT and p,p'-DDT), hexachlorocyclohexanes (HCHs: α -, β -, γ - and δ -HCH isomers),
135 chlordanes (trans- and cis-chlordane isomers, oxychlordane and trans-nonachlor), heptachlor and
136 its epoxide, aldrin, dieldrin, endosulfans (endosulfan I and II isomers and endosulfan sulfate),
137 hexachlorobenzene (HCBz), endrin, endrin aldehyde, endrin ketone and methoxychlor were
138 investigated. Separation of OCPs was accomplished with a DB-5 MS fused silica capillary
139 column (30 m $\times$ 0.25 mm i.d., 0.25 μ m film thickness, J&W Scientific). Abbreviations of OCPs
140 and more details on the instrumental analysis are given in Table (SI 3) and text (SI 3).

141 Procedural blanks, field blanks, matrix spikes and duplicate samples (20 % of the total
142 samples) were included with each sample batch, and were carried throughout the entire analytical
143 procedure in a manner identical to the samples. OCPs were completely absent from procedural
144 and field blanks indicating negligible contamination during transport, storage and analysis.
145 Limits of detection (LODs) were determined as the concentration of analytes in a sample giving
146 a peak with a signal-to-noise (S/N) of 3 (Doong et al., 2002). (QA/QC procedures are detailed in
147 text SI 4 and Table SI 3).

2.6. Physico-Chemical Properties

Internally consistent air-water partitioning coefficients (K_{AW}) and octanol-air partitioning coefficients (K_{OA}) were obtained from Schenker et al. (2005). Missing K_{OA} values were obtained by correlating available K_{OA} values from Schenker et al. (2005) against values obtained from EPI Suite (USEPA, 2011) (Figure SI 2). Missing K_{AW} values were calculated according to equation 1 (Reinhard and Drefahl, 1999):

$$K_{AW} = \frac{H_c}{RT} \quad (1)$$

where H_c is Henry's law constant (atm.L.mol^{-1}), R is the gas constant (0.08206 in L.atm/mol.K) and T is the absolute temperature in Kelvin. H_c for δ -HCH was obtained from Meylan and Howard (2005). Values for oxychlordane, trans-nonachlor, o,p'-DDD, o,p'-DDT and endosulfan sulfate were obtained from SPARC online calculator (V4.5). Values for endrin aldehyde, endrin ketone and methoxychlor were obtained from Mackay et al. (2006). Internally consistent values of the sub-cooled liquid vapor pressure (P_L/Pa) were obtained from Shen and Wania (2005). Missing P_L values were obtained by correlating values from Shen and Wania (2005) with those obtained from Mackay et al. (2006) (Figure SI 3). P_L value for trans-nonachlor was obtained from Shoeib and Harner (2002). Values for endrin aldehyde, endosulfan sulfate and endrin ketone were obtained from EPI Suite. Enthalpies of vaporization (ΔH_{vap} in KJ/mol) were obtained from Shoeib and Harner (2002). Missing ΔH_{vap} values were obtained by correlating available values against P_L (Figure SI 4). Sampler-water partitioning coefficients (K_{PE-W}) were calculated according to Lohmann (2012). Values of all the physico-chemical parameters are given in Table (SI 4).

Deleted:

171 2.7. Estimated atmospheric concentrations based on LDPE

172 PRCs, impregnated in the polyethylene samplers before their deployment, were used to
173 gauge whether OCPs had achieved equilibrium and to adjust for disequilibrium in polyethylene
174 (C_{LDPE}) assuming that uptake and elimination rates are equivalent. (Booij et al., 2002) (Text SI
175 5). Gaseous atmospheric concentrations (C_g , ng/l) were calculated from disequilibrium- corrected
176 polyethylene concentrations (C_{LDPE}) as follows:

$$177 \quad C_g = \frac{C_{LDPE}}{(1 - e^{-k_e t}) \cdot K_{PE-A}} \quad (2)$$

178 where C_{LDPE} is the OCP concentration in the LDPE (ng/kg), k_e is the loss rate constant of PRCs
179 (day^{-1}), t is the deployment period (days) and K_{PE-A} is the sampler-air partitioning coefficient
180 (L/kg). To estimate K_{PE-A} values for OCPs, a high volume sampler was left operating for a
181 period of 15 days (5 days per week) at one of the sampling sites (site 2) during the summer
182 season (See text SI 2 for more details). Detected OCP concentrations in the LDPE and PUF
183 samples were used to calculate the K_{PE-A} values according to equation 2. Partition coefficients
184 were adjusted for temperature according to a modified form of Van't Hoff equation (equation 3)

$$185 \quad K_{PE-A}(T_2) = K_{PE-A}(T_1) \times e^{\frac{-\Delta H_{\text{vap}}}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)} \quad (3)$$

186 where T_2 and T_1 are the mean temperatures of the deployment and at which the partitioning
187 properties were determined (K).

Deleted: and ΔH_{vap} is the enthalpy of vaporization (J mol^{-1})

189 2.8. Uncertainty associated with K_{PE-A} calculations

190 Calculated overall uncertainty combined the uncertainty of predicted partition
191 coefficients, equilibrium adjustment of polyethylene concentrations based on the use of PRCs

(the fraction equilibration remaining at time of PE sampler collection), and the uncertainty associated with the analysis of the OCPs (See text SI 6 for more details). K_{PE-A} values for undetected OCPs (β -HCH, δ -HCH, dieldrin, endosulfan sulfate and methoxychlor) in the field study were predicted from a simple linear relationship with P_L (see section 3.2, equation 5). Accordingly, we used the same relative uncertainties (RU) in predicted K_{PE-As} as given for the log P_L by Shen and Wania (2005). Since internally consistent P_L values did not exist for δ -HCH, endosulfan sulfate and methoxychlor, they were assigned the highest uncertainty estimate of P_L (500 %). The overall uncertainty (Table 1) ranged from ± 43.0 % to ± 504 % (0.20-0.80 log units), with higher uncertainty values (303-504 %) for predicted K_{PE-As} compared to the field measured ones (43.0-65.0 %) owing to the greater uncertainty estimates of their P_L .

Deleted:

Deleted:

Deleted: only as that of

Deleted: given

Deleted:

Deleted:

Table 1

3. Results and Discussion

3.1. Uptake of OCPs by the LDPE

The addition of PRCs to the samplers before deployment is useful to infer the exchange rate kinetics, assuming that the elimination rates (of PRCs) and uptake rates (of native compounds) are equivalent (Bartkow et al., 2004). Exchange kinetics were quantified by k_e (for details, see text SI 5). Four PRCs with different log K_{PE-A} values were used in this study. The least dissipated PRC in all the investigated samples was pentabromobiphenyl (PBB 103) (Table SI 5). A higher elimination rate was observed for PBB 103 during the winter (30-46 %) compared to the summer (25-33 %). The Kruskal-Wallis one way ANOVA statistical test indicated that loss rate constants decrease with increasing the sampler-air partitioning coefficient (log K_{PE-A}) (PBB 103 < OCN < PBB 52 < PBB 9) in both seasons ($H_{\text{summer}} = 39.1$, $H_{\text{winter}} = 38.6$

Deleted:

215 at $\alpha = 0.05$, $p < 0.001$). Moreover, loss rate constants for each of the PRCs were significantly
 216 higher in the winter compared to the summer season [Mann-Whitney rank sum test for PBB 9
 217 and PBB 103: $T = 66$ and 73 respectively at $p < 0.001$; t-test for PBB 52 ($t = -5.56$ at $p < 0.001$)
 218 and OCN ($t = -2.37$ at $p = 0.028$)], which was probably related to the increased wind speed
 219 during the winter season (Table SI 1). This indicates that the chemical exchange between the
 220 LDPE sampler and the air was limited by the air side boundary layer resistance ($k_e \propto 1/K_{PE-A}$)
 221 based on the following relation:

$$222 \quad k_e = \frac{k_a \cdot A_s}{V_s \cdot K_{PE-A}} \quad (4)$$

223 where k_a is the mass transfer coefficient through the air side boundary layer and A_s and V_s are
 224 the sampler surface area and volume respectively.

225 A correlation between log transformed k_e values and log K_{PE-A} of the PRCs was
 226 established to extrapolate k_e for all the investigated OCPs (Tables SI 6 and 7). At the end of the
 227 21 days deployment period, the average values of % equilibrium for OCPs (Figure 1) in the
 228 samples indicated that OCPs with higher P_L values were approaching equilibrium (α -HCH, γ -
 229 HCH, HCBz and aldrin) (> 80 % equilibrium), whereas OCPs like DDTs were still far from
 230 reaching equilibrium (< 60 %). To gain more information about the accumulation patterns of
 231 OCPs, three LDPEs were deployed in each season at one of the sites (site 11), and one was
 232 harvested after each week of exposure period (Figure 1). The equilibration of α - and γ -HCH,
 233 HCBz, aldrin, heptachlor, trans- and cis-chlordane after each week of deployment (week 2 -
 234 week 1 and week 3 - week 2) started to deviate from the linear uptake after three weeks
 235 deployment period in the summer season (Figure 1a), indicating that they were approaching
 236 equilibrium. Contrarily, endosulfan I and II, trans-nonachlor, endrin, oxychlordane, p,p'-DDE
 237 and p,p'-DDT increased linearly during the sampling period. During the winter season (Figure

1b), only p,p'-DDD/o,p'-DDT, o,p'-DDD and p,p'-DDT increased linearly during the sampling period, whereas all the other detected OCPs were approaching equilibrium. When the approached % equilibrium results of OCPs based on the PRCs at site 11 were compared to the variation in the accumulated amounts of OCPs in the LDPE (ng/g PE) with time (Figure SI 6), good agreement was generally observed for the predicted stage of exchange of OCPs between the LDPE sampler and air in both seasons. This implies that PRCs can be used to calibrate sampler/site specific mass transfer behavior, and thus accurately estimate the gaseous concentrations of OCPs (and other POPs) in the atmosphere.

Deleted:

Figure 1

3.2. Sampler-air partitioning constants (K_{PE-A})

PE-A partitioning constants were calculated directly for 17 OCPs which were detected both in the passive sampler (corrected for non-equilibrium) and the PUFs at site 2 during the field deployments (Table 1). To predict values for the other investigated OCPs, we examined correlation between our measured K_{PE-AS} and literature K_{OA} values, which has been used previously to predict K_{PE-AS} for PAHs (Bartkow et al., 2004; Khairy and Lohmann, 2012). As shown in Figure (SI 5), a highly significant and strong linear correlation can be observed for PAHs ($R^2 > 0.90$, $p < 0.001$), whereas for OCPs, the relation explained only 32 % of the total variability in the data.

Deleted: i

We observed a significant and strong correlation when K_{PE-AS} were regressed against P_L (Pa) ($p < 0.001$) (Figure 2a). The correlation explained 90.0 % of the total variability in the data. The slope was insignificantly different from -1 ($p < 0.001$; $R^2 = 0.90$), implying that P_L can be used as a good predictor of K_{PE-A} , and that the magnitude of K_{PE-A} for OCPs is dominated by their volatilities. The derived K_{PE-A} were calculated from two 3-week field deployments and

constrained by our ability to safely operate active sampling equipment during the entire field work. Yet the strong correlation with P_L implies that we were able to derive values that are well constrained by physico-chemical properties, validating the field-derived data. Accordingly, K_{PE-A} values for the non-quantified OCPs in the passive sampler were predicted (Table 2) based on the derived P_L - K_{PE-A} relation (equation 5):

$$\log K_{PE-A} = -0.77 \pm 0.07 \cdot \log P_L + 6.35 \pm 0.13 \quad (5)$$

(n = 17; $R^2 = 0.90$; SE = 0.19)

Table 2

Figure 2

We propose that K_{PE-A} for the non-quantified OCPs in the field calibration study can be best predicted from equation (5) despite of the higher degree of uncertainty for endrin, endrin aldehyde, endosulfan sulfate, endrin ketone and methoxychlor (Table 1) as internally consistent physico-chemical properties were missing. For further assessment, calculated K_{PE-A} values were compared with K_{PE-A} values derived using sampler-water partitioning coefficients (K_{PE-W}) and air-water partitioning coefficients (K_{AW}) according to equation 6:

$$K_{PE-A} = \frac{K_{PE-W}}{K_{AW}} \quad (6)$$

Calculated (from the field experiment) and predicted (K_{PE-W}/K_{AW}) K_{PE-A} values agreed well within 94 % (factor difference < 0.4 log units) for the majority of the investigated OCPs (Figure 2b). The good agreement observed for most OCPs supports the assumption that our field-derived K_{PE-A} values are good approximations of their real values, despite the limited deployment period in the field. Contrarily, lower degree of agreement was observed for endrin

286 aldehyde, endrin ketone, endosulfan sulfate, trans-nonachlor and oxychlordane (factor difference
287 ranged from 0.60-3.8 log units). One explanation for this lower agreement is related to the degree
288 of uncertainty associated with the partitioning coefficients (uncertainty from P_L and K_{AW} +
289 uncertainty from the K_{PE-A} and K_{PE-W} partitioning coefficients).

290

291 3.3. Spatial distribution and seasonal variations of OCPs

292
293 Estimated atmospheric OCP concentrations in Alexandria are given in Tables (3, SI 8 and

294 SI 9). Based on the average concentrations, the overall trend of OCPs showed the following

295 order: HCBz ($78 \pm 46 \text{ pg/m}^3$) > HCHs (sum of α - and γ - isomers: $67 \pm 33 \text{ pg/m}^3$) > chlordanes

296 (sum of cis- and trans- isomers, oxychlordane, trans-nonachlor, heptachlor and its epoxide: 60.5

297 $\pm 75 \text{ pg/m}^3$), aldrin ($59 \pm 48 \text{ pg/m}^3$) > endosulfans (sum of endosulfan I and II: $45 \pm 49 \text{ pg/m}^3$) >

298 DDTs (sum of o,p'-DDD, o,p'-DDT, p,p'-DDE, p,p'-DDD and p,p'-DDT: $34.5 \pm 43 \text{ pg/m}^3$) in

299 the summer season, and HCHs ($82 \pm 29 \text{ pg/m}^3$) > HCBz ($63 \pm 31 \text{ pg/m}^3$) > aldrin (30 ± 31

300 pg/m^3) > chlordanes ($19 \pm 15 \text{ pg/m}^3$) > endosulfans ($15 \pm 19 \text{ pg/m}^3$) > DDTs ($14 \pm 12 \text{ pg/m}^3$) in

301 the winter season. Endrin, endrin aldehyde (EA) and endrin ketone (EK) were only detected at

302 lower concentrations (<LOD- 28 pg/m^3) in the summer season and <LOD in all the winter

303 samples. For most of the investigated OCPs, highest concentrations were always observed at

304 sites 2 and 3 (Figure SI 7). These sites were located in the northeastern part of Alexandria,

305 closest to agricultural areas (Figure SI1). In the summer season, concentrations of chlordanes,

306 aldrin, endosulfans, endrin and its metabolites and DDTs were greater at site 2 and site 3,

307 whereas only slight between sample variations were observed for HCHs. A similar trend was

308 observed for aldrin, endosulfans and p,p'-DDE in the winter season. HCBz, in contrast, displayed

309 highest concentrations ($23\text{-}168 \text{ pg/m}^3$) in both seasons at site 2 and site 3, and other locations

Deleted:),

Deleted: which a

Deleted: and are the

Deleted: samples

Deleted: the

Deleted: metabolites

310 | characterized by high traffic and industrial activities (site 1, 4, 6 and 9). The average HCBz
311 | concentration in Alexandria was only slightly higher than its northern hemisphere average
312 | concentration (~ 50 pg/m³) (Jin et al., 2013), but much lower than measured concentrations in
313 | China (Liu et al., 2009) and Korea (Li et al., 2007) (Table SI 10). Although HCBz production
314 | has ceased in most countries, it is still being generated inadvertently as a by-product and/or
315 | impurity in several chemical processes, such as the manufacture of chlorinated solvents,
316 | chlorinated aromatics and pesticides. It is also released to the environment by incomplete
317 | combustion, and from old dumpsites (Barber et al., 2005).

Deleted: Average

Deleted: is

Deleted: in the northern hemisphere

318 | **Table 3**

319 | α - and γ -HCH were the only detected HCH isomers in the atmospheric environment of
320 | Alexandria (Figure SI 7). In all the samples, α -HCH concentrations (<LOD - 98.0 pg/m³) were
321 | much higher than the γ -HCH (<LOD - 36.0 pg/m³), especially in the winter season.

322 | Aldrin was detected in 82 % of the investigated PE samples with relatively high
323 | concentrations (<LOD - 147 pg/m³) compared to HCHs and HCBz (Figure SI 7). Chlordanes
324 | were detected in the majority of the investigated samples. Based on the average concentrations,
325 | trans-chlordane showed the highest concentrations in all the samples during both seasons
326 | (summer: 19 pg/m³; winter: 9 pg/m³) followed by cis-chlordane (8.0; 4.0 pg/m³). Oxychlordane,
327 | the metabolite of chlordane, was only observed in the summer season with an average
328 | concentration of 7.0 pg/m³. Heptachlor epoxide, the metabolite of heptachlor was also observed
329 | only in the summer season (higher temperature). The detection of both metabolites only in the
330 | summer suggests that they were produced from the photodegradation of parent compounds (Park
331 | et al., 2011).

332 Endosulfan is still in use globally for agriculture and also for the control of disease
333 vectors (Pozo et al., 2011). Concentrations of endosulfans ranged from 9.0-160 pg/m³ and
334 <LOD-65 pg/m³ in the summer and winter seasons respectively. In all the samples, endosulfan I
335 concentrations were higher than endosulfan II, and endosulfan sulfate was <LOD in all the
336 samples.

337 p,p'-DDT was widely used in Egypt for agricultural purposes and for the control of
338 disease vectors. Although officially banned in 1988, it was recently detected in sediments of the
339 Egyptian Mediterranean Sea coast (Khairy et al., 2012) and in coastal lakes (Barakat et al.,
340 2012a, b), because of its long environmental half-life time. In the current study, concentrations of
341 DDTs were lower than those of HCHs, endosulfans, chlordanes and HCBz (Figure SI 7) ranging
342 from 4 - 168 ng/m³ and 4-29 pg/m³ in the summer and winter seasons respectively. p,p'-DDE
343 was the most frequent and abundant isomer in all the samples comprising 48-96 % of the total
344 concentrations of DDTs in Alexandria.

345 No significant seasonal variations were observed for HCBz, chlordanes (trans- and cis-
346 isomers), aldrin and DDTs (p: 0.115-0.861), whereas concentrations of γ -HCH and endosulfans
347 were significantly higher in the summer (p = 0.009-0.048), and α -HCH was significantly higher
348 in the winter (p = 0.035) (Figure SI 7).

349 HCBz concentrations in Alexandria were higher than concentrations measured in Europe
350 but lower than in Asia (Table SI 10). Chlordane concentrations were among the highest when
351 compared to concentrations in other locations worldwide using PUF passive samplers). γ -HCH
352 and endosulfans on the other hand were lower than concentrations measured in parts of Europe,
353 Canada, India, China and in the other African countries such as Ghana and South Africa.

Deleted:

Deleted: their

Deleted: DDTs

Deleted: DDTs

Deleted: significance

Comment [RL1]: Didn't you already comment on HCB above? This last paragraph should either be deleted, or you paste the sentences in the relevant sections above.

355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377

3.4. Source identification of OCPs

Several isomeric ratios were applied (Figure SI 8) in the current study to determine the source of the atmospheric OCPs (HCHs), and whether OCPs were released to the atmosphere as a result of fresh application or an aged use (HCHs, chlordanes, endosulfans and DDTs).

In the current study, α -HCH/ γ -HCH ratio ranged from 1.3-2.9 (Figure SI 8) in the summer (hot) season, and 3.2-6.7 in the winter (cold) season indicating a possible seasonal re-volatilization of old lindane residues. Lower ratio values observed in the summer season could be related to the increased lindane volatilization from soil as a result of elevated atmospheric temperatures.

Comment [RL2]: Needs ref – maybe from Bidleman's work??

Technical chlordane has been widely used as agriculture pesticide and termiticide in all over the world (Syed et al., 2013). Calculated ratios of trans/cis chlordane in the current study ranged from 1.5-2.9 (Figure SI 8) in both seasons, indicating fresh application of technical chlordane (Hinkley et al., 1990, Jantunen et al., 2000). According to the UNEP, the use of chlordane is severely restricted and limited to non-agricultural uses in Egypt (UNEP, 2003).

Endosulfan comprises two isomers, endosulfan I and II, with a I/II ratio of about 2.33 (Daly et al. 2007) in the technical mixture. Endosulfan II is more reactive and so an elevated I/II ratio is representative of an aged signature, whereas ratios closer to the starting technical mixture suggest recent application. As shown in Figure (SI 8), calculated values in the current study were around the value of the technical mixture (2.3) indicating recent application of endosulfans.

In this study, the ratio of p,p'-DDE/p,p'-DDT ranged from 2-106, indicating an aged application of DDT in the area.

378

379

380 **3.5. Evaluation of the Predictive Ability of the LDPE Samplers**

381 Estimated gaseous OCP concentrations calculated from the LDPE passive samplers
382 corrected for non-equilibrium were compared with the gaseous concentrations actually measured
383 in the atmosphere using the high volume sampler. Results are given in Tables (SI 8 and 9).
384 Although the difference in the sampling periods between the passive (21 days) and active (2–3
385 days) sampling periods adds uncertainty to the results in Alexandria (Egypt), good agreement
386 was generally observed between estimated and measured OCP concentrations in all the
387 investigated samples (Figure 3 a & b). During the summer and winter seasons, the difference
388 ranged from a factor of 0.8 to 2.0. Additionally, slopes of both regression lines were not
389 significantly different from one (at $\alpha = 0.05$, $P < 0.001$) supporting this good agreement. The
390 significant correlations obtained here suggest that the gas-phase OCP concentrations in
391 Alexandria might not display significant short-term variations during our measurement periods.
392 This assumption is supported by the observed statistical insignificant difference ($p = 0.128$)
393 between the weekly measured gas-phase concentrations at site 2 (Figure SI 9) sampled with the
394 conventional active sampling method. Overall, LDPEs have proven to be a powerful tool for
395 predicting atmospheric OCP concentrations in the present study.

396

Figure 3

397 **4. Conclusions**

398 Our results indicated the feasibility of using LDPE as a practical and low-cost technique
399 for monitoring OCPs in Egypt, and to provide data on the spatial distribution and seasonal
400 variations of OCPs in an urban Egyptian city. Including PRCs enables the use of PE sheets as

passive air samplers for gas-phase OCPs in general. While our work demonstrated the importance of using PRCs in verifying the equilibrium status of OCPs in PE samplers, a cheaper alternative to circumvent the PRC approach is the use of LDPEs with different surface area-to-volume ratios, as presented by Bartkow et al. (2004). Different passive sampling materials are now popular and their utilization for monitoring organic contaminants in the atmosphere has greatly increased. Future work should concentrate on comparing different passive samplers as a function of deployment times, temperature and wind speed so that data derived from different studies are broadly comparable, and the levels of confidence around the data are known.

Estimated concentrations of atmospheric OCPs in Alexandria, Egypt were within the global range as measured in the global atmospheric passive sampling network (GAPS). Yet the detection of OCPs in the Egyptian environment, especially those with fresh sources as indicated by the isomeric ratios questions the efficacy of the official ban of most of the investigated OCPs in Egypt from decades. This could encourage decision makers in Egypt (and beyond) to utilize this cost-effective sampling tool for developing an extensive local air sampling network. With the use of passive samplers, data can be acquired about sources and concentrations of organic pollutants and to develop effective control measures.

Supplementary information

Supplementary data associated with this article can be found in the online version.

Acknowledgment

Authors would like to thank Mr. Dave Adelman (GSO, URI) for his effort in PE preparation. Dr. Mohammed Khairy acknowledges the Fulbright Foundation for offering him the opportunity to carry this research in USA. Dr. Rainer Lohmann acknowledges funding from EPA's Great Lakes Restoration Initiative Award GLAS # 00E00597-0 supporting passive sampler research at URI.

5. References

- Abou-Arab, A.A.K., Gomaa, M.N.E., Badawy, A., Naguib, K., 1995. Distribution of organochlorine pesticides in the Egyptian aquatic ecosystem. *Food Chemistry* 54, 141-146 .
- Baek, S.Y., Jurng, J., Chang, Y.S., 2013. Spatial distribution of polychlorinated biphenyls, organochlorine pesticides, and dechlorane plus in Northeast Asia. *Atmospheric Environment* 64, 40-46.
- Bailey, R., Barrie, L.A., Halsall, C.J., Fellin, P., Muir, D.C.G., 2000. Atmospheric organochlorine pesticides in the western Canadian Arctic: Evidence of transpacific transport. *Journal of Geophysical Research: Atmospheres* 105, 11805-11811.
- Barakat, A.O., Mostafa, A., Wade, T.L., Sweet, S.T., El Sayed, N.B., 2012a. Assessment of persistent organochlorine pollutants in sediments from Lake Manzala, Egypt. *Marine Pollution Bulletin* 64, 1713-1720.
- Barakat, A.O., Mostafa, A., Wade, T.L., Sweet, S.T., El Sayed, N.B., 2012b. Spatial distribution and temporal trends of persistent organochlorine pollutants in sediments from Lake Maryut, Alexandria, Egypt. *Marine Pollution Bulletin* 64, 395-404.
- Barakat, A.O., Khairy, M., Aukaily, I., 2013. Persistent organochlorine pesticide and PCB residues in surface sediments of Lake Qarun, a protected area of Egypt. *Chemosphere* 90, 2467-2476.
- Barber, J.L., Sweetman, A.J., van Wijk, D., Jones, K.C., 2005. Hexachlorobenzene in the global environment: Emissions, levels, distribution, trends and processes. *Science of the Total Environment* 349, 1-44.
- Bartkow, M.E., Hawker, D.W., Kennedy, K.E., Muller, J.F., 2004. Characterizing uptake kinetics of PAHs from the air using polyethylene-based passive air samplers of multiple surface

451 area-to-volume ratios. *Environmental Science and Technology* 38, 2701-2706.
 452 Booij, K., Smedes, F., van Weerlee, E.M., 2002. Spiking of performance reference compounds in
 453 low density polyethylene and silicone passive water samplers. *Chemosphere* 46, 1157-1161.
 454 Choi, S.D., Baek, S.Y., Chang, Y.S., Wania, F., Ikonomou, M.G., Yoon, Y.J., Park, B.K., Hong,
 455 S., 2008. Passive air sampling of polychlorinated biphenyls and organochlorine pesticides at the
 456 Korean Arctic and Antarctic research stations: Implications for long-range transport and local
 457 pollution. *Environmental Science and Technology* 42, 7125-7131.
 458 Daly, G.L., Lei, Y.D., Teixeira, C., Muir, D.C.G., Castillo, L.E., Jantunen, L.M.M., Wania, F.,
 459 2007. Organochlorine pesticides in the soils and atmosphere of Costa Rica. *Environmental*
 460 *Science and Technology* 41, 1124-1130.
 461 Devi, N.L., Qi, S., Chakraborty, P., Zhang, G., Yadav, I.C., 2011 Passive air sampling of
 462 organochlorine pesticides in a northeastern state of India, Manipur. *Journal of Environmental*
 463 *Sciences* 23, 808-815.
 464 Doong, R.A., Sun, Y.C., Liao, P.L., Peng, C.K., Wu, S.C., 2002. Distribution and fate of
 465 organochlorine pesticide residues in sediments from the selected rivers in Taiwan. *Chemosphere*
 466 48, 237-246.
 467 El Nemr, A., Abd-Allah, A.M.A., 2004. Organochlorine contamination in some marketable fish
 468 in Egypt. *Chemosphere* 54, 1401-1406.
 469 EPA, U.S. EPI Suite v4.10. 2011.
 470 Hinckley, D.A., Bidleman, T.F., Foreman, W.T., Tuschall, J.R., 1990 Determination of vapor
 471 pressures for nonpolar and semipolar organic compounds from gas chromatographic retention
 472 data. *Journal of Chemical Engineering Data* 35, 232-237.
 473 Hung, H., Kallenborn, R., Breivik, K., Su, Y., Brorstrom-Lunden, E., Olafsdottir, K., Thorlacius,

474 J.M., Leppanen, S., Bossi, R., Skov, H., Mano, S., Patton, G.W., Stern, G., Sverko, E., Fellin, P.,
 475 2010. Atmospheric monitoring of organic pollutants in the Arctic under the Arctic Monitoring
 476 and Assessment Programme (AMAP): 1993-2006. *Science of the Total Environment* 408, 2854-
 477 2873.

478 Jantunen, L.M.M., Bidleman, T.F., Harner, T., Parkhurst, W.J., 2000. Toxaphene, chlordane, and
 479 other organochlorine pesticides in Alabama air. *Environmental Science and Technology* 34,
 480 5097-5105.

481 Jaward, F.M., Farrar, N.J., Harner, T., Sweetman, A.J., Jones, K.C., 2004 Passive Air Sampling
 482 of PCBs, PBDEs, and Organochlorine Pesticides Across Europe. *Environmental Science and*
 483 *Technology* 38, 34-41.

484 Jaward, F.M., Zhang, G., Nam, J.J., Sweetman, A.J., Obbard, J.P., Kobara, Y., Jones, K.C.,
 485 2005. Passive air sampling of polychlorinated biphenyls, organochlorine compounds, and
 486 polybrominated diphenyl ethers across Asia. *Environmental Science and Technology* 39, 8638-
 487 8645.

488 Jin, G.Z., Kim, S.M., Lee, S.Y., Park, J.S., Kim, D.H., Lee, M.J., Sim, K.T., Kang, H.G., Kim,
 489 I.G., Shin, S.K., Seok, K.S., Hwang, S.R., 2013 Levels and potential sources of atmospheric
 490 organochlorine pesticides at Korea background sites. *Atmospheric Environment* 68, 333-342.

491 Kennedy, K.E., Hawker, D.W., Muller, J.F., Bartkow, M.E., Truss, R.W., 2007 A field
 492 comparison of ethylene vinyl acetate and low-density polyethylene thin films for equilibrium
 493 phase passive air sampling of polycyclic aromatic hydrocarbons. *Atmospheric Environment* 41,
 494 5778-5787.

495 Khairy, M.A., Lohmann, R., 2012. Field Validation of polyethylene passive air samplers for
 496 parent and alkylated PAHs in Alexandria, Egypt. *Environmental Science and Technology* 46,

497 3990-3998.

498 Khairy, M.A., Kolb, M., Mostafa, A., El-Fiky, A., Bahadir, M., 2012 Risk posed by chlorinated
 499 organic compounds in Abu Qir Bay, East Alexandria, Egypt. *Environmental Science and*
 500 *Pollution Research* 19, 794-811.

501 Lemaire, G.R., Terouanne, B.A., Mauvais, P., Michel, S., Rahmani, R., 2004. Effect of
 502 organochlorine pesticides on human androgen receptor activation in vitro. *Toxicology and*
 503 *Applied Pharmacology* 196, 235-246.

504 Li, J., Zhang, G., Guo, L., Xu, W., Li, X., Lee, C.S.L., Ding, A., Wang, T., 2007. Organochlorine
 505 pesticides in the atmosphere of Guangzhou and Hong Kong: Regional sources and long-range
 506 atmospheric transport. *Atmospheric Environment* 41, 3889-3903.

507 Liu, X., Zhang, G., Li, J., Yu, L.L., Xu, Y., Li, X.D., Kobara, Y., Jones, K.C., 2009. Seasonal
 508 patterns and current sources of DDTs, chlordanes, hexachlorobenzene, and endosulfan in the
 509 atmosphere of 37 Chinese Cities. *Environmental Science and Technology* 43, 1316-1321.

510 Lohmann, R., 2012 Critical review of low-density polyethylene's partitioning and diffusion
 511 coefficients for trace organic contaminants and implications for its use as a passive sampler.
 512 *Environmental Science and Technology* 46, 606-618.

513 Lohmann, R., Corrigan, B.P., Howsam, M., Jones, K.C., Ockenden, W.A., 2001. Further
 514 developments in the use of semipermeable membrane devices (SPMDs) as passive air samplers
 515 for persistent organic pollutants: field application in a spatial survey of PCDD/Fs and PAHs.
 516 *Environmental Science and Technology* 35, 2576-2582.

517 Mackay, D.S., Wan, W.Y., Ma, K.C., Lee, S.C., 2006 Handbook of physical-chemical properties
 518 and environmental fate of organic chemicals. Second edition ed.; CRC Press, Taylor & Francis
 519 Group, : Florida, USA, 2006.

520 Mansour, S.A., 2004. Pesticide exposure-Egyptian scene, *Toxicology* 198, 91-115.

521 Messing, P., Farenhorst, A., Waite, D., Sproull, J., 2013. Influence of usage and chemical-
 522 physical properties on the atmospheric transport and deposition of pesticides to agricultural
 523 regions of Manitoba, Canada. *Chemosphere* 90, 1997-2003.

524 Meylan, W.M., Howard, P.H., 2006. Estimating octanol-air partition coefficients with octanol-
 525 water partition coefficients and Henry's law constants. *Chemosphere* 61, 640-644.

526 Murvoll, K.M., Skaare, J.U., Moe, B., Anderssen, E., Jenssen, B.M., 2006. Spatial trends and
 527 associated biological responses of organochlorines and brominated flame retardants in hatchlings
 528 of North Atlantic kittiwakes (*Rissa tridactyla*). *Environmental Toxicology and Chemistry* 25,
 529 1648-1656.

530 Nakata, H., Kawazoe, M., Arizono, K., Abe, S., Kitano, T., Shimada, H., Li, W., Ding, X., 2002.
 531 Organochlorine pesticides and polychlorinated biphenyl residues in foodstuffs and human tissues
 532 from China: Status of contamination, historical trend, and human dietary exposure. *Archives of*
 533 *Environmental Contamination and Toxicology* 43, 0473-0480.

534 Park, J.S., Shin, S.K., Kim, W., Kim B.H., 2011. Residual levels and identify possible sources of
 535 organochlorine pesticides in Korea atmosphere. *Atmospheric Environment* 45, 7496-7502.

536 Pozo, K., Harner, T., Lee, S.C., Sinha, R.K., Sengupta, B., Loewen, M., Geethalakshmi, V.,
 537 Kannan, K., Volpi, V., 2011. Assessing seasonal and spatial trends of persistent organic
 538 pollutants (POPs) in Indian agricultural regions using PUF disk passive air samplers.
 539 *Environmental Pollution* 159, 646-653.

540 Reinhard, M., Drefahl, A., 1999. Handbook for estimating physicochemical properties for
 541 organic compounds; John Wiley & Sons, Inc.: New York, USA.

542 Sallam, K.I., Morshedy, A.E.M.A., 2008. Organochlorine pesticide residues in camel, cattle and

543 sheep carcasses slaughtered in Sharkia Province, Egypt. Food Chemistry 108, 154-164.

544 Schenker, U., MacLeod, M., Scheringer, M., Hungerbuhler, K., 2005. Improving data quality for
 545 environmental fate models: A least-squares adjustment procedure for harmonizing
 546 physicochemical properties of organic compounds. Environmental Science and Technology
 547 39, 8434-8441.

548 Shen, L., Wania, F., Lei, Y.D., Teixeira, C., Muir, D.C.G., Bidleman, T.F., 2004.
 549 Hexachlorocyclohexanes in the North American Atmosphere. Environmental Science and
 550 Technology 38, 965-975.

551 Shen, L., Wania, F., 2005. Compilation, evaluation, and selection of physical-chemical property
 552 data for organochlorine pesticides. Journal of Chemical and Engineering Data 50, 742-768.

553 Shen, L., Wania, F., Lei, Y.D., Teixeira, C., Muir, D.C.G., Bidleman, T.F., 2005. Atmospheric
 554 Distribution and Long-Range Transport Behavior of Organochlorine Pesticides in North
 555 America. Environmental Science and Technology 39, 409-420.

556 Shoeib, M., Harner, T., 2002. Using measured octanol-air partition coefficients to explain
 557 environmental partitioning of organochlorine pesticides. Environmental Toxicology and
 558 Chemistry 21, 984-990.

559 Souza, M.S., de Potas, G.M., de D'Angelo, A.M.P., 2004. Organophosphorous and
 560 organochlorine pesticides affect human placental phosphoinositides metabolism and PI-4 kinase
 561 activity. Journal of Biochemistry and Molecular Toxicology 18, 30-36.

562 Switch. Alexandria 2030:- Integrated urban water management (IUWM) strategic plan. Switch
 563 Central Management Unit, UNESCO-IHE, The Netherlands, 2011;

564 <http://switchurbanwater.lboro.ac.uk/outputs/pdfs/W1->
 565 [1_CALE_RPT_D1.1.6_Strategic_Planning_Process_-_Alexandria.pdf](http://switchurbanwater.lboro.ac.uk/outputs/pdfs/W1-1_CALE_RPT_D1.1.6_Strategic_Planning_Process_-_Alexandria.pdf).

566 Syed, L.H., Malik, R.N., Liu, D., Xu, Y., Wang, Y., Li, J., Zhang, G., Jones, K.C., 2013.
 567 Organochlorine pesticides in air and soil and estimated air-soil exchange in Punjab, Pakistan.
 568 Science of the Total Environment 444, 491-497.
 569 Tiido, T., Rignell-Hydbom, A., Jonsson, B., Giwercman, Y.L., Rylander, L., Hagmar, L.,
 570 Giwercman, A., 2005. Exposure to persistent organochlorine pollutants associates with human
 571 sperm Y:X chromosome ratio. Human Reproduction 20, 1903-1909.
 572 United Nation Environmental Program (UNEP), 2003. Draft technical guidelines on the
 573 environmentally sound management of persistent organic pollutants as wastes, Geneva.,
 574 Van Drooge, B.L., Grimalt, J.O., Torres Garcla, C.J.T., Cuevas, E., 2002. 'Semivolatile
 575 Organochlorine Compounds in the Free Troposphere of the Northeastern Atlantic.
 576 Environmental Science and Technology 36, 1155-1161.
 577 Wang, X.P., Gong, P., Yao, T.D., Jones, K.C., 2010. Passive air sampling of organochlorine
 578 pesticides, polychlorinated biphenyls, and polybrominated diphenyl ethers across the Tibetan
 579 Plateau. Environmental Science and Technology 44, 2988-2993.
 580 Zhang, G., Chakraborty, P., Li, J., Sampathkumar, P., Balasubramanian, T., Kathiresan, K.,
 581 Takahashi, S., Subramanian, A., Tanabe, S., Jones, K.C., 2008. Passive atmospheric sampling of
 582 organochlorine pesticides, polychlorinated biphenyls, and polybrominated diphenyl ethers in
 583 urban, rural, and wetland sites along the coastal length of India. Environmental Science and
 584 Technology 42, 8218-8223.
 585