

8-2013

Hemlock Woolly Adelgid and Elongate Hemlock Scale Induce Changes in Foliar and Twig Volatiles of Eastern Hemlock

Joshua Pezet

Joseph Elkinton

Sara Gomez

E. Alexa McKenzie

Michael Lavine

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/bio_facpubs

Citation/Publisher Attribution

Pezet, J., Elkinton, J., Gomez, S., McKenzie, E. A., Lavine, M., & Preisser, E. (2013). Hemlock woolly adelgid and elongate hemlock scale induce changes in foliar and twig volatiles of eastern hemlock. *Journal of Chemical Ecology*, 39(8), 1090-100.

Available at: <http://dx.doi.org/10.1007/s10886-013-0300-5>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Hemlock Woolly Adelgid and Elongate Hemlock Scale Induce Changes in Foliar and Twig Volatiles of Eastern Hemlock

Authors

Joshua Pezet, Joseph Elkinton, Sara Gomez, E. Alexa McKenzie, Michael Lavine, and Evan Preisser

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

1
2
3
4
5
6
7 HEMLOCK WOOLLY ADELGID AND ELONGATE HEMLOCK SCALE INDUCE CHANGES IN
8 FOLIAR AND TWIG VOLATILES OF EASTERN HEMLOCK
9

10
11 JOSHUA PEZET¹, JOSEPH ELKINTON^{1,2*}, SARA GOMEZ^{3,4},
12 E. ALEXA MCKENZIE¹, MICHAEL LAVINE⁵, and EVAN PREISSER³
13
14
15
16
17
18
19

20 ¹*Program in Organismic and Evolutionary Biology, University of Massachusetts, Amherst, MA 01003*
21

22 ²*Department of Environmental Conservation, University of Massachusetts, Amherst, MA 01003*
23

24 ³*Department of Biological Sciences, University of Rhode Island, Kingston, RI 02881*
25

26 ⁴*Department of Biology, Tufts University, Medford, MA 02155*
27

28 ⁵*Department of Mathematics and Statistics, University of Massachusetts, Amherst, MA 01003*
29
30
31
32
33
34
35
36
37

38 *To whom correspondence should be addressed. Email: elkinton@ent.umass.edu; phone: 413-545-4816;
39 fax: 413-545-5858.
40
41

42 **Abstract**— Eastern hemlock (*Tsuga canadensis*) is in rapid decline because of infestation by the invasive
43 hemlock woolly adelgid (*Adelges tsugae*; ‘HWA’) and, to a lesser extent, the invasive elongate hemlock
44 scale (*Fiorinia externa*; ‘EHS’). For many conifers, induced oleoresin-based defenses play a central role in
45 their response to herbivorous insects; however, it is unknown whether eastern hemlock mobilizes these
46 inducible defenses. We conducted a study to determine if feeding by HWA or EHS induced changes in the
47 volatile resin compounds of eastern hemlock. Young trees were experimentally infested for three years with
48 HWA, EHS, or neither insect. Twig and needle resin volatiles were identified and quantified by gas
49 chromatography/mass spectrometry. We observed a suite of changes in eastern hemlock’s volatile profile
50 markedly different from the largely terpenoid-based defense response of similar conifers. Overall, both
51 insects produced a similar effect: most twig volatiles decreased slightly, while most needle volatiles
52 increased slightly. Only HWA feeding led to elevated levels of methyl salicylate, a signal for systemic
53 acquired resistance in many plants, and benzyl alcohol, a strong antimicrobial and aphid deterrent. Green
54 leaf volatiles, often induced in wounded plants, were increased by both insects, but more strongly by EHS.
55 The array of phytochemical changes we observed may reflect manipulation of the tree’s biochemistry by
56 HWA, or simply the absence of functional defenses against piercing-sucking insects due to the lack of
57 evolutionary contact with these species. Our findings verify that HWA and EHS both induce changes in
58 eastern hemlock’s resin chemistry, and represent the first important step toward understanding the effects
59 of inducible chemical defenses on hemlock susceptibility to these exotic pests.

60

61 **Key Words**—*Adelges tsugae*; *Fiorinia externa*; *Tsuga canadensis*; plant-insect interactions; conifer
62 volatiles: induction.

63

INTRODUCTION

64

65 Conifers in the family Pinaceae are among the largest and longest-living organisms on earth. Their striking
66 longevity means that individual trees face an imposing array of biotic and abiotic challenges. They respond
67 to these challenges via complex constitutive and inducible defenses that enable them to survive under
68 highly diverse and taxing conditions and dominate vast areas of the earth's temperate and alpine forests
69 (Trapp and Croteau 2001, Dudareva et al. 2006).

70

71 Conifers commonly use oleoresin-based chemical defenses to combat herbivorous insects and
72 pathogens (Zulak and Bohlmann 2010). Oleoresin, or simply 'resin,' is a complex and species-specific
73 mixture of phytochemicals that is usually dominated by volatile monoterpenoids and non-volatile
74 diterpenoid acids but also contains smaller amounts of volatile organic chemicals such as sesquiterpenoids,
75 benzenoids (including phenolics), and fatty acid derivatives. These compounds are produced in resin-cells
76 of buds, needles and woody tissue, and in some conifers (such as *Pinus* species) they accumulate in
77 intercellular ducts or canals either constitutively or in response to trauma (Keeling and Bohlmann 2006).
78 Many conifers can respond to insect and microbial challenges via inducible increases in the biosynthesis
79 and accumulation of resin (Hudgins et al. 2004). These defenses variously act to physically engulf and
80 expel insects from the tree by the force of resin flow, seal off infected regions from surrounding tissue,
81 deter herbivory or oviposition, chemically interfere with insect developmental pathways, ATP production
82 and nervous system functioning, and disrupt microbial cell membranes causing cell leakage and death
83 (Langenheim 1994, Eyles et al. 2010). Herbivore attack can also induce the release of volatile resin
84 semiochemicals that attract predators of the colonizing plant-feeder (Mumm et al. 2003, Koepke 2010).

84

85 Over the last century, factors such as non-native pest introductions, forestry practices, and climate
86 change have sharply increased the amount of conifer mortality due to pests or pathogen (Trapp and Croteau
87 2001, Cudmore et al. 2010). The increasing frequency and severity of such outbreaks have spurred
88 intensive molecular and biochemical research into the factors underlying host susceptibility and
89 pest/pathogen defense in spruce (*Picea*; Bohlmann 2008), fir (*Abies*; Hain et al. 1991, Lewinsohn et al.
90 1993a), and pine (*Pinus*; Sampedro et al. 2011) species. Defense induction in conifers by mechanical
91 wounding (Lewinsohn et al. 1993a), experimental insect attack (Miller et al. 2005, Sampedro et al. 2011) or
'simulated' herbivory by application of chemical elicitors such as methyl jasmonate (Martin et al. 2002,

2003, Sampedro et al. 2010) leads to dramatic increases in bark and stem-wood terpenoid accumulation and volatile release from needles. An increasing number of the active genes and biosynthetic enzymes underlying defensive chemical outputs in these conifer systems have been identified, establishing strong evidence that resin-based—and primarily terpenoid-based—chemical defenses are central to the trees' evolved responses to insect or pathogen colonization (Franceschi et al. 2005, Keeling and Bohlmann 2006).

In eastern North America, the invasive twig-feeding hemlock woolly adelgid (*Adelges tsugae*; 'HWA') threatens to extirpate the native eastern hemlock (*Tsuga canadensis* Carr.; McClure and Cheah 1999). The first documented population of the adelgid in eastern North America was detected in the early 1950s, and appears to be of Japanese origin (Havill et al. 2006). The insect has now spread to the southern extent of eastern hemlock's range in northern Georgia, and is moving northward into Vermont, New Hampshire, and Maine (Preisser et al. 2008, Forest Health Protection Program 2011). The insect can take a year or two to reach high densities, but its effect on hemlocks is needle desiccation, branch mortality, and marked suppression of new spring growth, often leading to tree death in four years or less (McClure 1991a). As the only native shade-tolerant conifer in the eastern United States, eastern hemlock acts as a foundation species (sensu Ellison et al. 2005) that creates cool and moist microclimates in the midst of deciduous forests. The nearly complete removal of mature and seedling eastern hemlocks following HWA infestation (Preisser et al. 2011) substantially increases soil and stream temperatures, alters soil chemistry and nutrient cycling patterns, and favors fast-growing, early-successional trees—a series of changes that dramatically transforms the forest landscape (Orwig et al. 2008, Gandhi and Herms 2010). The elongate hemlock scale (*Fiorinia externa*; 'EHS') is another exotic pest of eastern hemlock; an armored scale introduced to the Northeastern United States in the early 20th century, this insect is also now present in much of the tree's range and continues to spread northward (Preisser et al. 2008). Reports seemingly based on observational, rather than experimental, evidence suggest that although EHS is usually not lethal, high densities can cause significant needle loss and contribute to the mortality of already stressed trees (McClure 1980, Abell and Van Driesche 2012).

Despite the existence of several studies documenting a correlation between terpenoid levels and herbivory in eastern hemlock, there has been no direct investigation into whether either of these exotic pests elicits resin defenses in eastern hemlock. One study reported a positive correlation between volatile

120 terpenoid levels and the fecundity of both EHS and a second armored scale pest of eastern hemlock
121 (McClure and Hare 1984). Lagalante and Montgomery (2003) compared the constitutive volatile terpenoid
122 profiles of HWA resistant and susceptible *Tsuga* species and suggested that several volatile terpenoids may
123 act as deterrents or attractants ('feedants') to HWA. In a follow up study focused on eastern hemlock,
124 Lagalante et al. (2006) measured spatial and temporal variability in resin volatiles and hypothesized that
125 these phytochemical fluctuations drive the HWA's unusual annual patterns of settlement, aestivation, and
126 feeding. European silver fir (*Abies alba*), a conifer of a genus related to *Tsuga*, showed increased levels of
127 monoterpenoid accumulation in bark naturally infested with *Adelges piceae*, the balsam woolly adelgid
128 (Hain et al. 1991). In addition, western hemlock (*Tsuga heterophylla*) responded to simulated herbivory
129 (treatment with methyl jasmonate) in a manner typical of the conifers of Pinaceae: traumatic resin ducts
130 formed and terpenoid concentrations increased (Hudgins et al. 2004). This evidence suggests resinosis may
131 also occur in species of *Tsuga*. However, despite the prevalence of research into herbivore-defense
132 responses of other conifers of Pinaceae, little is known about the inducible resin defenses of hemlocks.

133 There is growing evidence that HWA infestation induces changes in eastern hemlock chemistry
134 and physiology. Evidence of a localized and systemic hypersensitive response (a common plant defense
135 against pathogens and sessile herbivores leading to tissue necrosis at the infected site; Radville et al. 2011),
136 substantially higher foliar free amino acid concentrations (Gomez 2012), changes in woody plant anatomy
137 (Gonda-King et al. 2012) and a reduction of both new growth and percent total foliar nitrogen (Miller-
138 Pierce et al. 2010) have been reported in response to HWA feeding on eastern hemlocks. EHS, on the other
139 hand, appeared to produce only a localized hypersensitive response and did not significantly affect free
140 amino acid concentration, percent total foliar nitrogen, woody plant anatomy, or subsequent new growth.

141 We investigated whether HWA or EHS infestation induced oleoresin production in eastern
142 hemlock, an ecologically unique native U.S. conifer in rapid decline in many areas. Previous research has
143 suggested spatial and temporal fluctuations in volatile resin compounds can influence the establishment of
144 colonizing hemipteran herbivores (McClure and Hare 1984, Lagalante et al. 2006). To test this, we
145 measured levels of resin volatiles in both twigs and needles of eastern hemlocks experimentally infested
146 with HWA, EHS or neither insect in early summer and again in mid-autumn, each time following periods
147 of active feeding by both insects. We predicted that both insects would elicit changes in the concentrations

148 or composition of volatile resin compounds. We hypothesized that an agent as rapidly lethal as HWA
149 would elicit a defensive resinosis typical of many conifers of Pinaceae: pronounced increases in toxic or
150 deterrent phytochemicals, especially terpenoids. We also predicted that the much milder effects of EHS on
151 the host tree's physiology (Miller-Pierce et al. 2010, Radville et al. 2011, Gonda-king et al. 2012, Gomez et
152 al. 2012) would be accompanied by an induced resin response distinct from that of HWA.

153

154

METHODS AND MATERIALS

155 *Study System.* Eastern hemlock buds begin opening in May in the Northeastern United States, and the
156 young new growth shoots, at first green and pliant, complete their elongation at approximately the end of
157 summer. By that time the foliage has hardened and taken on the form and appearance of the fully mature,
158 previous year growth.

159 The HWA completes two clonal generations per year in eastern North America, as in its range of
160 origin in Japan (McClure and Cheah 1999). In the Northeastern United States, first instar nymphal crawlers
161 of the progredien generation settle in April (before bud-break) on already mature previous year's growth
162 just below the needle abscission layer and feed through a stylet bundle on xylem ray parenchyma cells in
163 the twig (Young et al. 1995). The sexuparae, a winged, sexually reproducing generation of HWA, hatch
164 concurrently with the clonal female progrediens and, in Japan, subsequently disperse to a spruce (*Picea*)
165 primary host to complete reproduction. Sexuparae in North America are unable to complete their life-cycle
166 due to the absence of a suitable spruce host and thus, only asexual reproduction occurs. The sessile
167 progredien adults complete egg laying in June, at which point the crawlers of the sisten generation emerge,
168 settle preferentially on the new, young current year's growth, and promptly enter aestivation. In early fall,
169 by the time the new growth has matured, the sisten nymphs resume feeding, completing development and
170 oviposition in April (McClure and Cheah 1999).

171 The EHS completes two full generations per year in its natural range in Japan, but in the
172 Northeastern United States it appears to lack a distinct and regular cycle of life stage development, and
173 completes between one and two generations annually (Abell and Driesche 2012). First instar nymphs begin
174 to hatch in early June and settle preferentially on the undersides of young hemlock needles. EHS is also a
175 sessile stylet feeder, inserting a thread-like stylet bundle and sucking fluid from needle mesophyll cells

176 (McClure 1980). Since generation times in the Northeast are irregular, life stages appear to overlap and
177 often two or more instars may be found developing concurrently on the same foliage (Abell and Driesche
178 2012).

179

180 *Experimental Design.* In April 2007, eastern hemlock saplings (0.7-1.0m) were removed from Cadwell
181 Experimental Forest (Pelham, MA, USA) and planted in an open field setting (East Farm, Kingston, RI,
182 USA) in a rectangular grid. The source forest was free of both HWA and EHS at the time of collection,
183 and careful inspection of the sapling trees revealed no prior infestation by either insect. Artificial
184 infestation with HWA, EHS, or neither insect was randomly applied to the saplings. Because both insects
185 are wind-dispersed during their first-instar crawler phase, each tree (including all uninfested controls) was
186 enclosed in a mesh cage annually from early spring to late fall to prevent cross-contamination. Each of the
187 1 m x 1 m x 2 m (length by width by height) cages consisted of a plastic PVC pipe frame covered by
188 mosquito netting (97 holes/cm² mesh size). Weed-inhibiting fabric (1 m²) was placed around the base of
189 each tree. By 2010, a combination of insect cross-contamination and tree death from transplantation-related
190 stress reduced the level of replication to nine trees in the HWA treatment, seven trees in the EHS treatment,
191 and eight trees in the control treatment.

192

193 *Insect Inoculations.* Insect inoculations were conducted following standard procedures (see Butin et al.
194 2007). Briefly, trees were inoculated with insects each spring from 2007 to 2010 to mimic natural
195 infestation cycles. Immediately prior to crawler emergence (May for HWA, June for EHS), naturally-
196 infested branches with comparable insect densities were collected from sites in southern New England and
197 attached to trees in the appropriate treatment group; control trees received uninfested branches. Individual
198 branches were placed in aquapics to slow needle desiccation and decrease insect mortality.

199

200 *Plant Material.* Plant tissue samples were collected from each tree in late June 2010 (fully mature, previous
201 year foliage segments) after the first-instar crawlers of both insects had settled and commenced feeding,
202 and again in mid-October 2010 (young, current year growth twigs) after settled HWA had ceased
203 aestivation and resumed feeding. An average of 10 cm of twig with foliage was clipped; in the case of the

204 insect treatments, infested foliage samples were selected. Each sample was placed in a polypropylene
205 cryovial, flash-frozen in liquid nitrogen, transported to the laboratory on dry ice and stored at -80°C until
206 extraction and analysis.

207

208 *Extraction of Resin Volatiles.* Extraction of resin volatiles was modified from a protocol developed by
209 Lewinsohn et al. (1993b). All reagents and reference standards were obtained from Sigma-Aldrich (St.
210 Louis, MO, USA). Solvents were HPLC or GC grade purity.

211 Needles were separated from twigs and ground to a homogenous powder using a mortar and pestle
212 under liquid nitrogen. Approximately 100-200 mg (dry weight) of needle tissue was combined with methyl
213 *tert*-butyl ether (MTBE; 1.3-1.5 mL) containing a known concentration of isobutylbenzene ($40\ \mu\text{g mL}^{-1}$) as
214 an internal standard in a pre-weighed 2 mL vial (glass with PTFE-coated screw cap, Sigma-Aldrich, St.
215 Louis, MO, USA). Needle samples were extracted overnight (20 h) with constant shaking at room
216 temperature. Each extract was transferred to a fresh glass vial and washed with aqueous $(\text{NH}_4)_2\text{CO}_3$ (0.3
217 mL, 1 M) to neutralize acidic impurities. The organic layer was then filtered through a Pasteur pipette
218 column packed with silica gel (0.3 g, Sigma-Aldrich, 60\AA) overlaid with MgSO_4 (0.2 g). Oxygenated
219 volatile compounds were subsequently eluted by washing the filter with diethyl ether (1 mL), and
220 combined eluates were collected in a GC vial (PTFE-coated screw cap, Agilent Technologies, Santa Clara,
221 CA, USA) and stored at -20°C until analysis.

222 Twig samples of approximately 10-50 mg (dry weight) were cooled with liquid nitrogen in a
223 mortar and pestle, ground to a coarse powder, and combined with MTBE (1.0 mL) containing
224 isobutylbenzene ($2\ \mu\text{g mL}^{-1}$) in a 2 mL glass vial. Twigs were extracted overnight (19 h) with constant
225 shaking at room temperature. Aqueous $(\text{NH}_4)_2\text{CO}_3$ (0.2 mL; 1 M) was added to each extract, followed by
226 thorough mixing. The organic layer was then transferred directly to a Pasteur pipette filter packed with
227 silica gel (0.2 g, 60\AA) overlaid with MgSO_4 (0.13 g). The filter was washed with diethyl ether (0.5 mL),
228 and combined eluates were collected and stored as described above. After extraction, each sample was
229 dried for at least 48 hours at $55\text{-}60^{\circ}\text{C}$ and weighed for the determination of tissue dry weight.

230

231 *Analysis of Resin Volatiles.* Needle volatile extracts were analyzed on a Hewlett-Packard (HP) 6890 GC
232 equipped with a flame ionization detector (FID). For all analyses, the injection volume was 1 μL , injector
233 temperature 220°C. Volatile compounds were separated on an Agilent DB-5, 0.25 mm i.d. x 30 m, 0.25 μm
234 coating thickness, fused silica capillary column. H_2 carrier gas flow was a constant 1.0 mL min^{-1} and the
235 split ratio was 20:1. The FID was heated to 250°C, with H_2 flow at 40 mL min^{-1} , air flow 350 mL min^{-1} ,
236 and constant make-up flow (N_2) at 45 mL min^{-1} . The GC oven was programmed with an initial temperature
237 of 60°C (no hold), an increase at 3°C min^{-1} to 156°C, then 50°C min^{-1} to 300°C (hold 3 min). GC-FID
238 generated peaks were integrated using HP ChemStation software (Agilent technologies). Datafiles for five
239 of the October needle samples were corrupted, reducing the level of replication to seven trees in the HWA
240 treatment, six trees in the EHS treatment, and six trees in the control treatment.

241 For all compound identifications, as well as all twig volatile quantification, analyses were
242 performed on a Shimadzu GC-2010 system equipped with a QP2010-Plus mass spectrometer (EI mode, 70
243 eV), running GCMSolution software (Shimadzu Corporation, Kyoto, Japan). Separations were performed
244 on the same column as described above for GC-FID. The injection volume was 1 μL and injector
245 temperature 220°C. Helium carrier gas flow was in constant linear velocity mode at 36.5 cm sec^{-1} , with
246 column flow set at 1.0 mL min^{-1} and a split ratio of 5:1. The GC oven was programmed with an initial
247 temperature of 60°C (no hold), an increase at 3°C min^{-1} to 175°C, then 30°C min^{-1} to 300°C (hold 5 min).
248 The interface and ion source temperatures were both set at 300°C, and the MS scan range was m/z 40-400.

249 Identification of each volatile compound was, wherever possible, based on comparison of the
250 experimental retention time and mass spectrum with those of an authentic standard (indicated in Table 1);
251 when a pure standard was unavailable, tentative identification was based on comparison with retention
252 index and mass spectral information reported in the literature (Adams 2001) and with mass spectra in the
253 NIST05 and NIST05s mass spectral libraries (Stein 2005). Concentrations of all compounds were
254 determined by normalizing integrated peak areas against that of the internal standard isobutylbenzene in
255 each chromatogram. Each tissue volatile concentration value was standardized to ' $\mu\text{g g}^{-1}$ dry weight' by
256 dividing by the sample dried weight.

257 Since both the HWA and the EHS are quite small and adhere tightly to their twig or needle feeding
258 sites, complete removal of insects and their ovisacs from infested samples prior to analysis was not

259 practical. To test whether detected volatiles could potentially be of insect, rather than hemlock, origin, we
260 obtained several samples of HWA-infested foliage of comparable size and insect density to our
261 experimental samples, collected the insects, eggs, and the wax of ovisacs into vials, and extracted and
262 analyzed the insect material using the plant-volatile protocol described herein.

263

264 *Statistical Analysis.* Resin volatile concentrations were log transformed prior to statistical analysis to
265 reduce heterogeneity of variance. Two-way mixed-model ANOVAs (Proc Mixed, SAS 9.3; SAS Institute
266 2011) were used to test twigs and needles separately for treatment-level differences in the concentration of
267 individual volatiles, total monoterpenoids, total sesquiterpenoids, total green leaf volatiles (needles only)
268 and total combined benzenoids (including phenolics; twigs only) with month (June vs. October) and
269 treatment (HWA, EHS and control) as fixed factors and tree as a random factor. Mixed-model analyses
270 were appropriate because we sampled from the same trees in both months. We also used ANOVA planned
271 contrasts to separately test HWA and EHS treatment means against the control mean, using treatment as the
272 fixed factor (R 2.14.0; R Development Core Team 2012).

273 Familywise error rate for the mixed model analyses was evaluated using a false discovery rate
274 (FDR) estimation method ('fdrtool' software package; R 2.14.0; Strimmer 2008). FDR techniques are now
275 used widely with multiple simultaneous hypothesis testing to estimate the proportion of tests with
276 incorrectly rejected null hypotheses among tests with statistically significant findings. This is in contrast to
277 traditional familywise error rate correction methods (e.g. the sequential Bonferroni) that estimate the
278 probability of a false rejection among all tests conducted and, arguably, unnecessarily sacrifice statistical
279 power.

280 As an additional measure of the overall strength of evidence for our mixed-model hypothesis test
281 findings, we used the following binomial equation (sensu Moran 2003) to calculate the overall probability
282 of obtaining K tests with P -values smaller than our specified α -level:

$$283 \quad P_B = [N!/(N-K)!K!] \times \alpha^K (1-\alpha)^{N-K}$$

284 where N = number of tests. This procedure allowed us to estimate the probability that so many statistically
285 significant treatment effects could arise by chance (i.e. could be 'false positives').

286

RESULTS

287
288 The overall effects of infestation with HWA and EHS were very similar in both June 2010 (mature
289 previous year's growth) and October 2010 (young current year's growth) samples. Both insects produced a
290 notable trend of decreases of most individual twig volatiles, though only a modest fraction of these
291 decreases carried statistical significance; the same trend of largely non-significant decreases was observed
292 for total twig volatile levels. Conversely, both insects produced a largely non-significant but also notable
293 trend of increases of most needle volatiles (Table 1), with the same non-significant trend of increases for all
294 measures of total needle volatiles. Across all treatments, the total twig or needle monoterpenoid levels were
295 25-40% higher in the young current year's growth than in the mature previous year's growth (Table 2A, B).
296 Total needle sesquiterpenoids followed a similar pattern, while current year's growth twigs had volatile
297 concentrations 65-70% higher than previous year's growth.

298 In twig tissue, 16 monoterpenoids, five sesquiterpenoids, and six benzenoid or phenolic
299 compounds were present in quantities sufficient for identification and quantification (Fig 1A); in needle
300 tissue, the corresponding numbers were 18 monoterpenoids, five sesquiterpenoids, one benzenoid, and
301 three fatty acid derivatives (i.e. green leaf volatiles or 'GLVs'; Fig 1B). Qualitatively, needle and twig
302 volatile profiles were overlapping but different (Table 1). Monoterpenoids dominated in terms of both
303 diversity and mass contribution, and had the greatest effect on the induced changes of total volatiles.
304 Sesquiterpenoids, present at somewhat lower abundance, generally increased in both twigs and needles.
305 GLVs were detected only in needle tissue, and were consistently increased by both insects, especially by
306 EHS—the total amount of these compounds had nearly doubled from June to October; Fig. 2.

307 The effects of insect feeding on volatile concentration were larger in twigs than in needles (Table
308 1, Online Resource 2). In twigs, the results of mixed-model ANOVAs (Online Resource 2A) show that
309 HWA feeding significantly ($P < 0.05$) or marginally significantly ($0.05 < P < 0.10$) decreased five of 17
310 individual monoterpenoids and two unidentified volatiles; additionally, the dramatic increases in the
311 benzenoid benzyl alcohol (more than 30-fold in June and about 10-fold in October; Fig. 3) and the
312 monophenolic phytohormone methyl salicylate ('MeSA'; two orders of magnitude in June and more than
313 10-fold in October; Fig. 4) were both significant. EHS feeding decreased two monoterpenoids significantly.
314 Both insects decreased the monophenolic raspberry ketone and several other monoterpenoids with marginal

315 significance. HWA feeding significantly decreased total monoterpenoids, while EHS feeding decreased
316 both total monoterpenoids and total volatiles with marginal significance. HWA feeding marginally
317 increased total benzenoids, while EHS feeding marginally decreased these compounds (Online Resource
318 2A).

319 In needles, (Online Resource 2B) EHS feeding increased *cis*-3-hexenal and total GLVs
320 significantly, and increased *trans*-2-hexenal and the benzenoid p-cymene with marginal significance.
321 There were no significant effects of HWA feeding on needle volatile levels.

322 Results of planned contrast ANOVA comparisons of average control versus treatment volatile
323 concentrations were quite similar to those we obtained using the mixed model analyses. Significance
324 values from these simpler analyses are indicated in Table 1.

325 The binomial probability that the 60 twig volatile mixed model tests we ran would generate *P*-
326 values smaller than the ones we observed was $P_B=0.00014$ if calculated at the $\alpha=0.05$ level, or 4.7×10^{-9} if
327 calculated at the $\alpha=0.10$ level. For the 54 needle volatile tests, the overall probability of no ‘real’ effect
328 was greater: $P_B=0.18$, and 0.12, respectively.

329 Estimated false discovery rate for twig volatile hypothesis tests is reported as *q*-value alongside
330 each test’s nominal *P*-value (Online Resource 2A). The *q*-value is the minimum FDR level that would be
331 needed to reject that hypothesis. Selection of an appropriate FDR level, in turn, depends on the proportion
332 of false rejections considered tolerable. We did not report FDR for needle volatile hypothesis tests (Online
333 Resource 2B). Since the method estimates the proportion of false rejections among only tests with
334 significant findings—and there was just one out of 54 needle volatile hypothesis tests that was statistically
335 significant—in that case an estimate of FDR was superfluous.

336

337

DISCUSSION

338 We found evidence of an induced response in eastern hemlock during infestation by both HWA and EHS ,
339 encompassing a number of feeding-elicited changes in the tree’s resin volatile profile. However, the
340 modest induction (mostly decreases) of resin metabolites in twig tissue and the non-significant trend of
341 modest increases in needle tissue produced by both insects, was conspicuously different from the profuse
342 resinosis observed in insect-infested pines, spruces, and firs (Trapp and Croteau 2001). In light of the

343 considerable evidence that HWA induces more extensive changes in eastern hemlock physiology than does
344 EHS (Miller-Pierce et al. 2010, Radville et al. 2011, Gonda-King et al. 2012, Gomez 2012), the observation
345 that HWA and EHS produced similar overall changes in the tree's volatiles was intriguing and ran counter
346 to our predictions.

347 In contrast to the modest changes in terpenoid levels, a number of the non-terpenoids were sharply
348 increased by HWA feeding, in what may reflect a hemlock defense response (Table 1). Benzyl alcohol was
349 induced in HWA-infested trees; this compound is a common plant volatile (Dudareva et al. 2008)
350 previously detected in the stem-wood of mountain hemlock (*T. mertensiana*; Shepherd et al. 2008) and in
351 volatiles released from mite-infested spruce foliage (Kannaste 2008; Fig. 3). In screening studies, benzyl
352 alcohol deterred feeding by the greenbug aphid *Schizaphis graminum*, reducing fecundity and causing
353 substantial mortality (Formusoh et al. 1997). MeSA, which was also induced by HWA (Fig. 4), has been
354 found in the volatile mix released after aphid feeding and identified as a deterrent to aphid settling and
355 fecundity in a number of plant-insect systems (Hardie et al. 1994, Quiroz et al. 1998).

356 The sharp increase of these two compounds in HWA-infested trees (Table 1) is notable in light of
357 the growing body of evidence that some plants respond to piercing-sucking hemiptera by activating
358 biosynthetic pathways similar or identical to those used in pathogen defense (Kaloshian and Walling 2005).
359 Benzyl alcohol is a strong antimicrobial agent against diverse microorganisms (Shenep et al. 2011), while
360 MeSA, the volatile methyl ester of salicylic acid (SA), activates a SA-dependent biosynthetic cascade in
361 numerous plants that leads to systemic acquired resistance (SAR) against pathogen infection (Durrant and
362 Dong 2004). For aphids, close relatives of adelgids, feeding has been shown in many studies to activate the
363 SA-dependent biosynthetic pathways normally associated with pathogen defense (Moran and Thompson
364 2001, Martinez de Ilarduya et al. 2003, Zhu-Salzman et al. 2004) or to induce pathogen-resistance outright
365 in their host plant (Russo et al. 1997). The elevated levels of these two compounds in HWA-induced
366 hemlock tissue is a sign that a SA-driven insect defense syndrome may be active in HWA-infested trees.

367 It is also possible that increased production of these volatiles reflects the tree's detection of a
368 microbial associate of HWA rather than of the insect itself. An endosymbiont was recently found
369 throughout the body of the HWA and appears essential to the insect's survival (Shields and Hirth 2005). It

370 is possible that the hemlocks may be responding to this bacterium, if it is introduced into the vascular tissue
371 of eastern hemlock during HWA feeding, by mobilizing a pathogen defense response.

372 Our results may help elucidate why HWA causes more extensive damage to eastern hemlock than
373 EHS. Radville et al. (2011) detected evidence of a local hypersensitive response (elevated hydrogen
374 peroxide levels) in both EHS- and HWA-infested trees, and showed that this hypersensitive response
375 occurs systemically in response to HWA-infestation. The hypersensitive response usually precedes the
376 development of SAR (Durrant and Dong 2004, Kaloshian and Walling 2005). Research on tobacco has
377 revealed that in pathogen-infected plant tissue SA is enzymatically converted to the volatile MeSA, which
378 acts as a mobile agent that is taken up by receptors on distant, uninfected tissue. There, the MeSA is
379 demethylated and transformed back to SA, which in turn activates an induced resistance response to the
380 invading organism (i.e. SAR; Shulaev et al. 1997, Park et al. 2007). Our discovery that MeSA levels were
381 elevated in only the HWA-infested trees suggests this compound could be a mobile signal that propagates
382 the ‘pathogen-like’ effects of the adelgid on uninfested foliage, extending the insect’s effects and
383 intensifying the overall damage to the tree. The observation that HWA elicited such a response, but EHS
384 did not, may reflect the species-specific nature of the hemlock defense elicitors carried in the insects’
385 salivary secretions, as has been observed in at least one other hemipteran-plant interaction (Ven et al.
386 2000).

387 The HWA-driven increases we observed in levels of benzyl alcohol and MeSA may also help
388 explain previously noted changes in the primary chemistry of the hemlock saplings of the present study
389 (Gomez et al. 2012). Although much of the biosynthetic pathway for the benzenoids has yet to be
390 determined, radio-labeling experiments show they are derived from L-phenylalanine (Dudareva et al.
391 2006). As with benzyl alcohol and MeSA, a marked increase in L-phenylalanine and many other free
392 amino acids occurred in trees infested with HWA, but not EHS (Gomez et al. 2012). Thus the increased
393 amino acid levels in HWA-infested trees may constitute an adaptive mobilization of precursors of defense-
394 related volatile compounds.

395 Alternatively, adelgid manipulation of host-plant biochemistry could explain a number of the
396 insect-induced changes in resin chemistry we have shown. HWA, like many adelgids, forms extensive galls
397 on the buds of its primary spruce host in its original range in Asia (Havill and Footitt 2007). Gallling insects

398 are known to be adept at manipulating host plant physiology to create a more nutritious and less defended
399 environment (Tooker and Moraes 2009). We have demonstrated a substantial decrease in monoterpenoids,
400 often compounds of direct defense against herbivory (Eyles et al. 2010, Schiestl 2010), in the tissue where
401 the adelgid feed. Our results also show a less pronounced elicitation of GLVs (typical wounding response
402 volatiles; Fig. 2; Shiojiri et al. 2006) in response to the feeding of HWA, relative to EHS, despite the
403 adelgid's much greater impacts on tree physiology (Miller Pierce et al. 2010, Radville et al. 2011, Gonda-
404 King et al. 2012). These observations, considered together with the noted increase in free amino acids only
405 in HWA-infested trees (Gomez et al. 2012), may constitute evidence that the host-manipulating capacity
406 conserved in adelgid biology may be an underlying mechanism in this system.

407 Lagalante et al. (2006) suggested that the lack of a co-evolutionary history between eastern
408 hemlock and sessile piercing-sucking insects resulted in the absence of biosynthetic pathways with which
409 eastern hemlock can defend against insects like HWA and EHS. This hypothesis is consistent with the
410 finding of little or no output of anatomical or chemical resin defenses. However, we did observe a resin
411 chemical response to HWA, and to the co-occurring EHS, though perhaps of a subtler nature than that often
412 seen in other conifers. It is possible that the resistance traits HWA elicits in eastern hemlock are simply not
413 well matched to the actual challenge of this introduced insect and do not confer resistance. A comparison of
414 the induced response of susceptible eastern hemlocks to those of HWA-resistant *Tsuga* species and strains
415 of eastern hemlock believed resistant to HWA, as well as conifers with putative resistance to EHS
416 (McClure and Fergione 1977), will test these hypotheses. Nonetheless, our findings establish that HWA
417 and EHS both induce changes in the resin chemistry of eastern hemlock, and constitute the first critical step
418 toward understanding the role inducible chemical defenses play in determining hemlock susceptibility to
419 these exotic hemipteran pests.

420

421

ACKNOWLEDGMENTS

422 This work was supported by USDA Forest Service cooperative agreements #103-CA-11244225- 130 and
423 #09-CA-11420004-360 to JSE, and NSF grant DEB-0715504 and NIFA grant #2011-67013-30142 to ELP.
424 We also gratefully acknowledge the conceptual and technical help of R. D'Alonzo, S. Eyles, J. Gershenson,

425 A. Kessler, C. Orians, R. Ragusso, and S. Teale; statistical guidance from J. Finn, K. McGarigal, P. Sievert,
426 and A. Whiteley; and assistance with fieldwork given by D. Cox, L. Glidden, C. Huffman, and L. Radville.

427

428

REFERENCES

429 ABELL, K. and DRIESCHE, R.V. 2012. Impact of latitude on synchrony of a scale (*Fiorinia externa*) (Hemiptera:
430 Diaspididae) and its parasitoid (*Encarsia citrina*) (Hymenoptera: Aphelinidae) in the Eastern United States.
431 *Biolog. Control* 63:339-347.

432 ADAMS, R.P. 2001. Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass
433 Spectroscopy. Allured Press, Carol Stream, Illinois.

434 BOHLMANN, J. 2008. Insect-Induced Terpenoid Defenses in Spruce, pp. 173-187, in: A. Schaller (ed.). Induced
435 Plant Resistance to Herbivory. Springer Press, Stuttgart, Germany.

436 BUTIN, E., PREISSER, E., and ELKINTON, J. 2007. Factors affecting settlement rate of the hemlock woolly
437 adelgid, *Adelges tsugae*, on eastern hemlock, *Tsuga canadensis*. *Agr. Forest Entomol.* 9:215-219.

438 CUDMORE, T.J., BJORKLUND, N., CARROLL, A.L., and LINDGREN, B.S. 2010. Climate change and range
439 expansion of an aggressive bark beetle: evidence of higher beetle reproduction in naive host tree
440 populations. *J. Appl. Ecol.* 47:1036-1043.

441 DUDAREVA, N., NEGRE, F., NAGEGOWDA, D.A., and ORLOVA, I. 2006. Plant Volatiles: Recent Advances
442 and Future Perspectives. *Crit. Rev. Plant Sci.* 25:417-440.

443 DURRANT, W. and DONG, X. 2004. Systemic Acquired Resistance. *Annu. Rev. Phytopathol.* 42:185-209.

444 ELLISON, A.M., BANK, M.S., CLINTON, B.D., COLBURN, E.A., ELLIOTT, K., FORD, C.R., FOSTER, D.R.,
445 KLOEPPPEL, B.D., KNOEPP, J.D., and LOVETT, G.M. 2005. Loss of foundation species: consequences
446 for the structure and dynamics of forested ecosystems. *Front. Ecol. Environ.* 3:479-486.

447 EYLES, A., BONELLO, P., GANLEY, R., and MOHAMMED, C. 2010. Induced resistance to pests and pathogens
448 in trees. *New Phytol.* 185:893-908.

449 FOREST HEALTH PROTECTION PROGRAM. 2011. Counties with established HWA populations 2010. USDA
450 Forest Service, Morgantown, WV.

451 FORMUSOH, E.S., REESE, J.C., and BRADFISCH, G. 1997. A miniaturized bioassay system for screening
452 compounds deleterious to greenbugs (Homoptera: Aphididae) on artificial diets. *J. Kansas Entomol. Soc.*

- 453 70:323-328.
- 454 FRANCESCHI, V.R., KROKENE, P., CHRISTIANSEN, E., and KREKLING, T. 2005. Anatomical and chemical
455 defenses of conifer bark against bark beetles and other pests. *New Phytol.* 167:353-376.
- 456 GANDHI, K.J.K. and HERMS, D.A. 2010. Direct and indirect effects of alien insect herbivores on ecological
457 processes and interactions in forests of eastern North America. *Biol. Invasions* 12:389-405.
- 458 GOMEZ, S., ORIAN, C.M., and PREISSER, E.L. 2012. Exotic herbivores on a shared host: foliar quality after
459 individual, simultaneous, and sequential attack. *Oecologia* 169:1015-1024.
- 460 GONDA-KING, L., RADVILLE, L., and PREISSER, E.L. 2012. False ring formation in eastern hemlock branches:
461 impacts of hemlock woolly adelgid and elongate hemlock scale. *Environ. Entomol.* 41:523-531.
- 462 HAIN, F.P., HOLLINGSWORTH, R.G., ARTHUR, F.H., SANCHEZ, F., and ROSS, R.K. 1991. Adelgid host
463 interactions with special reference to the balsam woolly adelgid in North America, pp. 271-287, in: Y.N.
464 Baranchikov, W.J. Mattson, F.P. Hain and T.L. Payne (eds.). *Forest Insect Guilds: Patterns of Interaction*
465 *with Host Trees*. USDA Forest Service General Technical Report NE-153.
- 466 HARDIE, J., ISAACS, R., PICKETT, J., WADHAMS, L., and WOODCOCK, C. 1994. Methyl salicylate and (-)-
467 (1R,5S)-myrtenal are plant-derived repellents for black bean aphid, *Aphis fabae* Scop. (Homoptera:
468 Aphididae). *J. Chem. Ecol.* 20:2847-2855.
- 469 HAVILL, N.P. and FOOTITT, R.G. 2007. Biology and evolution of Adelgidae. *Annu. Rev. Entomol.* 52:325-49.
- 470 HAVILL, N.P., MONTGOMERY, M.E., YU, G., SHIYAKE, S., and CACCONE, A. 2006. Mitochondrial DNA
471 from hemlock woolly adelgid (Hemiptera: Adelgidae) suggests cryptic speciation and pinpoints the source
472 of the introductions to eastern North America. *Ann. Entomol. Soc. Am.* 99:195-203.
- 473 HUDGINS, J.W., CHRISTIANSEN, E., and FRANCESCHI, V.R. 2004. Induction of anatomically based defense
474 responses in stems of diverse conifers by methyl jasmonate: a phylogenetic perspective. *Tree Phys.* 24:251-
475 264.
- 476 KALOSHIAN, I. and WALLING, L.L. 2005. Hemipterans as plant pathogens. *Annu. Rev. Phytopathol.* 43:491-521.
- 477 KANNASTE, A. 2008. Volatiles of Conifer Seedlings: Composition and Resistance Markers. PhD dissertation.
478 KTH School of Chemical Science and Engineering, Stockholm, Sweden.
- 479 KEELING, C.I. and BOHLMANN, J. 2006. Genes, enzymes and chemicals of terpenoid diversity in the constitutive
480 and induced defence of conifers against insects and pathogens. *New Phytol.* 170:657-675.

- 481 KOEPKE, D., BEYAERT, I., GERSHENZON, J., HILKER, M., and SCHMIDT, A. 2010. Species-specific
482 responses of pine sesquiterpene synthases to sawfly oviposition. *Phytochemistry* 71:909-917.
- 483 LAGALANTE, A.F., LEWIS, N., MONTGOMERY, M.E., and SHIELDS, K.S. 2006. Temporal and spatial
484 variation of terpenoids in eastern hemlock (*Tsuga canadensis*) in relation to feeding by *Adelges tsugae*. *J.*
485 *Chem. Ecol.* 32:2389-2403.
- 486 LAGALANTE, A.F. and MONTGOMERY, M.E. 2003. Analysis of terpenoids from hemlock (*Tsuga*) species by
487 solid-phase microextraction/gas chromatography/ion-trap mass spectrometry. *J. Agr. Food Chem.* 51:2115-
488 2120.
- 489 LANGENHEIM, J.H. 1994. Higher plant terpenoids: a phyto-centric overview of their ecological roles. *J. Chem.*
490 *Ecol.* 20:1224-1280.
- 491 LEWINSOHN, E., GIJZEN, M., MUZIKA, R.M., BARTON, K., and CROTEAU, R. 1993. Oleoresinosis in grand
492 fir (*Abies grandis*) saplings and mature trees. *Plant Physiol.* 101:1021-1028.
- 493 LEWINSOHN, E., THOMAS, J., GIJZEN, M., and CROTEAU, R. 1993. Simultaneous analysis of monoterpenes
494 and diterpenoids of conifer oleoresin. *Phytochem. Analysis* 4:220-225.
- 495 MARTIN, D., THOLL, D., GERSHENZON, J., and BOHLMANN, J. 2002. Methyl jasmonate induces traumatic
496 resin ducts, terpenoid resin biosynthesis, and terpenoid accumulation in developing xylem of Norway
497 spruce stems. *Plant Physiol.* 129:1003-18.
- 498 MARTIN, D.M. 2003. Induction of volatile terpene biosynthesis and diurnal emission by methyl jasmonate in
499 foliage of Norway spruce. *Plant Physiol.* 132:1586-1599.
- 500 MARTINEZ, D.E. ILARDUYA, O.M., XIE, Q., and KALOSHIAN, I. 2003. Aphid-induced defense responses in
501 *Mi-1* mediated compatible and incompatible tomato interactions. *Mol. Plant-Microbe In.* 16:699-708.
- 502 MCCLURE, M.S. 1980. Competition between exotic species: scale insects on hemlock. *Ecology* 61:1391-1401.
- 503 MCCLURE, M. 1991. Density-dependent feedback and population cycles in *Adelges tsugae* (Homoptera:
504 Adelgidae) on *Tsuga canadensis*. *Environ. Entomol.* 20:258-264.
- 505 MCCLURE, M.S. and CHEAH, C.A.S. 1999. Reshaping the ecology of invading populations of hemlock woolly
506 adelgid, *Adelges tsugae* (Homoptera: Adelgidae), in eastern North America. *Biol. Invasions* 1:247-254.
- 507 MCCLURE, M.S. and FERGIONE, M.B. 1977. *Fiorinia externa* and *Tsugaspidotus tsugae* (Homoptera:
508 Diaspididae): distribution, abundance, and new hosts of two destructive scale insects of eastern hemlock in

- 509 Connecticut. *Environ. Entomol.* 6:807-811.
- 510 MCCLURE, M.S. and HARE, J.D. 1984. Foliar terpenoids in *Tsuga* species and the fecundity of scale insects.
511 *Oecologia* 63:185-193.
- 512 MILLER, B., MADILAO, L.L., RALPH, S., and BOHLMANN, J. 2005. Insect-induced conifer defense: white pine
513 weevil and methyl jasmonate induce traumatic resinosis, de novo formed volatile emissions, and
514 accumulation of terpenoid synthase and putative octadecanoid pathway transcripts in Sitka spruce. *Plant*
515 *Physiol.* 137:369-382.
- 516 MILLER-PIERCE, M.R., ORWIG, D.A., and PREISSER, E.L. 2010. Effects of hemlock woolly adelgid and
517 elongate hemlock scale on eastern hemlock growth and foliar chemistry. *Environ. Entomol.* 39:513-519.
- 518 MORAN, M.D. 2003. Arguments for rejecting the sequential Bonferroni in ecological studies. *Oikos* 100:403-405.
- 519 MORAN, P.J. and THOMPSON, G.A. 2001. Molecular responses to aphid feeding in *Arabidopsis* in relation to
520 plant defense pathways. *Plant Physiol.* 125:1074-1085.
- 521 MUMM, R., SCHRANK, K., WEGENER, R., SCHULZ, S., and HILKER, M. 2003. Chemical analysis of volatiles
522 emitted by *Pinus sylvestris* after induction by insect oviposition. *J. Chem. Ecol.* 29:1235-1252.
- 523 ORWIG, D.A., COBB, R.C., D'AMATO, A.W., KIZLINSKI, M.L., and FOSTER, D.R. 2008. Multi-year
524 ecosystem response to hemlock woolly adelgid infestation in southern New England forests. *Can. J. Forest*
525 *Res.* 38:834-843.
- 526 PARK, S.-W., KAIMOYO, E., KUMAR, D., MOSHER, S., and KLESSIG, D.F. 2007. Methyl salicylate is a critical
527 mobile signal for plant systemic acquired resistance. *Science* 318:113-116.
- 528 PREISSER, E.L., LODGE, A.G., ORWIG, D.A., and ELKINTON, J.S. 2008. Range expansion and population
529 dynamics of co-occurring invasive herbivores. *Biol. Invasions* 10:201-213.
- 530 PREISSER, E.L., MILLER-PIERCE, M.R., VANSANT, J.L., and ORWIG, D.A. 2011. Eastern hemlock (*Tsuga*
531 *canadensis*) regeneration in the presence of hemlock woolly adelgid (*Adelges tsugae*) and elongate
532 hemlock scale (*Fiorinia externa*). *Canad. J. For. Res.* 41:2433-2439.
- 533 QUIROZ, A., PETTERSSON, J., PICKETT, J., WADHAMS, L., and NIEMEYER, H. 1998. Semiochemicals
534 mediating spacing behavior of bird cherry-oat aphids, *Rhopalosiphum padi*, feeding on cereals. *J. Chem.*
535 *Ecol.* 23:2599-2607.
- 536 R DEVELOPMENT CORE TEAM. 2012. R: a language and environment for statistical computing. R Foundation

- 537 for Statistical Computing, Vienna, Austria. <<http://www.R-project.org>>.
- 538 RADVILLE, L., CHAVES, A., and PREISSER, E.L. 2011. Variation in plant defense against invasive herbivores:
539 evidence for a hypersensitive response in eastern hemlocks (*Tsuga canadensis*). *J. Chem. Ecol.* 37:592-597.
- 540 RUSSO, V.M., RUSSO, B.M., PETERS, M., PERKINS-VEAZIE, P., and CARTWRIGHT, B. 1997. Interaction of
541 *Colletotrichum orbiculare* with thrips and aphid feeding on watermelon seedlings. *Crop Prot.* 16:581-584.
- 542 SAMPEDRO, L., MOREIRA, X., LLUSIA, J., PENUELAS, J., and ZAS, R. 2010. Genetics, phosphorus
543 availability, and herbivore-derived induction as sources of phenotypic variation of leaf volatile terpenes in a
544 pine species. *J. Exp. Biol.* 61:4437-4447.
- 545 SAMPEDRO, L., MOREIRA, X., and ZAS, R. 2011. Resistance and response of *Pinus pinaster* seedlings to
546 *Hylobius abietis* after induction with methyl jasmonate. *Plant Ecol.* 212:397-401.
- 547 SAS INSTITUTE INC. 2011. The SAS system for Windows, version 9.3. SAS Institute, Cary, NC.
- 548 SCHIESTL, F.P. 2010. The evolution of floral scent and insect chemical communication. *Ecol. Lett.* 13:643-56.
- 549 SHENEP, L.E., SHENEP, M.A., CHEATHAM, W., HOFFMAN, J.M., HALE, A., WILLIAMS, B.F., PERKINS,
550 R., HEWITT, C.B., HAYDEN, R.T., and SHENEP, J.L. 2011. Efficacy of intravascular catheter lock
551 solutions containing preservatives in the prevention of microbial colonization. *J. Hosp. Infect.* 79:317-322.
- 552 SHEPHERD, W.P., HUBER, D.P.W., SEYBOLD, S.J., and FETTIG, C.J. 2007. Antennal responses of the western
553 pine beetle, *Dendroctonus brevicomis* (Coleoptera: Curculionidae), to stem volatiles of its primary host,
554 *Pinus ponderosa*, and nine sympatric nonhost angiosperms and conifers. *Chemoecology* 17:209-221.
- 555 SHIELDS, K.S. and HIRTH, R.T. 2005. Bacterial endosymbionts of *Adelges tsugae* Annand: potential targets for
556 biocontrol?, pp. 357-59, in: B. Onken and R. Reardon, (eds.). Proceedings of the Fourth Symposium on the
557 Hemlock Woolly Adelgid in the Eastern United States. Forest Health Technology Enterprise Team,
558 Morgantown, WV: USDA, For. Serv.
- 559 SHIOJIRI, K., KISHIMOTO, K., OZAWA, R., KUGIMIYA, S., URASHIMO, S., ARIMURA, G., HORIUCHI, J.,
560 NISHIOKA, T., MATSUI, K., and TAKABAYASHI, J. 2006. Changing green leaf volatile biosynthesis in
561 plants: An approach for improving plant resistance against both herbivores and pathogens. *Proc. Natl.*
562 *Acad. Sci. USA* 103:16672-16676.
- 563 SHULAEV, V., SILVERMAN, P., and RASKIN, I. 1997. Airborne signalling by methyl salicylate in plant
564 pathogen resistance. *Nature* 385:718-721.

- 565 STEIN, S.E. 2005. NIST Standard Reference Database 1a. NIST/EPA/NIH Mass Spectral Database (NIST05) and
566 NIST Mass Spectral Search Program (NIST05s; Version 2.0g).
- 567 STRIMMER, K. 2008. Fdrtool: a versatile R package for estimating local and tail area-based false discovery rates.
568 *Bioinform. Applic. Note* 24:1461-1462.
- 569 TOOKER, J.F. and DE MORAES, C.M. 2009. A gall-inducing caterpillar species increases essential fatty acid
570 content of its host plant without concomitant increases in phytohormone levels. *Mol. Plant-Microbe In.*
571 22:551-9.
- 572 TRAPP, S. and CROTEAU, R. 2001. Defensive resin biosynthesis in conifers. *Annu. Rev. Plant Phys.* 52:689-724.
- 573 VEN, W.T.G. V.D., LEVESQUE, C.S., PERRING, T.M., and WALLING, L.L. 2000. Local and systemic changes
574 in squash gene expression in response to silverleaf whitefly feeding. *Plant Cell* 12:1409-1423.
- 575 YOUNG, R., SHIELDS, K., and BERLYN, G. 1995. Hemlock woolly adelgid (Homoptera:Adelgidae): Stylet
576 bundle insertion and feeding sites. *Ann. Entomol. Soc. Am.* 88:827-835.
- 577 ZHU-SALZMAN, K., SALZMAN, R.A., AHN, J., and KOIWA, H. 2004. Transcriptional regulation of sorghum
578 defense determinants against a phloem-feeding aphid. *Plant Physiol.* 134:420-431.
- 579 ZULAK, K.G. and BOHLMANN, J. 2010. Terpenoid biosynthesis and specialized vascular cells of conifer defense.
580 *J. Integr. Plant Biol.* 52:86-97.
- 581
- 582
- 583
- 584
- 585
- 586
- 587
- 588
- 589
- 590

591

FIGURES AND TABLES

592

Table 1 Resin volatile concentration relative change (treatment average/control average ratio) for eastern hemlock saplings treated with 3-year artificial infestation with hemlock woolly adelgid (HWA) or elongate hemlock scale (EHS)

593

594

	Twig Volatiles				Needle Volatiles				
	June ^a		October ^a		June		October		
	HWA	EHS	HWA	EHS	HWA	EHS	HWA	EHS	
Monoterpenoids^b									Monoterpenoids
Tricyclene	*0.49^c	0.80	*0.62	0.95	1.07	1.02	1.12	1.09	Tricyclene
α-Pinene ^d	0.78	0.93	1.08	0.71	1.08	1.05	1.15	1.09	α-Pinene ^d
Camphene ^d	*0.49	0.76	*0.68	0.91	1.04	1.05	1.08	1.13	Camphene ^d
β-Pinene ^d	0.76	1.07	1.22	0.75	1.14	0.98	1.19	0.91	Sabinene
Myrcene ^d	0.84	1.30	1.41	1.63	1.09	1.00	1.15	1.00	β-Pinene ^d
Limonene ^d	0.72	0.95	0.69	*1.47	1.05	1.02	1.08	1.08	Myrcene ^d
L-trans-Pinocarveol	0.74	0.87	0.71	*0.58	1.18	1.12	*1.34	1.09	α-Phellandrene ^d
cis-Verbenol	0.93	1.15	0.73	*0.46	1.19	1.15	1.12	1.07	Limonene ^d
trans-Verbenol	0.77	0.93	0.72	*0.58	0.88	0.83	1.14	0.92	γ-Terpinene ^d
Borneol ^d	0.73	0.87	*0.72	1.24	1.03	1.14	1.27	1.28	Terpinolene ^d
trans-Carveol ^d	0.73	0.88	0.72	*0.63	1.05	1.77	0.73	1.40	Camphor ^d
Myrtenol ^d	*0.61	*0.62	0.70	*0.56	0.05	0.10	2.30	1.68	Borneol ^d
α-Campholenal	*0.59	0.75	0.68	*0.61	0.83	0.82	0.93	0.91	4-Carvomenthenol ^d
Pinocarvone	*0.68	0.94	0.77	*0.56	0.84	0.74	1.15	1.08	p-Menth-1-en-9-ol
Verbenone ^d	*0.60	0.70	0.75	0.65	1.19	1.14	0.49	0.78	α-Terpineol ^d
Bornyl Acetate ^d	*0.45	*0.60	*0.71	0.82	0.70	*0.59	1.24	1.18	trans-Piperitol
total	*0.61	0.79	0.95	0.80	0.83	0.87	1.06	1.28	Piperitone ^d
Sesquiterpenoids					1.01	1.04	1.08	1.11	Bornyl Acetate ^d
β-Caryophyllene ^d	1.05	*0.00	2.38	1.25	1.04	1.03	1.11	1.10	total
α-Humulene ^d	1.89	*3.72	2.81	1.46					Sesquiterpenoids
Germacrene-D	1.05	1.34	1.17	0.55	1.01	1.07	1.24	1.13	β-Caryophyllene ^d
α-Amorphene	3.83	1.04	3.24	1.15	1.05	1.12	1.24	1.13	α-Humulene ^d
Caryophyllene Oxide ^d	0.95	0.85	0.88	*0.50	1.26	0.71	1.76	0.69	Germacrene-D
total	1.26	1.03	1.78	0.87	0.91	1.00	1.15	1.09	α-Amorphene
Benzenoids					0.97	1.02	1.24	1.13	δ-Cadinene
p-Cymene ^d	0.75	0.72	1.25	0.73	1.05	1.03	1.30	1.08	total
Benzyl Alcohol ^d	*31.43	0.53	*9.68	0.86					Benzenoids
p-Cymen-8-ol ^d	0.93	0.50	0.92	0.71	0.33	0.25	0.74	0.63	p-Cymene ^d
Methyl Salicylate ^d	*123.61	0.00	*12.05	1.10					Green leaf volatiles
3,4-Dimethoxyphenol	0.88	0.99	1.13	0.86	1.08	1.41	1.28	1.55	n-Hexanal ^d
Raspberry Ketone ^d	0.73	0.72	*0.66	*0.64	1.30	1.09	1.82	*2.55	trans-2-Hexenal ^d
total	*2.41	0.71	1.21	0.72	1.04	1.37	1.31	*2.55	cis-3-Hexenal
Unknown A	0.86	0.83	0.79	*0.50	1.22	1.19	*1.41	*1.87	total
Unknown B	*0.30	0.62	*0.67	0.59	1.03	1.03	1.13	1.12	Total Needle Volatiles
Unknown C	*0.30	0.80	*0.29	0.48					
Total Twig Volatiles	*0.77	0.78	0.98	0.79					

^a Foliage sampled in June was mature, previous year growth infested with EHS or progredien-generation HWA; foliage sampled in October was young, current year growth infested with EHS or sisten-generation HWA.

^b Compounds are ordered first by structural class, then by ascending order of elution from a non-polar DB-5 GC column. A summed total for each class of phytochemical is included.

^c Values >1 and <1 indicate an increase (dark gray shading) and decrease (light gray shading), respectively, from the control trees. Statistically significant differences from uninfested trees (planned contrast, P<0.05) are marked in bold text with asterisks. Marginally significant (0.05<P<0.10) values are marked in italics with asterisks.

^d Tentative identification based on comparison of GC retention time and mass spectrum with those of an authentic standard

595

597 **Fig. 1** (A) GC-FID total ion chromatogram showing volatiles tentatively identified in HWA-infested
 598 eastern hemlock needles: **1**, *cis*-3-hexenal; **2**, *n*-hexanal; **3**, *trans*-2-hexenal; **4**, tricyclene; **5**, α -pinene; **6**,
 599 camphene; **7**, sabinene; **8**, β -pinene; **9**, myrcene; **10**, α -phellandrene; **11**, isobutylbenzene (internal
 600 standard); **12**, *p*-cymene, **13**, D-limonene; **14**, γ -terpinene, **15**, terpinolene; **16**, camphor; **17**, borneol; **18**, 4-
 601 carvomenthenol; **19**, *p*-menth-1-en-9-ol; **20**, α -terpineol; **21**, *trans*-piperitol; **22**, piperitone; **23**, bornyl
 602 acetate; **24**, β -caryophyllene; **25**, α -humulene; **26**, germacrene-D; **27**, α -amorphene; **28**, δ -cadinene. (B)
 603 GC-MS total ion chromatogram showing volatiles in HWA-infested twigs: **1**, tricyclene; **2**, α -pinene; **3**,
 604 camphene; **4**, β -pinene; **5**, myrcene; **6**, isobutylbenzene (internal standard); **7**, *p*-cymene; **8**, D-limonene; **9**,
 605 benzyl alcohol; **10**, unknown; **11**, unknown; **12**, α -campholenal; **13**, *L-trans*-pinocarveol; **14**, *cis*-verbenol;
 606 **15**, *trans*-verbenol; **16**, pinocarvone; **17**, borneol; **18**, *p*-cymen-8-ol; **19**, methyl salicylate; **20**, myrtenol; **21**,
 607 verbenone; **22**, *cis*-carveol; **23**, bornyl acetate; **24**, unknown; **25**, β -caryophyllene; **26**, 3,4-
 608 dimethoxyphenol; **27**, α -humulene; **28**, germacrene-D; **29**, α -amorphene; **30**, raspberry ketone; **31**,
 609 caryophyllene dioxide.

610

611 **Fig. 2** Green leaf volatile ('GLV') content (average \pm SE) in needle tissue of control and insect-infested
 612 eastern hemlocks. 'HWA' or 'EHS' represents 3-year artificial infestation with hemlock woolly adelgid (*A.*
 613 *tsugae*) or elongate hemlock scale (*F. externa*). Data represents the average concentration ($\mu\text{g}\cdot\text{g dry wt}^{-1}$) of
 614 total GLVs in mature previous year growth (sampled 28 June) and young current year growth (sampled 19
 615 October), calculated from 6 to 9 trees per treatment group. *P*-values are shown when the difference
 616 between the treatment and control trees was significant ($P<0.05$), or marginally significant ($0.05<P<0.10$;
 617 planned contrast).

618

619 **Fig. 3** Benzyl alcohol content (average \pm SE) in twig tissue of control and insect-infested eastern hemlock
 620 trees. 'HWA' or 'EHS' represents 3-year artificial infestation with hemlock woolly adelgid (*A. tsugae*) or
 621 elongate hemlock scale (*F. externa*). Data represents the average concentration ($\mu\text{g}\cdot\text{g dry wt}^{-1}$) of benzyl
 622 alcohol in mature previous year growth (sampled 28 June) and young current year growth (sampled 19
 623 October), calculated from 7 to 9 trees per treatment group. *P*-values are shown when the difference

624 between the treatment and control trees was significant ($P<0.05$), or marginally significant ($0.05<P<0.10$;
625 planned contrast).

626

627 **Fig. 4** Methyl salicylate content (average \pm SE) in twig tissue of control and insect-infested eastern
628 hemlock trees. 'HWA' or 'EHS' represents 3-year artificial infestation with hemlock woolly adelgid (*A.*
629 *tsugae*) or elongate hemlock scale (*F. externa*). Data represents the average concentration ($\mu\text{g}\cdot\text{g dry wt}^{-1}$) of
630 methyl salicylate in mature previous year growth (sampled 28 June) and young current year growth
631 (sampled 19 October) calculated from 7 to 9 trees per treatment group. *P*-values are shown when the
632 difference between the treatment and control trees was significant ($P<0.05$), or marginally significant
633 ($0.05<P<0.10$; planned contrast).

634

635

636

637


638

639

640

641 **Fig. 1**

642



643

644 Fig. 2

645

646

647

648

649


650

651

652

653

654



655 Fig. 3

656

657

658

659


660

661

662


663

664




665 Fig. 4

666



667

ELECTRONIC SUPPLEMENTARY MATERIALS



Online Resource 1 Structures of representative volatile compounds of eastern hemlock (*Tsuga canadensis* Carr.): (A) monoterpenoids, (B) sesquiterpenoids, (C) green leaf volatiles (GLVs) and (D) benzenoids detected in twig or needle tissue of young trees.

Online Resource 2A Twig volatile concentrations (average \pm SE) with mixed model ANOVA results

	Previous year growth (June sampling)			Current year growth (October sampling)			HWA versus control analyses			EHS versus control analyses		
	Control	HWA	EHS	Control	HWA	EHS	Sample	Insect	q-value	Sample	Insect	q-value
	(N=8)	(N=9)	(N=7)	(N=8)	(N=9)	(N=7)	Date	Effect	F(P)	Date	Effect	F(P)
Tricyclene	15.55 (2.44)	7.70 (1.29)	12.49 (3.50)	26.32 (3.55)	16.45 (2.72)	25.04 (4.10)	41.56 ^b (0.00)	7.46 (0.01)	0.05	37.46 (0.00)	0.50 (0.49)	0.24
α -Pinene	29.48 (4.27)	22.96 (5.81)	27.45 (4.48)	345.86 (90.76)	372.80 (127.32)	245.02 (51.46)	163.97 (0.00)	1.10 (0.31)	0.18	218.00 (0.00)	1.02 (0.33)	0.19
Camphene	15.43 (2.06)	7.53 (1.24)	11.75 (2.87)	33.34 (3.65)	22.71 (3.85)	30.47 (4.50)	70.93 (0.00)	11.09 (0.00)	0.03	73.19 (0.00)	1.19 (0.29)	0.18
β -Pinene	2.74 (0.36)	2.09 (0.53)	2.94 (0.40)	35.65 (10.07)	43.59 (15.50)	26.84 (5.92)	145.38 (0.00)	0.50 (0.49)	0.24	205.05 (0.00)	0.29 (0.60)	0.28
Myrcene	3.31 (0.90)	2.78 (0.39)	4.30 (1.11)	31.51 (4.13)	44.46 (19.41)	51.46 (17.22)	61.94 (0.00)	0.32 (0.58)	0.28	91.03 (0.00)	0.72 (0.41)	0.22
Limonene	0.30 (0.18)	0.22 (0.11)	0.29 (0.20)	26.99 (2.13)	18.59 (4.38)	39.72 (5.49)	155.13 (0.00)	2.83 (0.11)	0.11	829.59 (0.00)	1.78 (0.20)	0.15
α -Campholenal	5.37 (0.89)	3.15 (0.73)	4.01 (0.48)	14.59 (2.53)	9.99 (1.52)	8.97 (0.88)	64.95 (0.00)	5.63 (0.03)	0.07	53.25 (0.00)	4.38 (0.06)	0.09
L-trans-Pinocarveol	3.31 (0.51)	2.44 (0.61)	2.90 (0.56)	11.08 (1.89)	7.87 (1.57)	6.47 (1.01)	63.33 (0.00)	4.04 (0.06)	0.09	34.49 (0.00)	4.48 (3.05)	0.09
cis-Verbenol	0.27 (0.09)	0.25 (0.08)	0.31 (0.15)	1.12 (0.15)	0.82 (0.17)	0.51 (0.25)	28.27 (0.00)	1.37 (0.26)	0.17	8.36 (0.01)	1.37 (0.11)	0.11
trans-Verbenol	4.76 (0.65)	3.65 (0.98)	4.45 (0.73)	21.63 (3.74)	15.58 (2.91)	12.45 (2.28)	70.76 (0.00)	3.46 (0.08)	0.10	58.47 (0.00)	3.56 (0.08)	0.10
Pinocarvone	3.62 (0.55)	2.47 (0.50)	3.40 (0.42)	9.66 (1.59)	7.43 (1.22)	5.45 (0.83)	69.94 (0.00)	3.42 (0.08)	0.10	26.51 (0.00)	4.22 (0.06)	0.09
Borneol	6.62 (1.37)	4.82 (1.28)	5.78 (1.11)	7.77 (1.00)	5.63 (0.86)	9.63 (2.41)	4.31 (0.05)	2.29 (0.15)	0.12	4.90 (0.04)	0.01 (0.93)	0.38
Myrtenol	5.09 (0.56)	3.12 (0.79)	3.17 (0.92)	22.18 (4.16)	15.59 (3.01)	12.46 (1.99)	137.21 (0.00)	4.61 (0.05)	0.09	50.32 (0.00)	8.48 (0.01)	0.05
Verbenone	9.74 (1.75)	5.85 (1.26)	6.87 (1.36)	15.40 (2.60)	11.49 (1.84)	10.06 (1.64)	14.32 (0.00)	3.50 (0.08)	0.10	7.82 (0.01)	3.02 (0.10)	0.11
trans-Carveol	0.72 (0.29)	0.52 (0.21)	0.63 (0.22)	4.77 (0.84)	3.45 (0.81)	2.98 (0.39)	84.00 (0.00)	1.14 (0.30)	0.18	76.51 (0.00)	1.10 (0.31)	0.18
Bornyl Acetate	32.87 (4.62)	14.86 (2.98)	19.63 (5.52)	78.83 (7.51)	56.32 (13.02)	64.61 (10.93)	86.15 (0.00)	9.76 (0.01)	0.04	74.67 (0.00)	3.69 (0.08)	0.10
β -Caryophyllene	0.50 (0.28)	0.52 (0.26)	0.00 (0.00)	2.88 (0.98)	6.85 (3.30)	3.61 (1.79)	12.20 (0.00)	0.28 (0.61)	0.28	11.83 (0.00)	0.65 (0.43)	0.23
α -Humulene	0.45 (0.24)	0.85 (0.34)	1.68 (0.57)	7.32 (2.57)	20.56 (10.00)	10.71 (3.80)	29.39 (0.00)	0.57 (0.46)	0.23	45.48 (0.00)	2.20 (0.16)	0.13
Germacrene-D	0.14 (0.09)	0.14 (0.10)	0.18 (0.14)	1.22 (0.28)	1.43 (0.37)	0.67 (0.32)	43.44 (0.00)	0.07 (0.79)	0.34	16.44 (0.00)	1.12 (0.31)	0.18
α -Amorphene	0.43 (0.39)	1.64 (1.14)	0.45 (0.45)	4.71 (2.90)	15.25 (7.88)	5.42 (4.01)	10.38 (0.01)	1.92 (0.18)	0.14	8.71 (0.01)	0.05 (0.83)	0.35
Caryophyllene Oxide	4.02 (0.79)	3.82 (0.86)	3.41 (0.96)	16.89 (3.93)	14.87 (3.81)	8.48 (1.29)	57.84 (0.00)	0.20 (0.66)	0.30	33.64 (0.00)	2.41 (0.14)	0.12
p-Cymene	4.91 (1.29)	3.69 (1.17)	3.55 (0.77)	26.64 (9.25)	33.32 (9.91)	19.54 (7.38)	87.06 (0.00)	0.02 (0.90)	0.37	55.30 (0.00)	0.61 (0.45)	0.23
Benzyl Alcohol	0.79 (0.33)	24.81 (10.70)	0.42 (0.42)	1.22 (0.24)	11.83 (3.94)	11.83 (3.94)	1.57 (0.23)	12.56 (0.00)	0.03	5.95 (0.03)	0.99 (0.34)	0.19
p-Cymen-8-ol	3.72 (1.01)	3.46 (0.82)	1.86 (0.95)	8.81 (1.99)	8.11 (1.56)	6.28 (1.86)	28.28 (0.00)	0.00 (0.99)	0.39	15.41 (0.00)	2.34 (0.15)	0.12
Methyl Salicylate ^c	0.02 (0.02)	2.91 (1.72)	0.00 (0.00)	0.08 (0.04)	0.96 (0.45)	0.09 (0.09)	3.69 (0.07)	8.03 (0.01)	0.05	2.34 (0.15)	0.10 (0.75)	0.33
3,4-Dimethoxyphenol	2.83 (0.41)	2.48 (0.44)	2.80 (0.52)	5.15 (0.76)	5.84 (0.83)	4.45 (0.92)	16.37 (0.00)	0.00 (0.96)	0.39	4.20 (0.06)	0.39 (0.54)	0.26
Raspberry Ketone	4.59 (0.62)	3.34 (0.59)	3.33 (0.70)	16.46 (2.28)	10.79 (1.70)	10.53 (3.30)	67.80 (0.00)	4.27 (0.06)	0.09	21.20 (0.00)	3.02 (0.10)	0.11
Unknown A	0.82 (0.35)	0.71 (0.22)	0.68 (0.50)	3.63 (0.52)	2.86 (0.45)	1.83 (0.62)	35.78 (0.00)	1.05 (0.32)	0.19	44.52 (0.00)	1.92 (0.19)	0.14
Unknown B	3.03 (0.44)	1.95 (0.44)	2.43 (0.67)	22.35 (4.30)	15.03 (3.92)	13.24 (1.72)	488.65 (0.00)	6.06 (0.03)	0.07	316.95 (0.00)	2.56 (0.13)	0.12
Unknown C	16.20 (2.77)	4.78 (0.70)	9.99 (1.79)	14.40 (4.10)	4.22 (0.67)	6.97 (1.30)	2.06 (0.17)	15.01 (0.00)	0.02	6.35 (0.02)	2.40 (0.14)	0.12
Total Monoterpenoids	139.16 (12.82)	84.40 (16.83)	110.36 (18.05)	686.71 (117.13)	652.75 (190.09)	552.14 (88.28)	103.68 (0.00)	5.60 (0.02)		118.05 (0.00)	3.14 (0.09)	
Total Sesquiterpenoids	5.53 (1.31)	6.98 (1.87)	5.71 (1.65)	33.04 (10.17)	58.96 (23.51)	28.88 (8.88)	35.13 (0.00)	0.44 (0.52)		41.86 (0.00)	0.16 (0.69)	
Total Benzenoids/Phenolics	16.87 (2.07)	40.69 (12.45)	11.94 (2.26)	58.36 (12.67)	70.84 (12.09)	41.94 (11.62)	33.17 (0.00)	3.07 (0.10)		46.06 (0.00)	3.35 (0.09)	
Total Twig Resin Volatiles	181.60 (15.13)	139.50 (22.67)	141.10 (20.22)	818.49 (141.03)	804.66 (228.07)	645.00 (103.38)	82.54 (0.00)	2.36 (0.14)		119.96 (0.00)	3.92 (0.06)	

^a Familywise error rate was estimated using a false discovery rate (FDR) method. FDR estimates the proportion of incorrectly rejected null hypotheses among tests with statistically significant findings. The q-value for each hypothesis test is the minimum FDR level that would be needed to reject that test's null hypothesis.

^b There were no significant sampling date x insect treatment interactions, except as noted.

^c The HWA versus control model includes a significant sampling date x treatment interaction term; F=1.83, P=0.018.

Online Resource 2B Needle volatile concentrations (average \pm SE) with mixed model ANOVA results

	Previous year growth (June sampling)			Current year growth (October sampling)			HWA versus control analyses			EHS versus control analyses		
	Control	HWA	EHS	Control	HWA	EHS	Sample	Insect	q-value	Sample	Insect	q-value
	(N=8)	(N=9)	(N=7)	(N=8)	(N=9)	(N=7)	Date	Effect	F(P)	Date	Effect	F(P)
<i>cis</i> -3-Hexenal	1.21 (0.64)	1.26 (0.31)	1.66 (0.48)	1.59 (0.31)	2.07 (0.31)	4.04 (1.13)	4.85 (0.05)	1.13 (0.31)	**	6.45 (0.03)	5.18 (0.04)	*
<i>n</i> -Hexanal	0.54 (0.15)	0.58 (0.25)	0.76 (0.31)	13.43 (2.80)	17.13 (1.19)	20.87 (4.32)	174.51 (0.00)	0.81 (0.39)	*	118.68 (0.00)	1.62 (0.23)	*
<i>trans</i> -2-Hexenal	3.27 (1.31)	4.26 (0.92)	3.58 (0.58)	4.81 (0.79)	8.76 (2.03)	12.25 (3.03)	6.23 (0.03)	2.78 (0.12)	*	14.47 (0.00)	3.29 (0.10)	*
Tricyclene	26.39 (2.29)	28.26 (2.35)	26.88 (4.66)	50.67 (3.01)	56.86 (4.61)	55.33 (8.66)	153.14 (0.00)	0.64 (0.44)	*	73.17 (0.00)	0.09 (0.77)	*
α -Pinene	91.56 (8.08)	98.81 (9.21)	96.22 (14.25)	186.27 (13.69)	213.45 (20.86)	202.57 (26.49)	177.07 (0.00)	0.63 (0.44)	*	63.48 (0.00)	0.00 (0.97)	*
Camphene	60.41 (5.13)	62.60 (5.52)	63.14 (9.42)	115.01 (6.74)	123.94 (10.52)	129.77 (16.85)	148.38 (0.00)	0.19 (0.67)	*	65.98 (0.00)	0.00 (0.95)	*
Sabinene	3.60 (0.35)	4.08 (0.38)	3.54 (0.57)	7.06 (0.82)	8.40 (0.77)	6.40 (0.86)	114.12 (0.00)	1.50 (0.24)	*	38.00 (0.00)	0.37 (0.56)	*
β -Pinene	12.92 (1.39)	14.11 (1.42)	12.91 (2.00)	29.00 (2.80)	33.21 (2.50)	29.00 (3.78)	215.16 (0.00)	0.91 (0.36)	*	70.42 (0.00)	0.05 (0.82)	*
Myrcene	15.26 (1.23)	16.05 (1.52)	15.57 (2.27)	29.20 (1.69)	31.65 (2.48)	31.40 (3.68)	139.98 (0.00)	0.33 (0.57)	*	63.15 (0.00)	0.01 (0.91)	*
α -Phellandrene	6.01 (1.08)	7.11 (1.31)	6.74 (1.19)	17.87 (2.45)	23.87 (2.07)	19.53 (3.06)	145.20 (0.00)	1.03 (0.33)	*	75.42 (0.00)	0.23 (0.64)	*
Limonene	25.18 (4.00)	29.89 (2.76)	29.08 (4.76)	58.90 (3.99)	66.21 (5.61)	63.23 (7.40)	31.97 (0.00)	1.55 (0.24)	*	23.70 (0.00)	0.39 (0.55)	*
γ -Terpinene	0.48 (0.08)	0.42 (0.11)	0.39 (0.15)	0.99 (0.05)	1.13 (0.12)	0.91 (0.20)	50.44 (0.00)	0.02 (0.89)	*	29.47 (0.00)	0.72 (0.41)	*
Terpinolene	2.03 (0.18)	2.10 (0.26)	2.33 (0.37)	4.75 (0.30)	6.01 (0.84)	6.07 (0.82)	130.28 (0.00)	0.70 (0.42)	*	77.55 (0.00)	0.81 (0.39)	*
Camphor	0.60 (0.15)	0.62 (0.32)	1.06 (0.29)	1.84 (0.44)	1.35 (0.69)	2.58 (1.07)	22.70 (0.00)	0.30 (0.59)	*	14.11 (0.00)	0.79 (0.39)	*
Borneol	1.52 (1.43)	0.08 (0.08)	0.16 (0.10)	0.09 (0.09)	0.21 (0.14)	0.15 (0.10)	0.28 (0.61)	0.62 (0.45)	*	0.71 (0.42)	0.33 (0.58)	*
4-Carvomenthenol	0.87 (0.11)	0.72 (0.11)	0.71 (0.15)	1.50 (0.17)	1.39 (0.18)	1.37 (0.31)	31.05 (0.00)	0.64 (0.44)	*	17.99 (0.00)	0.56 (0.47)	*
<i>p</i> -Menth-1-en-ol	1.72 (0.22)	1.44 (0.31)	1.27 (0.35)	2.73 (0.20)	3.13 (0.70)	2.94 (0.54)	65.17 (0.00)	0.36 (0.56)	*	61.76 (0.00)	1.20 (0.30)	*
α -terpineol	0.79 (0.06)	0.93 (0.08)	0.90 (0.04)	1.21 (0.48)	0.59 (0.11)	0.94 (0.39)	0.31 (0.59)	0.60 (0.45)	*	0.10 (0.76)	0.03 (0.86)	*
Piperitol	0.88 (0.08)	0.62 (0.14)	0.52 (0.19)	3.10 (0.62)	3.86 (0.42)	3.67 (0.70)	92.22 (0.00)	0.02 (0.90)	*	56.01 (0.00)	0.50 (0.49)	*
Piperitone	24.20 (4.60)	20.02 (2.30)	21.16 (4.72)	37.11 (4.67)	39.23 (4.83)	47.58 (6.37)	37.05 (0.00)	0.01 (0.91)	*	29.88 (0.00)	0.01 (0.94)	*
Bornyl Acetate	125.05 (10.85)	126.13 (11.00)	130.54 (18.86)	212.80 (16.12)	229.55 (27.16)	235.76 (29.51)	92.52 (0.00)	0.07 (0.79)	*	48.47 (0.00)	0.01 (0.91)	*
β -Caryophyllene	11.85 (0.91)	11.99 (0.72)	12.67 (2.12)	22.22 (2.09)	27.53 (2.66)	25.15 (3.69)	108.43 (0.00)	1.11 (0.31)	*	66.54 (0.00)	0.01 (0.92)	*
α -Humulene	13.51 (1.20)	14.16 (0.91)	15.12 (2.55)	26.54 (2.54)	32.89 (3.21)	30.09 (4.43)	121.09 (0.00)	1.42 (0.26)	*	70.01 (0.00)	0.05 (0.82)	*
Germacrene D	4.61 (0.91)	5.79 (1.19)	3.28 (1.12)	7.64 (2.38)	13.45 (2.99)	5.24 (2.06)	47.97 (0.00)	1.32 (0.27)	*	31.87 (0.00)	1.80 (0.21)	*
α -Amorphene	2.29 (0.22)	2.09 (0.22)	2.30 (0.34)	4.10 (0.55)	4.73 (0.69)	4.46 (0.64)	61.99 (0.00)	0.01 (0.91)	*	66.56 (0.00)	0.05 (0.83)	*
δ -Cadinene	2.34 (0.19)	2.28 (0.24)	2.40 (0.41)	4.16 (0.58)	5.14 (0.70)	4.72 (0.76)	58.84 (0.00)	0.26 (0.62)	*	60.82 (0.00)	0.06 (0.81)	*
<i>p</i> -Cymene	5.76 (3.66)	1.92 (0.28)	1.46 (0.27)	3.66 (0.79)	2.71 (0.26)	2.31 (0.35)	0.87 (0.37)	1.83 (0.20)	*	0.84 (0.38)	3.66 (0.08)	*
Total Green Leaf Volatiles	5.02 (1.89)	6.10 (1.23)	5.99 (0.81)	19.83 (2.84)	27.96 (3.11)	37.16 (4.49)	62.45 (0.00)	3.04 (0.11)	*	85.22 (0.00)	6.02 (0.03)	*
Total Monoterpenoids	399.47 (28.46)	413.99 (34.62)	413.11 (59.97)	760.09 (46.43)	844.04 (78.55)	839.20 (102.74)	22.89 (0.00)	1.12 (0.31)	*	18.00 (0.00)	0.40 (0.54)	*
Total Sesquiterpenoids	34.60 (2.66)	36.31 (1.87)	35.76 (5.67)	64.66 (7.03)	83.75 (9.16)	69.65 (9.99)	32.83 (0.00)	1.79 (0.21)	*	20.64 (0.00)	0.16 (0.69)	*
Total Needle Volatiles	444.84 (27.90)	458.33 (35.81)	456.32 (65.18)	848.23 (51.45)	958.46 (84.71)	948.32 (112.67)	183.62 (0.00)	0.44 (0.52)	*	75.63 (0.00)	0.00 (0.99)	*

* FDR results were not reported for needle volatile tests due to the scarcity of statistically significant findings. See text for a more detailed explanation.