

2016

N-Acyl Dehydrotyrosines, Tyrosinase Inhibitors from the Marine Bacterium, *Thalassotalea* sp. PP2-459

Robert W. Deering
University of Rhode Island, rdeering19@uri.edu

Jianwei Chen
University of Rhode Island

Jiadong Sun
University of Rhode Island

Hang Ma
University of Rhode Island, hang_ma@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/bps_facpubs

Javier Dubert

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Deering, R. W., Chen, J., Sun, J., Ma, H., Dubert, J., Barja, J. L., Seeram, N. P.,...Rowley D. C. (2016). N-Acyl Dehydrotyrosines, Tyrosinase Inhibitors from the Marine Bacterium *Thalassotalea* PP2-459. *Journal of Natural Products*, 79(2), 447-450. doi: 10.1021/acs.jnatprod.5b00972

Available at: <http://dx.doi.org/10.1021/acs.jnatprod.5b00972>

This Article is brought to you for free and open access by the Biomedical and Pharmaceutical Sciences at DigitalCommons@URI. It has been accepted for inclusion in Biomedical and Pharmaceutical Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors

Robert W. Deering, Jianwei Chen, Jiadong Sun, Hang Ma, Javier Dubert, Juan L. Barja, Navindra P. Seeram, Hong Wang, and David C. Rowley

***N*-Acyl Dehydrotyrosines, Tyrosinase Inhibitors from the Marine Bacterium *Thalassotalea* sp. PP2-459**

Robert W. Deering,^{†,⊥} Jianwei Chen,^{†,‡,⊥} Jiadong Sun,[†] Hang Ma,[†] Javier Dubert,[§]
Juan L. Barja,[§] Navindra P. Seeram,[†] Hong Wang,[‡] David C. Rowley^{†,*}

[†]Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, University of Rhode Island, Kingston, RI, United States

[‡]College of Pharmaceutical Science, Zhejiang University of Technology, Hangzhou 310014, China

[§]Department of Microbiology and Parasitology, CIBUS- Faculty of Biology and Aquaculture Institute, University of Santiago de Compostela, Santiago de Compostela 17582, Spain

Corresponding Author

*Email: drowley@uri.edu; Tel: 401-874-9228

Author Contributions

[⊥]R. W. Deering and J. Chen contributed equally to this work.

ABSTRACT

Thalassotalic acids A-C and thalassotalamides A and B are new *N*-acyl dehydrotyrosine derivatives produced by *Thalassotalea* sp. PP2-459, a Gram-negative bacterium isolated from a marine bivalve aquaculture facility. The structures were elucidated via a combination of spectroscopic analyses emphasizing two-dimensional NMR and high-resolution mass spectrometric data. Thalassotalic acid A (**1**) displays *in vitro* inhibition of the enzyme tyrosinase with an IC₅₀ value (130 μM) that compares favorably to the commercially-used control compounds kojic acid (46 μM) and arbutin (100 μM). These are the first natural products reported from a bacterium belonging to the genus *Thalassotalea*.

Marine bacteria continue to be a highly productive source of new natural products with 169 new compounds reported in 2013 alone.¹ While marine Gram-positive bacteria, especially actinomycetes, have garnered the most attention to date, genome mining and isolation-driven discovery efforts have raised awareness for the biosynthetic potential of marine Gram-negative bacteria (GNB).^{2,3} Recognition that marine GNB produce such molecules as the didemnins,⁴ and quite likely the bryostatins,⁵ highlights the potential for discovery of molecules with biomedical value. The marine γ -proteobacteria have been a particularly productive taxonomic group of GNB for natural product studies, yielding notable antibiotics such as andrimid and the moiramides, as well as the signaling molecule AI-2.^{6,7} These studies and others portend the future success of discovery efforts aimed at marine GNB.

Thalassotalea is a new Gram-negative, aerobic, γ -proteobacteria genus.⁸ This genus was first characterized in 2014 following reclassification of members of the genus *Thalassomonas*.⁸ There have been only 10 species of *Thalassotalea* discussed in the literature and they have been primarily studied for their agarolytic properties, commensalism with marine animals, and pathogenic contribution to coral diseases.⁸⁻¹³ To date, there have been no published secondary metabolite studies on this genus.⁸⁻¹¹

The enzyme tyrosinase is responsible for three transformations in the conversion of tyrosine to melanin, which is responsible for pigmentation of skin, hair, and eyes.¹⁴ It specifically catalyzes the conversion of L-tyrosine to 3,4-dihydroxyphenylalanine (L-DOPA), L-DOPA to dopaquinone, and 5,6-dihydroxyindole (DHI) to indole-quinone.¹⁴ Because of tyrosinase's role in pigmentary diseases such as albinism and melasma, as well as cosmetic relevance to increase or decrease skin or hair color and pigmentation, compounds that can modulate tyrosinase have potential therapeutic and cosmetic applications.¹⁵ Additionally,

tyrosinase is responsible for enzymatic browning of foods and inhibition of this process could lead to improved agricultural lifespans and nutrition.¹⁶

In this paper, we report new *N*-acyl dehydrotyrosine derivatives (**1-5**) produced during laboratory cultivation of *Thalassotalea* sp. PP2-459, formerly classified as *Thalassomonas* sp. PP2-459, a strain isolated from a carpet-shell clam (*Ruditapes decussata*) harvested in a bivalve hatchery located in Galicia (NW Spain).¹⁷ PP2-459 was previously determined to have quorum quenching ability against key bivalve aquaculture pathogens, including *Vibrio anguillarum*.¹⁷ Hence, PP2-459 has been proposed as a candidate for probiotic applications in aquaculture. Due to structural similarity with known tyrosinase inhibitors such as *N*-acylated tyrosines, **1-5** were evaluated for their inhibition of the tyrosinase enzyme.

Thalassotalea PP2-459 was cultured in YP seawater medium at 25 °C shaking and 200 rpm for two days. The cultures were centrifuged at 10,000 g for 10 min to substantially remove the cells, and the resulting supernatants were extracted with EtOAc. Purification efforts using reversed-phase (C18) chromatography methods yielded new compounds **1-5**.

Thalassotalic acid A (**1**) was isolated as a light brown powder. The molecular formula was determined to be C₁₉H₂₇NO₄ using HRESIMS (*m/z* 332.1866 [M-H]⁻), indicating seven degrees of unsaturation. The presence of a decanoyl chain was supported by ¹H NMR resonances at δ_H 0.90 (t, 3H, H-10'), 1.29-1.31 (m, 12H, H-4'-H-9'), 1.59 (m, 2H, H-3'), and 2.28 (t, 2H, H-2') that comprised a spin system in the COSY spectrum (Table 1). ¹³C NMR resonances at δ_C 172.0 (C-1') and 166.7 (C-1) indicated the presence of two carbonyl carbons, the former belonging to the decanoyl chain based on HMBC correlations from H-2' and H-3'. In the aromatic region, two doublets at δ_H 6.80 (H-6, *J* = 8.7 Hz) and 7.51 (H-5, *J* = 8.7 Hz) provided evidence for a para-substituted benzene ring. A HSQC correlation assigned a sharp singlet at δ_H

7.22 (H-3) as an olefinic methine attached to carbon at δ_C 132.2. HMBC correlations from H-5 to C-3 and H-3 to C-5 indicated that the C-3 methine was the benzylic position of the para-substituted aromatic ring. Consideration of the molecular formula and further HMBC correlations from the C-2-NH to C-1, C-2, and C-3, as well as correlations from H-3 to C-1 and C-2, established the α - β unsaturated tyrosine (Figure 1). The (*Z*)-configuration of the double bond was determined based on NOE correlations shown in Figure 1. TOF-MS/MS experiments further corroborated the proposed structure. Product ions were observed indicating the loss of CO₂H (m/z 288.2, [M-45]), the decanoyl chain (m/z 178.0, [M-155]), and both the carboxylic acid and the acyl chain (m/z 134.0, [M-199]).

The structure of thalassotalic acid B (**2**) was determined by comparison to spectroscopic data of **1**. An [M-H]⁻ ion at m/z 332.1872 indicated the same molecular formula as **1**: C₁₉H₂₇NO₄. A key difference in the ¹H NMR spectrum was the presence of a sharp doublet at δ_H 0.90 that integrated for six protons, suggesting a terminal branch in the aliphatic chain. Analysis of the HSQC spectrum revealed an aliphatic methine at δ_H 1.55 and δ_C 27.4, whose ¹H NMR resonance coupled to the C-8' methyl groups in the COSY spectrum, thus confirming the terminal (CH₃)₂CH- group. The remaining structure was determined to be identical to **1** based on comparison of ¹³C and 2D NMR data and TOF-MS/MS evidence.

Thalassotalic acid C (**3**) displayed a HRESIMS ion (m/z 318.1699 [M-H]⁻) consistent with a molecular formula of C₁₈H₂₅NO₄, which is one CH₂ unit fewer than **1** and **2**. Comparison of ¹H and ¹³C NMR data with **1** and **2** revealed that **3** shared the same core tyrosine structure, but incorporated an unbranched chain containing one fewer methylene unit than **1**. Thus, **3** was determined to be the nonanoyl analog of **1**. TOF-MS/MS further corroborated this structure with

the characteristic losses of CO₂H (m/z 274.2, [M-45]), the nonanoyl chain (m/z 178.1, [M-141]), and both the carboxylic acid and the acyl chain (m/z 134.1 [M-185]).

Thalassotalamide A (**4**) was isolated as an amorphous brown powder. HRESIMS indicated that the molecular formula was C₁₉H₂₈N₂O₃ based on an [M-H]⁻ ion (m/z 331.2021). Thus, **4** had an additional NH and one less O than **1**. Examination of the ¹H NMR spectrum of **4** indicated strong similarity to **1** based on resonances attributed to the *p*-substituted benzene ring (two doublets each integrating as 2H at δ_H 6.78 and 7.42) and unbranched aliphatic chain. The outstanding difference was two broad singlets at δ_H 7.05 and 7.28, indicating that **4** was likely the amide analog of **1**. TOF-MS/MS confirmed this suspicion as fragment ions of m/z 288.2 [M-44] and m/z 134.1 [M-198] were observed, indicating that the α - β unsaturated carbonyl was an amide and not a carboxylic acid. Analysis of ¹³C and 2D NMR and further confirmed the structure of **4**.

The structure of thalassotalamide B (**5**) was elucidated by comparison to spectroscopic data of **2** and **4**. HRESIMS indicated that the molecular formula of **5** was the same as **4** (C₁₉H₂₈N₂O₃) based on an [M-H]⁻ ion (m/z 331.2015). In the ¹H NMR spectrum, a doublet at δ_H 0.89 integrating 6H suggested that **5** had the same branched aliphatic chain as **2**. Comparison of ¹³C NMR data from the aliphatic chain of **2** to **5** confirmed they had the same 8-methyl nonanoyl aliphatic chain. Comparison of the remaining ¹³C NMR data to those of **4**, as well as the prominent loss of the amide in TOF-MS/MS (m/z 288.2), indicated that **5** was the amide derivative of **2**.

N-acyl tyrosines have previously been studied for their effects on pigmentation of skin and hair.¹⁸⁻²⁰ These compounds have been implicated in both increasing and decreasing pigmentation.¹⁸⁻²⁰ Because of the similarity in structure to *N*-acyl tyrosines, we evaluated compounds **1-5** for their effects on tyrosinase enzyme activity compared to the positive controls

arbutin (**6**) and kojic acid (**7**) that have been widely used in skin whitening products.^{21, 22}

Compounds **1**, **2**, and **3** exhibited tyrosinase enzyme inhibition with IC₅₀ values of 130 ± 10 μM, 470 ± 10 μM, and 280 ± 10 μM, respectively. The IC₅₀ value for **1** compared favorably to the **6** (100 ± 10 μM), but was less effective than **7** (IC₅₀ = 46 ± 2 μM). Compounds **4** and **5** were inactive at the maximum tested concentration (1 mM). These data support the requirement for a carboxylic acid and straight aliphatic chain for increasing enzyme inhibition within this structural class of tyrosinase inhibitors. While these *N*-acyl dehydrotyrosine compounds are not as potent as **6** or **7**, **1** displays comparable tyrosinase inhibition and could be useful as a whitening agent or as an agricultural additive to prevent the browning of foods.¹⁶

N-acyl derivatives of amino acids, including tyrosine, have been previously produced via heterologous expression of environmental DNA in *Escherichia coli*. These molecules are products of *N*-acyl amino acid synthases, which have been observed as single genes or as part of more complex biosynthetic sequences.^{23, 24} While the compounds discussed here are structurally similar, the dehydrogenation across the α-β carbons of the tyrosine, the terminal amide functional group in **4** and **5**, and the branched acyl chains in **2** and **5** represent biosynthetic variations not present in heterologously produced *N*-acyl tyrosines.²⁴ Additionally, these are the first natural products reported from a bacterium belonging to the genus *Thalassotalea*.

EXPERIMENTAL SECTION

General Experimental Procedures: All chemical reagents and solvents were purchased from Sigma-Aldrich and acquired via Wilkem Scientific. UV spectra were acquired with a DU 800 UV/Vis Spectrophotometer (Beckman Coulter). IR spectra were acquired with a Nicolet 380 FT-IR (Thermo Electron Corporation). NMR experiments were conducted using an Agilent

NMRS 500 MHz spectrometer with (CD₃)₂SO as the solvent (referenced to residual DMSO at δ_{H} 2.54 and δ_{C} 39.5) at 25 °C. Electrospray ionization mass spectra (ESIMS) were acquired using an AB Sciex TripleTOF 4600 spectrometer in the negative ion mode. Flash chromatography was completed with a Combiflash Rf200 equipped with an 86 g C₁₈ RediSepRf High Performance Gold column (Teledyne ISCO). HPLC experiments were performed on a Hitachi Elite LaChrom system equipped with a diode array detector (DAD) model L-2450, pump L-2130, and autosampler L-2200. Semi-preparative HPLC experiments were completed with a Waters XBridge Prep C18 5 μm , 10 \times 250 mm column.

Bacterial Strain and Culture Conditions: The bacteria strain (PP2-459) was isolated from a bivalve hatchery in Galicia, Spain and identified as a *Thalassotalea* sp. based on 16S rRNA sequence comparison (European Nucleotide Archive; accession number LN898116).¹⁷ It has been deposited in the Spanish Type Culture Collection (CECT) under restricted access. Cultures were maintained in YP seawater medium (1 g yeast extract, 5 g peptone, 22.5 g Instant Ocean (Spectrum Brands) per liter of deionized water) at 25 °C and shaking at 200 rpm.

Extraction and Isolation: A total of 17 L of bacterial cultures were cultivated and centrifuged at 10,000 g for 10 min to remove cells. The culture supernatants were extracted twice with equal volumes of EtOAc. The resultant organic layer was concentrated *in vacuo* to generate a total extract of 1.5 g.

The extract was adsorbed onto C₁₈ resin (BAKERBOND Octadecyl (C₁₈) 40 μm Prep LC Packing) and separated by reversed-phase flash chromatography. A linear gradient of 10 - 100% MeOH in H₂O, each solvent modified with 0.1% formic acid (FA), afforded five fractions. Fraction 4 (135.6 mg) was further purified by reversed-phase C₁₈ semi-preparative HPLC (25 min linear gradient from 40 - 78% MeOH in H₂O (0.1% FA) to yield five pure compounds: **1**

(28.3 mg, $t_R = 17.9$ min), **2** (10.1 mg, $t_R = 16.9$ min), **3** (7.8 mg, $t_R = 13.8$ min), **4** (2.3 mg, $t_R = 16.0$ min), and **5** (2.2 mg, $t_R = 15.1$ min).

Thalassotalic acid A (1): off-white amorphous powder; UV (MeOH) λ_{\max} (log ϵ) 201 (4.16) 298 (3.99) nm; IR (ZnSe) ν_{\max} 3308, 3067 (br), 2917, 2850, 1688, 1656, 1601, 1584, 1271, 1167 cm^{-1} ; ^1H and ^{13}C NMR data, Table 1; HRESIMS m/z 332.1866 $[\text{M-H}]^-$ (calcd for $\text{C}_{19}\text{H}_{26}\text{NO}_4$, 332.1867).

Thalassotalic acid B (2): brown amorphous powder; UV (MeOH) λ_{\max} (log ϵ) 201 (4.12) 299 (4.03) nm; IR (ZnSe) ν_{\max} 3326, 3118 (br), 2925, 2853, 1688, 1656, 1601, 1584, 1270, 1168 cm^{-1} ; ^1H and ^{13}C NMR data, Table 1; HRESIMS m/z 332.1872 $[\text{M-H}]^-$ (calcd for $\text{C}_{19}\text{H}_{26}\text{NO}_4$, 332.1867).

Thalassotalic acid C (3): brown amorphous powder; UV (MeOH) λ_{\max} (log ϵ) 202 (4.27) 299 (4.37) nm; IR (ZnSe) ν_{\max} 3317, 3067 (br), 2922, 2852, 1692, 1655, 1601, 1584, 1270, 1168 cm^{-1} ; ^1H and ^{13}C NMR data, Table 1; HRESIMS m/z 318.1699 $[\text{M-H}]^-$ (calcd for $\text{C}_{18}\text{H}_{24}\text{NO}_4$, 318.1711).

Thalassotalamide A (4): brown amorphous powder; ^1H and ^{13}C NMR data, Table 1; HRESIMS m/z 331.2021 $[\text{M-H}]^-$ (calcd for $\text{C}_{19}\text{H}_{27}\text{N}_2\text{O}_3$, 331.2027).

Thalassotalamide B (5): brown amorphous powder; UV (MeOH) λ_{\max} (log ϵ) 200 (4.26) 210 (4.19) 272 (3.59) nm; IR (ZnSe) ν_{\max} 3726, 3624, 3321, 2925, 2854, 1665, 1578, 1137 cm^{-1} ; ^1H and ^{13}C NMR data, Table 1; HRESIMS m/z 331.2015 $[\text{M-H}]^-$ (calcd for $\text{C}_{19}\text{H}_{27}\text{N}_2\text{O}_3$, 331.2027).

Tyrosinase Inhibition Assay: Inhibitory effects of compounds **1-5** on mushroom tyrosinase were evaluated spectrophotometrically using L-tyrosine as a substrate according to our

previously published method with minor modification.²¹ Tyrosinase inhibition assays were performed in 96-well microplate format using a SpectraMax M2 microplate reader (Molecular Devices). Briefly, compounds were dissolved at various concentrations in 10% MeOH in phosphate buffer solution (PBS; 0.1 M, pH 6.8). Each well contained 40 μ L of sample with 80 μ L of PBS, 40 μ L of tyrosinase (100 units/mL), and 40 μ L L-tyrosine (2.5 mM). The mixture was incubated for 30 min at 37 °C, and absorbance was measured at 490 nm. Each sample was accompanied by a blank containing all components except L-tyrosine. Arbutin and kojic acid were used as positive controls. The results were compared with a control consisting of 10% MeOH in place of the test compounds. Experiments were completed in triplicate for each compound. The percentage of tyrosinase inhibition was calculated as: $[(\Delta A_{\text{control}} - \Delta A_{\text{sample}}) / \Delta A_{\text{control}}] \times 100$.

ACKNOWLEDGEMENTS

Instruments used for mass spectrometric and other analyses were supported through an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institute of Health (grant number 2 P20 GM103430). NMR data were acquired on instruments supported in part by the National Science Foundation EPSCoR Cooperative Agreement #EPS-1004057. J.C. was supported in part by the China Scholarship Council (File no. 201408330156).

Supporting Information Available: 1D and 2D NMR and HRESI mass spectra for compounds **1-5**. The Supporting Information is available free of charge on the ACS Publications website at DOI:

NOTES

The authors declare no conflicts of interest. Thalassotalic acids were previously assigned the trivial names of Thalassomononic acids,²⁵ but have been renamed in the current study based on the reclassification of this bacterial genus.

REFERENCES

- (1) Blunt, J. W.; Copp, B. R.; Keyzers, R. A.; Munro, M. H.; Prinsep, M. R. *Nat. Prod. Rep.* **2015**, *32*, 116-211.
- (2) Machado, H.; Sonnenschein, E. C.; Melchiorson, J.; Gram, L. *BMC. Genomics* **2015**, *16*, 158.
- (3) Still, P. C.; Johnson, T. A.; Theodore, C. M.; Loveridge, S. T.; Crews, P. *J. Nat. Prod.* **2014**, *77*, 690-702.
- (4) Xu, Y.; Kersten, R. D.; Nam, S.-J.; Lu, L.; Al-Suwailem, A. M.; Zheng, H.; Fenical, W.; Dorrestein, P. C.; Moore, B. S.; Qian, P.-Y. *J. Am. Chem. Soc.* **2012**, *134*, 8625-8632.
- (5) Sudek, S.; Lopanik, N. B.; Waggoner, L. E.; Hildebrand, M.; Anderson, C.; Liu, H.; Patel, A.; Sherman, D. H.; Haygood, M. G. *J. Nat. Prod.* **2007**, *70*, 67-74.
- (6) Mansson, M.; Gram, L.; Larsen, T. O. *Mar. Drugs.* **2011**, *9*, 1440-68.
- (7) Isnansetyo, A.; Kamei, Y. *J. Ind. Microbiol. Biotechnol.* **2009**, *36*, 1239-48.
- (8) Zhang, Y.; Tang, K.; Shi, X.; Zhang, X.-H. *Int. J. Syst. Evol. Micr.* **2014**, *64*, 1223-1228.

- (9) Stelling, S. C.; Techtmann, S. M.; Utturkar, S. M.; Alshibli, N. K.; Brown, S. D.; Hazen, T. C. *Genome Announc.* **2014**, *2*, e01231-14.
- (10) Park, S.; Jung, Y.-T.; Kang, C.-H.; Park, J.-M.; Yoon, J.-H. *Int. J. Syst. Evol. Micr.* **2014**, *64*, 3676-3682.
- (11) Hou, T.-T.; Liu, Y.; Zhong, Z.-P.; Liu, H.-C.; Liu, Z.-P. *Int. J. Syst. Evol. Micr.* **2015**.
- (12) Thompson, F.; Barash, Y.; Sawabe, T.; Sharon, G.; Swings, J.; Rosenberg, E. *Int. J. Syst. Evol. Micr.* **2006**, *56*, 365-368.
- (13) Ohta, Y.; Hatada, Y.; Miyazaki, M.; Nogi, Y.; Ito, S.; Horikoshi, K. *Curr. Microbiol.* **2005**, *50*, 212-216.
- (14) Hearing, V. J.; Tsukamoto, K. *FASEB. J.* **1991**, *5*, 2902-2909.
- (15) Ando, H.; Kondoh, H.; Ichihashi, M.; Hearing, V. J. *J. Invest. Dermatol.* **2007**, *127*, 751-761.
- (16) Parvez, S.; Kang, M.; Chung, H.-S.; Bae, H. *Phytother. Res.* **2007**, *21*, 805-816.
- (17) Torres, M.; Romero, M.; Prado, S.; Dubert, J.; Tahrioui, A.; Otero, A.; Llamas, I. *Microbiol. Res.* **2013**, *168*, 547-554.
- (18) Guglielmini, G. *J. Appl. Cosmetol.* **2006**, *24*, 55-61.
- (19) Kafara, M.; Arct, J.; Dzierzgowski, S. *J. Appl. Cosmetol.* **2009**, *27*, 1-12.
- (20) Alvaro, M.; Cozzi, R.; Genovese, A.; Sedghi Zadeh, H. Patent, I. WO 2013/179098 A1, 2013.
- (21) Niesen, D. B.; Ma, H.; Yuan, T.; Bach 2nd, A.; Henry, G. E.; Seeram, N. P. *Nat. Prod. Commun.* **2015**, *10*, 491-493.
- (22) Wang, Y. H.; Avonto, C.; Avula, B.; Wang, M.; Rua, D.; Khan, I. A. *J. AOAC. Int.* **2015**, *98*, 5-12.

- (23) Brady, S. F.; Clardy, J. *Org. Lett.* **2005**, *7*, 3613-3616.
- (24) Brady, S. F.; Chao, C. J.; Clardy, J. *Appl. Environ. Microb.* **2004**, *70*, 6865-6870.
- (25) Deering, R.; Chen, J.; Ma, H.; Dubert, J.; Prado, S.; Seeram, N. P.; Rowley, D. C. *Natl. Meet.-Am. Chem. Soc.* **2015**, *250*, MEDI 153.

Table 1. NMR Spectroscopic data (^1H 500 MHz, ^{13}C 125 MHz, DMSO- d_6) for Thalassotalic acids A-C (**1-3**) and Thalassotalamides A (**4**) and B (**5**).

position	Thalassotalic acid A (1)		Thalassotalic acid B (2)		Thalassotalic acid C (3)		Thalassotalamide A (4)		Thalassotalamide B (5)	
	δ_{C} , type	δ_{H} (J in Hz)	δ_{C} , type	δ_{H} (J in Hz)	δ_{C} , type	δ_{H} (J in Hz)	δ_{C} , type	δ_{H} (J in Hz)	δ_{C} , type	δ_{H} (J in Hz)
1	166.7, C		166.7, C		166.8, C		167.2, C		167.2, C	
1-OH/NH ₂		12.35, br s		12.23, br s		12.26, br s		7.28/7.05, br s		7.28/7.05, br s
2	124.2, C		124.3, C		124.5, C		127.0, C		127.0, C	
2-NH		9.23, s		9.22, s		9.20, s		9.18, s		9.19, s
3	132.2, CH	7.22, s	132.0, CH	7.22, s	131.7, ^a CH	7.21, s	128.6, CH	7.06, s	128.6, CH	7.06, s
4	124.7, C		124.7, C		124.8, C		125.0, C		125.0, C	
5	131.7, CH	7.51, d (8.7)	131.7, CH	7.51, d (8.6)	131.7, ^a CH	7.50, d (8.5)	131.2, CH	7.42, d (8.7)	131.2, CH	7.42, d (8.5)
6	115.3, CH	6.80, d (8.7)	115.3, CH	6.80, d (8.5)	115.3, CH	6.79, d (8.5)	115.2, CH	6.78, d (8.7)	115.2, CH	6.78, d (8.5)
7	158.5, C		158.5, C		158.5, C		158.0, C		158.1, C	
7-OH		9.94, br s		10.02, br s		10.03, br s		10.00, br s		10.03, br s
1'	172.0, C		171.9, C		171.8, C		172.0, C		172.0, C	
2'	35.1, CH ₂	2.28, t (7.2)	35.2, CH ₂	2.28, t (7.4)	35.2, CH ₂	2.28, t (7.2)	35.2, CH ₂	2.32, t (7.5)	35.2, CH ₂	2.32, t (7.5)
3'	25.0, CH ₂	1.59, m	25.0, CH ₂	1.59, m	25.0, CH ₂	1.59, m	24.7, CH ₂	1.58, m	24.7, CH ₂	1.58, m
4'	28.6-29.0, CH ₂	1.31, m ^a	28.7, CH ₂	1.34, m ^a	28.6-28.8, CH ₂	1.31, m ^a	28.7-28.9, CH ₂	1.31, m ^a	28.8, CH ₂	1.32, m ^a
5'	28.6-29.0, CH ₂	1.31, m ^a	29.1, CH ₂	1.30, m ^a	28.6-28.8, CH ₂	1.31, m ^a	28.7-28.9, CH ₂	1.31, m ^a	29.1, CH ₂	1.28, m ^a
6'	28.6-29.0, CH ₂	1.31, m ^a	26.7, CH ₂	1.30, m ^a	28.6-28.8, CH ₂	1.31, m ^a	28.7-28.9, CH ₂	1.31, m ^a	26.7, CH ₂	1.30, m ^a
7'	28.6-29.0, CH ₂	1.31, m ^a	38.4, CH ₂	1.19, m	31.2, CH ₂	1.30, m ^a	28.7-28.9, CH ₂	1.31, m ^a	38.4, CH ₂	1.19, m
8'	31.3, CH ₂	1.29, m ^a	27.4, CH ₂	1.55, m	22.1, CH ₂	1.32, m ^a	31.3, CH ₂	1.29, m ^a	27.4, CH ₂	1.55, m
9'	22.1, CH ₂	1.30, m ^a	22.5, CH ₃	0.90, d (6.6)	13.9, CH ₃	0.90, t (7.0)	22.1, CH ₂	1.30, m ^a	22.5, CH ₃	0.89, d (6.5)
10'	13.9, CH ₃	0.90, t (7.0)					13.9, CH ₃	0.90, t (7.0)		

^aOverlapped signal





Figure 1. Key COSY, HMBC, and NOESY correlations for thalassotalic acid A (**1**).




Table of Contents/Abstract Graphic

