

2020

## 22. A glimpse into relativity

Gerhard Müller  
*University of Rhode Island, gmuller@uri.edu*

Robert Coyne  
*University of Rhode Island, robcoyne@uri.edu*

Follow this and additional works at: <https://digitalcommons.uri.edu/phy204-slides>

### Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/).

---

### Recommended Citation


Müller, Gerhard and Coyne, Robert, "22. A glimpse into relativity" (2020). *PHY 204: Elementary Physics II -- Slides*. Paper 47.  
<https://digitalcommons.uri.edu/phy204-slides/47>

This Course Material is brought to you for free and open access by the PHY 204: Elementary Physics II (2021) at DigitalCommons@URI. It has been accepted for inclusion in PHY 204: Elementary Physics II -- Slides by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

# Is There Absolute Motion?



Forces between two long, parallel, charged rods




- $\frac{F_E}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1\lambda_2}{d}$  (left),  $\frac{F_E^*}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1^*\lambda_2^*}{d}$ ,  $\frac{F_B}{L} = \frac{\mu_0}{2\pi} \frac{I_1I_2}{d}$ , (right)
- $\frac{F_E^* - F_B}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1^*\lambda_2^*}{d} \left(1 - \frac{v^2}{c^2}\right) = \frac{1}{2\pi\epsilon_0} \frac{\lambda_1\lambda_2}{d}$
- $c = \frac{1}{\sqrt{\epsilon_0\mu_0}} = 2.998 \times 10^8 \text{ms}^{-1}$  (speed of light)
- $\lambda_1^* = \frac{\lambda_1}{\sqrt{1 - v^2/c^2}}$ ,  $\lambda_2^* = \frac{\lambda_2}{\sqrt{1 - v^2/c^2}}$  (due to length contraction)

# Catching Up with a Photon? (1)




Forces between two long, parallel, charged rods in relative motion.



- Galilean kinematics predicts  $u = v_1 + v_2$ .
- Relativistic kinematics requires  $v_1 < c, v_2 < c, u < c$ .
- Relativistic dynamics requires  $F_E^* - F_B = F_E$ .
- Length-contracted charge densities:  $\lambda(v) = \frac{\lambda(0)}{\sqrt{1 - v^2/c^2}}$ .
- Electric currents:  $I(v) = \lambda(v)v$ .

## Catching Up with a Photon? (2)



- $\frac{F_E}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda(0)\lambda(v_1)}{d}, \quad \frac{F_E^*}{L} = \frac{1}{2\pi\epsilon_0} \frac{\lambda(v_2)\lambda(u)}{d}.$
- $\frac{F_B}{L} = \frac{\mu_0}{2\pi} \frac{[\lambda(v_2)v_2][\lambda(u)u]}{d} = \frac{1}{2\pi\epsilon_0} \frac{\lambda(v_2)\lambda(u)}{d} \frac{v_2 u}{c^2}.$
- $\frac{F_E^* - F_B}{L} = \frac{F_E}{L} \Rightarrow \frac{1}{2\pi\epsilon_0} \frac{\lambda(v_2)\lambda(u)}{d} \left(1 - \frac{v_2 u}{c^2}\right) = \frac{1}{2\pi\epsilon_0} \frac{\lambda(0)\lambda(v_1)}{d}$
- $\Rightarrow \frac{1}{\sqrt{1 - v_2^2/c^2}} \frac{1}{\sqrt{1 - u^2/c^2}} \left(1 - \frac{v_2 u}{c^2}\right) = \frac{1}{\sqrt{1 - v_1^2/c^2}}$  to be solved for  $u$ .
- Relativistic kinematics predicts  $u = \frac{v_1 + v_2}{1 + v_1 v_2 / c^2} < c.$



Four friends wish to cross a river at night on a narrow and treacherous bridge.

They have one flashlight.

To walk across the bridge it takes

- Gougouma 1minute,
- Ndakta 2 minutes,
- Maïtäina 5 minutes,
- Käissebo 10 minutes.

The bridge supports not more than two persons simultaneously.

The use of the flashlight is essential on every trip across the bridge.

The duration of any trip is dictated by the slower person.

Describe the sequence of trips that minimizes the total time for the four friends to make it to the other side of the river.