

3-1-2015

Hemlock Woolly Adelgid Alters Fine Root Bacterial Abundance and Mycorrhizal Associations in Eastern Hemlock

Justin F. Vendettuoli

David A. Orwig

Jennifer Adams Krumins

Matthew D. Waterhouse

Evan L. Preisser

University of Rhode Island, preisser@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/bio_facpubs

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

JF Vendettuoli, DA Orwig, J Adams Krumins, MD Waterhouse, and EL Preisser. (2015). "Hemlock woolly adelgid alters fine root bacterial abundance and mycorrhizal associations in eastern hemlock." *Forest Ecology and Management*. 339: 112-116. Available at: <http://www.sciencedirect.com/science/article/pii/S0378112714007178>.

This Article is brought to you for free and open access by the Biological Sciences at DigitalCommons@URI. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

1 Running head: Underground effects of adelgid

2

3 TITLE: Hemlock woolly adelgid alters fine root bacterial abundance and mycorrhizal
4 associations in eastern hemlock

5

6 Justin F. Vendettuoli¹, David A. Orwig², Jennifer Adams Krumins³, Matthew D.
7 Waterhouse⁴, and Evan L. Preisser^{5*}

8

9 ¹Department of Natural Resources Science, University of Rhode Island, Kingston, RI
10 02881

11 ²Harvard Forest, Harvard University, Petersham MA 01366

12 ³Department of Biology and Molecular Biology, Montclair State University, Montclair
13 NJ 07043

14 ⁴Department of Biology, University of British Columbia, Kelowna BC V1V 1V7

15 ⁵Department of Biological Sciences, University of Rhode Island, Kingston RI 02881

16

17 *Corresponding author: Evan Preisser

18 Department of Biological Sciences, University of Rhode Island, Kingston RI 02881

19 Ph: 401-874-2120 e-mail: preisser@uri.edu

20

21 **KEYWORDS**

22 *Adelges tsugae*, *Tsuga canadensis*, mycorrhizae, fine root bacteria, fine root

23 carbon:nitrogen

24 **ABSTRACT**

25 While the impact of aboveground herbivores on plant biomass and fitness has received
26 considerable attention, there has been far less research on the corresponding belowground
27 impacts. The belowground effects of aboveground feeding may be particularly noticeable for
28 invasive and/or outbreaking herbivore species that reach high densities and can cause major
29 damage and sometimes death. The hemlock woolly adelgid, *Adelges tsugae*, is an invasive pest
30 on the eastern seaboard of the United States that feeds on a native shade-tolerant conifer, the
31 eastern hemlock *Tsuga canadensis*. Trees rapidly decline and die following infestation, and the
32 invasion of this insect has devastated hemlock populations from Georgia in the south to Maine in
33 the north. Despite their substantial impact on tree health, we are unaware of any research into the
34 adelgid's effect on hemlock roots and the surrounding rhizosphere. We report the results of
35 research assessing ectomycorrhizal root colonization, rhizosphere bacterial abundance, and root
36 C:N ratios of infested and uninfested *T. canadensis*. We found that adelgid infestation decreased
37 the percentage of root material colonized by ectomycorrhizal fungi by more than 67%.
38 Rhizosphere bacterial abundance on fine roots was 25% lower on adelgid-infested versus
39 uninfested trees, and roots of adelgid-infested trees contained significantly less carbon. Our
40 results demonstrate that aboveground adelgid infestation can affect hemlock root composition
41 and alter belowground interactions with ectomycorrhizal fungi and bacteria. This information
42 demonstrates that above-belowground linkages can transmit the impact of herbivory far from the
43 site of localized damage.

44 **1.0 INTRODUCTION**

45 The impact of insect herbivores on plant growth and community structure can range from
46 inconsequential to major; in extreme cases the structure and functions of entire ecosystems can

47 be substantially altered (Lovett *et al.*, 2006). In some instances, insect herbivores can increase
48 biodiversity via preferential feeding on dominant species, allowing resources to be exploited by a
49 greater number of species (Carson and Root, 2000). In other cases where herbivores inflict
50 substantial damage, outbreaks of such species can devastate their hosts and cause major changes
51 to the environment (Smith and Schowalter, 2001; Gandhi and Herms, 2010) and economic loss
52 (Aukema *et al.*, 2011; Oliveira *et al.*, 2013).

53 Researchers are increasingly aware that herbivore grazing on aboveground green biomass
54 can have profound belowground impacts on the composition of organisms and subsequent
55 nutrient cycling in the rhizosphere (Bardgett and Wardle, 2010). The herbivore removal of
56 aboveground plant tissue can alter patterns of carbon and nutrient allocation in belowground
57 roots (Rasmann *et al.*, 2009). This can affect the composition and abundance of rhizosphere-
58 dwelling organisms. Cattle grazing, for instance, has been shown to alter microbial community
59 composition and food web structure in the root zone of grass (Hamilton and Frank, 2001; Veen
60 *et al.*, 2010). Moose and snowshoe hare grazing have also led to reduced ectomycorrhizal (EM)
61 colonization in roots of balsam poplar (*Populus balsamifera*) and willow (*Salix* spp.) (Rossow *et*
62 *al.*, 1997). These changes can in turn alter rates of nutrient cycling and availability in ways that
63 affect the grazed plant. This feedback loop has been well-documented and can be surprisingly
64 favorable for plant growth and recovery (Ruess and McNaughton, 1987; Krumins, 2014).

65 While the effect of folivory on aboveground-belowground interactions has been well-
66 studied in herbaceous plants, the impact of sap-feeding herbivores on aboveground-belowground
67 interactions in woody plant species has attracted less attention. This gap is notable in light of
68 work documenting that sap-feeder impacts on woody plant fitness equal or exceed those of
69 folivores (Zvereva *et al.*, 2010). Aphid infestations, for instance, indirectly reduce root growth in

70 Sitka spruce by limiting the tree's ability to provide carbon from photosynthesis (Day and
71 Cameron, 1997) and reduce root tissue density of Douglas fir by inducing the translocation of
72 additional carbohydrates from roots to shoots (Smith and Schowalter, 2001). Herbivory by the
73 needle-feeding scale *Matsucoccus acalyptus* reduced EM colonization of pinyon pine (Gehring
74 and Whitham, 1991); similarly, western spruce budworm (*Choristoneura occidentalis*)
75 defoliation reduced EM colonization in Douglas fir seedlings (Kolb *et al.*, 1999). More
76 generally, both folivores and sap feeders can also impact epiphytic microbial communities
77 through their production of nutrient-rich excrement (Stadler *et al.*, 2001).

78 The hemlock woolly adelgid (*Adelges tsugae*; 'adelgid') is an invasive sessile herbivore
79 that feeds exclusively on eastern hemlock (*Tsuga canadensis* (L.) Carr.) in the northeastern
80 United States. While it has minimal impact on hemlock health in its native range of Japan and
81 China (Havill *et al.*, 2006), it can kill even mature eastern hemlocks in as little as four years; few
82 heavily-infested trees survive longer than ten years (Orwig and Foster, 1998). While the impact
83 of the adelgid on hemlock physiology has been studied (Radville *et al.*, 2011; Gómez *et al.*,
84 2012; Gonda-King *et al.*, 2012; Domec *et al.*, 2013), its effects on hemlock root physiology and
85 the associated rhizosphere remain unexplored.

86 We report the results of work exploring the belowground impact of adelgid infestation on
87 *T. canadensis*. We measured root EM associations of infested and uninfested hemlock trees,
88 rhizosphere bacterial abundance, and root C:N ratios. Our findings demonstrate that
89 aboveground adelgid infestation of eastern hemlock has belowground consequences that likely
90 augment the ecosystem-level impact of this pest and may need to be addressed for the maximal
91 success of forest restoration efforts.

92 **2.0 METHODS**

93 As part of a long-term research program addressing the impacts of the hemlock woolly
94 adelgid on eastern hemlock, we characterized adelgid infestation level and stand vigor in 79
95 stands in CT and 63 stands in MA. Following their initial characterization (1997-1998 in CT,
96 2002-2004 in MA; Orwig *et al.*, 2002; Orwig *et al.*, 2012), these stands were repeatedly surveyed
97 for adelgid infestation and stand vigor in 2005, 2007, 2009, and 2011 (Preisser *et al.*, 2008;
98 Preisser *et al.*, 2011). We assessed the rhizosphere of eastern hemlocks in a subset of the stands
99 described above. In order to explore how adelgid infestation affects rhizosphere processes in
100 eastern hemlock, we characterized both fine-root EM colonization (Study #1) and bacterial
101 abundance (Study #2).

102 In Study #1, we quantified EM colonization of hemlock roots in three adelgid-infested
103 and two uninfested hemlock stands in 2003; we focused our research on ectomycorrhizae as
104 opposed to arbuscular (endo) mycorrhizae because conifers like hemlock are almost exclusively
105 colonized by ectomycorrhizae (Smith *et al.*, 1997). Infested stands were located in south-central
106 Connecticut, and had been colonized by adelgid for 3-10 years; uninfested forests were located
107 in central Massachusetts (Fig. 1). In Study #2, we quantified rhizosphere bacterial abundance
108 from ten infested and ten uninfested hemlock stands in central and northern Massachusetts in
109 2013 (Fig. 1). In both studies, trees sampled in the ‘infested’ treatment were heavily infested
110 themselves (>1 mature adelgid/cm foliage growth) and surrounded by other heavily-infested
111 trees. In contrast, trees sampled in the ‘uninfested’ treatments were in stands where no adelgid
112 had been detected during previous large-scale surveys; to ensure that the sampled trees had
113 maintained their uninfected status, we carefully surveyed each sampled tree and all trees within
114 10m of it to ensure the absence of adelgid. Because of logistical constraints, we were unable to
115 sample the same sites in both surveys. Regardless of location, all sampled stands consisted of

116 >50% hemlock canopy cover within 100m of the sampled trees, and 100% canopy cover within
117 10 m of the sampled trees.

118 *2.1. Study 1: Ectomycorrhizal colonization of hemlock roots*

119 From each of the five sites, we collected roots from 3–8 hemlock saplings (1.5-2m in
120 height); we sampled a total of 14 saplings from infested sites and 16 from uninfested sites. Soil
121 and stand traits were sampled in the course of several related studies (Cobb *et al.*, 2006; Orwig *et*
122 *al.*, 2013); infested sites averaged 708 ± 19 [SE] trees ha⁻¹, with a mean hemlock stand basal area
123 of 46.6 ± 2.7 m² ha⁻¹; uninfested sites averaged 1072 ± 97 [SE] trees ha⁻¹, with a mean hemlock
124 stand basal area of 45.6 ± 2.0 m² ha⁻¹. soils in infested sites had a mean organic (forest floor) soil
125 C:N ratio of $26.1 \pm 2.3\%$ [SE] and a mean mineral soil c:n ratio of $23.2 \pm 0.65\%$. Soils in
126 uninfested sites had a organic (forest floor) soil C:N ratio of 26.9% and a mineral soil c:n ratio of
127 24.1% (also see online appendix #1). Only trees growing under a hemlock-dominated canopy
128 were sampled. Each sapling was uprooted to expose the entire root system, and three 20cm root
129 samples per tree were collected and rinsed with deionized water (four of 30 sampled trees only
130 had sufficient roots for two 20cm root samples). Although we would have liked to sample mature
131 trees, the labor involved in uprooting multiple large hemlock trees (necessary to ensure that the
132 sampled roots in fact belonged to the chosen tree) necessitated using saplings. Each root sample
133 was assigned a random number to ensure an unbiased assessment and then trimmed down to a
134 5cm section. The grid intercept method (Giovanetti and Mosse, 1980) was used to assess the
135 percent EM colonization for each root sample. Roots with EM colonization were differentiated
136 on the basis of morphology, color, characteristics of the surface of the hyphal mantle, and planar
137 views of different mantle layers using standard methods (Agerer, 1992). Each root sample was
138 randomly dispersed in a 9cm diameter petri plate with 0.5cm grid lines. The intersection between

139 grid lines and roots were designated as either EM-colonized or non-mycorrhizal. The proportion
140 of root counts that were mycorrhizal was calculated for each root sample and averaged for each
141 sapling. We took a total of 5,902 counts, an average of 197 counts per tree (each tree had a
142 minimum of 100 counts).

143 2.2 Study 2: Bacterial abundance in the rhizosphere

144 We collected fine roots ($\leq 2\text{mm}$ diameter; Robertson *et al.*, 1999) from three understory
145 hemlocks (2 – 5m tall) at each of twenty sites (10 infested and 10 uninfested); all sampled trees
146 were growing under a hemlock-dominated canopy. Again, our choice of the sampled trees was
147 motivated by the difficulties inherent in uprooting multiple large hemlock trees (necessary to
148 ensure that the sampled roots in fact belonged to the chosen tree). Infested sites averaged $1312 \pm$
149 240 [SE] trees ha^{-1} , with a mean hemlock basal area of $33.2 \pm 1.5 \text{ m}^2 \text{ ha}^{-1}$; uninfested sites
150 averaged 859 ± 78 [SE] trees ha^{-1} , with a mean hemlock basal area of $38.2 \pm 1.7 \text{ m}^2 \text{ ha}^{-1}$. Soils in
151 infested sites had a mean organic (forest floor) soil C:N ratio of $26.9 \pm 1.4\%$ [SE] and a mean
152 mineral soil c:n ratio of $24.5 \pm 2.0\%$. Soils in uninfested sites had a mean organic soil C:N ratio
153 of $28.8 \pm 1.6\%$ and a mean mineral soil c:n ratio of $24.5 \pm 1.6\%$ (also see online appendix #1; for
154 a more detailed site description, see Orwig *et al.*, 2012). Roots were collected from each
155 hemlock by lightly scraping away leaf litter and organic soil from the base of the tree, extracting
156 the roots, and clipping three 12-15cm root samples. Root samples were combined in a single
157 plastic bag (one bag per tree), immediately returned to the lab, and refrigerated for <2 hours prior
158 to fixation. Before fixing each sample, loose debris and soil was manually shaken off; $\sim 0.1\text{g}$ of
159 fine roots from each tree was then placed in 5mL of phosphate buffer saline (PBS) solution and
160 vortexed for 2 minutes to suspend the bacteria. Each suspension was fixed with 1.5% filtered

161 formalin (1.5% final formalin concentration) and again vortexed. Fixed samples were stored at
162 4°C prior to staining and enumeration.

163 Staining took place within three days of the sample being fixed and collected; because the
164 samples were fixed immediately after collection, there were no time-related differences between
165 bacterial counts taken on different days. Bacterial abundance was determined using acridine
166 orange direct counts (AODC) (Kepner and Pratt, 1994). Cell concentration was optimized by
167 dilution to achieve countable samples. Between 0.1-0.5ml of each fixed sample was removed
168 (the extracted amount was supplemented with PBS to ensure a total volume of 1ml) and then
169 stained with 200µl of 0.1% acridine orange. Each sample was vacuum-captured onto a 0.2µm
170 black polycarbonate filter (EMD Millipore Corporation, MA) and fixed to a slide with
171 immersion oil. Enumeration was done using a Nikon Eclipse Ti inverted fluorescent microscope
172 fixed with a Nikon DS-Fi1 camera. We determined mean bacterial abundance per slide by
173 photographing ten randomly-chosen locations on each slide, using ImageJ (Schneider et al.
174 2012) to count bacterial densities, and averaging the ten counts. Each count thus reflects the
175 mean number of cells per g fresh weight of plant material. Because the extraction procedures
176 involved washing and removing fresh plant tissue, we could not calculate the dry weight of plant
177 material.

178 We used the procedures outlined above to collect an additional fine root sample in 2013
179 for carbon and nitrogen analysis from each tree used for bacterial abundance at the 20 sites. Fine
180 roots were rinsed with deionized water to remove soil particles and dried at 60°C in an oven for
181 48 hrs. Dried samples were ground into a fine powder using a grinding mill (Spex Mixer Mill
182 8000M, Metuchen, NJ) and analyzed for carbon and nitrogen content with a nutrient analyzer

183 (Elementar vario MICRO cube, Mount Laurel, NJ). Two replicates of each fine root sample were
184 analyzed, and the results averaged to calculate a site-level mean.

185 *2.3 Statistical Analysis:* To test the effect of adelgid infestation on EM colonization,
186 bacterial abundance, and root C:N, all samples were grouped by site and the infested and control
187 sites were compared. Because the data from Study #1 did not meet the assumptions of normality,
188 we analyzed it using a non-parametric median test; data from Study #2 was analyzed using one-
189 way ANOVA. EM colonization was calculated as percent colonization and bacterial abundance
190 was calculated as number of cells per gram fresh weight of root. All analyses were performed
191 using JMP 9.0 (SAS, Cary, NC).

192 **3.0 RESULTS**

193 The percentage of EM colonization was significantly lower ($10.6 \pm 2.4\%$ [SE]) on infested
194 hemlock roots compared to roots from uninfested trees ($32.4 \pm 10.2\%$) (Median Test; $p=0.046$).
195 Hemlock fine root bacterial abundance, measured in millions of cells per gram, was also lower
196 on infested versus uninfested trees ($F_{1,18}=2.22$, $p=0.044$; Fig. 2A). Chemical analysis of fine
197 roots from adelgid-infested versus uninfested trees revealed that percent carbon was significantly
198 lower in infested hemlock stands ($F_{1,18}=5.11$, $p=0.036$; Fig. 2B), but that adelgid infestation did
199 not affect percent nitrogen (Fig. 2C). Despite the differences in root C, roots from infested versus
200 uninfested stands did not differ in their root C:N ratio (Fig. 2D).

201 **4.0 DISCUSSION**

202 We found that aboveground infestation by hemlock woolly adelgid significantly affected
203 rhizosphere processes. The rhizosphere surrounding fine roots of adelgid-infested trees had less
204 ectomycorrhizal colonization and lower bacterial abundance (Fig. 2A), while the fine roots
205 themselves had lower carbon concentrations (Fig. 2B). The lower percentage of ectomycorrhizal

206 colonization found in our results is consistent with findings following herbivory from several
207 types of insects (Gehring and Whitham, 1994b). One of the most common causes of decreased
208 ectomycorrhizal colonization is herbivory-driven reductions in photosynthate availability; this
209 can disrupt carbohydrate supply to mycorrhizae and reduce mycorrhizal root tip abundance
210 (Gehring and Whitham, 1994b) and mycorrhizal inoculum potential (Lewis *et al.*, 2008). Aphid
211 outbreaks on other conifer species, for instance, can decrease photosynthetic efficiency (Day and
212 Cameron, 1997). The hemlock woolly adelgid has been shown to cause similar reductions in
213 photosynthesis (Nelson *et al.*, 2014), and continued feeding on hemlock likely leads to
214 continuous carbohydrate depletion. It is possible that the reduction in carbon found in the fine
215 roots of infested trees (Fig. 2B) and lower EM colonization result from a disruption of that
216 carbon source.

217 Environmental conditions can also play a significant role in the mycorrhizal response to
218 herbivory (Bardgett and Wardle, 2003; Gehring and Whitham, 2003). There has been a
219 documented decrease in EM colonization in sites with higher soil nutrient status compared with
220 water- and nutrient-stressed sites (Gehring and Whitham, 1994a). Several environmental
221 characteristics of chronic HWA infestation may have impacted the EM colonization response
222 seen in this study. First, Stadler *et al.* (2005; 2006) provided evidence that adelgid impacts the
223 composition of throughfall in infested stands. Their work showed higher inputs of N into the soil
224 under adelgid-infested trees. Second, in addition to inputs from throughfall, adelgid damage
225 often yields microenvironmental conditions that lead to increased soil N due to changes in
226 decomposition, N cycling and availability, and reduced tree uptake of nutrients (Kizlinski *et al.*
227 2002; Orwig *et al.* 2008). Enhanced soil N status resulting from either of these mechanisms may
228 also have led to reduced colonization of hemlock fine roots and associated bacterial levels.

229 Finally, altered mycorrhizal community structure resulting from increased N in the system may
230 have selected for ectomycorrhizal species that perform less of a service for their host but still
231 require the same carbon cost (Johnson, 1993).

232 We expected the abundance of bacterial cells colonizing the roots of adelgid-infested
233 trees to be higher. Experiments in grasslands ecosystems showed that grazed plants exude more
234 carbon into the rhizosphere, thus stimulating microbial growth and metabolism (Hamilton and
235 Frank, 2001). The priming effect (Kuzyakov *et al.*, 2000) stimulated by above-ground herbivory
236 fed back positively to the grasses, increasing nutrient availability. Rhizosphere microbes have the
237 primary responsibility for making nutrients available to plants through the decomposition and
238 mineralization of soil organic matter (Vessey, 2003). It is interesting to note that we did not find
239 increased microbial abundance in the trees we studied. This difference may be due to important
240 differences in the feeding behavior of herbivores and how they affect the host plant (Lovett *et al.*,
241 2006). Sucking insects, for instance, export far less wasted plant biomass (increasing soil organic
242 matter) and frass to the forest floor than do chewing insects or larger grazers (Zvereva *et al.*,
243 2010).

244 Bacterial and fungal communities are often tightly coupled in the rhizosphere. The
245 reduced bacterial abundance found in adelgid-infested hemlock stands may simultaneously be
246 linked to reduced ectomycorrhizal associations and changes in root and soil nutrient chemistry
247 associated with infestation. Indeed, some studies have found that soil communities experiencing
248 mycorrhizal loss also lose their fungally-associated bacteria (Hol *et al.*, 2014). However, the
249 significant decline in absolute numbers of bacteria suggests resource limitation from the root.
250 The implications of decreased bacterial load will be decreased mineral nutrient availability to the
251 tree (Vessey, 2003; Wardle *et al.*, 2004); this is consistent with our results (Fig. 2A). Knowing

252 both the load of EM and bacteria in the rhizosphere is, however, only the first critical step. Soil
253 communities are classically divided into bacterial- and fungal-based energy channels (Moore and
254 Hunt, 1988; De Ruiter *et al.*, 1995). It is the balance of these two energetic pathways, however,
255 that leads to stability and functioning of the rhizosphere community (Rooney *et al.*, 2006) and
256 overall plant health. Therefore, future work will test differences in community composition of
257 both mycorrhiza and bacteria colonizing the roots of trees in affected and unaffected sites. The
258 reduced colonization by EM and bacterial abundance found here may be caused by strong
259 competitors dominating the rhizosphere community.

260 Our work provides the first documentation of the below-ground consequences of above-
261 ground herbivory on eastern hemlock by an exotic herbivore. The impact of aboveground
262 feeding by the hemlock woolly adelgid on the rhizosphere processes of the hemlock and the
263 mycorrhizal and bacterial abundances illustrates the need for a greater understanding of how
264 herbivores impact all aspects of an ecosystem. From a management perspective, there is
265 increased recognition (Kardol and Wardle, 2010) of the importance of aboveground-
266 belowground linkages in determining the efficacy of management and restoration efforts.
267 Specifically, adelgid-mediated alterations in the belowground communities that facilitate
268 hemlock growth could make it more difficult to replant hemlocks in formerly-suitable areas.
269 Further research should investigate the soil community structure of infested hemlock stands
270 against uninfested hemlock stands, to see if there are shifts in the species found in addition to the
271 change in abundance of bacteria.

272 **5.0 ACKNOWLEDGEMENTS**

273 We thank M. Patel for her assistance with the lab work and C. Renaud for help
274 processing samples. The manuscript was greatly improved by by the comments and suggestions

275 of two anonymous reviewers. This work was funded by RI00HI-4004 to EP, NIFA 2011-67013-
276 30142 to EP and DO, NSF funding through its programs in Research Experience for
277 Undergraduates (DBI 10-03938), Long Term Ecological Research (DEB 12-37491), and is a
278 publication of the Harvard Forest Long Term Ecological Research (LTER) Site.

279 **6.0 LITERATURE CITED**

- 280 Agerer, R., 1992. Ectomycorrhizae of *Phellodon niger* on Norway spruce and their
281 chlamydospores. *Mycorrhiza* 2, 47-52.
- 282 Aukema, J., Leung, B., Kovacs, K., Chivers, C., Britton, K., Englin, J., Frankel, S., Haight, R.,
283 Holmes, T., Liebhold, A., McCullough, D., Von Holle, B., 2011. Economic impacts of non-
284 native forest insects in the continental United States. *PLoS ONE* 6, e24587.
- 285 Bardgett, R.D., Wardle, D.A., 2003. Herbivore-mediated linkages between aboveground and
286 belowground communities. *Ecology* 84, 2258-2268.
- 287 Bardgett, R.D., Wardle, D.A., 2010. Aboveground-Belowground Linkages: Biotic Interactions,
288 Ecosystem Processes, and Global Change. Oxford University Press.
- 289 Carson, W.P., Root, R.B., 2000. Herbivory and plant species coexistence: community regulation
290 by an outbreaking phytophagous insect. *Ecol Monogr* 70, 73-99.
- 291 Cobb, R., Orwig, D., Currie, S., 2006. Decomposition of green foliage in eastern hemlock forests
292 of southern New England impacted by hemlock woolly adelgid populations. *Can. J. For. Res.* 36,
293 1331-1341.
- 294 Day, K.R., Cameron, A., 1997. Effect of contemporary infestation by the spruce aphid
295 (*Elatobium abietinum*) on root growth in Sitka spruce transplants. *Forestry* 70, 1-5.
- 296 De Ruiter, P., Neutel, A.-M., Moore, J.C., 1995. Energetics, patterns of interaction strengths, and
297 stability in real ecosystems. *Science* 269, 1257-1260.

298 Domec, J.-C., Rivera, L.N., King, J.S., Peszlen, I., Hain, F., Smith, B., Frampton, J., 2013.
299 Hemlock woolly adelgid (*Adelges tsugae*) infestation affects water and carbon relations of
300 eastern hemlock (*Tsuga canadensis*) and Carolina hemlock (*Tsuga caroliniana*). *New Phytol*
301 199, 452-463.

302 Gandhi, K., Herms, D., 2010. Direct and indirect effects of alien insect herbivores on ecological
303 processes and interactions in forests of eastern North America. *Biol. Invasions* 12, 389-405.

304 Gehring, C.A., Whitham, T.G., 1991. Herbivore-driven mycorrhizal mutualism in insect-
305 susceptible pinyon pine. *Nature* 353, 556-557.

306 Gehring, C.A., Whitham, T.G., 1994a. Comparisons of ectomycorrhizae on pinyon pines (*Pinus*
307 *edulis*; Pinaceae) across extremes of soil type and herbivory. *Am. J. Bot.* 81, 1509-1516.

308 Gehring, C.A., Whitham, T.G., 1994b. Interactions between aboveground herbivores and the
309 mycorrhizal mutualists of plants. *Trends Ecol Evol* 9, 251-255.

310 Gehring, C.A., Whitham, T.G., 2003. Mycorrhizae-Herbivore Interactions: Population and
311 Community Consequences. In: van der Heijden, M.G.A., Sanders, I.R. (Eds.), *Mycorrhizal*
312 *Ecology*. Springer Berlin Heidelberg, pp. 295-320.

313 Giovanetti, M., Mosse, B., 1980. An evaluation of techniques for measuring vesicular-arbuscular
314 mycorrhizal infection in roots. *New Phytol* 84, 489-500.

315 Gómez, S., Orians, C., Preisser, E., 2012. Exotic herbivores on a shared native host: tissue
316 quality after individual, simultaneous, and sequential attack. *Oecologia* 169, 1015-1024.

317 Gonda-King, L., Radville, L., Preisser, E., 2012. False ring formation in eastern hemlock
318 branches: impacts of hemlock woolly adelgid and elongate hemlock scale. *Environ Entomol* 41,
319 523-531.

320 Hamilton, E.W., Frank, D.A., 2001. Can plants stimulate soil microbes and their own nutrient
321 supply? Evidence from a grazing tolerant grass. *Ecology* 82, 2397-2402.

322 Havill, N., Montgomery, M., Yu, G., Shiyake, S., Caccone, A., 2006. Mitochondrial DNA from
323 hemlock woolly adelgid (Hemiptera: Adelgidae) suggests cryptic speciation and pinpoints the
324 source of the introduction to eastern North America. *Ann. Entomol. Soc. Am.* 99, 195-203.

325 Hol, W.H.G., de Boer, W., Medina, A., 2014. Beneficial Interactions in the Rhizosphere. In:
326 Dighton, J., Krumins, J.A. (Eds.), *Interactions in Soil: Promoting Plant Growth*. Springer
327 Netherlands, pp. 59-80.

328 Johnson, N.C., 1993. Can fertilization of soil select less mutualistic mycorrhizae? *Ecol. Appl.* 3,
329 749-757.

330 Kardol, P., Wardle, D.A., 2010. How understanding aboveground–belowground linkages can
331 assist restoration ecology. *Trends Ecol Evol* 25, 670-679.

332 Kepner, R.L., Pratt, J.R., 1994. Use of fluorochromes for direct enumeration of total bacteria in
333 environmental samples: past and present. *Microbiol Rev* 58, 603-615.

334 Kolb, T.E., Dodds, K.A., Clancy, K.M., 1999. Effect of western spruce budworm defoliation on
335 the physiology and growth of potted Douglas-fir seedlings. *For Sci* 45, 280-291.

336 Krumins, J.A., 2014. The Positive Effects of Trophic Interactions in Soil. In: Dighton, J.,
337 Krumins, J.A. (Eds.), *Interactions in Soil: Promoting Plant Growth*. Springer Netherlands, pp.
338 81-94.

339 Kuzyakov, Y., Friedel, J.K., Stahr, K., 2000. Review of mechanisms and quantification of
340 priming effects. *Soil Biol. Biochem.* 32, 1485-1498.

341 Lewis, J., Licitra, J., Tuininga, A., Sirulnik, A., Turner, G., Johnson, J., 2008. Oak seedling
342 growth and ectomycorrhizal colonization are less in eastern hemlock stands infested with
343 hemlock woolly adelgid than in adjacent oak stands. *Tree Physiology* 28, 629-636.

344 Lovett, G.M., Canham, C.D., Arthur, M.A., Weathers, K.C., Fitzhugh, R.D., 2006. Forest
345 ecosystem responses to exotic pests and pathogens in eastern North America. *Bioscience* 56,
346 395-405.

347 Moore, J.C., Hunt, H.W., 1988. Resource compartmentation and the stability of real ecosystems.
348 *Nature* 333, 261-263.

349 Nelson, L.A., Dillaway, D.N., Rieske, L.K., 2014. Effect of an exotic herbivore, *Adelges tsugae*,
350 on photosynthesis of a highly susceptible *Tsuga* host, with notes on conspecifics. *Arthropod-
351 Plant Interactions* 8, 9-15.

352 Oliveira, C.M., Auad, A.M., Mendes, S.M., Frizzas, M.R., 2013. Economic impact of exotic
353 insect pests in Brazilian agriculture. *J Appl Entomol* 137, 1-15.

354 Orwig, D., Foster, D., 1998. Forest response to the introduced hemlock woolly adelgid in
355 southern New England, USA. *J Torrey Bot Soc* 125, 60-73.

356 Orwig, D., Foster, D., Mausel, D., 2002. Landscape patterns of hemlock decline in New England
357 due to the introduced hemlock woolly adelgid. *J Biogeogr* 29, 1475-1487.

358 Orwig, D., Plotkin, A., Davidson, E., Lux, H., Savage, K., Ellison, A., 2013. Foundation species
359 loss affects vegetation structure more than ecosystem function in a northeastern USA forest.
360 *PeerJ* 1, e41.

361 Orwig, D., Thompson, J., Povak, N., Manner, M., Niebyl, D., Foster, D., 2012. A foundation tree
362 at the precipice: *Tsuga canadensis* health after the arrival of *Adelges tsugae* in central New
363 England. *Ecosphere* 3, 10.

364 Preisser, E., Lodge, A., Orwig, D., Elkinton, J., 2008. Range expansion and population dynamics
365 of co-occurring invasive herbivores. *Biol. Invasions* 10, 201-213.

366 Preisser, E., Miller-Pierce, M., Vansant, J., Orwig, D., 2011. Eastern hemlock (*Tsuga*
367 *canadensis*) regeneration in the presence of hemlock woolly adelgid (*Adelges tsugae*) and
368 elongate hemlock scale (*Fiorinia externa*). *Can. J. For. Res.* 41, 2433-2439.

369 Radville, L., Chaves, A., Preisser, E., 2011. Variation in plant defense against invasive
370 herbivores: evidence for a hypersensitive response in eastern hemlocks (*Tsuga canadensis*). *J*
371 *Chem Ecol* 37, 592-597.

372 Rasmann, S., Agrawal, A.A., Cook, S.C., Erwin, A.C., 2009. Cardenolides, induced responses,
373 and interactions between above- and belowground herbivores of milkweed (*Asclepias* spp.).
374 *Ecology* 90:2393-2404. *Ecology* 90, 2393-2404.

375 Robertson, G.P., Coleman, D.C., Bledsoe, C.S., Sollins, P. (Eds.), 1999. *Standard Soil Methods*
376 *for Long-term Ecological Research*. Oxford University Press, New York.

377 Rooney, N., McCann, K., Gellner, G., Moore, J.C., 2006. Structural asymmetry and the stability
378 of diverse food webs. *Nature* 442, 265-269.

379 Rossow, L.J., Bryant, J.P., Kielland, K., 1997. Effects of above-ground browsing by mammals
380 on mycorrhizal infection in an early successional taiga ecosystem. *Oecologia* 110, 94-98.

381 Ruess, R.W., McNaughton, S.J., 1987. Grazing and the dynamics of nutrient and energy
382 regulated microbial processes in the Serengeti grasslands. *Oikos* 49, 101-110.

383 Smith, J., Schowalter, T., 2001. Aphid-induced reduction of shoot and root growth in Douglas-fir
384 seedlings. *Ecol Entomol* 26, 411-416.

385 Smith, S.E., Read, D.J., Harley, J.L., 1997. *Mycorrhizal Symbiosis*. Academic Press, San Diego
386 CA.

387 Stadler, B., Müller, T., Orwig, D., 2006. The ecology of energy and nutrient fluxes in hemlock
388 forests invaded by the hemlock woolly adelgid. *Ecology* 87, 1792-1804.

389 Stadler, B., Müller, T., Orwig, D., Cobb, R., 2005. Hemlock woolly adelgid in New England
390 forests: canopy impacts transforming ecosystem processes and landscapes. *Ecosystems* 8, 233-
391 247.

392 Stadler, B., Solinger, S., Michalzik, B., 2001. Insect herbivores and the nutrient flow from the
393 canopy to the soil in coniferous and deciduous forests. *Oecologia* 126, 104-113.

394 Veen, G.F., Olf, H., Duyts, H., van der Putten, W.H., 2010. Vertebrate herbivores influence soil
395 nematodes by modifying plant communities. *Ecology* 91, 828-835.

396 Vessey, J.K., 2003. Plant growth promoting rhizobacteria as biofertilizers. *Plant Soil* 255, 571-
397 586.

398 Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., van der Putten, W.H., Wall, D.H.,
399 2004. Ecological linkages between aboveground and belowground biota. *Science* 304, 1629-
400 1633.

401 Zvereva, E., Lanta, V., Kozlov, M., 2010. Effects of sap-feeding insect herbivores on growth and
402 reproduction of woody plants: a meta-analysis of experimental studies. *Oecologia* 163, 949-960.

403


404 **7.0 Figure Legends**

405 Figure 1. Sites in Connecticut and Massachusetts used for the mycorrhizal (Study #1) and
406 root bacterial abundance (Study #2) surveys.

407 Figure 2. Mean bacterial abundance (A), mean percent carbon (B), mean percent nitrogen
408 (C), and mean carbon:nitrogen ratio (D) of eastern hemlock fine roots in 10 uninfested hemlock
409 stands and 10 adelgid-infested hemlock stands.

410

7.1 Figure 1.



414

7.2 Figure 2.

415

