

2015

Testing sample stability using four storage methods and the macroalgae *Ulva* and *Gracilaria*

Autumn Oczkowski

Carol S. Thornber

University of Rhode Island, thornber@uri.edu

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/bio_facpubs

**The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.**

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Oczkowski, A., Thornber, C. S., Markham, E. E., Rossi, R., Ziegler, A. and Rinehart, S. (2015), Testing sample stability using four storage methods and the macroalgae *Ulva* and *Gracilaria*. *Limnology and Oceanography: Methods*, 13: 9–14. doi: 10.1002/lom3.10002. Available: <http://onlinelibrary.wiley.com/doi/10.1002/lom3.10002/abstract>

This Article is brought to you for free and open access by the Biological Sciences at DigitalCommons@URI. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors

Autumn Oczkowski, Carol S. Thornber, Erin E. Markham, Ryann Rossi, Amanda Ziegler, and Shelby Rinehart

1 Testing sample stability using four storage methods and the macroalgae *Ulva* and
2 *Gracilaria*

3

4 Autumn Oczkowski^{1*}, Carol S. Thornber², Erin E. Markham^{1,3}, Ryann Rossi^{1,4}, Amanda
5 Ziegler^{2,5}, Shelby Rinehart^{2,6}

6

7 ¹U.S. Environmental Protection Agency, Atlantic Ecology Division, 27 Tarzwell Drive,
8 Narragansett, Rhode Island 02882

9 ²Department of Biological Sciences, University of Rhode Island, Kingston, Rhode Island
10 02881

11 ³Present address: Graduate School of Oceanography, University of Rhode Island,
12 Narragansett, Rhode Island 02882

13 ⁴Present address: Department of Applied Ecology, North Carolina State University,
14 Raleigh, North Carolina 27695

15 ⁵Present address: University of Hawaii at Manoa, Department of Oceanography,
16 Honolulu, Hawaii 96822

17 ⁶Present address: San Diego State University, Department of Biology, San Diego,
18 California 92182

19

20 *Corresponding author: Oczkowski.Autumn@epa.gov

21

22

23 Running Head: Stability of macroalgal samples for stable isotope analysis

24 Acknowledgements

25

26 Thank you to R. McKinney for assistance with sample analysis, and to J. Carey, R.
27 Johnson, and M. Pelletier for providing useful edits of this manuscript. Partial funding
28 was provided by the University of Rhode Island through an undergraduate research grant
29 to A. Ziegler. This material is based, in part, upon work supported in part by the National
30 Science Foundation EPSCoR Cooperative Agreement #EPS-1004057 and the State of
31 Rhode Island. This is ORD Tracking Number ORD-008193 of the Atlantic Ecology
32 Division, National Health and Environmental Effects Research Laboratory, Office of
33 Research and Development, U. S. Environmental Protection Agency. Although the
34 research described in this article has been funded in part by the U.S. Environmental
35 Protection Agency, it has not been subjected to Agency review. Therefore, it does not
36 necessarily reflect the views of the Agency.

37

38

39

40

41

42

43

44

45

46

47 Abstract

48

49 Concern over the relative importance of different sample preparation and storage
50 techniques frequently used in stable isotope analysis of particulate nitrogen ($\delta^{15}\text{N}$) and
51 carbon ($\delta^{13}\text{C}$) prompted an experiment to determine how important such factors were to
52 measured values in marine organisms. We stored the marine macroalgae *Ulva* and
53 *Gracilaria* in four different ways and analyzed replicates every three months over the
54 course of a year to assess treatment effects on stability. Treatments consisted of algae
55 dried at 65°C, ground to a powder, and stored in a desiccator until analysis; algae left in a
56 drying oven or in a freezer and processed (dried and ground) just prior to analysis, as well
57 as some dried, ground samples kept out in the lab and re-analyzed quarterly for 12
58 months. Concurrently, to assess the ecological range in isotope values over the course of
59 a year, samples were freshly collected from the same location and analyzed along with
60 the other treatments at each time step. Neither storage technique nor time had an impact
61 on either $\delta^{15}\text{N}$ or $\delta^{13}\text{C}$ values or the %N and %C of the algae tissues. There were clear
62 and consistent differences between species and some large seasonal differences in the
63 freshly collected samples. The interspecies differences and seasonal ranges of values
64 underscore the stability associated with method and duration of sample storage.

65

66

67 Key words: stable isotope, *Ulva*, *Gracilaria*, nitrogen, carbon

68

69

70 Introduction

71

72 Oftentimes laboratory procedures, like legends, are passed down from one analyst
73 to the next, as previous experiences have determined the methods necessary to obtain the
74 best results. However, sometimes the reasoning behind these methods is lost and a
75 reassessment is needed. In using stable isotopes of nitrogen ($\delta^{15}\text{N}$) and carbon ($\delta^{13}\text{C}$) in
76 our own work, we have followed procedures developed by colleagues as well as adopted
77 practices described in the literature. As ecologists, we frequently collect plant and animal
78 tissues, as well as sediment, from coastal areas which are then cleaned with deionized
79 water, dried in a 65°C oven, ground to a powder, and then analyzed on an isotope ratio
80 mass spectrometer. While the paradigm has always been to analyze the samples quickly
81 after collection, it has not always been feasible. Though taught to store samples in a
82 desiccator prior to analysis, the sheer number of samples has precluded this practice for
83 all samples. From issues like these arose concern about the stability of the samples with
84 respect to storage time and method. We conducted an experiment to test the stability of
85 samples of macroalgae commonly found in our region (Southern New England, *Ulva* and
86 *Gracilaria*) over the course of a year, under four different storage methods.

87 Typically, published methods call for samples to be dried in an oven (~60°C) for
88 24 hours or until dry (Oczkowski et al. 2008; Wozniak et al. 2006). But, it is unclear
89 whether samples can be dried for ‘too long,’ where extensive exposure to heat (days,
90 weeks, or months) would eventually enhance tissue breakdown and alter results. In
91 addition to examining the effects of dried, ground samples left in a desiccator and on the
92 bench-top (in sealed scintillation vials), we included a drying treatment where samples

93 were left in open aluminum weigh pans in a drying oven for up to one year. Finally, to
94 approximate a fresh sample, subsamples were frozen and individually defrosted, dried,
95 and ground within a week of analysis. To assess stability over time, some subsamples
96 were analyzed after our initial collection and then periodically over the course of a year.
97 If sample degradation were to occur, we could observe an increase in isotope value over
98 time as the lighter isotope might be preferentially lost (e.g., Fry 2006). We further
99 hypothesized that samples left on the counter might contain more water compared to
100 those in a desiccator, which could both facilitate the decomposition of the sample and
101 possibly distort the masses weighed for individual sample analyses, thus distorting the
102 measured %N and %C values. Also, if the long-term heat of the drying oven aided in the
103 breakdown and volatilization of N, we might expect to see a change in the $\delta^{15}\text{N}$ values
104 and a decrease in the %N. Our results (thankfully) indicate that the isotope and N and C
105 contents of the two macroalgae genera examined were stable over time and among
106 treatments. Given the range of ecological data, sample storage technique may have an
107 inconsequential impact on analytical outcome.

108

109

110 Materials and Procedures

111

112 Sample collection and processing

113 We collected 75 samples each of *Ulva rigida* C. Agardh and *Gracilaria*
114 *vermiculophylla* (Ohmi) Papenfuss from Oakland Beach, RI (41.68399, -71.39787) on
115 October 23, 2011. All algal thalli (individuals) were brought back to the lab and

116 immediately sorted to the species level, obvious epiphytes were removed, and algae were
117 rinsed with deionized water. Samples were allocated as follows for *Ulva* and *Gracilaria*:
118 twenty individuals of each species were cleaned, placed in sealed zipper bags, and placed
119 into a freezer (-20°C) until later analysis (hereafter 'freezer' samples; see Fig. 1 for
120 sample breakdown). The remaining fifty-five individuals of each species were cleaned,
121 placed into separate aluminum weighing dishes, and then into a drying oven at 60°C.
122 Once these were dry (after 2 days), fifteen were promptly removed, ground individually
123 into a fine powder with a mortar and pestle, and 2 to 3 mg of tissue from each sample
124 were placed into individual capsules for mass spectrometry analysis (Nov 9, 2011).
125 These fifteen specimens were randomly allocated as the initial samples for one of three
126 storage treatments (five for freezer, five for drying oven, and five for desiccator) for *Ulva*
127 and *Gracilaria* (Fig. 1). In addition to serving as 'initial' data points for the different
128 treatments, the five initial desiccator samples were left out on the counter and re-analyzed
129 at each subsequent time step. While this allowed us to look for changes over time in
130 samples stored on the counter, they were treated separately in statistical analyses (as
131 described below).

132 For each species, the remaining forty samples were divided into two equally sized
133 treatments named 'desiccator' and 'drying oven'. Desiccator samples were removed from
134 the drying oven, immediately ground into powder, and stored in twenty scintillation vials
135 in a laboratory desiccator. Drying oven samples remained as intact thalli in the drying
136 oven. At set time points (February, June, August, and November 2012 -- based in part on
137 mass spectrometer availability), we removed five individuals from each of the three
138 treatments, for each species, and analyzed them in a mass spectrometer. Prior to analysis,

139 frozen specimens were dried and ground, and drying oven specimens were ground. At
140 each subsequent time step at approximately three-month intervals (January 22, May 14,
141 July 17, and October 19, 2012), we collected five fresh individuals from each species
142 from Oakland Beach, cleaned them in the lab, and then dried, ground, and analyzed them
143 (hereafter 'freshly collected').

144 To address some questions that arose regarding initial $\delta^{15}\text{N}$ isotope values, we
145 collected five additional *Ulva* and *Gracilaria* samples (hereafter called addendum
146 samples) on 13 July 2013 and analyzed them first on 31 July 2013 and then again 23
147 September 2013. As described above, samples were dried, ground, and stored in acid-
148 washed scintillation vials on the counter until initial and then final analysis for $\delta^{15}\text{N}$
149 values.

150

151 Sample analysis

152 Samples were weighed into small tin capsules and analyzed on an Isoprime 100
153 mass spectrometer interfaced with a Micro Vario Elemental Analyzer (Elementar
154 Americas, Mt. Laurel, NJ) for $\delta^{15}\text{N}$, %N, $\delta^{13}\text{C}$, and %C. The nitrogen isotope
155 composition was expressed as a part per thousand (permil, ‰) deviation from air, while
156 the carbon was referenced to PeeDee Belemnite (PDB) where $\delta X = [(R_{\text{sample}} - R_{\text{standard}}) /$
157 $R_{\text{standard}}] \times 10^3$, X is $\delta^{15}\text{N}$ or $\delta^{13}\text{C}$, and R is the ratio of heavy to light isotope ($^{15}\text{N}:^{14}\text{N}$,
158 $^{13}\text{C}:^{12}\text{C}$). Samples were analyzed in triplicate and in batches of approximately 30
159 samples. Internal standards were used for check for instrument drift in each run and to
160 correct for instrument offset. The %N and %C was calculated by comparing the peak

161 area of the unknown sample to a standard curve of peak area vs. standard %N or %C
162 content.

163

164 Statistics

165 We analyzed the changes among treatments, between species, and across time in
166 $\delta^{15}\text{N}$, %N, $\delta^{13}\text{C}$, and %C of desiccator, drying oven, and freezer samples via a three-way
167 fixed factor ANOVA using JMP v11 statistical software (www.jmp.com). We analyzed
168 changes in the same four parameters for the freshly collected samples between species
169 and across time via a two-way fixed factor ANOVA. Changes in the 'counter' samples
170 over time and between species were analyzed with a two-way repeated measures
171 ANOVA (using 3, 6, 9, and 12 month data). Addendum samples were analyzed similarly
172 for $\delta^{15}\text{N}$ with a repeated measures ANOVA (using initial and 2 month data). All data
173 were checked for normality and homogeneity of variances and transformed where
174 appropriate.

175

176

177 Assessment

178

179 $\delta^{15}\text{N}$

180 The $\delta^{15}\text{N}$ values for the oven, desiccator, and freezer samples were, on average
181 2‰ lower in *Ulva* than *Gracilaria* (Table 1; $F_{1,120} = 153.66$, $p < 0.001$; Fig. 2). However,
182 there were no significant differences in $\delta^{15}\text{N}$ values across treatments (Table 1; $F_{2,120} =$
183 0.45, $p = 0.64$). Surprisingly, it does not seem to matter if macroalgae are left uncovered

184 in a drying oven, dried and ground in a desiccator, or in a freezer, prior to analysis, at
185 least in a southern New England climate. We found similar isotope values for samples
186 dried, ground, and left on a counter (Fig. 2). Because the counter samples were
187 reanalyzed repeatedly using material from the same vial, they could not be treated with
188 the same statistical techniques as the drying oven, desiccator, and freezer treatments.
189 Despite the statistical limitations in our ability to directly compare the counter samples to
190 the other treatments, they do not appear distinct from the others.

191 There was, however, a statistically significant difference in $\delta^{15}\text{N}$ among analysis
192 dates ($F_{1,120} = 17.89$; $p < 0.0001$) for the oven, desiccator, and freezer samples. Due to the
193 lack of a significant treatment main effect or interactions, we removed treatment from the
194 analyses and re-ran the $\delta^{15}\text{N}$ analyses separately for each species (as there was a
195 significant species by time interaction). We used time as the main effect to determine
196 which analysis dates differed (Underwood 1997). For *Gracilaria*, $\delta^{15}\text{N}$ values for the
197 initial samples were significantly higher than those measured at 3, 6, 9, and 12 months
198 ($p < 0.05$). The initial *Ulva* samples were not statistically different from the later
199 measurements ($p = 0.15$). Our 'counter' samples did not exhibit significant variability in
200 $\delta^{15}\text{N}$ across the study period (3 to 12 months; $F_{1,6, 13.2} = 3.78$; $p = 0.06$).

201 The higher initial *Gracilaria* and slightly, but not statistically, higher *Ulva* values
202 may be reflecting some instrument instability during the initial (Nov. 2011)
203 measurements of $\delta^{15}\text{N}$. As part of our analysis, we used a series of check standards (a
204 homogenized blue mussel tissue that is periodically internally calibrated to standard
205 reference material) interspersed throughout the run. These standards are used to calibrate
206 the reference gas and to check for any instrument drift. Typically, standard deviations

207 around these check standards are well below 0.3‰ and generally <0.2‰. In our initial
208 sampling, the check standards had an average value of 11.68 ± 0.64 ‰ (S.D.). However,
209 the cystine standard that we use to calibrate our %N measurements had a high
210 reproducibility (9.66 ± 0.054 ‰ S.D., n=4) and the offset between the cystine $\delta^{15}\text{N}$ values
211 measured in this run and the actual (calibrated to reference material) was the same as for
212 the blue mussel check standard, lending strength to the check standard. But, overall,
213 variability appeared to be higher in this initial run. To address this drop in $\delta^{15}\text{N}$ values
214 between initial and subsequent sampling, we collected additional samples in July 2013
215 and analyzed them 2 weeks and then 10 weeks after collection. The $\delta^{15}\text{N}$ values in what
216 we termed the addendum samples did not change significantly over time ($F_{1,8} = 0.70$, $p =$
217 0.43), lending support to our supposition that the originally higher initial *Gracilaria* $\delta^{15}\text{N}$
218 values were due to instrument performance.

219 By contrast, there were clear seasonal differences in $\delta^{15}\text{N}$ in freshly collected
220 macroalgae. With a range of 2‰ for *Ulva* and 4‰ for *Gracilaria*, the highest values were
221 in the late fall and lowest in the winter and spring ($F_{3,31} = 62.32$, $p < 0.0001$; Table 1),
222 with a significant interaction ($F_{3,31} = 7.04$, $p = 0.001$, Fig. 2), although there was no
223 difference between species ($F_{1,31} = 0.20$, $p = 0.66$). The wide range in the values of
224 freshly collected algae underscores the stability of the algae collected initially (23
225 October 2011), regardless of storage technique.

226

227 %N

228 As with $\delta^{15}\text{N}$, there were no significant differences in %N among frozen, oven,
229 and desiccator treatments (Table 1; $F_{2,120} = 0.30$, $p = 0.74$; Fig. 3), although %N was

230 significantly higher in *Gracilaria* ($F_{1,120} = 233.77$; $p < 0.0001$) and varied significantly
231 among sampling dates ($F_{4,120} = 2.89$, $p = 0.0252$). However, when we removed all
232 treatment terms and re-ran the analyses (as for $\delta^{15}\text{N}$ above), post-hoc comparisons did not
233 yield any dates that significantly differed in %N. By contrast, *Gracilaria* left on the
234 counter varied significantly among analysis dates ($F_{1.6, 12.69} = 101.52$, $p < 0.0001$; Fig. 3),
235 although there was not a consistent trend over time. The lowest values (at 6 months) may
236 have been associated with samples which were weighed to one less decimal place than
237 usual, increasing the uncertainty of the %N (and %C) values.

238 Overall, *Gracilaria* had about a third more N in their tissues than did *Ulva*
239 ($\sim 3.75\%$ vs. $\sim 2.5\%$, $p < 0.0001$; Table 1; Fig. 3). A recent assessment of *Ulva* and
240 *Gracilaria* in Narragansett Bay found %N ranging from 1 to 5%, with differences in
241 newly formed vs. mature tissues (Thornber et al. 2008). By contrast, our %N values are
242 lower than reported in some other areas for both species (e.g., Abreu et al. 2011; Barr et
243 al. 2013). The freshly collected samples showed a distinct seasonal pattern, where %N
244 was lowest in the spring and summer and highest in the fall and winter months ($F_{3,31} =$
245 173.94 , $p < 0.0001$; Table 1, Fig. 3). While we suspect these values may be reflecting
246 spring and summer water column nutrient depletion and winter luxury uptake, they
247 nonetheless indicate a dynamic environment.

248

249 $\delta^{13}\text{C}$

250 We did not find significant differences in the $\delta^{13}\text{C}$ content of algae among oven,
251 freezer, desiccator treatments, or among analytical dates (Table 1), although *Ulva* had
252 much higher $\delta^{13}\text{C}$ values (~ -10 ‰) than *Gracilaria* (~ -15 ‰; $F_{1,120} = 153.37$, $p < 0.0001$;

253 Fig. 4). By contrast, we did find significant differences in $\delta^{13}\text{C}$ in our counter specimens
254 that were repeatedly sampled ($F_{1,3,10.8} = 37.46$, $p < 0.0001$; Fig. 4), with a significant time
255 by species interaction ($F_{1,3,10.8} = 11.79$, $p = 0.004$).

256 There has been substantial detailed work in cataloging and interpreting
257 differences in C isotopes among species, as these values can be indicative of how the
258 species acquire C from the environment as well as their photosynthetic performance (for
259 example, see Fry and Sherr 1984; Raven et al. 1995, 2002). While these discussions are
260 beyond the scope of this paper, it is useful to note that our measured values indicate that
261 these species are capable of taking up both CO_2 and HCO_3^- although isotope differences
262 between the two forms of inorganic carbon does not indicate proportional uptake of either
263 carbonate species (Raven et al. 2002). While variable, other measurements of $\delta^{13}\text{C}$ values
264 from macroalgae in Narragansett Bay have ranged from -26 to -12 ‰ (Oczkowski et al.
265 2008). And, our freshly collected *Ulva* samples similarly ranged from -22.23 to -9.5‰
266 throughout the year. In contrast, *Gracilaria* was more homogenous, with mean values
267 ranging only from -15.43 to -13.96‰; values were significantly higher (less negative) for
268 *Gracilaria* than *Ulva* ($F_{1,31} = 5.67$, $p = 0.24$; Table 1), with significant variation among
269 sampling dates ($p < 0.0001$) and a significant species by time interaction ($p < 0.0001$).

270 Overall, while our measured *Gracilaria* values are typical for this region, *Ulva* values
271 from the initial (October 2011) collection were slightly higher than previously measured,
272 but not uncharacteristically so for macroalgae (Raven et al. 2002; Oczkowski et al. 2008).

273

274

275

276 %C

277 As with the other parameters measured, the %C of the macroalgae ($23.8 \pm 0.33\%$)
278 did not vary significantly among freezer, oven, and desiccator treatments ($F_{2,120} = 0.38$, p
279 $= 0.68$; Table 1; Fig. 5), nor over analysis dates ($F_{4,120} = 1.92$, $p = 0.11$), although the %
280 C was significantly higher in *Gracilaria* than *Ulva* (27.3% vs. 20.2%C; $F_{1,120} = 86.73$, $p <$
281 0.0001). The %C of the freshly collected samples was significantly higher in *Gracilaria*
282 than *Ulva* ($p < 0.0001$, Table 1), where %C of *Gracilaria* ranged from 24.76% to 31.35%
283 and *Ulva* from 20.33 to 23.57%. Samples from January were the highest, followed by
284 samples from October 2012, and then May and July 2012 ($p < 0.0001$, Table 1, Tukey
285 post-hoc comparisons).

286

287

288 Discussion

289

290 We chose to conduct an experiment to assess sample stability using several
291 common sample storage techniques. Using macroalgae, our results clearly indicate that
292 sample storage method has no bearing on the resultant $\delta^{15}\text{N}$, %N, $\delta^{13}\text{C}$, and %C values.
293 This is particularly surprising for those samples left in open weighing tins in a 65°C
294 drying oven for up to a year prior to analysis. We speculate that these results are
295 transferrable to many other plant tissues and maybe even to some animal tissues as well.

296

297

298

299 References

300

301 Abreu, M. H., R. Pereira, A. H. Buschmann, I. Sousa-Pinto, and C. Yarish. 2011.

302 Nitrogen uptake responses of *Gracilaria vermiculophylla* (Ohmi) Papenfuss under

303 combined and single addition of nitrate and ammonium. *J. Exp. Mar. Biol. Ecol.*

304 407:190-199.

305 Barr, N. G., B. D. Dudley, K. M. Rogers, and C. D. Cornelisen. 2013. Broad-scale

306 patterns of tissue $\delta^{15}\text{N}$ and tissue-N indicated in frondose *Ulva* spp.; Developing a

307 national baseline indicator of nitrogen-loading in coastal New Zealand. *Mar.*

308 *Pollut. Bull.* 67:203-216.

309 Fry, B. 2006. *Stable Isotope Ecology*. Springer.

310 Fry, B., and E. B. Sherr. 1984. $\delta^{13}\text{C}$ measurements as indicators of carbon flow in

311 marine and freshwater ecosystems. *Contrib. Mar. Sci.* 27:15-47.

312 Oczkowski, A. J. and others 2008. Distribution and trophic importance of anthropogenic

313 nitrogen in Narragansett Bay: An assessment using stable isotopes. *Estuaries and*

314 *Coasts* 31:53-69.

315 Raven, J. A. and others 2002. Mechanistic interpretation of carbon isotope discrimination

316 by marine macroalgae and seagrasses. *Funct. Plant Biol.* 29:355-378.

317 Raven, J. A., D. I. Walker, A. M. Johnston, L. L. Handley, and J. E. Kubler. 1995.

318 Implications of ^{13}C natural abundance measurements for photosynthetic

319 performance by marine macrophytes in their natural environment. *Mar. Ecol.*

320 *Prog. Ser.* 123:193-205.

321 Thornber, C. S., P. Dimilla, S. W. Nixon, and R. A. Mckinney. 2008. Natural and
322 anthropogenic nitrogen uptake by bloom-forming macroalgae. *Mar. Pollut. Bull.*
323 56:261-269.

324 Underwood, A. J. 1997. *Experiments in Ecology*. Cambridge University Press.

325 Wozniak, A. S., C. T. Roman, S. C. Wainright, R. A. Mckinney, and M.-J. James-Pirri.
326 2006. Monitoring food web changes in tide-restored salt marshes: A carbon stable
327 isotope approach. *Estuaries and Coasts* 29:568-578.

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344 Figures

345

346


TREATMENT	<u>INITIAL</u> 9 Nov 2011	<u>3 MONTHS</u> 13 Feb 2012	<u>6 MONTHS</u> 13 Jun 2012	<u>9 MONTHS</u> 15 Aug 2012	<u>12 MONTHS</u> 14 Nov 2012
FREEZER	5	5	5	5	5
DRYING OVEN	5	5	5	5	5
DESICCATOR	5	5	5	5	5
COUNTER		5* →	5* →	5* →	5*
FRESHLY COLLECTED	5 23 Oct	5 22 Jan	5 14 May	5 17 Jul	5 19 Oct

347

348

349 Figure 1. Schematic of treatments for each species. For freezer, drying oven, and
350 desiccator treatments, 75 total individuals were collected in Fall 2011, and 15 were
351 analyzed at each time point (five per treatment). For the freshly collected samples, five
352 specimens were collected from the field at each time point. Dates listed indicate mass
353 spectrometer run dates. *Indicates repeated analysis on same samples ('counter'
354 treatment).

355



356

357

358 Figure 2. Mean (± 1 standard error) $\delta^{15}\text{N}$ values for *U. rigida* (top panel) and *G.*

359 *vermiculophylla* (bottom panel) over the length of the experiment. Shapes represent


360 storage techniques (desiccator, drying oven, freezer, and counter) where counter samples

361 were left on the bench-top and periodically reanalyzed, with separate replicates of

362 desiccator, drying oven, and freezer samples that were analyzed at each time step. Bars

363 represent samples freshly collected from the same location just prior to analysis and were

364 included to illustrate the range of values observed seasonally.



365

366

367 Figure 3. Mean (± 1 standard error) of %N values for *U. rigida* (top panel) and *G.*

368 *vermiculophylla* (bottom panel) over the length of the experiment. Results are presented


369 in the same manner as in Fig. 2.

370

371

372

373



374

375


376 Figure 4. Mean (± 1 standard error) of $\delta^{13}\text{C}$ values for *U. rigida* (top panel) and *G.*

377 *vermiculophylla* (bottom panel) over the length of the experiment. Results are presented

378 in the same manner as in Fig. 2.

379

380



381

382

383 Figure 5. Mean (± 1 standard error) of %C values for *U. rigida* (top panel) and *G.*

384 *vermiculophylla* (bottom panel) over the length of the experiment. Results are presented

385 in the same manner as in Fig. 2.

386

387

388

389

390 Tables

391

392 Table 1: Results from three way fixed factor ANOVAs for $\delta^{15}\text{N}$, %N, $\delta^{13}\text{C}$, %C for
393 frozen, oven, and desiccator samples of *Ulva* and *Gracilaria* and from two way fixed
394 factor ANOVAs from freshly collected samples.

395

Source	DF	$\delta^{15}\text{N}$			%N			$\delta^{13}\text{C}$			%C		
		SS	F	P	SS	F	P	SS	F	P	SS	F	P
Frozen, Oven, and Dessiccator Samples													
Species	1	153.66	153.66	<0.0001	13.5	233.77	<0.0001	264.41	153.37	<0.0001	209.74	86.73	<0.0001
Treatment	2	0.34	0.45	0.64	0.03	0.3	0.74	2.92	0.85	0.43	1.85	0.38	0.68
Species * Treatment	2	0.64	0.85	0.43	0.03	0.24	0.78	9.35	2.71	0.07	1.01	0.21	0.81
Time	4	26.94	17.89	<0.0001	0.67	2.89	0.02	15.08	2.19	0.07	18.62	1.92	0.11
Species * Time	4	7.28	4.84	0.001	0.58	2.52	0.04	13.2	1.91	0.11	34.27	3.54	0.01
Treatment * Time	8	4.38	1.45	0.18	0.68	1.48	0.17	7.09	0.51	0.84	46.59	2.41	0.02
Species * Treatment * Time	8	2.64	0.87	0.54	0.48	1.05	0.4	19.65	1.42	0.19	27.26	1.41	0.2
Error	120	45.18			6.94			206.88			290.19		
Freshly Collected Samples													
Species	1	0.04	0.2	0.66	8.67	89.39	<0.0001	16.62	5.66	0.024	303.71	122.41	<0.0001
Time	3	42.58	62.32	<0.0001	50.64	173.94	<0.0001	123.83	14.07	<0.0001	61.02	8.2	0.0004
Species * Time	3	4.81	7.04	0.001	2.42	8.31	0.0003	130.08	14.78	<0.0001	108.1	14.52	<0.0001
Error	31	7.06			3.01			90.96			76.91		

396
397