

2011

Dynamics of a two-dimensional system of rational difference equations of Leslie–Gower type

S. Kalabušić

Mustafa Kulenović

University of Rhode Island, mkulenovic@uri.edu

E. Pilav

Follow this and additional works at: https://digitalcommons.uri.edu/math_facpubs

Citation/Publisher Attribution

Kalabušić, A., Kulenović, M. R. S., & Pilav, E. (2011). Dynamics of a two-dimensional system of rational difference equations of Leslie-Gower type. *Advances in Difference Equations*, 2011:29. doi: 10.1186/1687-1847-2011-29

Available at: <https://doi.org/10.1186/1687-1847-2011-29>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Dynamics of a two-dimensional system of rational difference equations of Leslie–Gower type

Creative Commons License



This work is licensed under a [Creative Commons Attribution 2.0 License](https://creativecommons.org/licenses/by/2.0/).

RESEARCH

Open Access

Dynamics of a two-dimensional system of rational difference equations of Leslie–Gower type

S Kalabušić¹, MRS Kulenović^{2*} and E Pilav¹

* Correspondence:

mkulenovic@mail.uri.edu

²Department of Mathematics,
University of Rhode Island,
Kingston, RI 02881-0816, USA

Full list of author information is
available at the end of the article

Abstract

We investigate global dynamics of the following systems of difference equations

$$\begin{cases} x_{n+1} = \frac{\alpha_1 + \beta_1 x_n}{A_1 + \gamma_n} \\ \gamma_{n+1} = \frac{\gamma_2 \gamma_n}{A_2 + B_2 x_n + \gamma_n} \end{cases}, \quad n = 0, 1, 2, \dots$$

where the parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2, B_2$ are positive numbers, and the initial conditions x_0 and y_0 are arbitrary nonnegative numbers. We show that this system has rich dynamics which depends on the region of parametric space. We show that the basins of attractions of different locally asymptotically stable equilibrium points or non-hyperbolic equilibrium points are separated by the global stable manifolds of either saddle points or non-hyperbolic equilibrium points. We give examples of a globally attractive non-hyperbolic equilibrium point and a semi-stable non-hyperbolic equilibrium point. We also give an example of two local attractors with precisely determined basins of attraction. Finally, in some regions of parameters, we give an explicit formula for the global stable manifold.

Mathematics Subject Classification (2000)

Primary: 39A10, 39A11 Secondary: 37E99, 37D10

Keywords: Basin of attraction, Competitive map, Global stable manifold, Monotonicity, Period-two solution

1 Introduction

In this paper, we study the global dynamics of the following rational system of difference equations

$$\begin{cases} x_{n+1} = \frac{\alpha_1 + \beta_1 x_n}{A_1 + \gamma_n} \\ \gamma_{n+1} = \frac{\gamma_2 \gamma_n}{A_2 + B_2 x_n + \gamma_n} \end{cases}, \quad n = 0, 1, 2, \dots \quad (1)$$

where the parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2, B_2$ are positive numbers and initial conditions x_0 and y_0 are arbitrary nonnegative numbers.

System (1) was mentioned in [1] as one of three systems of Open Problem 3, which asked for a description of the global dynamics of some rational systems of difference equations. In notation used to label systems of linear fractional difference equations

used in [1], System (1) is referred to as (29, 38). This system is dual to the system where the roles of x_n and y_n are interchanged, which is labeled as (29, 38) in [1], and so all results proven here extend to the latter system. In this paper, we provide a precise description of the global dynamics of the System (1). We show that System (1) may have between zero and three equilibrium points, which may have different local character. If System (1) has one equilibrium point, then this point is either locally asymptotically stable or saddle point or non-hyperbolic equilibrium point. If System (1) has two equilibrium points, then they are either locally asymptotically stable and non-hyperbolic, or locally asymptotically stable and saddle point. If System (1) has three equilibrium points, then two of equilibrium points are locally asymptotically stable and the third point, which is between these two points in southeast ordering defined below, is a saddle point. The major problem for global dynamics of the System (1) is determining the basins of attraction of different equilibrium points. The difficulty in analyzing the behavior of all solutions of the System (1) lies in the fact that there are many regions of parameters where this system possesses different equilibrium points with different local character and that in several cases, the equilibrium point is non-hyperbolic. However, all these cases can be handled by using recent results from [2].

System (1) is a competitive system, and our results are based on recent results about competitive systems in the plane, see [2,3]. System (1) can be used as a mathematical model for competition in population dynamics. In fact, second equation in (1) is of Leslie-Gower type, and first equation can be considered to be of Leslie-Gower type with stocking which is represented with the term α_1 , see [4-6].

In the next section, we present some general results about competitive systems in the plane. Section 3 contains some basic facts such as the non-existence of period-two solution of System (1). Section 4 analyzes local stability which is fairly complicated for this system. Finally, Section 5 gives global dynamics for all values of parameters.

2 Preliminaries

A first-order system of difference equations

$$\begin{cases} x_{n+1} = f(x_n, y_n) \\ y_{n+1} = g(x_n, y_n) \end{cases}, \quad n = 0, 1, 2, \dots \quad (2)$$

where $S \subset \mathbb{R}^2$, $(f, g): S \rightarrow S$, f, g are continuous functions is *competitive* if $f(x, y)$ is non-decreasing in x and non-increasing in y , and $g(x, y)$ is non-increasing in x and non-decreasing in y . If both f and g are non-decreasing in x and y , the System (2) is *cooperative*. Competitive and cooperative maps are defined similarly. *Strongly competitive* systems of difference equations or strongly competitive maps are those for which the functions f and g are coordinate-wise strictly monotone.

Competitive and cooperative systems have been investigated by many authors, see [2,3,5-19]. Special attention to discrete competitive and cooperative systems in the plane was given in [2,3,5-7,10,12,17,20]. One of the reasons for paying special attention to two-dimensional discrete competitive and cooperative systems is their applicability and the fact that many examples of mathematical models in biology and economy which involve competition or cooperation are models which involve two species. Another reason is that the theory of two-dimensional discrete competitive and cooperative systems is very well developed, unlike such theory for three and higher

dimensional systems. Part of the reason for this situation is de Mottoni and Schiaffino theorem given below, which provides relatively simple scenarios for possible behavior of many two-dimensional discrete competitive and cooperative systems. However, this does not mean that one cannot encounter chaos in such systems as has been shown by Smith, see [17].

If $\mathbf{v} = (u, v) \in \mathbb{R}^2$, we denote with $\mathcal{Q}_\ell(\mathbf{v})$, $\ell \in \{1, 2, 3, 4\}$, the four quadrants in \mathbb{R}^2 relative to \mathbf{v} , i.e., $\mathcal{Q}_1(\mathbf{v}) = \{(x, y) \in \mathbb{R}^2: x \geq u, y \geq v\}$, $\mathcal{Q}_2(\mathbf{v}) = \{(x, y) \in \mathbb{R}^2: x \leq u, y \geq v\}$, and so on. Define the *South-East* partial order \preceq_{se} on \mathbb{R}^2 by $(x, y) \preceq_{se} (s, t)$ if and only if $x \leq s$ and $y \geq t$. Similarly, we define the *North-East* partial order \preceq_{ne} on \mathbb{R}^2 by $(x, y) \preceq_{ne} (s, t)$ if and only if $x \leq s$ and $y \leq t$. For $\mathcal{A} \subset \mathbb{R}^2$ and $x \in \mathbb{R}^2$, define the *distance from x to A* as $\text{dist}(x, \mathcal{A}) = \inf\{\|x-y\|: y \in \mathcal{A}\}$. By $\text{int } \mathcal{A}$, we denote the interior of a set \mathcal{A} .

It is easy to show that a map F is competitive if it is non-decreasing with respect to the South-East partial order, that is, if the following holds:

$$\begin{pmatrix} x^1 \\ y^1 \end{pmatrix} \preceq_{se} \begin{pmatrix} x^2 \\ y^2 \end{pmatrix} \Rightarrow F \begin{pmatrix} x^1 \\ y^1 \end{pmatrix} \preceq_{se} F \begin{pmatrix} x^2 \\ y^2 \end{pmatrix}. \tag{3}$$

For standard definitions of attracting fixed point, saddle point, stable manifold, and related notions see [11].

We now state three results for competitive maps in the plane. The following definition is from [17].

Definition 1 Let \mathcal{S} be a nonempty subset of \mathbb{R}^2 . A competitive map $T: \mathcal{S} \rightarrow \mathcal{S}$ is said to satisfy condition (O+) if for every x, y in \mathcal{S} , $T(x) \preceq_{ne} T(y)$ implies $x \preceq_{ne} y$, and T is said to satisfy condition (O-) if for every x, y in \mathcal{S} , $T(x) \preceq_{ne} T(y)$ implies $y \preceq_{ne} x$.

The following theorem was proved by de Mottoni and Schiaffino [20] for the Poincaré map of a periodic competitive Lotka-Volterra system of differential equations. Smith [14,15] generalized the proof to competitive and cooperative maps.

Theorem 1 Let \mathcal{S} be a nonempty subset of \mathbb{R}^2 . If T is a competitive map for which (O+) holds then for all $x \in \mathcal{S}$, $\{T^n(x)\}$ is eventually componentwise monotone. If the orbit of x has compact closure, then it converges to a fixed point of T . If instead (O-) holds, then for all $x \in \mathcal{S}$, $\{T^{2n}(x)\}$ is eventually componentwise monotone. If the orbit of x has compact closure in \mathcal{S} , then its omega limit set is either a period-two orbit or a fixed point.

The following result is from [17], with the domain of the map specialized to be the cartesian product of intervals of real numbers. It gives a sufficient condition for conditions (O+) and (O-).

Theorem 2 Let $\mathcal{R} \subset \mathbb{R}^2$ be the cartesian product of two intervals in \mathbb{R} . Let $T: \mathcal{R} \rightarrow \mathcal{R}$ be a C^1 competitive map. If T is injective and $\det J_T(x) > 0$ for all $x \in \mathcal{R}$ then T satisfies (O+). If T is injective and $\det J_T(x) < 0$ for all $x \in \mathcal{R}$ then T satisfies (O-).

The following result is a direct consequence of the Trichotomy Theorem of Dancer and Hess, see [3] and [21] and is helpful for determining the basins of attraction of the equilibrium points.

Corollary 1 If the nonnegative cone of \preceq is a generalized quadrant in \mathbb{R}^n , and if T has no fixed points in $[[u_1, u_2]]$ other than u_1 and u_2 , then the interior of $[[u_1, u_2]]$ is either a subset of the basin of attraction of u_1 or a subset of the basin of attraction of u_2 .

Next result is well-known global attractivity result that holds in partially ordered Banach spaces as well, see [21].

Theorem 3 *Let T be a monotone map on a closed and bounded rectangular region $\mathcal{R} \subset \mathbb{R}^2$. Suppose that T has a unique fixed point \bar{e} in \mathcal{R} . Then \bar{e} is a global attractor of T on \mathcal{R} .*

The following theorems were proved by Kulenović and Merino [2] for competitive systems in the plane, when one of the eigenvalues of the linearized system at an equilibrium (hyperbolic or non-hyperbolic) is by absolute value smaller than 1 while the other has an arbitrary value. These results are useful for determining basins of attraction of fixed points of competitive maps.

Theorem 4 *Let T be a competitive map on a rectangular region $\mathcal{R} \subset \mathbb{R}^2$. Let $\bar{x} \in \mathcal{R}$ be a fixed point of T such that $\Delta := \mathcal{R} \cap \text{int}(\mathcal{Q}_1(\bar{x}) \cup \mathcal{Q}_3(\bar{x}))$ is nonempty (i.e., \bar{x} is not the NW or SE vertex of \mathcal{R}), and T is strongly competitive on Δ . Suppose that the following statements are true.*

- a. *The map T has a C^1 extension to a neighborhood of \bar{x} .*
- b. *The Jacobian $J_T(\bar{x})$ of T at \bar{x} has real eigenvalues λ, μ such that $0 < |\lambda| < \mu$, where $|\lambda| < 1$, and the eigenspace E^λ associated with λ is not a coordinate axis.*

Then there exists a curve $C \subset \mathcal{R}$ through \bar{x} that is invariant and a subset of the basin of attraction of \bar{x} , such that C is tangential to the eigenspace E^λ at \bar{x} , and C is the graph of a strictly increasing continuous function of the first coordinate on an interval. Any endpoints of C in the interior of \mathcal{R} are either fixed points or minimal period-two points. In the latter case, the set of endpoints of C is a minimal period-two orbit of T .

The situation where the endpoints of C are boundary points of \mathcal{R} is of interest. The following result gives a sufficient condition for this case.

Theorem 5 *For the curve C of Theorem 4 to have endpoints in $\partial\mathcal{R}$, it is sufficient that at least one of the following conditions is satisfied.*

- i. *The map T has no fixed points nor periodic points of minimal period-two in Δ .*
- ii. *The map T has no fixed points in Δ , $\det J_T(\bar{x}) > 0$, and $T(\bar{x}) = \bar{x}$ has no solutions $x \in \Delta$.*
- iii. *The map T has no points of minimal period-two in Δ , $\det J_T(\bar{x}) < 0$, and $T(x) = \bar{x}$ has no solutions $x \in \Delta$.*

The next result is useful for determining basins of attraction of fixed points of competitive maps.

Theorem 6 (A) *Assume the hypotheses of Theorem 4, and let C be the curve whose existence is guaranteed by Theorem 4. If the endpoints of C belong to $\partial\mathcal{R}$, then C separates \mathcal{R} into two connected components, namely*

$$\mathcal{W}_- := \{x \in \mathcal{R} \setminus C : \exists y \in C \text{ with } x \prec_{se} y\} \quad \text{and} \quad \mathcal{W}_+ := \{x \in \mathcal{R} \setminus C : \exists y \in C \text{ with } y \prec_{se} x\}, \quad (4)$$

such that the following statements are true.

- (i) \mathcal{W}_- *is invariant, and $\text{dist}(T^n(x), \mathcal{Q}_2(\bar{x})) \rightarrow 0$ as $n \rightarrow \infty$ for every $x \in \mathcal{W}_-$.*
- (ii) \mathcal{W}_+ *is invariant, and $\text{dist}(T^n(x), \mathcal{Q}_4(\bar{x})) \rightarrow 0$ as $n \rightarrow \infty$ for every $x \in \mathcal{W}_+$.*

(B) If, in addition to the hypotheses of part (A), \bar{x} is an interior point of \mathcal{R} and T is C^2 and strongly competitive in a neighborhood of \bar{x} , then T has no periodic points in the boundary of $(Q_1(\bar{x}) \cup Q_3(\bar{x}))$ except for \bar{x} , and the following statements are true.

- (iii) For every $x \in \mathcal{W}_-$ there exists $n_0 \in \mathbb{N}$ such that $T^n(x) \in \text{int } Q_2(\bar{x})$ for $n \geq n_0$.
- (iv) For every $x \in \mathcal{W}_+$ there exists $n_0 \in \mathbb{N}$ such that $T^n(x) \in \text{int } Q_4(\bar{x})$ for $n \geq n_0$.

If T is a map on a set \mathcal{R} and if \bar{x} is a fixed point of T , the stable set $\mathcal{W}^s(\bar{x})$ of \bar{x} is the set $\{x \in \mathcal{R} : T^n(x) \rightarrow \bar{x}\}$ and unstable set $\mathcal{W}^u(\bar{x})$ of \bar{x} is the set

$$\{x \in \mathcal{R} : \text{there exists } \{x_n\}_{n=-\infty}^0 \subset \mathcal{R} \text{ s.t. } T(x_n) = x_{n+1}, x_0 = x, \text{ and } \lim_{n \rightarrow -\infty} x_n = \bar{x}\}$$

When T is non-invertible, the set $\mathcal{W}^s(\bar{x})$ may not be connected and made up of infinitely many curves, or $\mathcal{W}^u(\bar{x})$ may not be a manifold. The following result gives a description of the stable and unstable sets of a saddle point of a competitive map. If the map is a diffeomorphism on \mathcal{R} , the sets $\mathcal{W}^s(\bar{x})$ and $\mathcal{W}^u(\bar{x})$ are the stable and unstable manifolds of \bar{x} .

Theorem 7 In addition to the hypotheses of part (B) of Theorem 6, suppose that $\mu > 1$ and that the eigenspace E^μ associated with μ is not a coordinate axis. If the curve \mathcal{C} of Theorem 4 has endpoints in $\partial\mathcal{R}$, then \mathcal{C} is the stable set $\mathcal{W}^s(\bar{x})$ of \bar{x} , and the unstable set $\mathcal{W}^u(\bar{x})$ of \bar{x} is a curve in \mathcal{R} that is tangential to E^μ at \bar{x} and such that it is the graph of a strictly decreasing function of the first coordinate on an interval. Any endpoints of $\mathcal{W}^u(\bar{x})$ in \mathcal{R} are fixed points of T .

The following result gives information on local dynamics near a fixed point of a map when there exists a characteristic vector whose coordinates have negative product and such that the associated eigenvalue is hyperbolic. This is a well-known result, valid in much more general setting that we include it here for completeness. A point (x, y) is a subsolution if $T(x, y) \preceq_{se} (x, y)$, and (x, y) is a supersolution if $(x, y) \preceq_{se} T(x, y)$. An order interval $[[a, b), (c, d]]$ is the cartesian product of the two compact intervals $[a, c]$ and $[b, d]$.

Theorem 8 Let T be a competitive map on a rectangular set $\mathcal{R} \subset \mathbb{R}^2$ with an isolated fixed point $\bar{x} \in \mathcal{R}$ such that $\mathcal{R} \cap \text{int}(Q_2(\bar{x}) \cup Q_4(\bar{x})) \neq \emptyset$. Suppose T has a C^1 extension to a neighborhood of \bar{x} . Let $v = (v^{(1)}, v^{(2)}) \in \mathbb{R}^2$ be an eigenvector of the Jacobian of T at \bar{x} , with associated eigenvalue $\mu \in \mathbb{R}$. If $v^{(1)}v^{(2)} < 0$, then there exists an order interval \mathcal{J} which is also a relative neighborhood of \bar{x} such that for every relative neighborhood $\mathcal{U} \subset \mathcal{J}$ of \bar{x} the following statements are true.

- i. If $\mu > 1$, then $\mathcal{U} \cap \text{int } Q_2(\bar{x})$ contains a subsolution and $\mathcal{U} \cap \text{int } Q_4(\bar{x})$ contains a supersolution. In this case for every $x \in \mathcal{I} \cap \text{int}(Q_2(\bar{x}) \cup Q_4(\bar{x}))$ there exists N such that $T^n(x) \notin \mathcal{J}$ for $n \geq N$.
- ii. If $\mu < 1$, then $\mathcal{U} \cap \text{int } Q_2(\bar{x})$ contains a supersolution and $\mathcal{U} \cap \text{int } Q_4(\bar{x})$ contains a subsolution. In this case $T^n(x) \rightarrow \bar{x}$ for every $x \in \mathcal{J}$.

3 Some basic facts

In this section, we give some basic facts about the nonexistence of period-two solutions, local injectivity of the map T at the equilibrium point, and boundedness of solutions. See [22] for similar analysis.

3.1 Equilibrium points

The equilibrium points (\bar{x}, \bar{y}) of System (1) satisfy

$$\bar{x} = \frac{\alpha_1 + \beta_1 \bar{x}}{A_1 + \bar{y}}, \quad \bar{y} = \frac{\gamma_2 \bar{y}}{A_2 + B_2 \bar{x} + \bar{y}}. \tag{5}$$

Solutions of System (5) are:

(i) $\bar{y} = 0, \bar{x} = \frac{\alpha_1}{A_1 - \beta_1}, A_1 > \beta_1$, i.e. $E_1 = \left(\frac{\alpha_1}{A_1 - \beta_1}, 0\right)$. Thus, the equilibrium point

$E_1 = \left(\frac{\alpha_1}{A_1 - \beta_1}, 0\right)$ exists if $A_1 > \beta_1$.

(ii) If $\bar{y} \neq 0$, then using System (5), we obtain

$$\bar{y} = \gamma_2 - A_2 - B_2 \bar{x}, \quad \bar{x}^2 B_2 - \bar{x}(\gamma_2 + A_1 - A_2 - \beta_1) + \alpha_1 = 0. \tag{6}$$

Solutions of System (6) are:

$$\bar{x}_{3,2} = \frac{\gamma_2 + A_1 - A_2 - \beta_1 \pm \sqrt{D_0}}{2B_2}, \quad \bar{y}_{2,3} = \frac{\gamma_2 - A_2 - A_1 + \beta_1 \pm \sqrt{D_0}}{2}, \tag{7}$$

where $D_0 = (\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1$ which gives a pair of the equilibrium points $E_2 = (\bar{x}_2, \bar{y}_2)$ and $E_3 = (\bar{x}_3, \bar{y}_3)$.

The criteria for the existence of the three equilibrium points are summarized in Table 1.

3.2 Injectivity

Lemma 1 Assume that (\bar{x}, \bar{y}) is an equilibrium of the map T . Then the following holds:

1) If

$$B_2 > \frac{A_2 \beta_1}{\alpha_1}, \quad A_1(B_2 \alpha_1 - A_2 \beta_1) \gamma_2 - (B_2 \alpha_1 + (A_1 - A_2) \beta_1)(A_1 A_2 - \beta_1 A_2 + B_2 \alpha_1) = 0, \tag{8}$$

then $T\left(x, \frac{A_1 A_2 \beta_1 + x A_1 B_2 \beta_1}{B_2 \alpha_1 - A_2 \beta_1}\right) = (\bar{x}, \bar{y})$ for all $x \geq 0$, where

$$(\bar{x}, \bar{y}) = \left(\bar{x}, \frac{A_1 A_2 \beta_1 + \bar{x} A_1 B_2 \beta_1}{B_2 \alpha_1 - A_2 \beta_1}\right) = \left(\frac{B_2 \alpha_1 + A_2 \beta_1}{A_1 B_2}, \frac{-A_2 \beta_1^2 + A_1 A_2 \beta_1 + B_2 \alpha_1 \beta_1}{B_2 \alpha_1 - A_2 \beta_1}\right).$$

That is the line

$$\mathcal{I} = \left\{ \left(x, \frac{A_1 A_2 \beta_1 + x A_1 B_2 \beta_1}{B_2 \alpha_1 - A_2 \beta_1}\right) : x \geq 0 \right\}$$

is invariant, equilibrium $(\bar{x}, \bar{y}) \in \mathcal{I}$ and for $(x, y) \in \mathcal{I}$ the following holds $T(x, y) = (\bar{x}, \bar{y})$, that is every point of this line is mapped to the equilibrium point (\bar{x}, \bar{y}) .

- 1.i) If $(B_2 \alpha_1 - A_2 \beta_1)^2 - A_1^2 B_2 \alpha_1 > 0$ then $(\bar{x}, \bar{y}) = E_3$.
- 1.ii) If $(B_2 \alpha_1 - A_2 \beta_1)^2 - A_1^2 B_2 \alpha_1 < 0$ then $(\bar{x}, \bar{y}) = E_2$.
- 1.iii) If $(B_2 \alpha_1 - A_2 \beta_1)^2 - A_1^2 B_2 \alpha_1 = 0$ then $(\bar{x}, \bar{y}) = E_3 = E_2$.

2) If

Table 1 The equilibrium points of System (1)

E_1	$A_1 > \beta_1, A_2 < \gamma_2 < A_1 + A_2 - \beta_1, \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} < \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or $A_1 > \beta_1, A_2 > \gamma_2, \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}, A_1 + \gamma_2 \neq A_2 + \beta_1$ or $A_1 > \beta_1, A_2 = \gamma_2, \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or $A_1 > \beta_1, \alpha_1 > \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$
$E_1 \equiv E_2 \equiv E_3$	$A_1 > \beta_1, A_1 + A_2 = \beta_1 + \gamma_2, \alpha_1 = \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}$
$E_1 \equiv E_3, E_2$	$A_1 > \beta_1, A_1 + A_2 < \beta_1 + \gamma_2, A_2 < \gamma_2, \alpha_1 = \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}$
E_1, E_2, E_3	$A_1 > \beta_1, A_1 + A_2 < \beta_1 + \gamma_2, \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} < \alpha_1 < \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$
E_1, E_2	$A_1 > \beta_1, A_2 < \gamma_2, \alpha_1 < \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}$
$E_1 \equiv E_2$	$A_1 > \beta_1, A_2 < \gamma_2 < A_1 + A_2 - \beta_1, \alpha_1 = \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}$
$E_1, E_2 \equiv E_3$	$A_1 > \beta_1, A_1 + A_2 < \beta_1 + \gamma_2, \alpha_1 = \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$
E_2, E_3	$A_1 < \beta_1, A_1 + \gamma_2 > A_2 + \beta_1, \alpha_1 < \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or $A_1 = \beta_1, A_1 + A_2 < \beta_1 + \gamma_2, \alpha_1 < \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$
$E_2 = E_3$	$A_1 < \beta_1, A_1 + \gamma_2 > A_2 + \beta_1, \alpha_1 = \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or $A_1 = \beta_1, A_1 + A_2 < \beta_1 + \gamma_2, \alpha_1 = \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$
No equilibrium	$A_1 < \beta_1, A_2 < \gamma_2 < -A_1 + A_2 + \beta_1, \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or $A_1 < \beta_1, A_2 \geq \gamma_2, \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or $A_1 \leq \beta_1, \alpha_1 > \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or $A_1 = \beta_1, A_1 + A_2 > \gamma_2 + \beta_1, \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$

$$B_2 \leq \frac{A_2 \beta_1}{\alpha_1} \text{ or } A_1 (B_2 \alpha_1 - A_2 \beta_1) \gamma_2 - (B_2 \alpha_1 + (A_1 - A_2) \beta_1) (A_1 A_2 - \beta_1 A_2 + B_2 \alpha_1) \neq 0,$$

then the following holds.

$$T(x, \gamma) = (\bar{x}, \bar{\gamma}) \Rightarrow (x, \gamma) = (\bar{x}, \bar{\gamma}).$$

Proof $T(x, \gamma) = (\bar{x}, \bar{\gamma})$ is equivalent to

$$\left(\frac{\alpha_1 + \beta_1 x}{A_1 + \gamma}, \frac{\gamma_2 \gamma}{A_2 + B_2 x + \gamma} \right) = (\bar{x}, \bar{\gamma}). \quad (9)$$

Since $(\bar{x}, \bar{\gamma})$ is the equilibrium point of the map T then System (9) is equivalent to

$$\left(\frac{\alpha_1 + \beta_1 x}{A_1 + \gamma}, \frac{\gamma_2 \gamma}{A_2 + B_2 x + \gamma} \right) = \left(\frac{\alpha_1 + \beta_1 \bar{x}}{A_1 + \bar{\gamma}}, \frac{\gamma_2 \bar{\gamma}}{A_2 + B_2 \bar{x} + \bar{\gamma}} \right). \quad (10)$$

System (10) is equivalent to

$$-\gamma \alpha_1 + \bar{\gamma} \alpha_1 - \gamma \bar{x} \beta_1 + x \bar{\gamma} \beta_1 + x A_1 \beta_1 - \bar{x} A_1 \beta_1 = 0 \quad (11)$$

$$\gamma A_2 \gamma_2 - \bar{\gamma} A_2 \gamma_2 + \gamma \bar{x} B_2 \gamma_2 - x \bar{\gamma} B_2 \gamma_2 = 0. \quad (12)$$

Equation 11 implies

$$\gamma = \frac{\bar{\gamma}\alpha_1 + x\bar{\gamma}\beta_1 + xA_1\beta_1 - \bar{x}A_1\beta_1}{\alpha_1 + \bar{x}\beta_1}.$$

and Equation 12 is equivalent to

$$(x - \bar{x})(-\bar{\gamma}B_2\alpha_1 + \bar{\gamma}A_2\beta_1 + A_1A_2\beta_1 + \bar{x}A_1B_2\beta_1)\gamma_2 = 0. \tag{13}$$

We conclude the following: If $(-\bar{\gamma}B_2\alpha_1 + \bar{\gamma}A_2\beta_1 + A_1A_2\beta_1 + \bar{x}A_1B_2\beta_1) \neq 0$, then $x = \bar{x}$ and $\gamma = \bar{\gamma}$.

On the other hand, if $(-\bar{\gamma}B_2\alpha_1 + \bar{\gamma}A_2\beta_1 + A_1A_2\beta_1 + \bar{x}A_1B_2\beta_1) = 0$, since $(\bar{x}, \bar{\gamma})$ is the equilibrium of the map T , then

$$B_2 > \frac{A_2\beta_1}{\alpha_1}, \quad \bar{\gamma} = -\frac{A_1(A_2 + \bar{x}B_2)\beta_1}{A_2\beta_1 - B_2\alpha_1}$$

and

$$(\bar{x}, \bar{\gamma}) = \left(\frac{\alpha_1 + \beta_1\bar{x}}{A_1 + \bar{\gamma}}, \frac{\gamma_2\bar{\gamma}}{A_2 + B_2\bar{x} + \bar{\gamma}} \right).$$

Using these equations, we have

$$\bar{x} = \frac{B_2\alpha_1 - A_2\beta_1}{A_1B_2}, \quad \bar{\gamma} = \frac{\beta_1(-A_1A_2 + \beta_1A_2 - B_2\alpha_1)}{A_2\beta_1 - B_2\alpha_1}$$

and

$$A_1(B_2\alpha_1 - A_2\beta_1)\gamma_2 - (B_2\alpha_1 + (A_1 - A_2)\beta_1)(A_1A_2 - \beta_1A_2 + B_2\alpha_1) = 0, \tag{14}$$

which completes the proof of lemma.

3.3 Period-two solutions

In this section, we prove that System (1) has no minimal period-two solutions which will be essential for application of Theorem 4 and Corollary 6.

Lemma 2 *System (1) has no minimal period-two solution.*

Proof Period-two solution satisfies $T^2(x, y) = (x, y)$, that is

$$T^2(x, y) = \left(\frac{\alpha_1 + \frac{\beta_1(\alpha_1 + x\beta_1)}{y + A_1}}{A_1 + \frac{\gamma\gamma_2}{y + A_2 + xB_2}}, \frac{\gamma\gamma_2^2}{\frac{(y + A_2 + xB_2)((y + A_1)A_2 + B_2(\alpha_1 + x\beta_1))}{y + A_1} + \gamma\gamma_2} \right) = (x, y).$$

This is equivalent to

$$-\frac{(y + A_2 + xB_2)(-x\beta_1^2 - \alpha_1\beta_1 + (y + A_1)(xA_1 - \alpha_1)) + x\gamma(y + A_1)\gamma_2}{(y + A_1)(A_1(y + A_2 + xB_2) + \gamma\gamma_2)} = 0$$

and

$$\frac{\gamma((y + A_1)\gamma_2^2 - \gamma(y + A_1)\gamma_2 + (-y - A_2 - xB_2)((y + A_1)A_2 + B_2(\alpha_1 + x\beta_1)))}{(y + A_2 + xB_2)((y + A_1)A_2 + B_2(\alpha_1 + x\beta_1)) + \gamma(y + A_1)\gamma_2} = 0,$$

which is equivalent to

$$(y + A_2 + xB_2)(-x\beta_1^2 - \alpha_1\beta_1 + (y + A_1)(xA_1 - \alpha_1)) + x\gamma(y + A_1)\gamma_2 = 0 \tag{15}$$

$$\gamma((\gamma + A_1)\gamma_2^2 - \gamma(\gamma + A_1)\gamma_2 + (-\gamma - A_2 - xB_2)((\gamma + A_1)A_2 + B_2(\alpha_1 + x\beta_1))) = 0 \quad (16)$$

If $\gamma = 0$, we substitute in (15) to obtain the first fixed point, that is $x = \frac{\alpha_1}{A_1 - \beta_1}$ i
 $x = -\frac{A_2}{B_2}$. Assume

$$(\gamma + A_1)\gamma_2^2 - \gamma(\gamma + A_1)\gamma_2 + (-\gamma - A_2 - xB_2)((\gamma + A_1)A_2 + B_2(\alpha_1 + x\beta_1)) = 0. \quad (17)$$

From (17) we calculate x^2 . We have

$$x^2 = -\frac{(\gamma + A_1)A_2^2 + (\gamma^2 + A_1(\gamma + xB_2) + B_2(\alpha_1 + x(\gamma + \beta_1)))A_2}{B_2^2\beta_1} - \frac{x B_2^2 \alpha_1 + \gamma B_2(\alpha_1 + x\beta_1) + (\gamma + A_1)(\gamma - \gamma_2)\gamma_2}{B_2^2\beta_1}. \quad (18)$$

Put (18) into (15), we have that (15) is equivalent to

$$\gamma + A_1 = 0 \quad (19)$$

or

$$(A_1(\gamma + A_1) - \beta_1^2)\gamma_2^2 + \gamma(\beta_1^2 + xB_2\beta_1 - A_1(\gamma + A_1))\gamma_2 + (-\gamma - A_2 - xB_2)(-A_2\beta_1^2 + B_2\alpha_1\beta_1 + A_1((\gamma + A_1)A_2 + B_2\alpha_1)) = 0 \quad (20)$$

If (19) holds, then we obtain a negative solution. Now, assume that (20) holds. We have

$$x = \frac{(A_1(\gamma + A_1) - \beta_1^2)\gamma_2^2 - \gamma(A_1(\gamma + A_1) - \beta_1^2)\gamma_2}{B_2(A_2A_1^2 + (\gamma A_2 + B_2\alpha_1)A_1 - \beta_1(-B_2\alpha_1 + A_2\beta_1 + \gamma\gamma_2))} + \frac{(-\gamma - A_2)(-A_2\beta_1^2 + B_2\alpha_1\beta_1 + A_1((\gamma + A_1)A_2 + B_2\alpha_1))}{B_2(A_2A_1^2 + (\gamma A_2 + B_2\alpha_1)A_1 - \beta_1(-B_2\alpha_1 + A_2\beta_1 + \gamma\gamma_2))}. \quad (21)$$

Put (21) into (18), we obtain that (18) is equivalent to

$$-\gamma^2 + (-A_1 - A_2 + \beta_1 + \gamma_2)\gamma - B_2\alpha_1 + \beta_1(A_2 - \gamma_2) + A_1(\gamma_2 - A_2) = 0 \quad (22)$$

or

$$-(A_2 + \gamma_2)(A_1^2 + (\beta_1 - A_2)A_1 + \beta_1\gamma_2)\gamma^2 - (A_1 + \beta_1)(A_1 - A_2 + \beta_1 - \gamma_2) \times (B_2\alpha_1 + A_1(A_2 + \gamma_2) - \beta_1(A_2 + \gamma_2))\gamma + (A_1 + \beta_1)^2\gamma_2(B_2\alpha_1 + A_1(A_2 + \gamma_2) - \beta_1(A_2 + \gamma_2)) = 0. \quad (23)$$

If (22) holds, we obtain the fixed points. So, we assume that (23) holds. Set

$$\Delta := (A_1 + \beta_1)^2(B_2\alpha_1 + (A_1 - \beta_1)(A_2 + \gamma_2)) \times ((B_2\alpha_1 + (A_1 - \beta_1)(A_2 + \gamma_2))(A_1 - A_2 + \beta_1 - \gamma_2)^2 + 4\gamma_2(A_2 + \gamma_2)(A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2)). \quad (24)$$

If $\Delta \geq 0$ and $A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2 \neq 0$ hold, we obtain the real solution of the form

$$\gamma_1 = -\frac{(\Delta_1 - \sqrt{\Delta})}{2(A_2 + \gamma_2)(A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2)}$$

$$\gamma_2 = -\frac{(\Delta_1 + \sqrt{\Delta})}{2(A_2 + \gamma_2)(A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2)}$$

where

$$\Delta_1 := (A_1 + \beta_1)(A_1 - A_2 + \beta_1 - \gamma_2)(B_2\alpha_1 + (A_1 - \beta_1)(A_2 + \gamma_2)).$$

Substituting this into (21), we have that the corresponding solutions are

$$x_1 = \frac{(\Delta_2 - \sqrt{\Delta})}{2B_2(A_1 + \beta_1)(A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2)}$$

$$x_2 = \frac{(\Delta_2 + \sqrt{\Delta})}{2B_2(A_1 + \beta_1)(A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2)}$$

where

$$\Delta_2 := (A_1 + \beta_1)(-(A_1 + \beta_1)\gamma_2^2 - (A_1 + \beta_1)^2 + B_2\alpha_1)\gamma_2 + (-A_1 + A_2 - \beta_1)(A_2(A_1 + \beta_1) - B_2\alpha_1). \quad (25)$$

□

Claim 1 Assume $\Delta \geq 0$. Then we have:

- a) If $x_1 \geq 0$ then $y_1 < 0$.
- b) If $x_2 \geq 0$ then $y_2 < 0$.

Proof. Since $T : [0, \infty)^2 \rightarrow [0, \infty)^2$, $T(x_1, y_1) = (x_2, y_2)$ and $T(x_2, y_2) = (x_1, y_1)$, it is obvious that if $(x_i, y_i) \in [0, \infty)^2$ holds then $T(x_i, y_i) \in [0, \infty)^2$ for $i = 1, 2$. It is enough to show that the assumptions $(x_1, y_1), (x_2, y_2) \in [0, \infty)^2$ and $T(x_1, y_1) = (x_2, y_2) \neq (x_1, y_1)$ lead to a contradiction.

Indeed, if $A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2 > 0$ then $(x_1, y_1) \prec_{se} (x_2, y_2)$. Since T is strongly competitive map then $(x_2, y_2) = T(x_1, y_1) \prec_{se} T(x_2, y_2) = (x_1, y_1)$ which is impossible since $(x_1, y_1) \prec_{se} (x_2, y_2)$.

If $A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2 < 0$ then $(x_2, y_2) \prec_{se} (x_1, y_1)$. Similarly, we have the same conclusion if $A_1(A_1 - A_2 + \beta_1) + \beta_1\gamma_2 = 0$. □

3.4 Boundedness of solutions

Lemma 3 Assume that $y_0 = 0, x_0 \in \mathbb{R}^+$. Then the following statements are true.

- (i) If $A_1 > \beta_1$ then $y_n = 0, x_n \rightarrow \frac{\alpha_1}{A_1 - \beta_1}, n \rightarrow \infty$.
- (ii) If $A_1 < \beta_1$ then $y_n = 0, x_n \rightarrow \infty, n \rightarrow \infty$.
- (iii) If $A_1 = \beta_1$, then $x_n = x_0 + \frac{\alpha_1}{A_1}n$ and $y_n = 0, x_n \rightarrow \infty$.

Assume that $y_0 \neq 0$ and $(x_0, y_0) \in \mathbb{R}_2^+$. Then the following statements are true.

- (iv) $x_{n+1} \leq \frac{\alpha_1}{A_1} + \frac{\beta_1}{A_1}x_n$ for all $n = 0, 1, 2, \dots$
- (v) $y_n \leq \gamma_2, n \geq N, y_{n+1} \leq C\left(\frac{\gamma_2}{A_2}\right)^n$ and
- (a) $x_n \rightarrow \frac{\alpha_1}{A_1 - \beta_1}, A_1 > \beta_1$.
- (b) $x_n \leq \frac{\alpha_1}{A_1 - \beta_1} + \varepsilon, \varepsilon > 0, A_1 > \beta_1$.
- (c) If $\gamma_2 < A_2$ then $y_n \rightarrow 0, n \rightarrow \infty$

Proof. Take $y_0 = 0$ and $x_0 \in \mathbb{R}^+$. Then, we have $y_n = 0$, for all $n \in \mathbb{N}$, and

$$x_{n+1} = \frac{\alpha_1}{A_1} + \frac{\beta_1}{A_1} x_n. \tag{26}$$

Solution of Equation 26 is

$$x_n = c \left(\frac{\beta_1}{A_1} \right)^n + \frac{\alpha_1}{A_1 - \beta_1} \tag{27}$$

From $y_{n+1} = \frac{\gamma_2 y_n}{A_2 + B_2 x_n + \gamma_1}$ it follows that $y_{n+1} \leq \frac{\gamma_2}{A_2} y_n$, $y_{n+1} \leq \gamma_2$, $n \geq 0$. The proof of Lemma 3 follows from (27). \square

4 Linearized stability analysis

The map T associated to System (1) is given by

$$T(x, y) = \left(\frac{\alpha_1 + \beta_1 x}{A_1 + y}, \frac{\gamma_2 y}{A_2 + B_2 x + y} \right).$$

The Jacobian matrix of the map T has the form:

$$J_T = \begin{pmatrix} \frac{\beta_1}{A_1 + y} & -\frac{\alpha_1 + \beta_1 x}{(A_1 + y)^2} \\ -\frac{B_2 \gamma_2 y}{(A_2 + B_2 x + y)^2} & \frac{\gamma_2 A_2 + \gamma_2 B_2 x}{(A_2 + B_2 x + y)^2} \end{pmatrix}. \tag{28}$$

The value of the Jacobian matrix of T at the equilibrium point $E = (\bar{x}, \bar{y})$ is

$$J_T(\bar{x}, \bar{y}) = \begin{pmatrix} \frac{\beta_1}{A_1 + \bar{y}} & -\frac{\bar{x}}{A_1 + \bar{y}} \\ -\frac{B_2 \bar{y}}{A_2 + B_2 \bar{x} + \bar{y}} & \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2} \end{pmatrix}. \tag{29}$$

The determinant of (29) is given by

$$\det J_T(\bar{x}, \bar{y}) = \frac{\beta_1}{A_1 + \bar{y}} \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2} - \frac{\bar{x}}{A_1 + \bar{y}} \frac{B_2 \bar{y}}{A_2 + B_2 \bar{x} + \bar{y}}.$$

The trace of (29) is

$$\text{Tr} J_T(\bar{x}, \bar{y}) = \frac{\beta_1}{A_1 + \bar{y}} + \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2}.$$

The characteristic equation has the form

$$\lambda^2 - \lambda \left(\frac{\beta_1}{A_1 + \bar{y}} + \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2} \right) + \frac{\beta_1 (\gamma_2 A_2 + \gamma_2 B_2 \bar{x})}{(A_1 + \bar{y})(A_2 + B_2 \bar{x} + \bar{y})^2} - \frac{B_1 \bar{x} \bar{y}}{(A_1 + \bar{y})(A_2 + B_2 \bar{x} + \bar{y})} = 0.$$

Theorem 9 Assume that $A_1 > \beta_1$. Then there exists the equilibrium point E_1 and:

- (i) E_1 is locally asymptotically stable if $\gamma_2 - A_2 < \frac{B_2 \alpha_1}{A_1 - \beta_1}$.
- (ii) E_1 is a saddle point if $\gamma_2 - A_2 > \frac{B_2 \alpha_1}{A_1 - \beta_1}$. The eigenvalues are

$$\lambda_1 = \frac{\beta_1}{A_1}, \quad \lambda_2 = \frac{(A_1 - \beta_1) \gamma_2}{B_2 \alpha_1 + A_2 (A_2 - \beta_1)}.$$

The corresponding eigenvectors, respectively, are

$$v_1 = (1, 0), \quad v_2 = \left(\frac{\alpha_1}{A_1(A_1 - \beta_1) \left(\frac{\beta_1}{\alpha_1} - \frac{(A_1 - \beta_1)}{A_1 A_2 + B_2 \alpha_1 - A_2 \beta_1} \right)}, 1 \right).$$

(iii) E_1 is non-hyperbolic if $\gamma_2 - A_2 = \frac{B_2 \alpha_1}{A_1 - \beta_1}$. The eigenvalues are $\lambda_1 = \frac{\beta_1}{A_1}$, $\lambda_2 = 1$. The corresponding eigenvectors are $\left(-\frac{\alpha_1}{(A_1 - \beta_1)^2}, 1 \right)$ and $(1, 0)$.

Proof. Evaluating Jacobian (29) at the equilibrium point $E_1(\alpha_1/(A_1 - \beta_1), 0)$,

$$J_T(E_1) = \begin{pmatrix} \frac{\beta_1}{A_1} & -\frac{\alpha_1}{A_1(A_1 - \beta_1)} \\ 0 & \frac{(A_1 - \beta_1)\gamma_2}{A_2(A_1 - \beta_1) + B_2 \alpha_1} \end{pmatrix}. \tag{30}$$

The determinant of (30) is given by

$$\det J_T(\bar{x}, \bar{y}) = \frac{\beta_1 \gamma_2 (A_1 - \beta_1)}{A_1 [A_2 (A_1 - \beta_1) + B_2 \alpha_1]}.$$

The trace of (30) is

$$\text{Tr} J_T(\bar{x}, \bar{y}) = \frac{\beta_1}{A_1} + \frac{(A_1 - \beta_1)\gamma_2}{A_2 (A_1 - \beta_1) + B_2 \alpha_1}.$$

The characteristic equation associated to System (1) at E_1 has the form

$$\left(\frac{\beta_1}{A_1} - \lambda \right) \left(\frac{(A_1 - \beta_1)\gamma_2}{A_2 (A_1 - \beta_1) + B_2 \alpha_1} - \lambda \right) = 0. \tag{31}$$

From Equation 31 we have

$$\lambda_1 = \frac{\beta_1}{A_1}, \quad \lambda_2 = \frac{(A_1 - \beta_1)\gamma_2}{A_2 (A_1 - \beta_1) + B_2 \alpha_1}.$$

- (i) If $A_1 > \beta_1$ and $\gamma_2 - A_2 < \frac{B_2 \alpha_1}{A_1 - \beta_1}$ then $\lambda_1 < 1$ and $\lambda_2 < 1$. Hence, E_1 is a sink.
- (ii) If $A_1 > \beta_1$ and $\gamma_2 - A_2 > \frac{B_2 \alpha_1}{A_1 - \beta_1}$. Then $\lambda_1 < 1$, and $\lambda_2 > 1$. Hence, E_1 is a saddle.
- (iii) If $A_1 > \beta_1$ and $\gamma_2 - A_2 = \frac{B_2 \alpha_1}{A_1 - \beta_1}$. Then, using Equation 31, we have that $\lambda_1 < 1$ and $\lambda_2 = 1$.

From (30) we obtain the eigenvectors that correspond to these eigenvalues. \square

We now perform a similar analysis for the other cases in table.

Theorem 10 Assume

$$A_1 > \beta_1, \quad A_1 + A_2 < \beta_1 + \gamma_2, \quad \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} < \alpha_1 < \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}.$$

Then E_1, E_2, E_3 exist and:

- (i) Equilibrium E_1 is locally asymptotically stable.

(ii) *Equilibrium E_3 is a saddle point. The eigenvalues are*

$$\lambda_1 = \frac{-\bar{\gamma}_3 (A_1 + \bar{\gamma}_3) + \gamma_2 (A_1 + \beta_1 + \bar{\gamma}_3) - \sqrt{\mathcal{D}}}{2\gamma_2 (A_1 + \bar{\gamma}_3)}$$

and

$$\lambda_2 = \frac{-\bar{\gamma}_3 (A_1 + \bar{\gamma}_3) + \gamma_2 (A_1 + \beta_1 + \bar{\gamma}_3) + \sqrt{\mathcal{D}}}{2\gamma_2 (A_1 + \bar{\gamma}_3)},$$

and $|\lambda_1| < 1$, $|\lambda_2| > 1$, where

$$\mathcal{D} = \bar{\gamma}_3^2 (A_1 + \bar{\gamma}_3)^2 - 2\gamma_2 \bar{\gamma}_3 (A_1 - \beta_1 - 2B_2 \bar{x}_3 + \bar{\gamma}_3) (A_1 + \bar{\gamma}_3) + \gamma_2^2 (A_1 - \beta_1 + \bar{\gamma}_3)^2.$$

The corresponding eigenvectors, respectively, are

$$v_1 = \left(-\bar{\gamma}_3 (A_1 + \bar{\gamma}_3) + \gamma_2 (A_1 - \beta_1 + \bar{\gamma}_3) + \sqrt{\mathcal{D}}, \quad 2B_2 \bar{\gamma}_3 (A_1 + \bar{\gamma}_3) \right)$$

$$v_2 = \left(-\bar{\gamma}_3 (A_1 + \bar{\gamma}_3) + \gamma_2 (A_1 - \beta_1 + \bar{\gamma}_3) - \sqrt{\mathcal{D}}, \quad 2B_2 \bar{\gamma}_3 (A_1 + \bar{\gamma}_3) \right).$$

(iii) *Equilibrium E_2 is locally asymptotically stable.*

Proof. By Theorem 9 (i) holds.

Equilibrium E_3 is a saddle if and only if the following conditions are satisfied

$$|\text{Tr} J_T(\bar{x}, \bar{y})| > |1 + \det J_T(\bar{x}, \bar{y})| \quad \text{and} \quad \text{Tr}^2 J_T(\bar{x}, \bar{y}) - 4 \det J_T(\bar{x}, \bar{y}) > 0.$$

The first condition is equivalent to

$$\frac{\beta_1}{A_1 + \bar{y}} + \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2} > \left| 1 + \frac{\beta_1}{(A_1 + \bar{y})} \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2} - \frac{B_2 \bar{x} \bar{y}}{(A_1 + \bar{y})(A_2 + B_2 \bar{x} + \bar{y})} \right|$$

which is equivalent to

$$\beta_1 (A_2 + B_2 \bar{x} + \bar{y})^2 + (A_1 + \bar{y})(\gamma_2 A_2 + \gamma_2 B_2 \bar{x})$$

$$> (A_1 + \bar{y})(A_2 + B_2 \bar{x} + \bar{y})^2 + \beta_1 \gamma_2 (A_2 + B_2 \bar{x}) - B_2 \bar{x} \bar{y} (A_2 + B_2 \bar{x} + \bar{y}).$$

This is equivalent to

$$(A_2 + B_2 \bar{x} + \bar{y})^2 (\beta_1 - A_1 - \bar{y}) + \gamma_2 (A_2 + B_2 \bar{x})(A_1 + \bar{y} - \beta_1) > -B_2 \bar{x} \bar{y} (A_2 + B_2 \bar{x} + \bar{y})$$

$$\gamma_2^2 (\beta_1 - A_1 - \bar{y}) + \gamma_2 (A_2 + B_2 \bar{x})(A_1 + \bar{y} - \beta_1) > -B_2 \gamma_2 \bar{x} \bar{y}$$

$$(A_1 - \beta_1 + \bar{y})(A_2 + B_2 \bar{x} - \gamma_2) > -B_2 \bar{x} \bar{y}$$

$$(\beta_1 - A_1 - \bar{y})(A_2 + B_2 \bar{x} - \gamma_2) < B_2 \bar{x} \bar{y}.$$

We have to prove that $(\beta_1 - A_1 - \bar{\gamma}_3)(A_2 + B_2 \bar{x}_3 - \gamma_2) < B_2 \bar{x}_3 \bar{\gamma}_3$. Notice that

$$\beta_1 - A_1 - \bar{\gamma}_3 = -B_2 \bar{x}_2 \quad \text{and} \quad A_2 + B_2 \bar{x}_3 - \gamma_2 = -\bar{\gamma}_3.$$

Now,

$$(\beta_1 - A_1 - \bar{\gamma}_3)(A_2 + B_2 \bar{x}_3 - \gamma_2) < B_2 \bar{x}_3 \bar{\gamma}_3$$

is equivalent to $B_2 \bar{x}_2 \bar{\gamma}_3 < B_2 \bar{x}_3 \bar{\gamma}_3$. This implies $\bar{x}_2 < \bar{x}_3$ which is true. Condition

$$\text{Tr}^2 J_T(\bar{x}, \bar{y}) - 4 \det J_T(\bar{x}, \bar{y}) > 0$$

is equivalent to

$$\left(\frac{\beta_1}{A_1 + \bar{y}} - \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2} \right)^2 + \frac{4B_2 \bar{x} \bar{y}}{(A_1 + \bar{y})(A_2 + B_2 \bar{x} + \bar{y})} > 0$$

which is clearly satisfied. Hence, E_3 is a saddle.

Now, we prove that E_2 is locally asymptotically stable. Notice that

$$|\text{Tr} J_T(\bar{x}, \bar{y})| < 1 + \det J_T(\bar{x}, \bar{y}) < 2$$

implies $\bar{x}_3 > \bar{x}_2$ which is true.

The second condition is equivalent to

$$\frac{\beta_1}{(A_1 + \bar{y})} \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{y})^2} - \frac{B_2 \bar{x} \bar{y}}{(A_1 + \bar{y})(A_2 + B_2 \bar{x} + \bar{y})} < 1.$$

This implies the following

$$\beta_1 \gamma_2 (A_2 + B_2 \bar{x}) - B_2 \bar{x} \bar{y} (A_2 + B_2 \bar{x} + \bar{y}) < (A_1 + \bar{y})(A_2 + B_2 \bar{x} + \bar{y})^2.$$

Now, using Equation 5, we obtain

$$\begin{aligned} \beta_1 \gamma_2 (\gamma_2 - \bar{y}) - B_2 \bar{x} \bar{y} \gamma_2 &< (A_1 + \bar{y}) \gamma_2^2 \\ -(\beta_1 \bar{y} + B_2 \bar{x} \bar{y}) &< (A_1 - \beta_1 + \bar{y}) \gamma_2 \end{aligned}$$

which is true, since the left side is always negative, while the right side is always positive.

Theorem 11 *Assume*

$$A_1 > \beta_1, \quad A_1 + A_2 < \beta_1 + \gamma_2, \quad \alpha_1 = \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}.$$

Then $E_1(\alpha_1/(A_1 - \beta_1), 0)$ and $E_2 = E_3 = \left(\frac{\gamma_2 - A_2 + A_1 - \beta_1}{2B_2}, \frac{\gamma_2 - A_2 - A_1 + \beta_1}{2} \right)$ exist and

- (i) *Equilibrium E_1 is locally asymptotically stable.*
- (ii) *Equilibrium E_2 is non-hyperbolic. The eigenvalues are*

$$\lambda_1 = 1, \quad \lambda_2 = \frac{A_1^2 - A_2^2 + 2A_2\beta_1 - \beta_1^2 + 2A_2\gamma_2 + 2\beta_1\gamma_2 - \gamma_2^2}{2\gamma_2(A_1 - A_2 + \beta_1 + \gamma_2)}.$$

The corresponding eigenvectors are

$$\left(-1/B_2, 1 \right), \quad \left(\frac{2\gamma_2(A_1 - A_2 - \beta_1 + \gamma_2)}{B_2(-A_1 - A_2 + \beta_1 + \gamma_2)(A_1 - A_2 + \beta_1 + \gamma_2)}, 1 \right).$$

Proof. By Theorem 9, E_1 is locally asymptotically stable.

Now, we prove that E_2 is non-hyperbolic.

Evaluating Jacobian (29) at the equilibrium point $E_2 = \left(\frac{\gamma_2 - A_2 + A_1 - \beta_1}{2B_2}, \frac{\gamma_2 - A_2 - A_1 + \beta_1}{2} \right)$,

$$J_T(E_2) = \begin{pmatrix} \frac{\beta_1}{A_1 + \bar{y}} - \frac{\bar{x}}{A_1 + \bar{y}} & -\frac{\bar{x}}{A_1 + \bar{y}} \\ -\frac{B_2 \bar{y}}{\gamma_2} & \frac{A_2 + B_2 \bar{x}}{\gamma_2} \end{pmatrix} = \begin{pmatrix} \frac{2\beta_1}{A_1 + \gamma_2 - A_2 + \beta_1} & \frac{-\gamma_2 + A_2 - A_1 + \beta_1}{B_2(A_1 + \gamma_2 - A_2 + \beta_1)} \\ -\frac{B_2(\gamma_2 - A_2 - A_1 + \beta_1)}{2\gamma_2} & \frac{A_2 + \gamma_2 + A_1 - \beta_1}{2\gamma_2} \end{pmatrix}. \quad (32)$$

The eigenvalues of (32) are

$$\lambda_1 = 1, \quad \text{and} \quad \lambda_2 = \frac{A_1^2 - A_2^2 + 2A_2\beta_1 - \beta_1^2 + 2A_2\gamma_2 + 2\beta_1\gamma_2 - \gamma_2^2}{2\gamma_2(A_1 - A_1 + \beta_1 + \gamma_2)}.$$

Notice that $|\lambda_2| < 1$. Hence, E_2 is non-hyperbolic.

Theorem 12 *Assume*

$$\begin{aligned} A_1 > \beta_1, \quad A_2 < \gamma_2 < A_1 + A_2 - \beta_1, \quad \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} < \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2} \\ A_1 > \beta_1, \quad A_2 > \gamma_2, \quad \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}, \quad A_1 + \gamma_2 \neq A_2 + \beta_1 \\ A_1 > \beta_1, \quad A_2 = \gamma_2, \quad \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2} \\ A_1 > \beta_1, \quad \alpha_1 > \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2} \end{aligned}$$

Then there exists a unique equilibrium $E_1 (\alpha_1/(A_1 - \beta_1), 0)$ which is locally asymptotically stable.

Proof. Observe that the assumption of Theorem 12 implies that the y coordinates of the equilibrium E_2 and E_3 are less than zero. By Theorem 9 E_1 is locally asymptotically stable.

Theorem 13 *Assume*

$$A_1 > \beta_1, \quad A_2 < \gamma_2, \quad \alpha_1 < \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}.$$

Then there exist two equilibrium points E_1 and E_2 . E_1 is a saddle point. The eigenvalues are

$$\lambda_1 = \frac{\beta_1}{A_1}, \quad \lambda_2 = \frac{(A_1 - \beta_1)\gamma_2}{B_2\alpha_1 + A_2(A_2 - \beta_1)}.$$

The corresponding eigenvectors, respectively, are

$$v_1 = (1, 0), \quad v_2 = \left(\frac{\alpha_1}{A_1(A_1 - \beta_1) \left(\frac{\beta_1}{\alpha_1} - \frac{(A_1 - \beta_1)}{A_1 A_2 + B_2 \alpha_1 - A_2 \beta_1} \right)}, 1 \right).$$

The equilibrium E_2 is locally asymptotically stable.

Proof. By Theorem 9 (ii), E_1 is a saddle point.

Now, we check the sign of coordinates of the equilibrium point E_2 . We have that $\bar{x}_2 > 0$, since all parameters are positive. Consider \bar{y}_2 . Since

$$\frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2} - \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} = \frac{(A_1 + A_2 - \beta_1 - \gamma_2)^2}{4B_2} > 0,$$

we have that $(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4\alpha_1 B_2 > 0$.

$$\bar{y}_1 > 0 \Leftrightarrow \gamma_2 - A_2 + \beta_1 - A_1 + \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4\alpha_1 B_2} > 0.$$

This implies

$$\sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4\alpha_1 B_2} > (A_1 - \beta_1) - (\gamma_2 - A_2). \tag{33}$$

From Equation 33, we see that inequality is always true if $A_1 - \beta_1 < \gamma_2 - A_2$. If $A_1 - \beta_1 > \gamma_2 - A_2$, then

$$\begin{aligned} (\gamma_2 - A_2)^2 + 2(\gamma_2 - A_2)(A_1 - \beta_1) + (A_1 - \beta_1)^2 - 4\alpha_1 B_1 &> (A_1 - \beta_1)^2 - 2(A_1 - \beta_1)(\gamma_2 - A_2) \\ (\gamma_2 - A_2)(A_1 - \beta_1) &> \alpha_1 B_2 \end{aligned}$$

which is true, since $A_1 - \beta_1 > \frac{B_2 \alpha_1}{\gamma_2 - A_2}$. So, in both cases $\bar{x}_2 > 0$ and $\bar{y}_2 > 0$.

Notice, that $\bar{x}_3 > 0$. Now, we check the sign of \bar{y}_3 . Assume that $\bar{y}_3 > 0$. Then, we have

$$\begin{aligned} \bar{y}_2 > 0 &\Leftrightarrow (\gamma_2 - A_2) - (A_1 - \beta_1) > \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4\alpha_1 B_2}. \\ &\Leftrightarrow (\gamma_2 - A_2)(A_1 - \beta_1) < \alpha_1 B_2. \end{aligned}$$

This is a contradiction with the assumption of theorem and so E_3 is not in considered domain.

By Theorem 10, E_2 is a locally asymptotically stable.

Theorem 14 Assume

$$A_1 > \beta_1, \quad A_1 + A_2 < \beta_1 + \gamma_2, \quad \alpha_1 = \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}.$$

Then there exist two equilibrium points $E_1 \equiv E_3 = \left(\frac{\alpha_1}{A_1 - \beta_1}, 0\right)$ and $E_2 = \left(\frac{\alpha_1}{\gamma_2 - A_2}, \frac{\gamma_2 - A_2 - A_1 + \beta_1}{2}\right)$, and $E_1 \equiv E_3$ is non-hyperbolic. The eigenvalues are $\lambda_1 = \frac{\beta_1}{A_1}$, $\lambda_2 = 1$. The corresponding eigenvectors are $\left(-\frac{\alpha_1}{(A_1 - \beta_1)^2}, 1\right)$ and $(1, 0)$. The equilibrium point E_2 is locally asymptotically stable.

Proof. By Theorem 10, E_2 is locally asymptotically stable. By Theorem 9 (iii), E_1 is non-hyperbolic.

Now, we consider the special case of System (1) when $A_1 = \beta_1$.

In this case, System (1) becomes

$$\begin{cases} x_{n+1} = \frac{\alpha_1 + A_1 x_n}{A_1 + \gamma_n} \\ \gamma_{n+1} = \frac{\gamma_2 \gamma_n}{A_2 + B_2 x_n + \gamma_n} \end{cases}, \quad n = 0, 1, 2, \dots \tag{34}$$

The map T associated to System (34) is given by

$$T(x, \gamma) = \left(\frac{\alpha_1 + A_1 x}{A_1 + \gamma}, \frac{\gamma_2 \gamma}{A_2 + B_2 x + \gamma} \right).$$

The Jacobian matrix of the map T has the form:

$$J_T = \begin{pmatrix} \frac{A_1}{A_1 + \gamma} & -\frac{\alpha_1 + A_1 x}{(A_1 + \gamma)^2} \\ -\frac{\beta_2 \gamma_2 \gamma}{(A_2 + B_2 x + \gamma)^2} & \frac{\gamma_2 A_2 + \gamma_2 B_2 x}{(A_2 + B_2 x + \gamma)^2} \end{pmatrix}. \tag{35}$$

The value of the Jacobian matrix of T at the equilibrium point $E = (\bar{x}, \bar{\gamma})$ is

$$J_T(\bar{x}, \bar{\gamma}) = \begin{pmatrix} \frac{A_1}{A_1 + \bar{\gamma}} & -\frac{\bar{x}}{A_1 + \bar{\gamma}} \\ -\frac{B_2 \bar{\gamma}}{A_2 + B_2 \bar{x} + \bar{\gamma}} & \frac{\gamma_2 A_2 + \gamma_2 B_2 \bar{x}}{(A_2 + B_2 \bar{x} + \bar{\gamma})^2} \end{pmatrix} = \begin{pmatrix} \frac{A_1}{A_1 + \bar{\gamma}} & -\frac{\bar{x}}{A_1 + \bar{\gamma}} \\ -\frac{B_2 \bar{\gamma}}{\gamma_2} & \frac{A_2 + B_2 \bar{x}}{\gamma_2} \end{pmatrix}. \tag{36}$$

The characteristic equation of T at $(\bar{x}, \bar{\gamma})$ has the form

$$\lambda^2 - \lambda \left(\frac{A_1}{A_1 + \bar{\gamma}} + \frac{A_2 + B_2\bar{x}}{\gamma_2} \right) + \frac{A_1}{A_1 + \bar{\gamma}} \frac{A_2 + B_2\bar{x}}{\gamma_2} - \frac{B_2\bar{x}\bar{\gamma}}{(A_1 + \bar{\gamma})\gamma_2} = 0.$$

Equilibrium points satisfy the following System

$$\begin{aligned} \bar{x} &= \frac{\alpha_1 + A_1\bar{x}}{A_1 + \bar{\gamma}} \\ \bar{\gamma} &= \frac{\gamma_2\bar{\gamma}}{A_2 + B_2\bar{x} + \bar{\gamma}} \quad n = 0, 1, \dots \end{aligned} \tag{37}$$

Notice, if $\bar{\gamma} = 0$, then using the first equation of System (37) we obtain $\alpha_1 = 0$ which is impossible. If $\bar{\gamma} \neq 0$ then, using System (37), we obtain

$$\begin{aligned} \bar{\gamma} &= \gamma_2 - A_2 - B_2\bar{x} \\ 0 &= B_2\bar{x}^2 - \bar{x}(\gamma_2 - A_2) + \alpha_1. \end{aligned}$$

and the equilibrium points are:

$$\begin{aligned} E_3 &= \left(\frac{\gamma_2 - A_2 + \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2B_2}, \frac{\gamma_2 - A_2 - \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2} \right), \\ E_2 &= \left(\frac{\gamma_2 - A_2 - \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2B_2}, \frac{\gamma_2 - A_2 + \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2} \right). \end{aligned}$$

We prove the following.

Theorem 15 *Assume*

$$A_1 = \beta_1.$$

Then the following statements hold.

(i) *If $\gamma_2 > A_2$, $(\gamma_2 - A_2)^2 - 4B_2\alpha_1 > 0$ then System (34) has two positive equilibrium points*

$$E_3 = \left(\frac{\gamma_2 - A_2 + \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2B_2}, \frac{\gamma_2 - A_2 - \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2} \right)$$

and

$$E_2 = \left(\frac{\gamma_2 - A_2 - \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2B_2}, \frac{\gamma_2 - A_2 + \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2} \right).$$

E_3 is a saddle point. The eigenvalues are

$$\lambda_1 = \frac{-\bar{\gamma}_3(A_1 + \bar{\gamma}_3) + \gamma_2(2A_1 + \bar{\gamma}_3) - \sqrt{F}}{2\gamma_2(A_1 + \bar{\gamma}_3)}, \quad |\lambda_1| < 1$$

$$\lambda_2 = \frac{-\bar{\gamma}_3(A_1 + \bar{\gamma}_3) + \gamma_2(2A_1 + \bar{\gamma}_3) + \sqrt{F}}{2\gamma_2(A_1 + \bar{\gamma}_3)}, \quad \lambda_2 > 1,$$

where

$$F = \bar{\gamma}_3^2(A_1 + \bar{\gamma}_3)^2 - 2\gamma_2\bar{\gamma}_3(\bar{\gamma}_3 - 2B_2\bar{x}_3)(A_1 + \bar{\gamma}_3) + \gamma_2^2\bar{\gamma}_3^2.$$

The corresponding eigenvectors are

$$v_1 = (-\bar{\gamma}_3(A_1 + \bar{\gamma}_3) + \gamma_2\bar{\gamma}_3 + \sqrt{F}, 2B_2\bar{\gamma}_3(A_1 + \bar{\gamma}_3)),$$

$$v_2 = (-\bar{\gamma}_3(A_1 + \bar{\gamma}_3) + \gamma_2\bar{\gamma}_3 - \sqrt{F}, 2B_2\bar{\gamma}_3(A_1 + \bar{\gamma}_3)).$$

The equilibrium E_2 is locally asymptotically stable.

(ii) If $\gamma_2 > A_2$, $(\gamma_2 - A_2)^2 - 4B_2\alpha_1 > 0$ then System (34) has a unique equilibrium point

$E = \left(\frac{\gamma_2 - A_2}{2B_2}, \frac{\gamma_2 - A_2}{2}\right)$ which is non-hyperbolic. The eigenvalues are $\lambda_1 = 1$ and

$\lambda_2 = \frac{2A_1A_2 - A_2^2 + 2A_1\gamma_2 + 2A_2\gamma_2 - \gamma_2^2}{2\gamma_2(2A_1 - A_2 + \gamma_2)}$. The corresponding eigenvectors are: $(-1/B_2, 1)$ and

$$\left(\frac{2\gamma_2}{B_2(2A_1 - A_2 + \gamma_2)}, 1\right).$$

(iii) If $\gamma_2 < A_2$ and $(\gamma_2 - A_2)^2 - 4B_2\alpha_1 \geq 0$ or $(\gamma_2 - A_2)^2 - 4B_2\alpha_1 > 0$ then System (34) has no equilibrium points.

Proof. (i) First, notice that under these assumptions, E_3 and E_2 are positive. Now, we prove that E_3 is a saddle point.

The equilibrium point E_3 is a saddle if and only if the following conditions are satisfied $|\text{Tr}J_T(\bar{x}, \bar{\gamma})| > |1 + \det J_T(\bar{x}, \bar{\gamma})|$ and $\text{Tr}^2 J_T(\bar{x}, \bar{\gamma}) - 4 \det J_T(\bar{x}, \bar{\gamma}) > 0$.

The first condition is equivalent to

$$\frac{A_1}{A_1 + \bar{\gamma}} + \frac{A_2 + B_2\bar{x}}{\gamma_2} > \left| 1 + \frac{A_1(A_2 + B_2\bar{x})}{\gamma_2(A_1 + \bar{\gamma})} - \frac{B_2\bar{x}\bar{\gamma}}{\gamma_2(A_1 + \bar{\gamma})} \right|,$$

which is equivalent to

$$A_1\gamma_2 + (A_1 + \bar{\gamma})(A_2 + B_2\bar{x}) > \gamma_2(A_1 + \bar{\gamma}) + A_1(A_1 + B_2\bar{x}) - B_2\bar{x}\bar{\gamma},$$

and this is equivalent to

$$\gamma_2 - A_2 < 2B_2\bar{x}.$$

In the case of equilibrium E_3 , this condition becomes

$$\gamma_2 - A_2 < 2B_2\bar{x}_3 \Leftrightarrow \gamma_2 - A_2 < \gamma_2 - A_2 + \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1} \Leftrightarrow \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1} > 0,$$

which is true.

The second condition becomes

$$\left(\frac{A_1}{A_1 + \bar{\gamma}} + \frac{A_2 + B_2\bar{x}}{\gamma_2}\right)^2 - 4\frac{A_1(A_2 + B_2\bar{x})}{\gamma_2(A_1 + \bar{\gamma})} + 4\frac{B_2\bar{x}\bar{\gamma}}{\gamma_2(A_1 + \bar{\gamma})} = \left(\frac{A_1}{A_1 + \bar{\gamma}} - \frac{A_2 + B_2\bar{x}}{\gamma_2}\right)^2 + 4\frac{B_2\bar{x}\bar{\gamma}}{\gamma_2(A_1 + \bar{\gamma})}$$

which is greater than zero. Hence, E_3 is a saddle.

Now, we prove that E_2 is locally asymptotically stable. The equilibrium point E_2 is locally asymptotically stable if the following is satisfied

$$|\text{Tr}J_T(\bar{x}, \bar{\gamma})| < 1 + \det J_T(\bar{x}, \bar{\gamma}) < 2.$$

The first condition is equivalent to

$$\frac{A_1}{A_1 + \bar{y}} + \frac{A_2 + B_2\bar{x}}{\gamma_2} < 1 + \frac{A_1(A_2 + B_2\bar{x})}{\gamma_2(A_1 + \bar{y})} - \frac{B_2\bar{x}\bar{y}}{\gamma_2(A_1 + \bar{y})}.$$

This implies

$$A_1\gamma_2 + (A_1 + \bar{y})(A_2 + B_2\bar{x}) < \gamma_2(A_1 + \bar{y}) + A_1(A_2 + B_2\bar{x}) - B_2\bar{x}\bar{y}$$

which is equivalent to $\gamma_2 - A_2 > 2B_2\bar{x}$. In the case of the equilibrium point E_2 , we have

$$\gamma_2 - A_2 > \gamma_2 - A_2 - \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1} \Leftrightarrow -\sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1} < 0$$

which is true.

The second condition is equivalent to

$$\frac{A_1(A_2 + B_2\bar{x})}{\gamma_2(A_1 + \bar{y})} - \frac{B_2\bar{x}\bar{y}}{\gamma_2(A_1 + \bar{y})} < 1.$$

This implies

$$A_1(A_2 + B_2\bar{x}) - B_2\bar{x}\bar{y} < \gamma_2(A_1 + \bar{y}) \Leftrightarrow A_1(A_2 - \gamma_2 + B_2\bar{x}) < \bar{y}(\gamma_2 + B_2\bar{x}).$$

Notice that

$$A_2 - \gamma_2 + B_2\bar{x}_2 = \frac{A_2 - \gamma_2 - \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2} = -\frac{\gamma_2 - A_2 + \sqrt{(\gamma_2 - A_2)^2 - 4B_2\alpha_1}}{2} = -\bar{y}_2.$$

Now, condition $A_1(A_2 - \gamma_2 + B_2\bar{x}) < \bar{y}(\gamma_2 + B_2\bar{x})$ becomes $-A_1\bar{y}_2 < \bar{y}_2(\gamma_2 + B_2\bar{x}_2) \Leftrightarrow -A_1 < \gamma_2 + B_2\bar{x}_2$ which is true. Hence, E_2 is locally asymptotically stable.

(ii) The characteristic equation associated to System (37) at E has the form

$$\lambda^2 - \lambda \left(\frac{2A_1}{2A_1 + \gamma_2 - A_2} + \frac{A_2 + \gamma_2}{2\gamma_2} \right) + \frac{A_1}{\gamma_2} - \frac{(\gamma_2 - A_2)^2}{2\gamma_2(2A_1 + \gamma_2 - A_2)} = 0. \quad (38)$$

Solutions of Equation (38) are $\lambda_1 = 1$ and $\lambda_2 = \frac{2A_1A_2 - A_2^2 + 2A_1\gamma_2 + 2A_2\gamma_2 - \gamma_2^2}{2\gamma_2(2A_1 - A_2 + \gamma_2)}$.

The corresponding eigenvectors are $(-1/B_2, 1)$ and $(\frac{2\gamma_2}{B_2(2A_1 - A_2 + \gamma_2)}, 1)$.

If $\gamma_2 < A_2$ and $(\gamma_2 - A_2)^2 - 4B_2\alpha_1 \geq 0$ then $\bar{x}_2 < 0$ and $\bar{x}_3 < 0$.

Theorem 16 Assume

$$A_1 < \beta_1, \gamma_2 > A_2, \gamma_2 - A_2 > \beta_1 - A_1 \quad \text{and} \quad (\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1 > 0.$$

Then there exist two positive equilibrium points

$$E_2 = \left(\frac{\gamma_2 - A_2 + A_1 - \beta_1 - \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2B_2}, \frac{\gamma_2 - A_2 - A_1 + \beta_1 + \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2} \right)$$

and

$$E_3 = \left(\frac{\gamma_2 - A_2 + A_1 - \beta_1 + \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2B_2}, \frac{\gamma_2 - A_2 + \beta_1 - A_1 - \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2} \right).$$

E_2 is locally asymptotically stable and E_3 is a saddle. The eigenvalues of characteristic equation at E_3 are

$$\lambda_1 = \frac{-\bar{\gamma}_3 (A_1 + \bar{\gamma}_3) + \gamma_2 (A_1 + \beta_1 + \bar{\gamma}_3) \mp \sqrt{D}}{2\gamma_2 (A_1 + \bar{\gamma}_3)},$$

where

$$D = \bar{\gamma}_3^2 (A_1 + \bar{\gamma}_3)^2 - 2\gamma_2 \bar{\gamma}_3 (A_1 - \beta_1 - 2B_2 \bar{x}_3 + \bar{\gamma}_3) (A_1 + \bar{\gamma}_3) + \gamma_2^2 (A_1 - \beta_1 + \bar{\gamma}_3)^2.$$

The corresponding eigenvectors are

$$v_{1,2} = \left(-\bar{\gamma}_3 (A_1 + \bar{\gamma}_3) + \gamma_2 (A_1 - \beta_1 + \bar{\gamma}_3) \pm \sqrt{D}, 2B_2 \bar{\gamma}_3 (A_1 + \bar{\gamma}_3) \right).$$

Proof. Now, we prove that E_2 is a sink. We check the condition $|\text{Tr} J_T(\bar{x}, \bar{y})| < 1 + \det J_T(\bar{x}, \bar{y}) < 2$. The first condition is equivalent to

$$\frac{\beta_1}{A_1 + \bar{y}} + \frac{A_2 + B_2 \bar{x}}{\gamma_2} < 1 + \frac{\beta_1 (A_2 + B_2 \bar{x})}{\gamma_2 (A_1 + \bar{y})} - \frac{B_2 \bar{x} \bar{y}}{\gamma_2 (A_1 + \bar{y})}.$$

This implies

$$\begin{aligned} \beta_1 \gamma_2 + (A_1 + \bar{y})(A_2 + B_2 \bar{x}) &< \gamma_2 (A_1 + \bar{y}) + \beta_1 (A_2 + B_2 \bar{x}) - B_2 \bar{x} \bar{y} \\ \gamma_2 (\beta_1 - A_1 - \bar{y}) + (A_2 + B_2 \bar{x})(A_1 + \bar{y} - \beta_1) &< -B_2 \bar{x} \bar{y} \\ (A_1 - \beta_1 + \bar{y})(A_2 + B_2 \bar{x} - \gamma_2) &< -B_2 \bar{x} \bar{y} \\ \bar{y}(A_1 - \beta_1 + \bar{y}) &> B_2 \bar{x} \bar{y} \\ (A_1 - \beta_1 + \bar{y}) &> B_2 \bar{x}. \end{aligned}$$

So, we have to prove

$$(A_1 - \beta_1 + \bar{y}_2) > B_2 \bar{x}_2. \tag{39}$$

Note that

$$\begin{aligned} A_1 - \beta_1 + \bar{y}_2 &= A_1 - \beta_1 + \frac{\gamma_2 - A_2 + \beta_1 - A_1 + \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2 \alpha_1}}{2} \\ &= \frac{A_1 - \beta_1 + \gamma_2 - A_2 + \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2 \alpha_1}}{2B_2} B_2 \\ &= B_2 \bar{x}_3. \end{aligned}$$

Now, (39) becomes $B_2 \bar{x}_3 > B_2 \bar{x}_2 \Rightarrow \bar{x}_3 > \bar{x}_2$ which is true.

The second condition is equivalent to

$$\frac{\beta_1 (A_2 + B_2 \bar{x})}{\gamma_2 (A_1 + \bar{y})} - \frac{B_2 \bar{x} \bar{y}}{\gamma_2 (A_1 + \bar{y})} < 1.$$

This implies $\beta_1 (\gamma_2 - \bar{y}) - B_2 \bar{x} \bar{y} < \gamma_2 (A_1 + \bar{y})$. Using equations of equilibrium points, we obtain $\bar{y}_2 (\beta_1 + B_2 \bar{x}_2) > \gamma_2 (\beta_1 - A_1 - \bar{y}_2)$ and

$$\begin{aligned} \beta_1 + B_2\bar{x}_2 &= \beta_1 + \frac{\gamma_2 - A_2 + A_1 - \beta_1 - \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2} \\ &= \frac{\gamma_2 - A_2 + A_1 + \beta_1 - \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2} > 0. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} (\beta_1 - A_1 - \bar{\gamma}_2 &= \beta_1 - A_1 - \frac{\gamma_2 - A_2 + \beta_1 - A_1 + \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2} \\ &= \frac{\beta_1 - A_1 + A_2 - \gamma_2 - \sqrt{(\gamma_2 - A_2 + A_1 - \beta_1)^2 - 4B_2\alpha_1}}{2} < 0 \end{aligned}$$

since $\gamma_2 - A_2 > \beta_1 - A_1$. Hence, E_2 is locally asymptotically stable.

Now, we prove that E_3 is a saddle.

The equilibrium point E_3 is a saddle if and only if the following conditions are satisfied

$$|\text{Tr}J_T(\bar{x}, \bar{\gamma})| > |1 + \det J_T(\bar{x}, \bar{\gamma})| \quad \text{and} \quad \text{Tr}^2 J_T(\bar{x}, \bar{\gamma}) - 4 \det J_T(\bar{x}, \bar{\gamma}) > 0.$$

Note that the first condition is equivalent to $B_2\bar{x}_3 > B_2\bar{x}_2 \Rightarrow \bar{x}_3 > \bar{x}_2$ which is true.

The second condition becomes

$$\left(\frac{\beta_1}{A_1 + \bar{\gamma}} + \frac{A_2 + B_2\bar{x}}{\gamma_2}\right)^2 - 4\frac{\beta_1(A_2 + B_2\bar{x})}{(A_1 + \bar{\gamma})\gamma_2} + 4\frac{B_2\bar{x}\bar{\gamma}}{\gamma_2(A_1 + \bar{\gamma})} = \left(\frac{\beta_1}{A_1 + \bar{\gamma}} - \frac{A_2 + B_2\bar{x}}{\gamma_2}\right)^2 + 4\frac{B_2\bar{x}\bar{\gamma}}{\gamma_2(A_1 + \bar{\gamma})} > 0.$$

Hence, E_3 is a saddle. \square

Theorem 17 Assume

$$A_1 < \beta_1, \quad \gamma_2 > A_2, \quad \gamma_2 - A_2 > \beta_1 - A_1 \quad \text{and} \quad (\gamma_2 + A_1 - A_2 - \beta_1)^2 - 4\alpha_1 B_2 = 0.$$

Then there exists a unique equilibrium point $E_2 \equiv E_3 = E = \left(\frac{\gamma_2 - A_2 + A_1 - \beta_1}{2B_2}, \frac{\gamma_2 - A_2 + \beta_1 - A_1}{2}\right)$ which is non-hyperbolic. The eigenvalues are:

$$\lambda_1 = 1 \quad \text{and} \quad \lambda_2 = \frac{A_1^2 - A_2^2 + 2A_2\beta_1 - \beta_1^2 + 2A_2\gamma_2 + 2\beta_1\gamma_2 - \gamma_2^2}{2\gamma_2(A_1 - A_2 + \beta_1 + \gamma_2)}.$$

The corresponding eigenvectors are:

$$\left(-\frac{1}{B_2}, 1\right), \quad \text{and} \quad \left(\frac{2\gamma_2(A_1 - A_2 - \beta_1 + \gamma_2)}{B_2(-A_1 - A_2 + \beta_1 + \gamma_2)(A_1 - A_2 + \beta_1 + \gamma_2)}, 1\right).$$

Proof. The value of the Jacobian matrix of T at the equilibrium point $E = (\bar{x}, \bar{\gamma})$ is

$$J_T(\bar{x}, \bar{\gamma}) = \begin{pmatrix} \frac{2\beta_1}{\gamma_2 - A_2 + \beta_1 + A_1} & \frac{\gamma_2 - A_2 + A_1 - \beta_1}{B_2(A_1 + \gamma_2 - A_2 + \beta_1)} \\ -\frac{B_2(\gamma_2 - A_2 + \beta_1 - A_1)}{2\gamma_2} & \frac{A_2 + \gamma_2 + A_1 - \beta_1}{2\gamma_2} \end{pmatrix}. \tag{40}$$

The characteristic equation is given by

$$\begin{aligned} \lambda^2 - \lambda \left(\frac{2\beta_1}{\gamma_2 - A_2 + \beta_1 + A_1} + \frac{A_2 + \gamma_2 + A_1 - \beta_1}{2\gamma_2} \right) + \frac{\beta_1(A_2 + \gamma_2 + A_1 - \beta_1)}{\gamma_2(\gamma_2 - A_2 + \beta_1 + A_1)} \\ - \frac{(\gamma_2 - A_2 + \beta_1 - A_1)(\gamma_2 - A_2 + A_1 - \beta_1)}{2\gamma_2(A_1 + \gamma_2 - A_2 + \beta_1)} = 0. \end{aligned} \tag{41}$$

Solutions of Equation (41) are:

$$\lambda_1 = 1 \quad \text{and} \quad \lambda_2 = \frac{A_1^2 - A_2^2 + 2A_2\beta_1 - \beta_1^2 + 2A_2\gamma_2 + 2\beta_1\gamma_2 - \gamma_2^2}{2\gamma_2(A_1 - A_2 + \beta_1 + \gamma_2)}.$$

By using (40), we obtain the corresponding eigenvectors.

5 Global behavior

Theorem 18 Table 2 describes the global behavior of System (1).

Proof. Throughout the proof of theorem \preceq will denote \preceq_{se} .

($\mathcal{R}_i, i = \overline{1, 4}$) By Theorem 9, E_1 is locally asymptotically stable. Consider $M(t) = (0, t)$ for $t \geq \gamma_2 - A_2$. Since $M(t) - T(M(t)) = \left(-\frac{\alpha_1}{t+A_1}, \frac{t(t+A_2-\gamma_2)}{t+A_2}\right)$, we have $M(t) \preceq T(M(t))$ for $t \geq \gamma_2 - A_2$. By induction, $T^n M(t) \preceq T^{n+1}(M(t)) \preceq E_1$ for all $n = 0, 1, 2, \dots$ because $M(t) \preceq E_1$ for all $t \geq 0$. By monotonicity and boundedness, the sequence $\{T^n(M(t))\}$ has to converge to the unique equilibrium E_1 . Consider $N(u) = (u, 0)$ for $u \geq 0$. Lemma 3 implies $T^n(N(u)) \rightarrow E_1$ as $n \rightarrow \infty$. Take any point $(x, y) \in [0, +\infty) \times [0, +\infty)$. Then there exists $t^*, u^* \geq 0$, such that $M(t^*) \preceq (x, y) \preceq N(u^*)$. The monotonicity of the map T implies $T^n M(t^*) \preceq T^n((x, y)) \preceq T^n(N(u^*))$. Since $T^n M(t^*), T^n(N(u^*)) \rightarrow E_1$ as $n \rightarrow \infty$, then $T^n((x, y)) \rightarrow E_1$. This completes the proof.

(\mathcal{R}_5) The first part of this theorem is proven in Theorem 9. The rest of the proof is similar to the proof of part ($\mathcal{R}_i, i = \overline{1, 4}$).

(\mathcal{R}_6) By Lemma 3 $y_0 = 0$ implies $y_n = 0, \forall n \in \mathbb{N}$, and $x_n \rightarrow \frac{\alpha_1}{A_1 - \beta_1}, n \rightarrow \infty$, which shows that x -axis is a subset of the basin of attraction of E_1 .

Furthermore, every solution of (1) enters and stays in the box $\mathcal{B}(E_2)$ and so we can restrict our attention to solutions that starts in $\mathcal{B}(E_2)$. Clearly, the set $Q_2(E_2) \cap \mathcal{B}(E_2)$ is an invariant set with a single equilibrium point E_2 and by Theorem 3, every solution that starts there is attracted to E_2 . In view of Corollary 1, the interior of rectangle $\llbracket E_2, E_1 \rrbracket$ is attracted to either E_1 or E_2 , and because E_2 is the local attractor, it is attracted to E_2 . If $(x, y) \in \mathcal{A} = \mathcal{B} \setminus (\llbracket E_2, E_1 \rrbracket \cup (Q_2(E_2) \cap \mathcal{B}) \cup \{(x, 0) : x \geq 0\})$, then there exist the points $(x_w, y_w) \in \mathcal{A} \cap Q_4(E_2)$ and $(x_b, y_b) \in Q_2(E_2) \cap \mathcal{B}$ such that $(x_b, y_b) \preceq_{se} (x, y) \preceq_{se} (x_w, y_w)$. Consequently, $T^n((x_b, y_b)) \preceq_{se} T^n((x, y)) \preceq_{se} T^n((x_w, y_w))$ for all $n = 1, 2, \dots$ and so $T^n((x, y)) \rightarrow E_2$ as $n \rightarrow \infty$, which completes the proof.

(\mathcal{R}_7) The first part of this Theorem is proven in Theorem 13.

Now, we prove a global result.

$$J_T(E_1) = \begin{pmatrix} \frac{\beta_1}{A_1} & -\frac{\alpha_1}{A_1(A_1 - \beta_1)} \\ 0 & \frac{(A_1 - \beta_1)\gamma_2}{A_1 A_2 - \beta_1 A_2 + B_2 \alpha_1} \end{pmatrix} \quad (42)$$

The eigenvalues of $J_T(E_1)$ are given by $\lambda_1 = \frac{\beta_1}{A_1}$ and $\lambda_2 = \frac{(A_1 - \beta_1)\gamma_2}{A_1 A_2 - \beta_1 A_2 + B_2 \alpha_1}$ and so

$$A_2 < \gamma_2, \quad \alpha_1 < \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} \Rightarrow \lambda_2 > 1, \quad A_1 > \beta_1 \Rightarrow \lambda_1 < 1.$$

The eigenvector of T at E_1 that corresponds to the eigenvalue $\lambda_1 < 1$ is $(1, 0)$.

The rest of the proof is similar to the proof of part (\mathcal{R}_6).

($\mathcal{R}_8, \mathcal{R}_9$) The first part of theorem follows from Theorems 15 and 16. If parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2$ and B_2 do not satisfy the condition (8) of Lemma 1, then the map T defined on the set $\mathcal{R} = \mathbb{R}_+^2$, satisfies all conditions of Theorems 4, 6-8. This implies that

Table 2 Global behavior of System (1)

Region	Global behavior
\mathcal{R}_1 $A_1 > \beta_1, A_2 < \gamma_2 < A_1 + A_2 - \beta_1,$ or $\frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} < \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ \mathcal{R}_2 $A_1 > \beta_1, A_2 > \gamma_2, \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2},$ $A_1 + \gamma_2 \neq A_2 + \beta_1,$ or \mathcal{R}_3 $A_1 > \beta_1, A_2 = \gamma_2, \alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$ or \mathcal{R}_4 $A_1 > \beta_1, \alpha_1 > \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$	There exists a unique equilibrium E_1 , and it is globally asymptotically stable (G.A.S.). The basin of attraction of E_1 is given by $\mathcal{B}^s(E_1) = [0, +\infty)^2$
\mathcal{R}_5 $A_1 > \beta_1, \gamma_2 + \beta_1 \leq A_1 + A_2, \alpha_1 = \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}$	There exists a unique equilibrium $E_1 = E_2$ which is non-hyperbolic. Furthermore, this equilibrium is the global attractor. Its basin of attraction is given by $\mathcal{B}^s(E_1) = [0, +\infty)^2$. This is an example of globally attractive non-hyperbolic equilibrium point
\mathcal{R}_6 $A_1 > \beta_1, A_1 + A_2 < \beta_1 + \gamma_2, \alpha_1 = \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}$	There exist two equilibrium points $E = E_1 = E_3$ which is non-hyperbolic, and E_2 , which is locally asymptotically stable. Furthermore, the x -axis is the basin of attraction of E_1 . The equilibrium point E_2 is globally asymptotically stable with the basin of attraction $\mathcal{B}^s(E_2) = [0, +\infty) \times [0, +\infty)$
\mathcal{R}_7 $A_2 > \beta_1, A_2 < \gamma_2, \alpha_1 < \frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2}$	There exist two equilibrium points E_1 , which is a saddle, and E_2 , which is a locally asymptotically stable equilibrium point. Furthermore, the x -axis is the global stable manifold of $\mathcal{W}^s(E_1)$. The equilibrium point E_2 is globally asymptotically stable with the basin of attraction $\mathcal{B}^s(E_2) = [0, +\infty) \times [0, +\infty)$
\mathcal{R}_8 $A_1 < \beta_1, A_1 + \gamma_2 > A_2 + \beta_1, \alpha_1 < \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$	There exist two equilibrium points E_3 , which is a saddle, and E_2 , which is locally asymptotically stable. Furthermore, there exists the global stable manifold $\mathcal{B}^s(E_3)$ that separates the positive quadrant so that all orbits below this manifold are asymptotic to $(+\infty, 0)$, and all orbits above this manifold are asymptotic to the equilibrium point E_2 . All orbits that starts on $\mathcal{B}^s(E_3)$ are attracted to E_3

Table 2 Global behavior of System (1) (Continued)

or

$$\mathcal{R}_9 A_1 = \beta_1, A_1 + A_2 < \beta_1 + \gamma_2,$$

$$\alpha_1 < \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

$$\mathcal{R}_{10} A_1 > \beta_1, A_1 + A_2 < \beta_1 + \gamma_2,$$

$$\frac{(A_1 - \beta_1)(\gamma_2 - A_2)}{B_2} < \alpha_1 < \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2},$$

$$\mathcal{R}_{11} \quad A_1 > \beta_1, A_1 + A_2 < \beta_1 + \gamma_2, \alpha_1 = \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

$$\mathcal{R}_{12} \quad A_1 < \beta_1, A_1 + \gamma_2 > A_2 + \beta_1, \alpha_1 = \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

$$\mathcal{R}_{13} A_1 = \beta_1, A_1 + A_2 < \beta_1 + \gamma_2,$$

or

$$\alpha_1 = \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

$$\mathcal{R}_{14} A_1 < \beta_1, A_2 < \gamma_2 < -A_1 + A_2 + \beta_1,$$

$$\alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

$$\mathcal{R}_{15} A_1 < \beta_1, A_2 \geq \gamma_2,$$

or

$$\alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

or

$$\mathcal{R}_{16} \quad A_1 \leq \beta_1, \alpha_1 > \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

$$\mathcal{R}_{17} A_1 = \beta_1, A_1 + A_2 > \gamma_2 + \beta_1,$$

or

$$\alpha_1 \leq \frac{(A_1 - A_2 - \beta_1 + \gamma_2)^2}{4B_2}$$

There exist three equilibrium points E_1, E_2 , and E_3 , where E_1 and E_2 are locally asymptotically stable and E_3 is a saddle. There exists the global stable manifold $\mathcal{W}^s(E_3)$ that separates the positive quadrant so that all orbits below this manifold are attracted to the equilibrium point E_1 , and all orbits above this manifold are attracted to the equilibrium point E_2 . All orbits that start on $\mathcal{W}^s(E_3)$ are attracted to E_3 . The global unstable manifold $\mathcal{W}^u(E_3)$ is the graph of a continuous strictly decreasing function, and $\mathcal{W}^u(E_3)$ has endpoints E_2 and E_1 .

There exist two equilibrium points $E = E_2 = E_3$ and E_1 . E_1 is locally asymptotically stable and E is non-hyperbolic. There exists a continuous increasing curve \mathcal{W}_E which is a subset of the basin of attraction of E . All orbits that start below this curve are attracted to E_1 . All orbits that start above this curve are attracted to E .

There exists a unique equilibrium point $E = E_2 = E_3$ which is non-hyperbolic. There exists a continuous increasing curve \mathcal{W}_E which is a subset of basin of attraction of E . All orbits that start below this curve are attracted to $(+\infty, 0)$. All orbits that start above this curve are attracted to E . This is an example of semi-stable non-hyperbolic equilibrium point.

System (1) does not possess an equilibrium point. Its behavior is as follows $x_n \rightarrow \infty, y_n \rightarrow \infty, n \rightarrow \infty$.

there exists the global stable manifold $\mathcal{W}^s(E_3)$ that separates the first quadrant into two invariant regions $\mathcal{W}(E_3)$ (above the stable manifold) and $\mathcal{W}^+(E_3)$ (below the stable manifold) which are connected. Now, we show that each orbit starting in the region $\mathcal{W}^+(E_3)$ is asymptotic to $(\infty, 0)$. Indeed, set $T_1(x, y) = \frac{\alpha_1 + \beta_1 x}{A_1 + y}$, $T_2(x, y) = \frac{\gamma_2 y}{A_2 + B_2 x + y}$. Take $x = (x_0, y_0) \in \mathcal{W}^+(E_3) \cap \mathcal{R}(+, -)$, where $\mathcal{R}(+, -) = \{(x, y) \in \mathcal{R} : T_1(x, y) > x, T_2(x, y) < y\}$. As is known, see [12], the set $\mathcal{R}(+, -)$ is invariant. We have

$$T_1(x_0, y_0) = \frac{\alpha_1 + \beta_1 x_0}{A_1 + y_0} > x_0, \quad T_2(x_0, y_0) = \frac{\gamma_2 y_0}{A_2 + B_2 x_0 + y_0} < y_0,$$

which implies the following

$$(x_0, y_0) \preceq_{se} (T_1(x_0, y_0), T_2(x_0, y_0)) \Leftrightarrow (x_0, y_0) \preceq_{se} T(x_0, y_0).$$

By monotonicity, $T(x_0, y_0) \preceq_{se} T^2(x_0, y_0)$ and by induction, $T^n(x_0, y_0) \preceq_{se} T^{n+1}(x_0, y_0)$. This implies that sequence $\{x_n\}$ is non-decreasing and $\{y_n\}$ is non-increasing. Since, $\{y_n\}$ is bounded from above, hence it must converges. Now $\lim_{n \rightarrow \infty} y_n = 0$ since otherwise (x_n, y_n) will converge to another limit which is strictly south-east of E_3 , which is impossible. By Lemma 3, $x_n \rightarrow \infty$. By Theorems 6-8 for all $(x, y) \in \mathcal{W}^+(E_3)$, there exists $n_0 > 0$ such that $T^n((x, y)) \in \text{int}(Q_4(E_3) \cap \mathcal{R})$, $n > n_0$. We see that for all $(x, y) \in \text{int}(Q_4(E_3) \cap \mathcal{R})$, there exists $(x_l, y_l) \in \mathcal{W}^+(E_3) \cap \mathcal{R}(+, -)$ such that $(x_l, y_l) \preceq (x, y)$. By monotonicity $T^n((x_l, y_l)) \preceq T^n((x, y)) \preceq (\infty, 0)$. This implies $T^n((x, y)) \rightarrow (\infty, 0)$ as $n \rightarrow \infty$.

Now, we show that each orbit starting in the region $\mathcal{W}(E_3)$ converges to E_2 . By Theorem 6, for all $(x, y) \in \mathcal{W}(E_3)$, there exists $n_0 > 0$ such that, $T^n((x, y)) \in \text{int}(Q_2(E_3) \cap \mathcal{R})$, $n > n_0$. Set $M(t) = (0, t)$ By part $((\mathcal{R}_i, i = \overline{1, 4}))$, for $t \geq \gamma_2 - A_2$, we have $M(t) \preceq T(M(t)) \preceq E_2$. By using monotonicity, $T^n(M(t)) \rightarrow E_2$ as $n \rightarrow \infty$. By Corollary 1, the interior of rectangle $\llbracket E_2, E_3 \rrbracket$ is attracted to either E_2 or E_3 , and because E_2 is local attractor, it is attracted to E_2 . If $(x, y) \in \text{int}(Q_2(E_3) \cap \mathcal{R})$, then there exist the points $(x_r, y_r) \in \llbracket E_2, E_3 \rrbracket$ and $t^* \geq \gamma_2 - A_2$, such that $M(t^*) \preceq_{se} (x, y) \preceq_{se} (x_r, y_r)$. Consequently, $T^n(M(t^*)) \preceq_{se} T^n((x, y)) \preceq_{se} T^n((x_r, y_r))$ for all $n = 1, 2, \dots$ and so $T^n((x, y)) \rightarrow E_2$ as $n \rightarrow \infty$.

Now, assume that parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2$, and B_2 satisfy the condition (8) and inequality 1.i) of Lemma 1. Then the set

$$\mathcal{I} = \left\{ \left(x, \frac{A_1 A_2 \beta_1 + x A_1 B_2 \beta_1}{B_2 \alpha_1 - A_2 \beta_1} \right) : x \geq 0 \right\}$$

is invariant and contains the equilibrium point E_3 , and $T(x, y) = E_3$ for $(x, y) \in \mathcal{I}$. In view of the uniqueness of global stable manifold, we conclude that $\mathcal{W}^s(E_3) = \mathcal{I}$. Take any point $(x, y) \in \mathcal{W}^+(E_3)$. Then there exists the point $(x_l, y_l) \in \mathcal{I}$ such that $(x_l, y_l) \ll_{se} (x, y)$. Since, the map T is strongly competitive, then $E_3 = T(x_l, y_l) \ll_{se} T(x, y)$. This implies $T(x, y) \in \text{int}(Q_4(E_3) \cap \mathcal{R})$. Similarly, if $(x, y) \in \mathcal{W}(E_3)$, then $T(x, y) \in \text{int}(Q_2(E_3) \cap \mathcal{R})$. The rest of the proof is similar to the proof of the first case. This completes the proof.

(\mathcal{R}_{10}) The first part of the theorem follows from Theorem 10. If parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2$, and B_2 do not satisfy the condition (8) of Lemma 1, then the map T , defined on the set $\mathcal{R} = \mathbb{R}_+^2$, satisfies all conditions of Theorems 4, 6-8. This implies that there exists the global stable manifold $\mathcal{W}^s(E_3)$ that separates the first quadrant into two

invariant regions $\mathcal{W}^+(E_3)$ (below the stable manifold) and $\mathcal{W}^-(E_3)$ (above the stable manifold) which are connected.

Using Theorems 6, 7, and 8, we have that for all $(x, y) \in \mathcal{W}^+(E_3)$, there exists $n_0 > 0$ such that for $n > n_0$, $T^n((x, y)) \in \text{int}(Q_4(E_3) \cap \mathcal{R})$, and for all $(x, y) \in \mathcal{W}^-(E_3)$, there exists $n_1 > 0$ such that for all $n > n_1$, $T^n((x, y)) \in \text{int}(Q_2(E_3) \cap \mathcal{R})$. Now, we show that each orbit starting in the region $\text{int}(Q_4(E_3))$ converges to E_1 , and each orbit starting in the region $\text{int}(Q_2(E_3))$ converges to E_2 .

Take $0 \leq t \leq (\gamma_2 - A_2)/B_2$, $U(t) = (t, -A_2 - tB_2 + \gamma_2)$. It is easy to see that the following holds

$$U(\bar{x}) = E = E_2 = E_3 \preceq E_1 \quad \text{where } \bar{x} = x_2 = x_3 \quad \text{and}$$

$$U(t) - T(U(t)) = \left(-\frac{(-A_1 + A_2 + 2tB_2 + \beta_1 - \gamma_2)^2}{4B_2(A_1 - A_2 - tB_2 + \gamma_2)}, 0 \right).$$

Since x_2 and x_3 are solutions of the equation $B_2t^2 + (-A_1 + A_2 + \beta_1 - \gamma_2)t + \alpha_1 = 0$ and the following inequality holds $A_2 + tB_2 - \gamma_2 < 0$, we have that $U(t) \preceq_{se} T(U(t))$ for $0 \leq t \leq x_2$ and $x_3 \leq t \leq (\gamma_2 - A_2)/B_2$ and $T(U(t)) \preceq_{se} U(t)$ for $x_2 \leq t \leq x_3$.

By using monotonicity of T , we have that for $0 \leq t < x_2$, $T^n(U(t)) \preceq T^{n+1}(U(t)) \preceq E_2$, and for $x_2 \leq t < x_3$, $E_2 \preceq T^{n+1}(U(t)) \preceq T^n(U(t)) \preceq E_3$. This implies $T^n(U(t)) \rightarrow E_2$ as $n \rightarrow \infty$. Similarly, for $x_3 \leq t \leq (\gamma_2 - A_2)/B_2$, we have $E_3 \preceq T^n(U(t)) \preceq T^{n+1}(U(t)) \preceq E_1$. This implies $T^n(U(t)) \rightarrow E_1$ as $n \rightarrow \infty$. By using the property of points $U(t)$ and $N(u)$, we have that for each point $(x, y) \in \text{int}(Q_4(E_3) \cap \mathcal{R})$, there exists $x_3 < t^* < (\gamma_2 - A_2)/B_2$ and $u^* > 0$ such that $U(t^*) \preceq (x, y) \preceq N(u^*)$. By using monotonicity of T , we have $T^n(U(t^*)) \preceq T^n((x, y)) \preceq T^n(N(u^*))$. This implies $T^n((x, y)) \rightarrow E_1$ as $n \rightarrow \infty$. Furthermore, for each point $(x, y) \in \text{int}(Q_2(E_3) \cap \mathcal{R})$, there exist $t_1 > 0$ and t_2 , $x_2 < t_2 < x_3$ such that $M(t_1) \preceq (x, y) \preceq U(t_2)$. By using monotonicity of T , we have $T^n(M(t_1)) \preceq T^n((x, y)) \preceq T^n(U(t_2))$. This implies $T^n((x, y)) \rightarrow E_2$ as $n \rightarrow \infty$.

Now, assume that parameters $\alpha_1, A_1, \gamma_2, A_2$, and B_2 satisfy the condition (8) and inequality 1.i) of Lemma 1. Then the set

$$\mathcal{I} = \left\{ \left(x, \frac{A_1A_2\beta_1 + xA_1B_2\beta_1}{B_2\alpha_1 - A_2\beta_1} \right) : x \geq 0 \right\}$$

is invariant and contains the equilibrium point E_3 and $T(x, y) = E_3$ for $(x, y) \in \mathcal{I}$. In view of the uniqueness of global stable manifold, we conclude that $\mathcal{W}^s(E_3) = \mathcal{I}$. Take any point $(x, y) \in \mathcal{W}^+(E_3)$, then there exists the point $(x_b, y_b) \in \mathcal{I}$ such that $(x_b, y_b) \ll_{se} (x, y)$. Since, the map T is strongly competitive, then $E_3 = T(x_b, y_b) \ll_{se} T(x, y)$. This implies $T(x, y) \in \text{int}(Q_4(E_3) \cap \mathcal{R})$. Similarly, if $(x, y) \in \mathcal{W}^-(E_3)$, then $T(x, y) \in \text{int}(Q_2(E_3) \cap \mathcal{R})$. The rest of the proof is similar to the proof of the first case. This completes the proof.

(\mathcal{R}_{11}) The first part of theorem follows from Theorems 15 and 16. If parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2$, and B_2 do not satisfy the condition (8) of Lemma 1, then the map T , defined on the set $\mathcal{R} = \mathbb{R}_{+,n}^2$, satisfies all conditions of Theorems 4, 6, and 8. This implies that there exists an invariant curve \mathcal{C} , which is a subset of the basin of attraction of the equilibrium point E , which separates the first quadrant into two invariant regions, $\mathcal{W}^+(E)$ (below the stable manifold) and $\mathcal{W}^-(E)$ (above the stable manifold) which are connected.

By Theorems 6 and 7 and 8 for all $(x, y) \in \mathcal{W}^+(E)$, there exists $n_0 > 0$ such that $T^n((x, y)) \in \text{int}(Q_4(E) \cap \mathcal{R})$ for $n > n_0$. For all $(x, y) \in \mathcal{W}^-(E)$, there exists $n_1 > 0$ such that for all $n > n_1$, $T^n((x, y)) \in \text{int}(Q_2(E) \cap \mathcal{R})$. Now, we show that each orbit starting in the region $\text{int}(Q_4(E))$ converges to E_1 , and each orbit starting in the region $\text{int}(Q_2(E))$ converges to E .

Now, for $0 \leq t \leq (\gamma_2 - A_2)/B_2$, take $U(t) = (t, -A_2 - tB_2 + \gamma_2)$. Since $\alpha_1 = (A_1 - A_2 - \beta_1 + \gamma_2)^2/(4B_2)$, it is easy to see that the following holds

$$U(\bar{x}) = E = E_2 = E_3 \preceq E_1 \quad \text{where } \bar{x} = x_2 = x_3 \quad \text{and}$$

$$U(t) - T(U(t)) = \left(-\frac{(-A_1 + A_2 + 2tB_2 + \beta_1 - \gamma_2)^2}{4B_2(A_1 - A_2 - tB_2 + \gamma_2)}, 0 \right).$$

Since $A_2 + tB_2 - \gamma_2 < 0$, we have $U(t) \preceq_{se} T(U(t)) \preceq$ for $0 \leq t \leq (\gamma_2 - A_2)/B_2$.

By using monotonicity of T , we have that $T^n(U(t)) \preceq T^{n+1}(U(t)) \preceq E$ for $0 \leq t < \bar{x}$. This implies $T^n(U(t)) \rightarrow E$ as $n \rightarrow \infty$. Similarly, for $\bar{x} \leq t < (\gamma_2 - A_2)/B_2$, $E \preceq T^n(U(t)) \preceq T^{n+1}(U(t)) \preceq E_1$. This implies $T^n(U(t)) \rightarrow E_1$ as $n \rightarrow \infty$. By using the property of the points $U(t)$ and $N(u)$, we have that for each point $(x, y) \in \text{int}(Q_4(E) \cap \mathcal{R})$, there exist $\bar{x} < t^* < (\gamma_2 - A_2)/B_2$ and $u^* > 0$ such that $U(t^*) \preceq (x, y) \preceq N(u^*)$. By using monotonicity of T , we have that $T^n(U(t^*)) \preceq T^n((x, y)) \preceq T^n(N(u^*))$. This implies $T^n((x, y)) \rightarrow E_1$ as $n \rightarrow \infty$. Furthermore, for each point $(x, y) \in \text{int}(Q_2(E) \cap \mathcal{R})$ there exists $t_1 > 0$ such that $M(t_1) \preceq (x, y) \preceq E$. By using monotonicity of T , we have $T^n(M(t_1)) \preceq T^n((x, y)) \preceq E$. This implies $T^n((x, y)) \rightarrow E$ as $n \rightarrow \infty$.

Now, assume that parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2$, and B_2 satisfy the condition (8) and inequality 1.i) of Lemma 1. The proof of Theorem is similar to the proof of Theorem in the regions (\mathcal{R}_9) and (\mathcal{R}_{10}) .

$(\mathcal{R}_{12}, \mathcal{R}_{13})$ The first part of theorem follows from Theorems 15 and 17. If parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2$, and B_2 do not satisfy (8) of Lemma 1, then the map T , defined on the set $\mathcal{R} = \mathbb{R}_+^2$, satisfies all conditions of Theorems 4,6, and 8. This implies that there exists an invariant curve \mathcal{C} , which is a subset of the basin of attraction of the equilibrium point E , and which separates the first quadrant into two invariant regions, $\mathcal{W}^+(E)$ (below the stable manifold) and $\mathcal{W}^-(E)$ (above the stable manifold) which are connected.

By Theorems 6 and 8 for all $(x, y) \in \mathcal{W}^+(E)$, there exists $n_0 > 0$ such that $T^n((x, y)) \in \text{int}(Q_4(E) \cap \mathcal{R})$ for $n > n_0$, and for all $(x, y) \in \mathcal{C}(E)$, there exists $n_1 > 0$ such that $T^n((x, y)) \in \text{int}(Q_2(E) \cap \mathcal{R})$ for all $n > n_1$. Now, we show that each orbit starting in the region $\text{int}(Q_4(E))$ is asymptotic to $(\infty, 0)$ and each orbit starting in the region $\text{int}(Q_2(E))$ converges to E .

Since $\alpha_1 = (A_1 - A_2 - \beta_1 + \gamma_2)^2/(4B_2)$, for $0 \leq t \leq (\gamma_2 - A_2)/B_2$, we have $U(t) = (t, -A_2 - tB_2 + \gamma_2)$. It is easy to see

$$U(\bar{x}) = E = E_2 = E_3 \quad \text{where } \bar{x} = x_2 = x_3 \quad \text{and}$$

$$U(t) - T(U(t)) = \left(-\frac{(-A_1 + A_2 + 2tB_2 + \beta_1 - \gamma_2)^2}{4B_2(A_1 - A_2 - tB_2 + \gamma_2)}, 0 \right).$$

Since $A_2 + tB_2 - \gamma_2 < 0$, for $0 \leq t \leq (\gamma_2 - A_2)/B_2$, we have $U(t) \preceq_{se} T(U(t))$.

By using monotonicity of T , we have $E \preceq T^n(U(t)) \preceq T^{n+1}(U(t)) \preceq E_1$ for $0 \leq t < \bar{x}$. This implies $T^n(U(t)) \rightarrow E$ as $n \rightarrow \infty$. Similarly, $E \preceq T^n(U(t)) \preceq T^{n+1}(U(t)) \preceq (\infty, 0)$ for $\bar{x} < t^* < (\gamma_2 - A_2)/B_2$. This implies $T^n(U(t)) \rightarrow (\infty, 0)$ as $n \rightarrow \infty$. For each point (x, y)

$\in \text{int}(Q_4(E_3) \cap \mathcal{R})$, there exists $\bar{x} < t^* < (\gamma_2 - A_2)/B_2$ such that $0 \leq t < \bar{x}$. $0 \leq t < \bar{x}$. By monotonicity of T , we have $T^n(U(t^*)) \preceq T^n((x, y)) \preceq (\infty, 0)$. This implies $T^n((x, y)) \rightarrow (\infty, 0)$ as $n \rightarrow \infty$. Furthermore, for each point $(x, y) \in \text{int}(Q_2(E_3) \cap \mathcal{R})$, there exists $t_1 > 0$ such that $U(t^*) \preceq (x, y) \preceq N(u^*)$. By monotonicity of T , we have $T^n(M(t_1)) \preceq T^n((x, y)) \preceq E$. This implies $T^n((x, y)) \rightarrow E$ as $n \rightarrow \infty$.

If parameters $\alpha_1, \beta_1, A_1, \gamma_2, A_2$, and B_2 satisfy the condition (8) and inequality 1.i) of Lemma 1, then the proof of Theorem is similar to the proof of parts (\mathcal{R}_9) and (\mathcal{R}_{10}) . This completes the proof of Theorem in the regions $\mathcal{R}_{12}, \mathcal{R}_{13}$. This is an example of semistable non-hyperbolic equilibrium point.

$(\mathcal{R}_i, i = \overline{14, 17})$ Assumptions of this theorem imply that equilibrium does not exist.

Set $M(t) = (0, t)$ for $t \geq \gamma_2 - A_2$. Since $M(t) - T(M(t)) = \left(-\frac{\alpha_1}{t + A_1}, \frac{t(t + A_2 - \gamma_2)}{t + A_2} \right)$, we have $M(t) \preceq T(M(t))$ for $t \geq \gamma_2 - A_2$. By using monotonicity $T^n(M(t)) \preceq T^{n+1}(M(t))$, for all $n = 0, 1, 2, \dots$. Set $(x_n^*, \gamma_n^*) = T^n(M(t))$. This implies that $\{\gamma_n^*\}$ is non-increasing and bounded, hence it has to converge. Set $\lim_{n \rightarrow \infty} \gamma_n^* = \gamma^*$. Since $\{x_n^*\}$ is unbounded and non-decreasing, we have that $x_n^* \rightarrow \infty$. By using the second equation of the System (1), we see that $\gamma^* = 0$. Take any point $(x, y) \in [0, \infty)^2$. Then there exists t^* , such that $M(t^*) \preceq (x, y) \preceq (\infty, 0)$. By using monotonicity, $T^n(M(t^*)) \preceq T^n((x, y)) \preceq (\infty, 0)$ as $T^n(M(t^*)) \rightarrow (\infty, 0)$ as $n \rightarrow \infty$, we obtain $T^n((x, y)) \rightarrow (\infty, 0)$ as $n \rightarrow \infty$, as which completes the proof of theorem.

Author details

¹Department of Mathematics, University of Sarajevo, Sarajevo, Bosnia and Herzegovina ²Department of Mathematics, University of Rhode Island, Kingston, RI 02881-0816, USA

Authors' contributions

All authors contributed equally to the manuscript and read and approved the final draft.

Competing interests

The authors declare that they have no competing interests.

Received: 26 January 2011 Accepted: 23 August 2011 Published: 23 August 2011

References

1. Camouzis, E, Kulenović, MRS, Ladas, G, Merino, O: Rational systems in the plane. *J Differ Equ Appl.* **15**, 303–323 (2009). doi:10.1080/10236190802125264
2. Kulenović, MRS, Merino, O: Invariant manifolds for competitive discrete systems in the plane. *Int J Bifurcat Chaos.* **20**, 2471–2486 (2010). doi:10.1142/S0218127410027118
3. Kulenović, MRS, Merino, O: Global bifurcation for competitive systems in the plane. *Discret Cont Dyn Syst B.* **12**, 133–149 (2009)
4. AlSharawi, Z, Rhouma, M: Coexistence and extinction in a competitive exclusion Leslie/Gower model with harvesting and stocking. *J Differ Equ Appl.* **15**, 1031–1053 (2009). doi:10.1080/10236190802459861
5. Cushing, JM, Leverage, S, Chitnis, N, Henson, SM: Some discrete competition models and the competitive exclusion principle. *J Differ Equ Appl.* **10**, 1139–1152 (2004). doi:10.1080/10236190410001652739
6. Kulenović, MRS, Nurkanović, M: Asymptotic behavior of a linear fractional system of difference equations. *J Inequal Appl.* 127–143 (2005)
7. Clark, D, Kulenović, MRS, Selgrade, JF: Global asymptotic behavior of a two dimensional difference equation modelling competition. *Nonlinear Anal TMA.* **52**, 1765–1776 (2003). doi:10.1016/S0362-546X(02)00294-8
8. Hirsch, MW: Systems of differential equations which are competitive or cooperative. I *Limits Sets SIAM J Math Anal.* **13**(2):167–179 (1982)
9. Hirsch, M, Smith, H: Monotone dynamical systems. In: Canada A, Drabek P, Fonda A (eds.) *Handbook of Differential Equations: Ordinary Differential Equations*, vol. II, pp. 239–357. Elsevier, Amsterdam (2005)
10. Kalabušić, S, Kulenović, MRS, Pilav, E: Global dynamics of a competitive system of rational difference equations in the plane. *Adv Differ Equ* **30** (2009). Article ID 132802
11. Kulenović, MRS, Merino, O: *Discrete Dynamical Systems and Difference Equations with Mathematica*. Chapman & Hall/CRC Press, Boca Raton (2002)
12. Kulenović, MRS, Merino, O: Competitive-exclusion versus competitive-coexistence for systems in the plane. *Discret Cont Dyn Syst Ser B.* **6**, 1141–1156 (2006)

13. Leonard, WJ, May, R: Nonlinear aspects of competition between species. *SIAM J Appl Math.* **29**, 243–275 (1975). doi:10.1137/0129022
14. Smith, HL: Invariant curves for mappings. *SIAM J Math Anal.* **17**, 1053–1067 (1986). doi:10.1137/0517075
15. Smith, HL: Periodic competitive differential equations and the discrete dynamics of competitive maps. *J Differ Equ.* **64**, 165–194 (1986). doi:10.1016/0022-0396(86)90086-0
16. Smith, HL: Periodic solutions of periodic competitive and cooperative systems. *SIAM J Math Anal.* **17**, 1289–1318 (1986). doi:10.1137/0517091
17. Smith, HL: Planar competitive and cooperative difference equations. *J Differ Equ Appl.* **3**, 335–357 (1998). doi:10.1080/10236199708808108
18. Smith, HL: Non-monotone systems decomposable into monotone systems with negative feedback. *J Math Biol.* **53**, 747–758 (2006). doi:10.1007/s00285-006-0004-3
19. Takáč, P: Domains of attraction of generic ω -limit sets for strongly monotone discrete-time semigroups. *J Reine Angew Math.* **423**, 101–173 (1992)
20. de Mottoni, P, Schiaffino, A: Competition systems with periodic coefficients: a geometric approach. *J Math Biol.* **11**, 319–335 (1981). doi:10.1007/BF00276900
21. Hess, P: Periodic-Parabolic Boundary Value Problems and Positivity. In Pitman Research Notes in Mathematics Series, vol. 247, Longman Scientific & Technical, Harlow (1991)
22. Kocic, V, Ladas, G: Global Behavior of Nonlinear Difference Equations of Higher Order with Applications. Kluwer, Dordrecht (1993)

doi:10.1186/1687-1847-2011-29

Cite this article as: Kalabušić *et al.*: Dynamics of a two-dimensional system of rational difference equations of Leslie–Gower type. *Advances in Difference Equations* 2011 **2011**:29.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
