1-25-2015

Manipulation of Host Quality and Defense by a Plant Virus Improves Performance of Whitefly Vectors

Qi Su

Evan L. Preisser

University of Rhode Island, preisser@uri.edu

Wen Xie

Shao Li Wang

Qin Jun Wu

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/bio_facpubs

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Qi Su, Evan L. Preisser, Xiao Mao Zhou, Wen Xie, Bai Ming Liu, Shao Li Wang, Qing Jun Wu, and You Jun Zhang. (2015). "Manipulation of host quality and defense by a plant virus improves performance of whitefly vectors." *Journal of Economic Entomology*. 108(1): 11-19. Available at: <http://dx.doi.org/10.1093/jee/tou012>

This Article is brought to you for free and open access by the Biological Sciences at DigitalCommons@URI. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Authors

Qi Su, Evan L. Preisser, Wen Xie, Shao Li Wang, Qin Jun Wu, and You Jun Zhang

Running title: Plant-virus-whitefly interactions

**TITLE: Manipulation of host quality and defense by a plant virus improves
performance of whitefly vectors**

Qi Su¹, Evan L. Preisser², Wen Xie¹, Shao Li Wang¹, Qin Jun Wu¹, You Jun Zhang^{1*}

¹*Department of Plant Protection, Institute of Vegetables and Flowers, Chinese Academy
of Agricultural Sciences, Beijing 100081, China*

²*Department of Biological Sciences, University of Rhode Island, Kingston RI 02881 USA*

* Corresponding Author:

Dr. Youjun Zhang

Department of Entomology, Institute of Vegetables and Flowers

Chinese Academy of Agricultural Sciences

12 Zhongguancun Nandajie, Haidian District

Beijing 100081, China

Phone: 86-10-82109518

Fax: 86-10-82109518

Email: zhangyoujun@caas.cn

Total words in text (excluding references, tables, and figure legends): 4,490

Abstract

Pathogen-mediated interactions between insect vectors and their host plants can affect herbivore fitness and the epidemiology of plant diseases. While the role of plant quality and defense in mediating these tripartite interactions has been recognized, there are many ecologically- and economically-important cases where the nature of the interaction has yet to be characterized. The *Bemisia tabaci* cryptic species MED is an important vector of *tomato yellow leaf curl virus* (TYLCV), and performs better on virus-infected tomato than on uninfected controls. We assessed the impact of TYLCV infection on plant quality and defense, and the direct impact of TYLCV infection on MED feeding. We found that although TYLCV infection has a minimal direct impact on MED, the virus alters the nutritional content of leaf tissue and phloem sap in a manner beneficial to MED. TYLCV infection also suppresses herbivore-induced production of plant defensive enzymes and callose deposition. The strongly positive net effect on TYLCV on MED is consistent with previously-reported patterns of whitefly behavior and performance, and provides a foundation for further exploration of the molecular mechanisms responsible for these effects and the evolutionary processes that shape them.

Keywords

Tomato yellow leaf curl virus, *Bemisia tabaci* MED, *Solanum lycopersicum*, persistent transmission, plant defense, mutualism, plant–virus–vector interactions

Introduction

Phloem-feeding insects are major pests of many agricultural crops. In addition to the feeding-related damage they cause, these insects can also serve as vectors for a wide variety of economically-important plant viruses (Jones 2003). Because virtually all of these viruses require a vector for between-host dispersal, insect behavior can affect pathogen success; as a result, there is likely to be strong selection for viral traits capable of manipulating plant-insect interactions in a manner that optimizes pathogen transmission (Hogenhout et al. 2008). Research testing this hypothesis has found that viruses can alter plant defense, nutritional composition, and other traits in ways that the preference, performance, and dispersal of viral vectors (Eigenbrode et al. 2002, Belliure et al. 2005, Ingwell et al. 2012, Mauck et al. 2012, Liu et al. 2013b, Moreno-Delafuente et al. 2013). The resulting changes in plant-insect interactions can improve viral transmission and alter epidemiological patterns (Sisterson 2008, Ingwell et al. 2012, Roosien et al. 2013).

Because of the ecological and agricultural importance of virus-vector-host interactions, there has been a surge of interest in the biochemical and physiological mechanisms underlying virally-mediated changes to host-vector interactions (Luan et al. 2014). In the case of the persistently-transmitted *tomato yellow leaf curl China virus* (TYLCCNV), for example, viral infection of tobacco (*Nicotiana tabacum*) improves nutritional assimilation by the whitefly vector *Bemisia tabaci* MEAM1 (formerly called the ‘B’ biotype) and suppresses both terpenoid and jasmonic acid defenses against MEAM1 (Wang et al. 2012, Zhang et al. 2012, Luan et al. 2013b). Interactions between *B. tabaci*, host plants, and persistently-transmitted begomoviruses have been of particular interest since this whitefly ‘species’ (actually a species complex that encompasses MEAM1 and several other cryptic but genetically-distinct species; De Barro et al.

2011) is a major agricultural pest and viral vector that has been named one of the world's '100 worst invasive species' (Lowe et al. 2000).

A recent review (Luan et al. 2014) documented considerable progress in addressing plant-mediated whitefly–begomovirus interactions, especially in regards to the highly-invasive MEAM1. Less is known, however, about vector-virus-host interactions involving *B. tabaci* MED (formerly the 'Q' biotype). In their review, Luan et al (2014) documented nine studies assessing MEAM1 performance on infected versus control plants, but only four studies involving MED; they also found four and zero studies addressing the direct effect of begomovirus infection on MEAM1 and MED, respectively. The lack of research on MED is noteworthy because this cryptic species is also invasive and a major pest; although generally competitively inferior to MEAM1, MED is more tolerant of insecticides and has displaced MEAM1 throughout China and other Asian countries (Crowder et al. 2010, Pan et al. 2011, Pan et al. in review).

MED and MEAM1 also differ in their relationship to *tomato yellow leaf curl virus* (TYLCV), a complex of circular, single-stranded DNA plant geminiviruses that infects tomato (*Solanum lycopersicum*) and is transmitted by *B. tabaci* in a persistent and circulative manner (Hogenhout et al. 2008, Ghanim 2014). Recent research has revealed that TYLCV-infected plants have different effects on MEAM1 and MED feeding and host preference (Fang et al. 2013, Liu et al. 2013b) and induce salicylic-acid defenses against MEAM1 but not MED (Shi et al. 2013). Perhaps as a result, TYLCV appears to have a mutualistic/neutral relationship with MED (Matsuura and Hoshino 2009, Li et al. 2011, Pan et al. 2013) but a neutral/parasitic relationship with MEAM1 (Liu et al. 2009, Pan et al. 2013).

We report the results of research investigating the biochemical and physiological mechanisms underlying the virus-plant-host relationship in order to address the direct and

indirect impacts of TYLCV infection on *S. lycopersicum*, MED, and the insect-plant interaction. We found that although TYLCV infection has a minimal direct impact on MED, the virally-mediated improvement in plant nutritional traits and reductions in host plant defenses is so beneficial that the net interaction is strongly mutualistic.

Materials and Methods

Tomato (*S. lycopersicum* Miller, cv Zhongza 9) and cotton (*Gossypium hirsutum* L., cv DP99B) plants were grown in potting mix and raised individually in 1.5 L pots. They were enclosed in whitefly-proof screen cages under natural lighting and controlled temperature ($26 \pm 2^{\circ}\text{C}$) in a glasshouse. Cotton plants were used at the 6–7 true-leaf stage; tomato plants were used at the 3–4 true-leaf stage for viral inoculation and 6–7 true-leaf stage for all other experiments. Plants were watered every 3–4 days as necessary.

The MED used in this study originated from the Haidian District of Beijing, where it was collected in 2009 from poinsettia (*Euphorbia pulcherrima* Wild. ex Klotz.). It was reared on *S. lycopersicum* in screen cages under natural lighting and controlled temperature in a glasshouse.

TYLCV inoculation: We infected tomato plants with TYLCV using *Agrobacterium tumefaciens*-mediated inoculation; an infectious clone (pBINPLUS-SH2-1.4A) of TYLCV- Israel [CN: SH2] was constructed using *A. tumefaciens* strain EHA105 (Zhang et al. 2009). TYLCV-infected plants were produced by inoculation at the 3–4 true-leaf stage (Zhang et al. 2009). Infection was determined visually and confirmed via PCR validation with primers TYLCV-473 and TYLCV-61 (Ghanim et al. 2007). Control plants were mock-inoculated using the *A. tumefaciens* strain EHA105 empty vector to account for mechanical inoculation.

Viral transmission assays: To assess the likelihood of whitefly infection with TYLCV, we allowed 20 MED to feed on a TYLCV-infected tomato plant for 10 h. **Because multiple**

previous studies (reviewed in Ghanim 2014) have shown that MED is both able to acquire TYLCV from infected tomato plants and transmit it to uninfected plants, we conducted both this work and the research described in the following paragraph solely to ensure that our lines were performing as expected. We extracted DNA from each whitefly as per White et al. (2009), and TYLCV presence was verified via PCR validation as above. We repeated this procedure using 20 MED and an uninfested tomato plant.

To assess the likelihood of TYLCV transmission to neighboring plants, we established whitefly colonies on tomato that had been infected with TYLCV for three weeks. Following whitefly colonization, two of the TYLCV-infected plants were placed in an arena containing five healthy plants. Whiteflies from the TYLCV-infected plants were allowed to move throughout the arena for two weeks. DNA was then extracted from the apical leaves of each healthy plant and TYLCV infection assessed using PCR (Ghanim et al. 2007).

Impact of TYLCV on MED mass, fecundity, and survival: To assess the impact of TYLCV on MED mass, 40 two-day-old adult whiteflies were first weighed singly and then placed individually into a clip cage attached to either the third- or fifth-to-bottom leaf of either a TYLCV-inoculated (n=10) or mock-inoculated (n=10) plant (two whiteflies per plant). After seven days, we recorded the final weight of each whitefly.

To assess the impact of TYLCV on MED fecundity and survival, 300 two-day-old mated female whiteflies were collected from uninfected tomato plants. Ten whiteflies were placed in a clip-cage (3 cm diameter × 4 cm height) attached to the fifth-from-bottom leaf of either a mock-inoculated (n=15) or TYLCV-inoculated (n=15) tomato plant. After seven days, we counted whitefly eggs and live adults within the clip-cage on each replicate.

Impact of TYLCV on MED nutritional assimilation: TYLCV could affect MED nutritional assimilation directly (via changes in the insect itself) and/or indirectly (via changes in the infected host plant that alter its nutritional quality for whiteflies). Assessing the direct impact of TYLCV infection by feeding viruliferous and uninfected MED on uninfected tomato plants could infect the plant and potentially alter plant nutritional quality; as a result, tomato plants cannot be used to isolate the direct effect of TYLCV infection on MED nutrient assimilation. We overcame this obstacle by allowing viruliferous and uninfected MED to feed on cotton, a non-host plant of TYLCV, and analyzing their excreted honeydew.

Viruliferous and uninfected whiteflies were obtained by allowing newly-emerged adults to feed on TYLCV-infected or mock-inoculated tomato plants for one day; after one day spent feeding on virus-infected plants, a PCR analysis of 20 randomly-selected whiteflies detected TYLCV in all of them (Su et al. 2013a). A group of 200 viruliferous whiteflies and another group of 200 non-viruliferous whiteflies were placed in two separate clip cages on different leaves of the same cotton plant; this procedure was replicated for six different cotton plants. A 16 cm² tin foil square was placed beneath each leaf to collect honeydew. Whiteflies were removed after one week, and each clip cage and its corresponding tin-foil square were rinsed with 1ml of deionized water and stored at -20°C for analysis. After one week of feeding on cotton, a PCR analysis of 20 randomly-selected whiteflies found that all of them were still viruliferous (Su et al. 2013a). The honeydew was analyzed for amino acid composition, percent amino acids, and the sugar:amino acid ratio using the procedures detailed in the following section.

Impact of TYLCV on plant nutritional composition: We analyzed the epidermis, mesophyll tissue, and phloem sap of 6-7 true-leaf stage plants that were either TYLCV-infected (n=6) or mock-inoculated (n=6). We sampled the two most-recently expanded leaves (fifth- and

sixth-from-bottom) from each plant between 8:00-12:00 am. Phloem sap was sampled from the sixth expanded leaf, which was cut four cm down the petiole and placed into 1.5 ml of 20 mM EDTA solution (pH 7.0). Leaves were chilled in an ice bath housed in a dark box (to prevent transpiration) following collection, then stored at -20°C until analysis. Epidermal and mesophyll tissue was sampled from the fifth leaf using a two-cm diameter cork borer that allowed sampling between major veins. Five leaf discs per plant were collected, weighed, flash-frozen in 1.5 ml Eppendorf tubes, and held at -80°C until analysis.

Carbohydrate and amino acid determination: Leaf disc samples were ground in liquid nitrogen and extracted using 1 ml of pH = 3.0 extraction liquid (ethanol/distilled water/HCl, 2:1:0.004 v/v) spiked with internal standards of the metabolites of interest. Phloem sap and honeydew samples were combined with the internal standard and 0.3 ml of chloroform, vortexed, and centrifuged at 12,000 g for two minutes before the organic material was removed. The aqueous fraction containing amino acids and carbohydrates was placed in an Eppendorf tube and dried in a Speed-vac; the dry extracts were suspended in 0.5 ml of double-distilled water.

Carbohydrates were purified using 3.5 g g^{-1} plant material ion exchange resins. Samples were concentrated to 0.4 ml and filtered through a $0.45\text{-}\mu\text{m}$ filter; $20\text{ }\mu\text{l}$ was injected and analyzed by HPLC using a Hi-Plex H column ($300\text{ mm} \times 7.7\text{ mm}$ column; Agilent, Palo Alto, CA, USA) flushed with 0.6 ml min^{-1} double-distilled water at 85°C with a refractive index detector (Waters). Carbohydrates were identified using reference sugars, and quantified with standard curves.

Amino acids were analyzed by reverse-phase HPLC with pre-column derivatization using o-phthaldialdehyde (OPA) and 9-fluorenylmethyloxycarbonyl (FMOC). Amino acids were quantified using the AA-S-17 (Agilent) reference amino-acid mixture, supplemented with

asparagine, glutamine and tryptophan (Sigma-Aldrich Co., St. Louis, MO, USA). Analyses were performed using an Agilent 1100 HPLC; a reverse-phase Agilent Zorbax Eclipse C18 column AAA (5 μ m, 250 $\times$ 4.6 mm) and fluorescence detector were used for chromatographic separation. Amino acids were quantified by comparing peak areas to the standard curve of each reference amino acid. Peak areas were converted to ng amounts relative to the known internal standard added to each sample, and corrected for leaf tissue weight. Peak areas for both phloem sap and honeydew were similarly converted to ng amounts based on the internal standard. Total sugar contents were expressed in terms of total monosaccharide contents to calculate the sugar:amino acid ratio of the epidermis and mesophyll tissue, phloem sap, and honeydew.

Impact of TYLCV infection and MED infestation on plant defensive enzymes and callose deposition: Fifty adult whiteflies were placed in a clip cage attached to a leaf of either TYLCV-infected (n=6) or mock-inoculated (n=6) plants. In the whitefly-free treatment, empty clip cages were attached to a leaf of either TYLCV-infected (n=6) or mock-inoculated (n=6) plant. Whiteflies were removed after two days and the plant tissue within each clip cage from all four treatments was immediately harvested and stored in liquid nitrogen.

Quantification of enzyme activity: The defensive enzymes phenylalanine ammonia lyase (PAL), peroxidase (POD), polyphenol oxidase (PPO), and superoxide dismutase (SOD) were extracted from 0.5 g frozen tissue by grinding in a 50 mM Tris-HCl buffer (pH = 7.5, 3 ml g⁻¹ of leaf tissue) containing 7% polyvinyl polypyrrolidone (PVPP), 1.67mM phenylthiourea, 0.3 M KCl, and 0.4 mM ascorbic acid. The thawed extract was centrifuged at 13,000 g for 10 min and enzyme activity measured in the supernatant. PAL, POD, and PPO activity was quantified according to Guo et al. (2012); SOD activity was quantified according to Zhang et al. (2008). Soluble protein was quantified by the dye-binding method (Bradford 1976) with bovine serum

albumin as a standard. Results are expressed in units per mg protein; treatment values are the mean of six replicates. Analyses were conducted using a fluorescence microplate reader (SpectraMax M2e, Molecular Devices) at room temperature ($25 \pm 2^{\circ}\text{C}$).

Callose visualization: Leaf samples were placed in 70% ethanol for one hour, 95% ethanol with chloroform overnight, and 100% ethanol for two hours to clear the chlorophyll. Samples were next washed in 0.1M phosphate buffer (pH 7.0) and incubated for 15 min in the same buffer containing 0.005% calcofluor (Fluorescent Brightener, Sigma, USA) and 0.01% aniline blue. Leaves were then washed repeatedly in 0.01% aniline blue in 0.1M phosphate buffer (pH 7.0), mounted in glycerol (Ton and Mauch-Mani 2004), then examined with a Leica DM RA2 microscope with an A4 fluorescence cube. Callose deposits were counted using QUANTITY ONE software (Bio-Rad). Counts from five adjacent fields of view along the length of the leaf (not including the mid-vein or leaf edge) were averaged to generate a mean leaf value. Mean values from 4-6 leaves were averaged to generate a mean treatment value.

Statistical analysis

Prior to analysis, data were checked for normality and homogeneity of variance. In cases where we took multiple samples from, or had multiple whiteflies feed on, a single plant, we averaged the data from that plant to generate a single mean per-plant response. We used one-way ANOVA to compare MED mass, fecundity, survival, and fecundity on TYLCV-infected versus control plants. Because of the large number of amino acids that we quantified, the p values for these analyses were corrected for multiple comparisons at $\alpha = 0.05$ using step-up false discovery rate (FDR), a sequential Bonferroni-type procedure (Benjamini and Hochberg 1995). We used two-way ANOVA to analyze the impact of TYLCV infection (present/absent) and MED infestation (present/absent) on plant defensive enzymes and callose deposition. We performed

means separation tests, where appropriate, using Tukey's HSD. All data were analyzed using JMP 9.0.0 (SAS Institute, Cary NC USA).

Results

TYLCV rapidly infects MED, and MED effectively transmits TYLCV: After feeding on a TYLCV-infected plant for 10 h, PCR validation revealed that all (20/20) of the initially-uninfected MED had become viruliferous. When uninfected MED were allowed to feed on both TYLCV-infected and healthy plants for two weeks, PCR validation revealed that all (5/5) of the previously-uninfected plants tested positive for TYLCV.

TYLCV-infected plants increase MED mass, fecundity, and survival: Whiteflies feeding on TYLCV-infected plants gained 68% more weight than those feeding on mock-inoculated plants (51.3 ± 3.1 [SE] μg and 30.5 ± 2.6 μg , respectively; $F_{1,18} = 33.4$, $p < 0.001$). Fecundity was also higher: whiteflies reared on TYLCV-infected plants laid 81% more eggs (43.9 ± 3.1 eggs and 24.2 ± 2.6 eggs, respectively; $F_{1,28} = 24.1$, $p < 0.001$). Whiteflies also survived 17% longer on TYLCV-infected plants. After one week, $83.3 \pm 3.6\%$ of whitefly adults survived on TYLCV-infected plants, versus $70.7 \pm 3.6\%$ on uninfected plants ($F_{1,28} = 6.2$, $p = 0.019$).

TYLCV alters MED nutritional assimilation: Honeydew excreted by viruliferous whiteflies had a sugar:amino acid ratio half that of honeydew from uninfected whiteflies (0.34 ± 0.04 vs. 0.68 ± 0.05 ; $F_{1,10} = 27.1$, $p < 0.001$). Their honeydew did not differ in the percentage of essential amino acids ($F_{1,10} = 1.56$, $p = 0.24$), however, nor in any of 16 individual amino acids ($F_{1,10}$, all $p > 0.05$ after correction for multiple comparisons).

TYLCV improves plant nutritional composition for MED: Free amino acid concentrations were 55% lower in the epidermis and mesophyll tissues of infected versus uninfected plants (573 ± 35 ng mg^{-1} tissue vs. 1267 ± 66 ng mg^{-1} tissue; $F_{1,10} = 85.2$, $p < 0.001$).

The concentrations of all 20 amino acids were lower in infected plants; 13 of these differences were significant after correction for multiple comparisons (Fig. 1A). There were, however, no differences in simple carbohydrates (sucrose, glucose, and fructose; Fig. 1B); as a result, the sugar:amino acid ratio in the epidermis and mesophyll tissues of infected plants was 2.5x higher than that of uninfected plants (6.42 ± 0.34 vs. 2.53 ± 0.25 ; $F_{1,10} = 83.6$, $p < 0.001$).

In contrast to the epidermis and mesophyll tissue, free amino acid concentrations were 92% higher in the phloem sap of infected plants (4985 ± 170 ng sample⁻¹ vs. 2596 ± 70 ng sample⁻¹ in uninfected plants; $F_{1,10} = 168.8$, $p < 0.001$). Concentrations of all 20 individual amino acids were higher in the phloem of infected plants; 17 of these differences were significant after correction for multiple comparisons (Fig. 1C). Because infected plants also had higher concentrations of simple carbohydrates (Fig. 1D), however, there were no between-treatment differences in the sugar:amino acid ratio of their phloem sap (3.12 ± 0.11 in infected vs. 2.91 ± 0.17 in uninfected; $F_{1,10} = 1.1$, $p = 0.33$).

TYLCV infection reduces plant defensive response to MED infestation: Both TYLCV infection and MED infestation significantly altered PAL (Fig. 2A), POD (Fig. 2B), PPO (Fig. 2C), and SOD (Fig. 2D) concentrations relative to uninfested tomato plants (main effects of ‘TYLCV’ and ‘MED’ all $p < 0.05$). For PAL, PPO, and SOD, infestation with MED only increased enzyme concentrations in uninfected plants (Tukeys’ HSD, $p < 0.05$). In the case of POD, both MED infestation and TYLCV infection increased enzyme concentrations, but there was no interaction between the two factors (Fig. 2C; TYLCV*MED; $F_{1,20} = 0.92$, $p = 0.35$). In contrast, callose formation was decreased 44% by TYLCV infection and increased 250% by MED infestation (Fig. 2E; TYLCV: $F_{1,16} = 45.9$, $p < 0.001$; MED: $F_{1,16} = 104.4$, $p < 0.001$).

Importantly, infection with TYLCV also reduced MED-induced callose formation by 52% (TYLCV*MED: $F_{1,16} = 26.1$, $p < 0.001$).

Discussion

We found that TYLCV-mediated alterations to plant nutritional traits and defensive responses improve the growth and reproduction of its MED vector. Our results provide a mechanistic basis for the results of several recently-published papers that found MED was preferentially attracted to TYLCV-infected plants (Fang et al. 2013) and performed better on infected versus uninfected plants (Liu et al. 2013b, Pan et al. 2013). Preferential feeding on infected plants improves the likelihood of viral acquisition, and our viral transmission assays confirm that MED can both rapidly acquire and effectively transmit TYCLV. As a result, the beneficial impact of TYLCV infection on MED fitness should favor improved viral transmission. An array of persistently-transmitted viruses have been found to similarly manipulate plant-herbivore relationships; many of these manipulations improve plant resource quality for their insect vectors (Mauck et al. 2012).

Our analysis of honeydew excreted by viruliferous and uninfected whiteflies feeding on cotton found minimal direct impacts of TYLCV infection. This is consistent with work documenting that TYLCV infection does not directly affect MED fitness (Li et al. 2011, Pan et al. 2013). Although honeydew from viruliferous whiteflies had a lower sugar:amino acid ratio, there was no difference in either the percentage of essential amino acids or in any of 16 individual amino acids. The absence of a direct impact of TYLCV on its vector is somewhat surprising in light of the fact that the closely-related *tomato yellow leaf curl China virus* had a negative direct effect on the fecundity and longevity of MEAM1 (Jiu et al. 2007); in their recent

review, Luan et al. (2014) found that four of six studies addressing the direct effects of viral infection on *Bemisia* species noted deleterious impacts.

Our analyses of plant nutritional composition found that TYLCV infection alters the concentrations of simple carbohydrates, amino acids, and the sugar:amino acid ratio in both the epidermis/mesophyll (Fig. 1A,B) and phloem (Fig. 1C,D). Viral manipulation of the epidermis/mesophyll is especially interesting, since whitefly attraction to suitable host plants is mediated by gustatory cues encountered during shallow probes of leaf tissue (Powell et al. 2006). Lower amino acid concentrations in the epidermis/mesophyll increase the sugar:amino acid ratio in infected tissues; higher values of this ratio have been shown to stimulate aphid feeding (Mauck et al. 2014). Aphids, whiteflies, and other phloem-feeding insects use small amounts of watery saliva to dissolve surface chemicals, determine physical features, and taste the chemical defenses of the phylloplane; this pre-phloem assessment of cellular contents plays a critical role in subsequent feeding, oviposition, and dispersal decisions (Walling 2008, Liu et al. 2013a). Phylloplane manipulation by TYLCV provides a basis for virally-mediated changes to the plant's volatile profile (Fang et al. 2013), and may help explain why MED prefers TYLCV-infected plants over healthy ones (Fang et al. 2013, Liu et al. 2013b).

While the nutritional content of the epidermis/mesophyll plays an important role in whitefly perceptions of plant quality, *Bemisia* performance is determined by the phloem on which they feed. While the sugars in phloem sap provide an abundant source of energy, amino acid concentrations (and thus N) are often relatively low; many whiteflies and other phloem-feeding insects overcome this limitation by hosting a complement of nutrient-overproducing bacterial symbionts (Douglas 2006, Su et al. 2013b). The better performance of MED on TYLCV-infected plants (discussed in detail below) suggest that it may be a better food source;

consistent with this hypothesis, we found that the phloem of infected plants had higher concentrations of both sugar and amino acids (Fig. 1C,D). Because whiteflies often use gradients of sucrose or other carbohydrates to locate phloem (Powell et al. 2006), this may help explain why MED locates phloem faster and begins ingesting sap more quickly on virus-infected versus healthy tomato plants (Liu et al. 2013b). This response is also consistent with work by Colvin et al (2006): they found that phloem sap from cassava (*Manihot esculenta* Crantz) infected with *East African cassava mosaic virus*-Uganda had higher concentrations of four essential amino acids, and that *Bemisia* Asia I did better on infected versus uninfected plants.

Because TYLCV does not directly affect MED, the preference for and improved performance of MED of TYLCV-infected plants is almost certainly due to virally-mediated changes in plant physiology (Figs. 1, 2). In addition to the changes in plant nutritional composition (Fig. 1), we also found that TYLCV infection decreased the ability of plants to mount a defensive response to whitefly feeding. The production of reactive oxygen species (ROS) is a rapid and generalized defensive response in plants that can also trigger subsequent defensive reactions (Low and Merida 1996). Plants produce an array of materials that scavenge ROS and protect the plant against ROS-induced oxidative bursts; these include small molecular antioxidants and enzymes such as PAL, POD, PPO, and SOD (Asada 2006). In uninfected plants, whitefly feeding induced increases in POD and SOD (involved in ROS synthesis) as well as in PAL and PPO (involved in phenol oxidation). In contrast, TYLCV infection blocked herbivore-induced increases in PAL, PPO, and SOD production (Fig. 2A-D). This is consistent with work (Luan et al. 2013a) on MEAM1 whiteflies feeding on TYLCV-infected and uninfected tobacco plants. They found that genes involved in both detoxification and redox activity were downregulated in MEAM1 allowed to feed on TYLCV-infected plants, and speculated that this

was in response to infected plants decreasing their *Bemisia*-induced production of ROS and other defenses (Luan et al. 2013a). In summarizing their work, they say that “Reduced detoxification activity is likely to attenuate energy costs, thereby enhancing the performance of whiteflies on virus-infected plants” (p. 597 in Luan et al. 2013a); this statement accords with both our findings of reduced plant defense and improved MED performance.

Callose deposition, a key plant defense that prevents phloem feeding by repairing punctured sieve elements (Walling 2008), increased in response to MED infestation in both infected and uninfected plants (Fig. 2E). The magnitude of the increase was much smaller in TYLCV-infested plants (80%) than in uninfected plants (196%), however, again demonstrating a TYLCV-mediated suppression of this defensive response. Decreased callose deposition may help explain why the mean duration of MED feeding bouts (i.e., the time spent ingesting sap from a single sieve element) was much higher in TYLCV-infected versus control plants (Liu et al. 2013b). This may in turn play a role in MED’s preference for, and better performance on, TYLCV-infected plants (Fang et al. 2013, Pan et al. 2013). This latter finding is also documented in our work (but see Li et al. 2011 and Matsuura and Hoshino 2009 for work finding no impact of TYLCV on MED fitness).

Our research into the mechanistic underpinnings of the MED-TYLCV-tomato interaction helps explicate recent research into this tripartite interaction while complementing similar MEAM1-focused work (reviewed in Luan et al. 2014). We found that the better performance of MED on TYLCV-infected plants is likely linked to improved plant nutritional quality and suppressed plant defenses. The host-mediated benefits of TYLCV infection to MED may explain this whitefly’s attraction to TYLCV-infected plants (Fang et al. 2013), a phenomena that should increase both viral acquisition and transmission. More generally, our findings provide insight

into how virally-induced changes in host plant biochemistry and physiology alter this ecologically- and economically-important interaction. Since begomovirus are among the most widely distributed plant viruses, and plants in natural settings are frequently infected (Hogenhout et al. 2008), future research addressing these tripartite interactions is likely to provide specific benefits while fostering our general understanding of plant–virus–vector interactions.

Funding

This work was supported by the National Science Fund for Distinguished Young Scholars (31025020), the 973 Program (2013CB127602), the National Natural Science Foundation of China (31171857), the Beijing Natural Science Foundation (6131002), and the Beijing Key Laboratory for Pest Control and Sustainable Cultivation of Vegetables. The granting agencies had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank Dr. Xueping Zhou (Institute of Biotechnology, Zhejiang University, Hangzhou, China) for providing the infectious TYLCV clone.

Literature Cited

Asada, K. 2006. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* 141: 391-396.

Belliure, B., A. Janssen, P. C. Maris, D. Peters, and M. W. Sabelis. 2005. Herbivore arthropods benefit from vectoring plant viruses. *Ecol Lett* 8: 70-79.

Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254.

Colvin, J., C. Omongo, M. Govindappa, P. Stevenson, M. Maruthi, G. Gibson, S. Seal,
and V. Muniyappa. 2006. Host-plant viral infection effects on arthropod-vector population
growth, development and behaviour: management and epidemiological implications. *Adv. Virus*
Res. 67: 419-452.

Crowder, D. W., A. R. Horowitz, P. J. De Barro, S. S. Liu, A. M. Showalter, S.
Kontsedalov, V. Khasdan, A. Shargal, J. Liu, and Y. Carrière. 2010. Mating behaviour, life
history and adaptation to insecticides determine species exclusion between whiteflies. *J Anim*
Ecol 79: 563-570.

De Barro, P., S. Liu, L. Boykin, and A. Dinsdale. 2011. *Bemisia tabaci*: A statement of
species status. *Annu Rev Entomol* 56: 1-19.

Douglas, A. E. 2006. Phloem-sap feeding by animals: problems and solutions. *J Exp Bot*
57: 747-754.

Eigenbrode, S., H. Ding, P. Shiel, and P. Berger. 2002. Volatiles from potato plants
infected with potato leafroll virus attract and arrest the virus vector, *Myzus persicae* (Homoptera:
Aphididae). *Proceedings of the Royal Society B: Biological Sciences* 269: 2603-2603.

Fang, Y., X. G. Jiao, W. Xie, S. L. Wang, Q. J. Wu, X. J. Shi, G. Chen, Q. Su, X. Yang,
H. P. Pan, and Y. J. Zhang. 2013. *Tomato yellow leaf curl virus* alters the host preferences of its
vector *Bemisia tabaci*. *Sci. Rep.* 3: 2876.

Ghanim, M. 2014. A review of the mechanisms and components that determine the
transmission efficiency of *tomato yellow leaf curl virus* (Geminiviridae; *Begomovirus*) by its
whitefly vector. *Virus Res.* 186: 47-54.

Ghanim, M., I. Sobol, M. Ghanim, and H. Czosnek. 2007. Horizontal transmission of
begomoviruses between *Bemisia tabaci* biotypes. *Arthropod-Plant Interactions* 1: 195-204.

407 **Guo, H., Y. Sun, Q. Ren, K. Zhu-Salzman, L. Kang, C. Wang, C. Li, and F. Ge. 2012.**
 408 Elevated CO₂ reduces the resistance and tolerance of tomato plants to *Helicoverpa armigera* by
 409 suppressing the JA signaling pathway. PLoS ONE 7: e41426.

410 **Hogenhout, S. A., E.-D. Ammar, A. E. Whitfield, and M. G. Redinbaugh. 2008.** Insect
 411 vector interactions with persistently transmitted viruses. Annu Rev Phytopathol 46: 327-359.

412 **Ingwell, L. L., S. D. Eigenbrode, and N. A. Bosque-Pérez. 2012.** Plant viruses alter insect
 413 behavior to enhance their spread. Sci. Rep. 2: 578.

414 **Jiu, M., X. P. Zhou, L. Tong, J. Xu, X. Yang, F. H. Wan, and S. S. Liu. 2007.** Vector-
 415 virus mutualism accelerates population increase of an invasive whitefly. PLoS ONE 2: e182.

416 **Jones, D. 2003.** Plant viruses transmitted by whiteflies. Eur J Plant Pathol 109: 195-219.

417 **Li, M., J. Liu, and S. S. Liu. 2011.** *Tomato yellow leaf curl virus* infection of tomato does
 418 not affect the performance of the Q and ZHJ2 biotypes of the viral vector *Bemisia tabaci*. Insect
 419 Sci 18: 40-49.

420 **Liu, B. M., E. L. Preisser, X. Jiao, H. P. Pan, W. Xie, S. L. Wang, Q. J. Wu, and Y. J.**
 421 **Zhang. 2013a.** Plant-mediated changes in the feeding behavior of an invasive whitefly. Environ
 422 Entomol 42: 980-986.

423 **Liu, B. M., E. L. Preisser, D. Chu, H. P. Pan, W. Xie, S. L. Wang, Q. J. Wu, X. G. Zhou,**
 424 **and Y. J. Zhang. 2013b.** Multiple forms of vector manipulation by a plant-infecting virus:
 425 *Bemisia tabaci* and *tomato yellow curl leaf virus*. J. Virol. 87: 4929-4937.

426 **Liu, J., H. Zhao, K. Jiang, X. P. Zhou, and S. S. Liu. 2009.** Differential indirect effects of
 427 two plant viruses on an invasive and an indigenous whitefly vector: implications for competitive
 428 displacement. Ann Appl Biol 155: 439-448.

Low, P. S., and J. R. Merida. 1996. The oxidative burst in plant defense: Function and signal transduction. *Physiol Plant* 96: 533-542.

Lowe, S., M. Browne, S. Boudjelas, and M. De Poorter (eds.). 2000. 100 of the world's worst invasive alien species: A selection from the Global Invasive Species Database. The IUCN Invasive Species Specialist Group, New York NY.

Luan, J. B., X. W. Wang, J. Colvin, and S. S. Liu. 2014. Plant-mediated whitefly–begomovirus interactions: research progress and future prospects. *Bull Entomol Res* 104: 267-276.

Luan, J. B., Y. L. Wang, J. Wang, X. W. Wang, and S. S. Liu. 2013a. Detoxification activity and energy cost is attenuated in whiteflies feeding on *Tomato yellow leaf curl China virus*-infected tobacco plants. *Insect Mol Biol* 22: 597-607.

Luan, J. B., D. M. Yao, T. Zhang, L. L. Walling, M. Yang, Y. J. Wang, and S. S. Liu. 2013b. Suppression of terpenoid synthesis in plants by a virus promotes its mutualism with vectors. *Ecol Lett* 16: 390-398.

Matsuura, S., and S. Hoshino. 2009. Effect of tomato yellow leaf curl disease on reproduction of *Bemisia tabaci* Q biotype (Hemiptera: Aleyrodidae) on tomato plants. *Applied Entomology & Zoology* 44: 143-148.

Mauck, K., N. A. Bosque-Pérez, S. D. Eigenbrode, C. M. De Moraes, and M. C. Mescher. 2012. Transmission mechanisms shape pathogen effects on host–vector interactions: evidence from plant viruses. *Funct Ecol* 26: 1162-1175.

Mauck, K. E., C. M. De Moraes, and M. C. Mescher. 2014. Biochemical and physiological mechanisms underlying effects of *Cucumber mosaic virus* on host-plant traits that mediate transmission by aphid vectors. *Plant, Cell Environ.* 37: 1427-1439.

452 **Moreno-Delafuente, A., E. Garzo, A. Moreno, and A. Fereres. 2013.** A plant virus
 453 manipulates the behavior of its whitefly vector to enhance its transmission efficiency and spread.
 454 PLoS ONE 8: e61543.

455 **Pan, H. P., E. L. Preisser, D. Chu, S. L. Wang, Q. J. Wu, X. G. Zhou, and Y. J. Zhang.**
 456 **in review.** Insecticides promote viral outbreaks by altering herbivore competition. Ecol Appl.

457 **Pan, H. P., D. Chu, B. M. Liu, X. B. Shi, W. Xie, Y. Carriere, X. C. Li, and Y. J. Zhang.**
 458 **2013.** Differential effects of virus on its two closely-related vectors, *Bemisia tabaci* B and Q. Sci.
 459 Rep. 3: 2230.

460 **Pan, H. P., D. Chu, D. Q. Ge, S. L. Wang, Q. J. Wu, W. Xie, X. G. Jiao, B. M. Liu, X.**
 461 **Yang, N. N. Yang, Q. Su, B. Y. Xu, and Y. J. Zhang. 2011.** Further spread of and domination by
 462 *Bemisia tabaci* (Hemiptera: Aleyrodidae) biotype Q on field crops in China. J Econ Entomol
 463 104: 978-985.

464 **Powell, G., C. R. Tosh, and J. Hardie. 2006.** Host plant selection by aphids: behavioral,
 465 evolutionary, and applied perspectives. Annu Rev Entomol 51: 309-330.

466 **Roosien, B. K., R. Gomulkiewicz, L. L. Ingwell, N. A. Bosque-Pérez, D. Rajabaskar,**
 467 **and S. D. Eigenbrode. 2013.** Conditional vector preference aids the spread of plant pathogens:
 468 results from a model. Environ Entomol 42: 1299-1308.

469 **Shi, X., H. Pan, W. Xie, Q. Wu, S. Wang, Y. Liu, Y. Fang, G. Chen, X. Gao, and Y.**
 470 **Zhang. 2013.** Plant virus differentially alters the plant's defense response to its closely related
 471 vectors. PLoS ONE 8: e83520.

472 **Sisterson, M. S. 2008.** Effects of insect-vector preference for healthy or infected plants on
 473 pathogen spread: insights from a model. J Econ Entomol 101: 1-8.

474 **Su, Q., H. P. Pan, B. M. Liu, D. Chu, W. Xie, Q. J. Wu, S. L. Wang, B. Y. Xu, and Y. J.**
475 **Zhang. 2013a.** Insect symbiont facilitates vector acquisition, retention, and transmission of plant
476 virus. *Sci. Rep.* 3: 1367.

477 **Su, Q., K. M. Oliver, H. P. Pan, X. Jiao, B. M. Liu, W. Xie, S. L. Wang, Q. Wu, B. Xu,**
478 **J. A. White, X. Zhou, and Y. J. Zhang. 2013b.** Facultative symbiont *Hamiltonella* confers
479 benefits to *Bemisia tabaci* (Hemiptera: Aleyrodidae), an invasive agricultural pest worldwide.
480 *Environ Entomol* 42: 1265-1271.

481 **Ton, J., and B. Mauch-Mani. 2004.** Beta-amino-butyric acid-induced resistance against
482 necrotrophic pathogens is based on ABA-dependent priming for callose. *The Plant journal : for*
483 *cell and molecular biology* 38: 119-130.

484 **Walling, L. L. 2008.** Avoiding effective defenses: strategies employed by phloem-feeding
485 insects. *Plant Physiol.* 146: 859-866.

486 **Wang, J., X.-L. Bing, M. Li, G.-Y. Ye, and S.-S. Liu. 2012.** Infection of tobacco plants by
487 a begomovirus improves nutritional assimilation by a whitefly. *Entomol Exp Appl* 144: 191-201.

488 **White, J. A., S. E. Kelly, S. J. Perlman, and M. S. Hunter. 2009.** Cytoplasmic
489 incompatibility in the parasitic wasp *Encarsia inaron*: disentangling the roles of *Cardinium* and
490 *Wolbachia* symbionts. *Heredity* 102: 483-489.

491 **Zhang, H., H. Gong, and X. Zhou. 2009.** Molecular characterization and pathogenicity of
492 *tomato yellow leaf curl virus* in China. *Virus Genes* 39: 249-255.

493 **Zhang, S.-Z., B.-Z. Hua, and F. Zhang. 2008.** Induction of the activities of antioxidative
494 enzymes and the levels of malondialdehyde in cucumber seedlings as a consequence of *Bemisia*
495 *tabaci* (Hemiptera: Aleyrodidae) infestation. *Arthropod-Plant Interactions* 2: 209-213.

496 **Zhang, T., J. B. Luan, J. F. Qi, C. J. Huang, M. Li, X. P. Zhou, and S. S. Liu. 2012.**
497 Begomovirus-whitefly mutualism is achieved through repression of plant defences by a virus
498 pathogenicity factor. *Mol Ecol* 21: 1294-1304.
499

Figure Legends

Figure 1. Impact of TYLCV infection on (A) amino acids in the epidermis and mesophyll tissue; (B) simple carbohydrates in the epidermis and mesophyll tissue; (C) amino acids in the phloem sap; and (D) simple carbohydrates in phloem sap. Gray bars: TYLCV-infected; white bars: uninfected. Values are mean $\pm$ SE; * differences significant at $\alpha = 0.05$.

Figure 2. Impact of TYLCV infection, MED infestation, and their interaction on plant defenses. A: concentration of phenylalanine ammonia lyase (PAL); B: concentration of peroxidase (POD); C: concentration of polyphenol oxidase (PPO); D: concentration of superoxide dismutase (SOD); E: callose deposition. Gray bars: TYLCV-infected; white bars: uninfected. Hatched bars: MED-infested; open bars: uninfested. Values are mean $\pm$ SE; lower-case letters indicate differences significant at $\alpha = 0.05$ using Tukey's HSD test.

512 Figure 1.