

1994

A fibered polynomial hull without an analytic selection

J. William Helton

Orlando Merino

University of Rhode Island, merino@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/math_facpubs

Citation/Publisher Attribution

Helton, J. William; Merino, Orlando. A fibered polynomial hull without an analytic selection. *Michigan Math. J.* 41 (1994), no. 2, 285–287. doi:10.1307/mmj/1029004995. <https://projecteuclid.org/euclid.mmj/1029004995>

Available at: <http://dx.doi.org/10.1307/mmj/1029004995>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

A fibered polynomial hull without an analytic selection

Terms of Use

All rights reserved under copyright.

A Fibered Polynomial Hull without an Analytic Selection

J. WILLIAM HELTON & ORLANDO MERINO

This note shows that certain polynomial hulls in \mathbf{C}^3 have no analytic selection, thus settling a standing question about such hulls.

Recall that the *polynomial hull* $\mathcal{P}(S)$ of a set S in \mathbf{C}^M is the set

$$\mathcal{P}(S) = \{w \in \mathbf{C}^M : |p(w)| \leq \max_{v \in S} |p(v)| \text{ for all polynomials } p \text{ on } \mathbf{C}^M\}.$$

We shall be considering sets S in \mathbf{C}^{1+N} that are fibered over the unit circle $\partial\mathbf{D}$ in \mathbf{C} the complex plane. Thus S has the form

$$\{(e^{i\theta}, S_\theta) : 0 \leq \theta \leq 2\pi\},$$

where each S_θ is a subset of \mathbf{C}^N . Let \mathbf{D} denote the unit disk in the complex plane and let H_N^∞ denote the \mathbf{C}^N -valued functions bounded and analytic on \mathbf{D} . Clearly (from the maximum principle) if f is any H_N^∞ function satisfying

$$f(e^{i\theta}) \in S_\theta \text{ for almost all } \theta,$$

then the graph $\{z, f(z) : z \in \mathbf{D}\}$ of f lies in $\mathcal{P}(S)$. Such a function f is called an *analytic selection* of $\mathcal{P}(S)$. A significant question about polynomial hulls is which hulls have analytic selections.

An obvious necessary condition is that $\mathcal{P}(S)$ in \mathbf{C}^{1+N} be a set whose projection onto the first coordinate is a set containing \mathbf{D} . We shall refer to such $\mathcal{P}(S)$ as having *nontrivial fiber over the unit disk*. Also, if the S_θ are not connected then it is easy to make up examples where $\mathcal{P}(S)$ has no analytic selection.

QUESTION (Q). Are these conditions sufficient for $\mathcal{P}(S)$ to have an analytic selection?

By giving a highly pathological example (for $N=1$), Wermer [Wr] showed that in general the answer is No. However, when the S_θ are nicely behaved the story is different. For $N=1$ Slodkowski [Sl] and independently Wegert [Wg] and Helton–Marshall [HM] showed that the answer is Yes. This article gives a simple very well-behaved S in \mathbf{C}^3 for which the answer to (Q) is No.

Now we write down the S that provides our example. Write \mathbf{C}^2 as $\{z = (x_1, y_1, x_2, y_2) = (z_1, z_2)\}$. Let \mathcal{C} denote the semicircle

$$\mathcal{C} = \{(z_1, z_2) : |z_1| = 1, \text{Im } z_1 \geq 0, z_2 = 0\}$$

embedded in the first complex coordinate of \mathbf{C}^2 . Our example is based on rotating \mathcal{C} inside of real 3-dimensional space in a way that varies with θ . Here \mathbf{R}^3 is embedded in \mathbf{C}^2 in the usual way ($y_2 = 0$). Let R_θ^1 denote the map which acts on \mathbf{C}^2 by the rotation of z in the z_1 plane by θ (z_2 is kept fixed), that is,

$$R_\theta^1(z_1, z_2) = (e^{i\theta} z_1, z_2).$$

Also, let R_θ^2 denote rotation in the (y_1, x_2) plane by θ .

THEOREM 1. *The polynomial hull $\mathcal{P}(\mathcal{S})$ of \mathcal{S} with fibers*

$$\mathcal{S}_\theta = R_{\theta/2}^1(R_{\theta/2}^2(\mathcal{C}))$$

has nontrivial fibers over the disk, since $\mathcal{P}(\mathcal{S})$ contains the graph

$$\{(z, \pm z^{1/2}, 0) : z \in \mathbf{D}\}$$

of $z^{1/2}$. Moreover, each \mathcal{S}_θ is connected, and the sets \mathcal{S}_θ vary continuously with θ . However, $\mathcal{P}(\mathcal{S})$ contains no analytic selection.

First we need a lemma. For $k = 1, 2$, let Proj_k be the map on \mathbf{C}^3 given by $\text{Proj}_k(\zeta, z_1, z_2) = z_k$.

LEMMA. $\mathcal{P}(\mathcal{S}) \subset \mathbf{D} \times \mathcal{P}(\text{Proj}_1(\mathcal{S})) \times \mathcal{P}(\text{Proj}_2(\mathcal{S}))$.

Proof. $\mathcal{S} \subset \partial\mathbf{D} \times \text{Proj}_1(\mathcal{S}) \times \text{Proj}_2(\mathcal{S})$. Thus

$$\mathcal{P}(\mathcal{S}) \subset \mathcal{P}(\partial\mathbf{D} \times \text{Proj}_1(\mathcal{S}) \times \text{Proj}_2(\mathcal{S})) \subset \mathbf{D} \times \mathcal{P}(\text{Proj}_1(\mathcal{S})) \times \mathcal{P}(\text{Proj}_2(\mathcal{S})). \quad \square$$

Proof of Theorem 1. Suppose that $f = (f^1, f^2)$ in H_2^∞ has its graph contained in $\mathcal{P}(\mathcal{S})$. By the lemma, the function f^2 is in $\mathcal{P}(\text{Proj}_2(\mathcal{S})) \subset [-1, 1]$ in \mathbf{R} . In particular $\text{Im } f^2$ is 0; therefore f^2 is constant, but $f^2(1) = 0$ since $f(1) \in \mathcal{C}$. We conclude that f^2 is 0.

The value of $f = (f^1, 0)$ at θ is in $\mathcal{S}_\theta \cap \{(z_1, 0)\} = \{(\pm e^{i\theta/2}, 0)\}$ except possibly for $\theta = 0$ or 2π . That is, $f^1 = \pm e^{i\theta/2}$ almost everywhere. No such function exists in H_1^∞ , so we have a contradiction.

Now we check that the \mathcal{S}_θ are smoothly varying. If $0 < \theta < 2\pi$, then the maps R_θ^1 and R_θ^2 on \mathbf{C}^2 are jointly C^∞ in θ and (z_1, z_2) . The only potential difficulty is at $\theta = 0$ or $\theta = 2\pi$. One can easily visualize \mathcal{S}_θ since it lies in $\mathbf{R}^3 \subset \mathbf{C}^2$. The key is the set of points $\pm z_\theta = (\pm e^{i\theta/2}, 0)$ which are the image of $\pm i$ under the map $R_{\theta/2}^1 R_{\theta/2}^2$ (and which consequently lie in \mathcal{S}_θ). Observe that $\pm i = \pm z_0 = \mp z_{2\pi}$, implying that as θ moves from 0 to 2π the points $(+1, 0)$ and $(-1, 0)$ rotate into each other. Thus $\mathcal{S}_{2\pi}$ is a rotation of \mathcal{C} in the $(z_1, 0)$ plane which has the same endpoints as \mathcal{C} ; we have $\mathcal{S}_{2\pi} = \mathcal{C}$ or $\mathcal{S}_{2\pi} = -\mathcal{C}$. However,

$$R_{2\pi/2}^1(R_{2\pi/2}^2(1, 0)) = R_\pi^1((-1, 0)) = (1, 0).$$

We have established that the \mathcal{S}_θ vary continuously in θ . □

Lest one complain that our example hinges on degeneracy of \mathcal{S}_θ , we now give an example where \mathcal{S} is the closure of an open set in \mathbf{C}^3 in addition to maintaining other nice properties.

THEOREM 2. *For $n \in \mathbf{N}$, let \mathcal{S}^n be the set in \mathbf{C}^3 with fibers $\mathcal{S}_\theta + (1/n)B$, where B is the unit ball of \mathbf{C}^2 . Then there exists $n_0 \in \mathbf{N}$ such that the polynomial hull of \mathcal{S}^{n_0} has no analytic selection.*

Proof. Suppose not. Then for each n there exists f_n in H_2^∞ whose graph lies in \mathcal{S}^n . Let f_0 be a normal families limit of $\{f_n\}$. By polynomial convexity of each \mathcal{S}_θ and by Corollary 2 in [HM], the function f_0 has graph lying in $\mathcal{O}(\mathcal{S})$. Thus $\mathcal{O}(\mathcal{S})$ has a selection, contrary to Theorem 1. \square

References

- [HM] J. W. Helton and D. E. Marshall, *Frequency domain design and analytic selections*, Indiana Univ. Math. J. 39 (1990), 157–184.
- [Sl] Z. Slodowski, *Polynomial hulls for \mathbf{C}^2 and quasicircles*, preprint.
- [Wg] E. Wegert, preprint.
- [Wr] J. Wermer, *Polynomially convex hulls and analyticity*, Ark. Mat. 20 (1982), 129–135.

J. W. Helton
Department of Mathematics
University of California – San Diego
La Jolla, CA 92093-0112

O. Merino
Department of Mathematics
University of Rhode Island
Kingston, RI 02881

