

2013

Global Dynamics of Three Anticompetitive Systems of Difference Equations in the Plane

M. DiPippo

Mustafa Kulenovic

University of Rhode Island, mkulenovic@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/math_facpubs

Citation/Publisher Attribution

M. DiPippo and M. R. S. Kulenović, "Global Dynamics of Three Anticompetitive Systems of Difference Equations in the Plane," *Discrete Dynamics in Nature and Society*, vol. 2013, Article ID 751594, 11 pages, 2013. <https://doi.org/10.1155/2013/751594>.

Available at: <http://dx.doi.org/10.1155/2013/751594>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Global Dynamics of Three Anticompetitive Systems of Difference Equations in the Plane

Creative Commons License



This work is licensed under a [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/).

Research Article

Global Dynamics of Three Anticompetitive Systems of Difference Equations in the Plane

M. DiPippo¹ and M. R. S. Kulenović²

¹ Department of Mathematics, Rhode Island College, Providence, RI 02881-0816, USA

² Department of Mathematics, University of Rhode Island, Kingston, RI 02881-0816, USA

Correspondence should be addressed to M. R. S. Kulenović; mkulenovic@mail.uri.edu

Received 11 September 2012; Revised 16 January 2013; Accepted 22 January 2013

Academic Editor: Raghieb Abu-Saris

Copyright © 2013 M. DiPippo and M. R. S. Kulenović. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the global dynamics of several anticompetitive systems of rational difference equations which are special cases of general linear fractional system of the forms $x_{n+1} = (\alpha_1 + \beta_1 x_n + \gamma_1 y_n)/(A_1 + B_1 x_n + C_1 y_n)$, $y_{n+1} = (\alpha_2 + \beta_2 x_n + \gamma_2 y_n)/(A_2 + B_2 x_n + C_2 y_n)$, $n = 0, 1, \dots$, where all parameters and the initial conditions x_0, y_0 are arbitrary nonnegative numbers, such that both denominators are positive. We find the basins of attraction of all attractors of these systems.

1. Introduction

A first order system of difference equations:

$$\begin{aligned} x_{n+1} &= f(x_n, y_n), & y_{n+1} &= g(x_n, y_n), \\ n &= 0, 1, \dots, & (x_0, y_0) &\in \mathcal{R}, \end{aligned} \quad (1)$$

where $\mathcal{R} \subset \mathbb{R}^2$, $(f, g) : \mathcal{R} \rightarrow \mathcal{R}$, f, g are continuous functions, is *competitive* if $f(x, y)$ is nondecreasing in x and nonincreasing in y , and $g(x, y)$ is nonincreasing in x and nondecreasing in y .

System (1) where the functions f and g have monotonic character opposite the monotonic character in competitive system is called *anticompetitive*; see [1, 2].

In this paper, we consider the following anticompetitive systems of difference equations:

$$x_{n+1} = \frac{\gamma_1 y_n}{B_1 x_n + y_n}, \quad y_{n+1} = \frac{\beta_2 x_n}{B_2 x_n + y_n}, \quad n = 0, 1, \dots, \quad (2)$$

where all parameters are positive numbers and the initial conditions (x_0, y_0) are arbitrary nonnegative numbers such that $x_0 + y_0 > 0$. In the classification of all linear fractional

systems in [3], System (2) was mentioned as system (18, 18). We also consider systems

$$x_{n+1} = \frac{\gamma_1 y_n}{B_1 x_n + y_n}, \quad y_{n+1} = \frac{\alpha_2 + \beta_2 x_n}{y_n}, \quad n = 0, 1, \dots, \quad (3)$$

$$x_{n+1} = \frac{\gamma_1 y_n}{A_1 + x_n}, \quad y_{n+1} = \frac{\alpha_2 + \beta_2 x_n}{y_n}, \quad n = 0, 1, \dots, \quad (4)$$

with $x_0 > 0, y_0 > 0$, which were labeled as systems (18, 23) and (16, 23), respectively, in [3]. Three systems have interesting and different dynamics. While System (2) has all bounded solutions, most of solutions of Systems (3) and (4) are unbounded. Another major difference is the existence of the unique period-two solution for (2) and, in a special case, the abundance of such solutions, while neither (3) nor (4) has period-two solutions. We show that every solution of System (3) converges to the unique equilibrium or is approaching $(0, \infty)$ and so System (3) gives an example of a semistable equilibrium point. Finally, we show that all solutions of (4) which start on the stable set converge to the unique equilibrium, while all solutions which start off the stable set are approaching $(0, \infty)$ or $(\infty, 0)$. We also get that

for some special values of parameters, Systems (4) and (3) can be decoupled and explicitly solved.

Competitive systems of the form (1) were studied by many authors such as Clark and Kulenović [4], Hess [5], Hirsch and Smith [6], Kulenović and Merino [7], Kulenović and Nurkanović [8], Garić-Demirović et al. [9, 10], and Smith [11, 12]. Precise results about the basins of attraction of the equilibrium points have been obtained in [13].

The study of anticompetitive systems started in [1] and has advanced since then; see [2]. The principal tool of study of anticompetitive systems is the fact that the second iterate of the map associated with anticompetitive system is competitive map and so elaborate theory for such maps developed recently in [6, 14, 15] can be applied.

The major result on a global behavior of System (2) is the following theorem.

Theorem 1. (a) Assume that $B_1 < B_2$. Then the unique positive equilibrium point $E(\bar{x}, \bar{y})$ of System (2) is locally asymptotically stable with the basin of attraction $\mathcal{B}(E) = (0, \infty)^2$. The unique period-two solution $\{P_1, P_2\} = \{(\gamma_1, 0), (0, \beta_2/B_2)\}$ is a saddle point and its basin of attraction is the union of coordinate axes without the origin, that is, $\mathcal{B}(\{P_1, P_2\}) = ((x, 0) \mid x > 0) \cup (0, y) \mid y > 0$.

(b) Assume that $B_1 > B_2$. Then every solution of System (2) converges to the period-two solution $\{P_1, P_2\}$ or to the equilibrium E . More precisely, there exists a set $\mathcal{C} = \{(x, (\bar{y}/\bar{x})x) : x > 0\} \subset \mathcal{R} = (0, \infty)^2$ which is the basin of attraction of E . The set \mathcal{C} has the property that for

$$\begin{aligned} \mathcal{W}_- &:= \{x \in \mathcal{R} \setminus \mathcal{C} : \exists y \in \mathcal{C} \text{ with } x \leq_{se} y\} \\ &= \left\{ (x, y) : y > \frac{\bar{y}}{\bar{x}}x \geq 0 \right\}, \\ \mathcal{W}_+ &:= \{x \in \mathcal{R} \setminus \mathcal{C} : \exists y \in \mathcal{C} \text{ with } y \leq_{se} x\} \\ &= \left\{ (x, y) : 0 \leq y < \frac{\bar{y}}{\bar{x}}x \right\}, \end{aligned} \quad (5)$$

the following holds.

(i) If $(x_0, y_0) \in \mathcal{W}_+$, then

$$\lim_{n \rightarrow \infty} (x_{2n}, y_{2n}) = P_1, \quad \lim_{n \rightarrow \infty} (x_{2n+1}, y_{2n+1}) = P_2. \quad (6)$$

(ii) If $(x_0, y_0) \in \mathcal{W}_-$,

$$\lim_{n \rightarrow \infty} (x_{2n}, y_{2n}) = P_2, \quad \lim_{n \rightarrow \infty} (x_{2n+1}, y_{2n+1}) = P_1. \quad (7)$$

(c) Assume that $B_1 = B_2$. Then every solution of System (2) is equal to either the period-two solution $\{B_1 y_0 / (B_1 x_0 + y_0), \beta_2 x_0 / (B_1 x_0 + y_0)\}$, $\{B_1 D x_0 / (B_1 y_0 + D x_0), \beta_2 y_0 / (B_1 y_0 + D x_0)\}$ or to the equilibrium $E(\bar{x}, \bar{y})$, where $D = \beta_2 / \gamma_1$ and $\bar{x} = \sqrt{\beta_2 \gamma_1} / (B_1 + 1)$.

2. Preliminaries

We now give some basic notions about competitive systems and maps in the plane of the form (1), where f and g are continuous functions, $f(x, y)$ is nondecreasing in x

and nonincreasing in y , and $g(x, y)$ is nonincreasing in x and nondecreasing in y in some domain A with nonempty interior.

Consider a map $T = (f, g)$ on a set $\mathcal{R} \subset \mathbb{R}^2$, and let $E \in \mathcal{R}$. The point $E \in \mathcal{R}$ is called a *fixed point* if $T(E) = E$. An *isolated fixed point* is a fixed point that has a neighborhood with no other fixed points in it. A fixed point $E \in \mathcal{R}$ is an *attractor* if there exists a neighborhood \mathcal{U} of E such that $T^n(\mathbf{x}) \rightarrow E$ as $n \rightarrow \infty$ for $\mathbf{x} \in \mathcal{U}$; the *basin of attraction* is the set of all $\mathbf{x} \in \mathcal{R}$ such that $T^n(\mathbf{x}) \rightarrow E$ as $n \rightarrow \infty$. A fixed point E is a global attractor on a set \mathcal{X} if E is an attractor and \mathcal{X} is a subset of the basin of attraction of E . If T is differentiable at a fixed point E and if the Jacobian $J_T(E)$ has one eigenvalue with modulus less than one and a second eigenvalue with modulus greater than one, E is said to be a *saddle*. See [16] for additional definitions.

Definition 2. Let $T = (f, g)$ be a continuously differentiable vector function and let U be a neighborhood of a saddle point (\bar{x}, \bar{y}) of (1). The local stable manifold \mathcal{W}_{loc}^s is the set

$$\begin{aligned} \mathcal{W}_{loc}^s((\bar{x}, \bar{y})) &= \left\{ (x, y) : T^n(x, y) \in U (\forall n \geq 0) \right. \\ &\quad \left. \wedge \lim_{n \rightarrow \infty} T^n(x, y) = (\bar{x}, \bar{y}) \right\}. \end{aligned} \quad (8)$$

The global stable manifold W^s of a saddle point (\bar{x}, \bar{y}) is the set

$$\mathcal{W}^s((\bar{x}, \bar{y})) = \left\{ (x, y) : \lim_{n \rightarrow \infty} T^n(x, y) = (\bar{x}, \bar{y}) \right\}. \quad (9)$$

The map T may be viewed as a monotone map if we define a partial order on \mathbb{R}^2 so that the positive cone in this new partial order is the fourth (resp. first) quadrant. Define a south-east (resp. north-east) partial order \leq_{se} (resp. \leq_{ne}) on \mathbb{R}^2 so that the positive cone is the fourth quadrant (resp. first quadrant), that is, $(x^1, y^1) \leq_{se} (x^2, y^2)$ (resp. $(x^1, y^1) \leq_{ne} (x^2, y^2)$) if and only if $x^1 \leq x^2$ and $y^1 \geq y^2$ (resp. $x^1 \leq x^2$ and $y^1 \leq y^2$). For $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ the *order interval* $[\mathbf{x}, \mathbf{y}]$ is the set of all \mathbf{z} such that $\mathbf{x} \leq \mathbf{z} \leq \mathbf{y}$. The map T is called *competitive* (resp. *cooperative*) on a set S if $\mathbf{v} \leq_{se} \mathbf{w}$ (resp. $\mathbf{v} \leq_{ne} \mathbf{w}$) implies $T(\mathbf{v}) \leq_{se} T(\mathbf{w})$ (resp. $T(\mathbf{v}) \leq_{ne} T(\mathbf{w})$).

Two points $\mathbf{v}, \mathbf{w} \in \mathbb{R}_+^2$ are said to be *related* if $\mathbf{v} \leq \mathbf{w}$ or $\mathbf{w} \leq \mathbf{v}$. Also, a strict inequality between points may be defined as $\mathbf{v} < \mathbf{w}$ if $\mathbf{v} \leq \mathbf{w}$ and $\mathbf{v} \neq \mathbf{w}$. A stronger inequality may be defined as $\mathbf{v} \ll \mathbf{w}$ if $v_1 < w_1$ and $w_2 < v_2$. A map $f : \text{Int } \mathbb{R}_+^2 \rightarrow \text{Int } \mathbb{R}_+^2$ is *strongly monotone* if $\mathbf{v} < \mathbf{w}$ implies that $f(\mathbf{v}) \ll f(\mathbf{w})$ for all $\mathbf{v}, \mathbf{w} \in \text{Int } \mathbb{R}_+^2$. Clearly, being related is invariant under the iteration of a strongly monotone map. Differentiable strongly monotone maps have Jacobian with constant sign configuration

$$\begin{bmatrix} + & - \\ - & + \end{bmatrix}. \quad (10)$$

The mean value theorem and the convexity of \mathbb{R}_+^2 may be used to show that T is monotone, as in [17].

For $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2$, define $Q_l(\mathbf{x})$ for $l = 1, \dots, 4$ to be the usual four quadrants based at \mathbf{x} and numbered in

a counterclockwise direction, for example, $Q_1(\mathbf{x}) = \{y = (y_1, y_2) \in \mathbb{R}^2 : x_1 \leq y_1, x_2 \leq y_2\}$. We now state three results for competitive maps in the plane.

The following definition is from [11].

Definition 3. Let \mathcal{S} be a nonempty subset of \mathbb{R}^2 . A competitive map $T : \mathcal{S} \rightarrow \mathcal{S}$ is said to satisfy condition (O+) if for every x, y in \mathcal{S} , $T(x) \leq_{ne} T(y)$ implies $x \leq_{ne} y$, and T is said to satisfy condition (O-) if for every x, y in \mathcal{S} , $T(x) \leq_{ne} T(y)$ implies $y \leq_{ne} x$.

The following theorem was proved by de Mottoni-Schiaffino for the Poincaré map of a periodic competitive Lotka-Volterra system of differential equations. Smith generalized the proof to competitive and cooperative maps [11, 12].

Theorem 4. Let \mathcal{S} be a nonempty subset of \mathbb{R}^2 . If T is a competitive map for which (O+) holds, then for all $x \in \mathcal{S}$, $\{T^n(x)\}$ is eventually componentwise monotone. If the orbit of x has compact closure, then it converges to a fixed point of T . If instead (O-) holds, then for all $x \in \mathcal{S}$, $\{T^{2n}\}$ is eventually componentwise monotone. If the orbit of x has compact closure in \mathcal{S} , then its omega limit set is either a period-two orbit or a fixed point.

The following result is from [11], with the domain of the map specialized to be the cartesian product of intervals of real numbers. It gives a sufficient condition for conditions (O+) and (O-).

Theorem 5. Let $\mathcal{R} \subset \mathbb{R}^2$ be the cartesian product of two intervals in \mathbb{R} . Let $T : \mathcal{R} \rightarrow \mathcal{R}$ be a C^1 competitive map. If T is injective and $\det J_T(x) > 0$ for all $x \in \mathcal{R}$, then T satisfies (O+). If T is injective and $\det J_T(x) < 0$ for all $x \in \mathcal{R}$, then T satisfies (O-).

The following result is a direct consequence of the Trichotomy Theorem of Dancer and Hess, see [5, 15, 18], and is helpful for determining the basins of attraction of the equilibrium points.

Corollary 6. If the nonnegative cone with respect to the partial order \leq is a generalized quadrant in \mathbb{R}^2 and if the competitive map $T : \mathcal{R} \rightarrow \mathcal{R}$ has no fixed points in $\llbracket u_1, u_2 \rrbracket$ other than u_1 and u_2 , then the interior of $\llbracket u_1, u_2 \rrbracket$ is either a subset of the basin of attraction of u_1 or a subset of the basin of attraction of u_2 .

The next two results are from [15, 18].

Theorem 7. Let T be a competitive map on a rectangular region $\mathcal{R} \subset \mathbb{R}^2$. Let $\bar{x} \in \mathcal{R}$ be a fixed point of T such that $\Delta := \mathcal{R} \cap \text{int}(Q_1(\bar{x}) \cup Q_3(\bar{x}))$ is nonempty (i.e., \bar{x} is not the NW or SE vertex of \mathcal{R}), and T is strongly competitive on Δ . Suppose that the following statements are true.

- (a) The map T has a C^1 extension to a neighborhood of \bar{x} .
- (b) The Jacobian matrix of T at \bar{x} has real eigenvalues λ, μ such that $0 < |\lambda| < \mu$, where $|\lambda| < 1$, and the

eigenspace E^λ associated with λ is not a coordinate axis.

Then there exists a curve $\mathcal{C} \subset \mathcal{R}$ through \bar{x} that is invariant and a subset of the basin of attraction of \bar{x} , such that \mathcal{C} is tangential to the eigenspace E^λ at \bar{x} , and \mathcal{C} is the graph of a strictly increasing continuous function of the first coordinate on an interval. Any endpoints of \mathcal{C} in the interior of \mathcal{R} are either fixed points or minimal period-two points. In the latter case, the set of endpoints of \mathcal{C} is a minimal period-two orbit of T .

Theorem 8. Let $\mathcal{I}_1, \mathcal{I}_2$ be intervals in \mathbb{R} with endpoints a_1, a_2 and b_1, b_2 with endpoints, respectively, with $a_1 < a_2$ and $b_1 < b_2$, where $-\infty \leq a_1 < a_2 \leq \infty$ and $-\infty \leq b_1 < b_2 \leq \infty$. Let T be a competitive map on a rectangle $\mathcal{R} = \mathcal{I}_1 \times \mathcal{I}_2$ and $\bar{x} \in \text{int}(\mathcal{R})$. Suppose that the following hypotheses are satisfied.

- (1) $T(\text{int}(\mathcal{R})) \subset \text{int}(\mathcal{R})$ and T is strongly competitive on $\text{int}(\mathcal{R})$.
- (2) The point \bar{x} is the only fixed point of T in $(Q_1(\bar{x}) \cup Q_3(\bar{x})) \cap \text{int}(\mathcal{R})$.
- (3) The map T is continuously differentiable in a neighborhood of \bar{x} , and \bar{x} is the saddle point.
- (4) At least one of the following statements is true.
 - (a) T has no minimal period-two orbits in $(Q_1(\bar{x}) \cup Q_3(\bar{x})) \cap \text{int}(\mathcal{R})$.
 - (b) $\det J_T(\bar{x}) > 0$ and $T(\mathbf{x}) = \bar{x}$ only for $\mathbf{x} = \bar{x}$.

Then the following statements are true.

- (i) The stable manifold $\mathcal{W}^s(\bar{x})$ is connected and it is the graph of a continuous increasing curve with endpoints in $\partial\mathcal{R}$. $\text{int}(\mathcal{R})$ is divided by the closure of $\mathcal{W}^s(\bar{x})$ into two invariant connected regions \mathcal{W}_+ ("below the stable set"), and \mathcal{W}_- ("above the stable set"), where
$$\mathcal{W}_+ := \{\mathbf{x} \in \mathcal{R} \setminus \mathcal{W}^s(\bar{x}) : \exists \mathbf{x}' \in \mathcal{W}^s(\bar{x}) \text{ with } \mathbf{x} \leq_{se} \mathbf{x}'\},$$

$$\mathcal{W}_- := \{\mathbf{x} \in \mathcal{R} \setminus \mathcal{W}^s(\bar{x}) : \exists \mathbf{x}' \in \mathcal{W}^s(\bar{x}) \text{ with } \mathbf{x}' \leq_{se} \mathbf{x}\}.$$
(11)
- (ii) The unstable manifold $\mathcal{W}^u(\bar{x})$ is connected and it is the graph of a continuous decreasing curve.
- (iii) For every $\mathbf{x} \in \mathcal{W}_+$, $T^n(\mathbf{x})$ eventually enters the interior of the invariant set $Q_4(\bar{x}) \cap \mathcal{R}$, and for every $\mathbf{x} \in \mathcal{W}_-$, $T^n(\mathbf{x})$ eventually enters the interior of the invariant set $Q_2(\bar{x}) \cap \mathcal{R}$.
- (iv) Let $\mathbf{m} \in Q_2(\bar{x})$ and $\mathbf{M} \in Q_4(\bar{x})$ be the endpoints of $\mathcal{W}^u(\bar{x})$, where $\mathbf{m} \leq_{se} \bar{x} \leq_{se} \mathbf{M}$. For every $\mathbf{x} \in \mathcal{W}_-$ and every $\mathbf{z} \in \mathcal{R}$ such that $\mathbf{m} \leq_{se} \mathbf{z}$, there exists $m \in \mathbb{N}$ such that $T^m(\mathbf{x}) \leq_{se} \mathbf{z}$, and for every $\mathbf{x} \in \mathcal{W}_+$ and every $\mathbf{z} \in \mathcal{R}$ such that $\mathbf{z} \leq_{se} \mathbf{M}$, there exists $m \in \mathbb{N}$ such that $\mathbf{M} \leq_{se} T^m(\mathbf{x})$.




FIGURE 1: Global dynamics of System (2).

3. Global Dynamics of System (2) (See Figure 1)

The equilibrium point $E(\bar{x}, \bar{y})$ of System (2) satisfies the following system of equations:

$$\bar{x} = \frac{\gamma_1 \bar{y}}{B_1 \bar{x} + \bar{y}}, \quad \bar{y} = \frac{\beta_2 \bar{x}}{B_2 \bar{x} + \bar{y}}. \quad (12)$$

It is easy to see that System (12) has unique equilibrium point E in the first quadrant, for all values of the parameters. Indeed, the positive equilibrium point is an intersection of the following two curves:

$$y = \frac{B_1 x^2}{\gamma_1 - x}, \quad (13)$$

$$x = \frac{y^2}{\beta_2 - B_2 y}. \quad (14)$$

It is clear that at the point of intersection E curve (13) is steeper than curve (14), that is,

$$\left. \frac{dy}{dx} \right|_{(13)} (E) > \left. \frac{dy}{dx} \right|_{(14)} (E), \quad (15)$$

which gives

$$\frac{B_1 \bar{x} (2\gamma_1 - \bar{x})}{(\gamma_1 - \bar{x})^2} > \frac{(\beta_2 - B_2 \bar{y})^2}{\bar{y} (2\beta_2 - B_2 \bar{y})}. \quad (16)$$

This inequality is equivalent to the following inequality:

$$\left(\beta_2 + \frac{\bar{y}^2}{\bar{x}} \right) \left(\gamma_1 + B_1 \frac{\bar{x}^2}{\bar{y}} \right) > B_1 \bar{x} \bar{y}, \quad (17)$$

which is always satisfied.

3.1. Linearized Stability Analysis of System (2). In this section we present the linearized stability analysis of the equilibrium E of System (2).

Theorem 9. (i) If $B_1 < B_2$, then E is locally asymptotically stable.

(ii) If $B_1 = B_2$, then E is a nonhyperbolic equilibrium point.

(iii) If $B_1 > B_2$, then E is a saddle point.

Proof. The map T associated to System (2) is

$$T(x, y) = \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix} = \begin{pmatrix} \frac{\gamma_1 y}{B_1 x + y} \\ \frac{\beta_2 x}{B_2 x + y} \end{pmatrix}. \quad (18)$$

The Jacobian matrix of map (18) is

$$J_T(x, y) = \begin{pmatrix} -\frac{B_1 \gamma_1 y}{(B_1 x + y)^2} & \frac{B_1 \gamma_1 x}{(B_1 x + y)^2} \\ \frac{\beta_2 y}{(B_2 x + y)^2} & -\frac{\beta_2 x}{(B_2 x + y)^2} \end{pmatrix}, \quad (19)$$

and evaluated at the equilibrium point $E = (\bar{x}, \bar{y})$ is

$$J_T(\bar{x}, \bar{y}) = \begin{pmatrix} -\frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} & \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^2} \\ \frac{\bar{y}^3}{\beta_2 \bar{x}^2} & -\frac{\bar{y}^2}{\beta_2 \bar{x}} \end{pmatrix}. \quad (20)$$

The characteristic equation has the following form:

$$\lambda^2 + \left(\frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} + \frac{1}{\beta_2} \frac{\bar{y}^2}{\bar{x}} \right) \lambda = 0, \quad (21)$$

and the characteristic roots are

$$\lambda_1 = 0, \quad \lambda_2 = -\frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} - \frac{1}{\beta_2} \frac{\bar{y}^2}{\bar{x}}. \quad (22)$$

A straightforward calculation shows that the conditions $\lambda_2 \in (-1, 0)$, $\lambda_2 = -1$, and $\lambda_2 \in (-\infty, -1)$, respectively, are equivalent to the conditions

$$\left(\frac{\bar{x}}{\bar{y}}\right)^2 < \frac{\beta_2}{\gamma_1}, \quad \left(\frac{\bar{x}}{\bar{y}}\right)^2 = \frac{\beta_2}{\gamma_1}, \quad \left(\frac{\bar{x}}{\bar{y}}\right)^2 > \frac{\beta_2}{\gamma_1}. \quad (23)$$

On the other hand, by dividing two equilibrium equations (12) we obtain

$$\left(\frac{\bar{x}}{\bar{y}}\right)^2 = \frac{\beta_2 B_1 \bar{x} + \bar{y}}{\gamma_1 B_2 \bar{x} + \bar{y}}, \quad (24)$$

which implies that the condition (23) is equivalent to the conditions

$$B_1 < B_2, \quad B_1 = B_2, \quad B_1 > B_2, \quad (25)$$

which completes the proof of the theorem. \square

3.2. *Global Results for System (2).* In this section we present the proof of Theorem 1 on the global dynamics of System (2).

$$T^2(x, y) = \left[x\beta_2\gamma_1 \frac{y + xB_1}{xy\beta_2 + x^2\beta_2B_1 + y^2\gamma_1B_1 + xy\gamma_1B_1B_2} \quad y\beta_2\gamma_1 \frac{y + xB_2}{\beta_2B_1x^2 + \gamma_1xyB_2^2 + \beta_2xy + \gamma_1y^2B_2} \right]. \quad (28)$$

A period-two solution of System (2) satisfies $T^2(x, y) = (x, y)$, which immediately leads to the following equations:

$$(B_2 - B_1)(\beta_2\gamma_1x - B_2\gamma_1xy - \gamma_1y^2) = 0, \quad (29)$$

$$(B_2 - B_1)(\beta_2\gamma_1y - B_2xy - \gamma_1B_1x^2) = 0, \quad (30)$$

$J_{T^2}(x, y)$

$$= \begin{bmatrix} y\beta_2\gamma_1^2B_1 \frac{B_1B_2x^2 + 2B_1xy + y^2}{(xy\beta_2 + x^2\beta_2B_1 + y^2\gamma_1B_1 + xy\gamma_1B_1B_2)^2} & -x\beta_2\gamma_1^2B_1 \frac{B_1B_2x^2 + 2B_1xy + y^2}{(xy\beta_2 + x^2\beta_2B_1 + y^2\gamma_1B_1 + xy\gamma_1B_1B_2)^2} \\ -y\beta_2^2\gamma_1 \frac{B_1B_2x^2 + 2B_1xy + y^2}{(\beta_2B_1x^2 + \gamma_1xyB_2^2 + \beta_2xy + \gamma_1y^2B_2)^2} & x\beta_2^2\gamma_1 \frac{B_1B_2x^2 + 2B_1xy + y^2}{(\beta_2B_1x^2 + \gamma_1xyB_2^2 + \beta_2xy + \gamma_1y^2B_2)^2} \end{bmatrix} \quad (31)$$

The Jacobian matrix of T^2 evaluated at P_1 is

$$J_{T^2}(P_1) = \begin{bmatrix} 0 & -\frac{1}{\beta_2}\gamma_1B_2 \\ 0 & \frac{1}{B_1}B_2 \end{bmatrix} \quad (32)$$

First, we prove the following result on existence and local behavior of the period-two solutions.

Lemma 10. *System (2) has the minimal period-two solution*

$$\{P_1, P_2\} = \left\{ (\gamma_1, 0), \left(0, \frac{\beta_2}{B_2} \right) \right\} \quad (26)$$

for all values of parameters. If $B_1 \neq B_2$, then the solution (26) is the unique period-two solution and when $B_1 = B_2$ there are infinitely many period-two solutions. The set

$$\mathcal{B} = \{(x, y) \mid x > 0, y = 0 \text{ or } x = 0, y > 0\} \quad (27)$$

is a subset of the basin of attraction of the solution (26). The period-two solution (26) is locally stable if $B_1 > B_2$ and a saddle point if $B_1 < B_2$. Finally the period-two solution (26) is nonhyperbolic if $B_1 = B_2$.

Proof. The second iterate of the map T is given as

which have either unique solution if $B_1 \neq B_2$ or it has infinitely many solutions if $B_1 = B_2$. In the first case, (29) gives immediately (13) and (30) gives (14), which means that in this case the only minimal period-two solution is (26).

A straightforward calculation shows that $T(P_1) = P_2$, $T(P_2) = P_1$, which shows that $\{P_1, P_2\}$ is a minimal period-two solution. Moreover, $T(a, 0) = P_2$, $T(0, b) = P_1$ for every $a > 0, b > 0$, which shows that the set \mathcal{B} is a subset of the basin of attraction of $\{P_1, P_2\}$. The Jacobian matrix of T^2 is

and the Jacobian matrix of T^2 evaluated at P_2 is

$$J_{T^2}(P_2) = \begin{bmatrix} \frac{1}{B_1}B_2 & 0 \\ B_2 - \frac{1}{\beta_2\gamma_1} \left(\frac{\beta_2^2}{B_2} + \beta_2\gamma_1B_2 \right) & 0 \end{bmatrix}. \quad (33)$$

In both cases, the eigenvalues of the Jacobian matrix of T^2 are $\lambda_1 = 0$, $\lambda_2 = B_2/B_1$, which implies the result on local stability of the minimal period-two solution $\{P_1, P_2\}$. \square

Proof of Theorem 1. First, observe that the rectangle $\mathcal{R} = [0, \gamma_1] \times [0, \beta_2/B_2] \setminus \{0, 0\} = \llbracket P_2, P_1 \rrbracket \setminus \{0, 0\}$ is an invariant and attracting set for the map T and so is for the map T^2 . More precisely, $(x_n, y_n) \in \mathcal{R}$ for $n \geq 1$. The map T^2 is a competitive map on \mathcal{R} . \square

Case a. Assume that $B_1 < B_2$. Then in view of Theorem 9 and Lemma 10, the map T^2 has three equilibrium points P_1 , P_2 , and E , where $P_2 \preceq_{se} E \preceq_{se} P_1$. The equilibrium points P_1 and P_2 are saddle points and E is a local attractor. The ordered intervals $\llbracket P_2, E \rrbracket$ and $\llbracket E, P_1 \rrbracket$ are both invariant sets of T^2 and in view of Corollary 6, their interiors are attracted to E . If we take the point $(x, y) \in \mathcal{R} \setminus \llbracket P_2, E \rrbracket \cup \llbracket E, P_1 \rrbracket$, we can find the points $(x_l, y_l) \in \text{int} \llbracket P_2, E \rrbracket$ and $(x_u, y_u) \in \text{int} \llbracket E, P_1 \rrbracket$, such that $(x_l, y_l) \preceq_{se} (x, y) \preceq_{se} (x_u, y_u)$. Consequently, since T^2 is competitive $T^{2n}((x_l, y_l)) \preceq_{se} T^{2n}((x, y)) \preceq_{se} T^{2n}((x_u, y_u))$ for $n \geq 1$ and so $\lim_{n \rightarrow \infty} T^{2n}((x, y)) = E$, which by continuity of T implies that

$$\begin{aligned} \lim_{n \rightarrow \infty} T^{2n+1}((x, y)) &= \lim_{n \rightarrow \infty} T(T^{2n}((x, y))) \\ &= T\left(\lim_{n \rightarrow \infty} T^{2n}((x, y))\right) = T(E) = E, \end{aligned} \quad (34)$$

and so $\lim_{n \rightarrow \infty} T^n((x, y)) = E$.

Case b. Assume that $B_1 > B_2$. Then in view of Theorem 9 and Lemma 10, the map T^2 has three equilibrium points P_1 and P_2 which are local attractors and E which is a saddle point. The ordered intervals $\llbracket P_2, E \rrbracket$ and $\llbracket E, P_1 \rrbracket$ are both invariant sets for T^2 and in view of Corollary 6, their interiors are attracted to P_2 and P_1 , respectively. In view of Theorems 7 and 8, there is the set \mathcal{C} with described properties. Direct calculation shows that the half-line $y = (\bar{y}/\bar{x})x$, $x > 0$ is an invariant set which in view of a uniqueness of stable manifold implies that this half-line is exactly stable manifold mentioned in Theorems 7 and 8. It should be observed that because of the fact that one of the characteristic values at the equilibrium point E is 0, this equilibrium is superattractive, that is, $T(x_0, y_0) = (\bar{x}, \bar{y})$, for every $(x_0, y_0) \in \mathcal{C}$.

Case c. Assume that $B_1 = B_2$. Then by dividing two equations of System (2) we obtain that the solution of (2) satisfies

$$\frac{y_{n+1}}{x_{n+1}} = \frac{\gamma_2 x_n}{\beta_1 y_n} = \frac{\gamma_2}{\beta_1} \frac{1}{y_n/x_n}. \quad (35)$$

This means that y_n/x_n satisfies first order difference equation $u_{n+1} = D/u_n$, where $D = \gamma_2/\beta_1$. All nonconstant solutions of $u_{n+1} = D/u_n$ are period-two solutions $\{u_0, D/u_0\}$.

Thus $y_n = \{u_0 x_n, (D/u_0)x_n\}$. In this case, System (2) becomes

$$\begin{aligned} x_{n+1} &= \frac{B_1 u_0}{B_1 + u_0}, & y_{n+1} &= \frac{\beta_2}{B_1 + u_0}, & n &= 1, 2, \dots, \\ x_{n+1} &= \frac{B_1 D}{D + B_1 u_0}, & y_{n+1} &= \frac{\beta_2 u_0}{D + B_1 u_0}, & n &= 1, 2, \dots, \end{aligned} \quad (36)$$

which completes the proof of Case (c).

Remark 11. System (2) is an example of the homogeneous system which is a special case of a general System (1), where both functions f and g are homogeneous functions of the same degree k , that is, $f(tu, tv) = t^k f(u, v)$, $g(tu, tv) = t^k g(u, v)$ for all u, v in intersection of domains of f and g and all $t \neq 0$. In that case, the ratio $z_n = y_n/x_n$ of every solution of (1) satisfies the first order difference equation:

$$z_{n+1} = \frac{f(1, z_n)}{g(1, z_n)} = F(z_n), \quad n = 0, 1, \dots \quad (37)$$

whose analysis gives valuable information about the dynamics of System (1), but does not provide the global dynamics. In particular, this approach cannot determine precisely the basins of attraction of different types of attractors such as equilibrium points and periodic solutions. This approach in the case of the system of linear fractional equations:

$$\begin{aligned} x_{n+1} &= \frac{\alpha_1 + \beta_1 x_n + \gamma_1 y_n}{A_1 + B_1 x_n + C_1 y_n}, & y_{n+1} &= \frac{\alpha_2 + \beta_2 x_n + \gamma_2 y_n}{A_2 + B_2 x_n + C_2 y_n}, \\ & & n &= 0, 1, \dots, \end{aligned} \quad (38)$$

where all parameters and the initial conditions (x_0, y_0) are arbitrary nonnegative numbers such that $A_i + B_i x_0 + C_i y_0 > 0$, $i = 1, 2$, was first used in [19] and was systematically developed in the recent paper [20]. In [20], the authors studied all possible homogeneous systems of the form (38) and they proved that every bounded solution converges to either an equilibrium solution or to period-two solution. They were able to find a part of the basin of attraction of the period-two solution but not the complete basin of attraction. In the case of system (2) the auxiliary equation for $z_n = y_n/x_n$ is

$$z_{n+1} = \frac{\beta_2 (B_1 + z_n)}{\gamma_1 (B_2 + z_n) z_n} = F(z_n), \quad n = 0, 1, \dots \quad (39)$$

Since F is decreasing, every solution of the auxiliary equation is approaching not necessarily minimal period-two solution. Further analysis can be continued either by checking negative feedback condition for F^2 or by using Theorem 3.2 from [20]. In neither case the complete description of the basins of attraction of the equilibrium and the period-two solution is possible. We prefer our approach because it is more precise and also apply equally well to anticompetitive systems which are not homogeneous. The approach which is making use of homogeneous properties of functions is applicable also to the systems which are neither competitive nor anticompetitive.




FIGURE 2: Global dynamics of System (3).

4. Global Dynamics of System (3) (See Figure 2)

The equilibrium point $E(\bar{x}, \bar{y})$ of System (3) satisfies the following system of equations:

$$\bar{x} = \frac{\gamma_1 \bar{y}}{B_1 \bar{x} + \bar{y}}, \quad \bar{y} = \frac{\alpha_2 + \beta_2 \bar{x}}{\bar{y}}. \quad (40)$$

It is easy to see that System (40) has the unique equilibrium point E in the first quadrant, for all values of the parameters. Indeed, the positive equilibrium point is an intersection of the following two curves:

$$y = \frac{B_1 x^2}{\gamma_1 - x}, \quad (41)$$

$$x = \frac{y^2 - \alpha_2}{\beta_2}. \quad (42)$$

It is clear that at the point of intersection E curve (41) is steeper than curve (42), that is,

$$\left. \frac{dy}{dx} \right|_{(41)} (E) > \left. \frac{dy}{dx} \right|_{(42)} (E), \quad (43)$$

which gives

$$\frac{B_1 \bar{x} (2\gamma_1 - \bar{x})}{(\gamma_1 - \bar{x})^2} > \frac{\beta_2}{2\bar{y}}. \quad (44)$$

This inequality is equivalent to the following inequality:

$$2\beta_1 \bar{x} \bar{y} (2\gamma_1 - \bar{x}) > \beta_2 (\gamma_1 - \bar{x})^2, \quad (45)$$

which in turn is equivalent to

$$2 + 2\frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} > \beta_2 \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^3}. \quad (46)$$

4.1. *Linearized Stability Analysis of System (3).* In this section we prove the following result.

Theorem 12. *The unique equilibrium E of System (3) is a saddle point.*

Proof. The map S associated to System (3) is

$$S(x, y) = \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix} = \begin{pmatrix} \frac{\gamma_1 y}{B_1 x + y} \\ \frac{\alpha_2 + \beta_2 x}{y} \end{pmatrix}. \quad (47)$$

The Jacobian matrix of map (47) is

$$J_S(x, y) = \begin{pmatrix} -\frac{B_1 \gamma_1 y}{(B_1 x + y)^2} & \frac{B_1 \gamma_1 x}{(B_1 x + y)^2} \\ \frac{\beta_2}{y} & -\frac{\alpha_2 + \beta_2 x}{y^2} \end{pmatrix}, \quad (48)$$

and evaluated at the equilibrium point $E = (\bar{x}, \bar{y})$ is

$$J_S(\bar{x}, \bar{y}) = \begin{pmatrix} -\frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} & \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^2} \\ \frac{\beta_2}{\bar{y}} & -1 \end{pmatrix}. \quad (49)$$

The characteristic equation has the following form:

$$\lambda^2 + \left(1 + \frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}}\right) \lambda + \frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} - \beta_2 \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^2} = 0. \quad (50)$$

Set

$$P = 1 + \frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}}, \quad Q = \frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} - \beta_2 \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^2}. \quad (51)$$

Then the necessary and sufficient condition for (50) to have one root inside the unit circle and one root outside the unit circle is $|P| > |1 + Q|$, $P^2 - 4Q > 0$; see [7, 21]. The condition $|P| > |1 + Q|$ leads to $P > -1 - Q$ which is equivalent to

$$2 + 2\frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} > \beta_2 \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^3}, \quad (52)$$

which is condition (46).

The condition $P^2 - 4Q > 0$ becomes

$$\left(1 + \frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}}\right)^2 - 4\left(\frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}} - \beta_2 \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^3}\right) > 0, \quad (53)$$

which is equivalent to

$$\left(1 - \frac{B_1 \bar{x}^2}{\gamma_1 \bar{y}}\right)^2 + 4\beta_2 \frac{B_1 \bar{x}^3}{\gamma_1 \bar{y}^3} > 0, \quad (54)$$

which is clearly satisfied. \square

4.2. Global Results for System (3)

Lemma 13. System (3) has no minimal period-two solution.

Proof. The second iterate of the map S is given as

$$S^2(x, y) = \left(\gamma_1(y + xB_1) \frac{\alpha_2 + x\beta_2}{\beta_2 B_1 x^2 + \beta_2 xy + \alpha_2 B_1 x + \gamma_1 B_1 y^2 + \alpha_2 y}, \frac{y}{(y + xB_1)(\alpha_2 + x\beta_2)} (\gamma\alpha_2 + y\beta_2\gamma_1 + x\alpha_2 B_1) \right). \quad (55)$$

Period-two solution satisfies $S^2(x, y) = (x, y)$ which reduces to the following two equations:

$$\begin{aligned} \gamma_1(y + xB_1)(\alpha_2 + x\beta_2) - x(\beta_2 B_1 x^2 + \beta_2 xy + \alpha_2 B_1 x + \gamma_1 B_1 y^2 + \alpha_2 y) &= 0, \\ y\alpha_2 + y\beta_2\gamma_1 + x\alpha_2 B_1 - (y + xB_1)(\alpha_2 + x\beta_2) &= 0. \end{aligned} \quad (56)$$

Equation (57) leads immediately to $y\gamma_1 - yx - x^2 B_1 = 0$ which is exactly the equilibrium equation (41). Using (41) in (56), we obtain after some elementary simplification that period-two solution satisfies (42). This shows that System (3) has no minimal period-two solution. \square

Lemma 14. The maps S and S^2 associated with System (3) have the following properties.

- (i) The maps S and S^2 are injective.
- (ii) $\det J_{S^2}(x, y) > 0$ for all (x, y) , $y > 0$.

Consequently, S^2 satisfies (O_+) condition and so $\{S^{2n}(x_0, y_0)\}$ is eventually componentwise monotone.

Proof. (i) We will prove that S is injective and the injectivity of S^2 will follow immediately. The condition

$$S(x_1, y_1) = S(x_2, y_2) \quad (58)$$

is reduced to the following two conditions:

$$x_2 y_1 = x_1 y_2, \quad \alpha_2(y_2 - y_1) = \beta_2(x_2 y_1 - x_1 y_2), \quad (59)$$

which immediately implies $y_1 = y_2$ and so $x_1 = x_2$.

- (ii) A direct calculation shows that

$$\det J_{S^2} = \frac{B_1^2 \alpha_2^2 \gamma_1^2 y^2}{(\alpha_2 + x\beta_2)(\gamma\alpha_2 + xy\beta_2 + x\alpha_2 B_1 + x^2 \beta_2 B_1 + y^2 \gamma_1 B_1)^2}, \quad (60)$$

which implies our statement.

The statement on (O_+) condition follows from Theorem 5. \square

Theorem 15. Consider System (3). Then there exists a set $\mathcal{C} \subset \mathcal{D} = [0, \infty) \times (0, \infty)$ which is invariant subset of the basin of attraction of E . The set \mathcal{C} is a graph of a strictly increasing continuous function of the first variable on an interval and separates \mathcal{D} into two connected and invariant components, namely,

$$\begin{aligned} \mathcal{W}_- &:= \{x \in \mathcal{D} \setminus \mathcal{C} : \exists y \in \mathcal{C} \text{ with } x \leq_{se} y\}, \\ \mathcal{W}_+ &:= \{x \in \mathcal{D} \setminus \mathcal{C} : \exists y \in \mathcal{C} \text{ with } y \leq_{se} x\}, \end{aligned} \quad (61)$$

which satisfy the following.

- (i) If $(x_0, y_0) \in \mathcal{W}_+$, then

$$\lim_{n \rightarrow \infty} (x_{2n}, y_{2n}) = E, \quad \lim_{n \rightarrow \infty} (x_{2n+1}, y_{2n+1}) = (0, \infty). \quad (62)$$

- (ii) If $(x_0, y_0) \in \mathcal{W}_-$,

$$\lim_{n \rightarrow \infty} (x_{2n}, y_{2n}) = (0, \infty), \quad \lim_{n \rightarrow \infty} (x_{2n+1}, y_{2n+1}) = E. \quad (63)$$

Proof. Clearly, the rectangle $[0, \gamma_1] \times (0, \infty)$ is an invariant and attracting set for the map S . In particular, $x_n \leq \gamma$ for $n \geq 1$.

Then by Lemma 14 and Theorem 4, every solution $\{(x_n, y_n)\}$ of System (3) has eventually monotone components $\{(x_{2n}, y_{2n})\}$ and $\{(x_{2n+1}, y_{2n+1})\}$, which shows that every bounded solution converges to period-two solution. In view of Lemma 13, there are no minimal period-two solutions and so every bounded solution of System (3) converges to the equilibrium E . In view of Theorems 7 and 8, there is a set \mathcal{C} with described properties. Consequently, every solution with an initial point $(x_0, y_0) \in \mathcal{W}_+$ converges to E , while every solution which starts in \mathcal{W}_- approaches $(0, \infty)$ and is asymptotic to the global unstable manifold $W^u(E)$. \square

5. Global Dynamics of System (4) (See Figure 3)

The equilibrium point $E(\bar{x}, \bar{y})$ of System (4) satisfies the following system of equations:

$$\bar{x} = \frac{\gamma_1 \bar{y}}{A_1 + \bar{x}}, \quad \bar{y} = \frac{\alpha_2 + \beta_2 \bar{x}}{\gamma}. \quad (64)$$

It is easy to see that System (64) has the unique equilibrium point E in the first quadrant, which is an intersection of two parabolas:

$$y = \frac{x(B_1 + x)}{\gamma_1}, \quad (65)$$

$$x = \frac{y^2 - \alpha_2}{\beta_2}. \quad (66)$$

It is clear that at the point of intersection E , curve (65) is steeper than curve (66), that is,

$$\left. \frac{dy}{dx} \right|_{(65)}(E) > \left. \frac{dy}{dx} \right|_{(66)}(E), \quad (67)$$

which gives

$$2\bar{y}(A_1 + 2\bar{x}) > \beta_2 \gamma_1. \quad (68)$$




FIGURE 3: Global dynamics of System (4).

5.1. *Linearized Stability Analysis of System (4).* In this section we prove the following result.

Theorem 16. *The unique equilibrium E of System (4) is a saddle point.*

Proof. The map S associated to System (4) is

$$U(x, y) = \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix} = \begin{pmatrix} \frac{\gamma_1 y}{A_1 + x} \\ \frac{\alpha_2 + \beta_2 x}{y} \end{pmatrix}. \quad (69)$$

The Jacobian matrix of map (69) has the form:

$$J_U(x, y) = \begin{pmatrix} -\frac{\gamma_1 y}{(A_1 + x)^2} & \frac{\gamma_1}{A_1 + x} \\ \frac{\beta_2}{y} & -\frac{\alpha_2 + \beta_2 x}{y^2} \end{pmatrix}, \quad (70)$$

which evaluated at the equilibrium point $E = (\bar{x}, \bar{y})$ is

$$J_U(\bar{x}, \bar{y}) = \begin{pmatrix} -\frac{\bar{x}}{A_1 + \bar{x}} & \frac{\gamma_1}{A_1 + \bar{x}} \\ \frac{\beta_2}{\bar{y}} & -1 \end{pmatrix}. \quad (71)$$

The characteristic equation of System (4) has the following form:

$$\lambda^2 + \left(1 + \frac{\bar{x}}{A_1 + \bar{x}}\right)\lambda + \frac{\bar{x}}{A_1 + \bar{x}} - \frac{\beta_2 \gamma_1}{\bar{y}(A_1 + \bar{x})} = 0. \quad (72)$$

Set

$$P = 1 + \frac{\bar{x}}{A_1 + \bar{x}}, \quad Q = \frac{\bar{x}}{A_1 + \bar{x}} - \frac{\beta_2 \gamma_1}{\bar{y}(A_1 + \bar{x})}. \quad (73)$$

The necessary and sufficient condition for (72) to have one root inside the unit circle and one root outside the unit circle

is $|P| > |1 + Q|$, $P^2 - 4Q > 0$; see [7, 21]. In view of the fact that $P > 1 + Q$, the condition $|P| > |1 + Q|$ leads to $P > -1 - Q$ which is equivalent to

$$2 + 2\frac{\bar{x}}{A_1 + \bar{x}} > \frac{\beta_2 \gamma_1}{\bar{y}(A_1 + \bar{x})}, \quad (74)$$

which is equivalent to the condition (68).

The condition $P^2 - 4Q > 0$ becomes

$$\left(1 + \frac{\bar{x}}{A_1 + \bar{x}}\right)^2 - 4\left(\frac{\bar{x}}{A_1 + \bar{x}} - \frac{\beta_2 \gamma_1}{\bar{y}(A_1 + \bar{x})}\right) > 0, \quad (75)$$

which is equivalent to

$$\left(1 - \frac{\bar{x}}{A_1 + \bar{x}}\right)^2 + 4\frac{\beta_2 \gamma_1}{\bar{y}(A_1 + \bar{x})} > 0, \quad (76)$$

which is clearly satisfied. \square

5.2. *Global Results for System (4)*

Lemma 17. *System (4) has no minimal period-two solution.*

Proof. The second iterate of the map U is given as

$$\begin{aligned} U^2(x, y) &= \left(\frac{1}{y}\gamma_1(x + A_1)\frac{\alpha_2 + x\beta_2}{A_1^2 + xA_1 + y\gamma_1}, \frac{y}{(x + A_1)(\alpha_2 + x\beta_2)}(x\alpha_2 + \alpha_2 A_1 + y\beta_2 \gamma_1)\right). \end{aligned} \quad (77)$$

Period-two solution satisfies $U^2(x, y) = (x, y)$ which reduces to the following two equations:

$$xy(A_1 x + y) = \gamma_1(\alpha_2 + \beta_2 x), \quad (78)$$

$$\gamma_1 y = x(A_1 + x). \quad (79)$$

Equation (79) is exactly the equilibrium equation (65). Using (65) in (78) we obtain after some elementary simplification that period-two solution satisfies (66). This shows that System (4) has no minimal period-two solution. \square

Lemma 18. *The maps U and U^2 associated with System (4) have the following properties.*

- (i) *If $A_1 \beta_2 \neq \alpha_2$, then the maps U and U^2 are injective.*
- (ii) *If $A_1 \beta_2 \neq \alpha_2$, then $\det J_{U^2}(x, y) > 0$ for all (x, y) , $y > 0$.*

Consequently, U^2 satisfies (O+) condition, when $A_1 \beta_2 \neq \alpha_2$.

Proof. (i) We will prove that U is injective which will imply the injectivity of U^2 . The condition

$$U(x_1, y_1) = U(x_2, y_2) \quad (80)$$

is reduced to the following two conditions:

$$\begin{aligned} A_1(y_2 - y_1) &= x_2y_1 - x_1y_2, \\ \alpha_2(y_2 - y_1) &= \beta_2(x_2y_1 - x_1y_2), \end{aligned} \tag{81}$$

which implies that $y_2 - y_1 = (1/A_1)(x_2y_1 - x_1y_2) = (\beta_2/\alpha_2)(x_2y_1 - x_1y_2)$

$$\left(\frac{1}{A_1} - \frac{\beta_2}{\alpha_2}\right)(x_2y_1 - x_1y_2) = 0, \tag{82}$$

and $x_2y_1 - x_1y_2 = 0$, when $A_1\beta_2 \neq \alpha_2$. This implies $y_1 = y_2$ and so $x_1 = x_2$.

(ii) A direct calculation shows that

$$\det J_{U^2} = (\alpha_2 - A_1\beta_2)^2 \frac{\gamma_1}{(\alpha_2 + \beta_2x)(A_1^2 + A_1x + \gamma_1y)^2}, \tag{83}$$

which proves our statement.

The statement on $(O+)$ condition follows from Theorem 5. \square

Theorem 19. Consider System (4). Assume that $A_1\beta_2 \neq \alpha_2$. Then there exists a set $\mathcal{C} \subset \mathcal{D} = [0, \infty) \times (0, \infty)$ which is invariant subset of the basin of attraction of E . The set \mathcal{C} is a graph of a strictly increasing continuous function of the first variable on an interval and separates \mathcal{D} into two connected and invariant components, namely,

$$\begin{aligned} \mathcal{W}_- &:= \{x \in \mathcal{D} \setminus \mathcal{C} : \exists y \in \mathcal{C} \text{ with } x \leq_{se} y\}, \\ \mathcal{W}_+ &:= \{x \in \mathcal{D} \setminus \mathcal{C} : \exists y \in \mathcal{C} \text{ with } y \leq_{se} x\}, \end{aligned} \tag{84}$$

which satisfy the following.

(i) If $(x_0, y_0) \in \mathcal{W}_+$, then

$$\lim_{n \rightarrow \infty} (x_{2n}, y_{2n}) = (\infty, 0), \quad \lim_{n \rightarrow \infty} (x_{2n+1}, y_{2n+1}) = (0, \infty). \tag{85}$$

(ii) If $(x_0, y_0) \in \mathcal{W}_-$,

$$\lim_{n \rightarrow \infty} (x_{2n}, y_{2n}) = (0, \infty), \quad \lim_{n \rightarrow \infty} (x_{2n+1}, y_{2n+1}) = (\infty, 0). \tag{86}$$

Proof. Assume that $A_1\beta_2 \neq \alpha_2$. Then by Lemma 18 and Theorem 4, every solution $\{(x_n, y_n)\}$ of System (4) has eventually monotone components $\{(x_{2n}, y_{2n})\}$ and $\{(x_{2n+1}, y_{2n+1})\}$, which shows that every bounded solution converges to a period-two solution. In view of Lemma 17, there are no minimal period-two solutions and so every bounded solution of System (4) converges to the equilibrium E . In view of Theorems 7 and 8, there is a set \mathcal{C} with described properties. Consequently, the regions \mathcal{W}_- and \mathcal{W}_+ are invariant for U^2 and every solution in \mathcal{W}_- (\mathcal{W}_+) is asymptotic to the unstable manifold $\mathcal{W}^u(E)$ and so the statement of the theorem follows. \square

Theorem 20. Consider System (4). Assume that $A_1\beta_2 = \alpha_2$. Then System (4) can be decoupled and written as

$$\begin{aligned} x_{n+1} &= \frac{\beta_2\gamma_1^2}{x_n(A_1 + x_n)}, \quad y_{n+1} = \beta_1 \frac{\beta_2\gamma_1 + A_1y_n}{y_n^2}, \\ n &= 0, 1, \dots \end{aligned} \tag{87}$$

Every solution $\{(x_n, y_n)\}$ of System (87) has eventually monotone subsequences $\{(x_{2n}, y_{2n})\}$ and $\{(x_{2n+1}, y_{2n+1})\}$. Every bounded solution converges to the unique positive equilibrium. Every unbounded solution $\{(x_n, y_n)\}$ approaches either $(\infty, 0)$ or $(0, \infty)$.

Proof. Using the condition $A_1\beta_2 = \alpha_2$ in the second equation of System (4) gives

$$y_{n+1} = \beta_2 \frac{A_1 + x_n}{y_n}, \tag{88}$$

and so $x_{n+1}y_{n+1} = \beta_2\gamma_1$ which shows that System (4) has an invariant of the form:

$$x_n y_n = \beta_2 \gamma_1, \quad n = 1, 2, \dots \tag{89}$$

Using (89), System (4) is reduced to System (87). Both equations of System (87) are first order difference equations with decreasing functions and so by Theorem 1.19 [7], the subsequences of even and odd indexes are eventually monotonic and so every bounded solution converges to a period-two solution. An immediate checking shows that neither one of the two equations of System (87) has period-two solutions. For example, the unique equilibrium \bar{x} of the first equation of System (87) satisfies

$$x^3 + A_1x^2 - \beta_2\gamma_1^2 = 0, \tag{90}$$

while period-two solution satisfies $f^2(x) = x$ which becomes $x^2((A+x)^2/(A^2x + Ax^2 + C)) = x$ and so is reduced to (90). Thus every bounded solution converges to the unique equilibrium.

The result for unbounded solutions follows immediately from (89). \square

Acknowledgment

The authors are grateful to one of the anonymous referees for pointing out the equation of the stable manifold for (2) and for observing the alternative method of the proof of Theorem 1 mentioned in Remark 11.

References

- [1] S. Kalabušić and M. R. S. Kulenović, "Dynamics of certain anti-competitive systems of rational difference equations in the plane," *Journal of Difference Equations and Applications*, vol. 17, no. 11, pp. 1599–1615, 2011.
- [2] M. R. S. Kulenović and M. Nurkanović, "Basins of attraction of an anti-competitive system of difference equations in the plane," *Communications on Applied Nonlinear Analysis*, vol. 19, no. 2, pp. 41–53, 2012.

- [3] E. Camouzis, M. R. S. Kulenović, G. Ladas, and O. Merino, "Rational systems in the plane—open problems and conjectures," *Journal of Difference Equations and Applications*, vol. 15, no. 3, pp. 303–323, 2009.
- [4] D. Clark and M. R. S. Kulenović, "A coupled system of rational difference equations," *Computers & Mathematics with Applications*, vol. 43, no. 6-7, pp. 849–867, 2002.
- [5] P. Hess, *Periodic-Parabolic Boundary Value Problems and Positivity*, vol. 247 of *Pitman Research Notes in Mathematics Series*, Longman Scientific Technical, Harlow, UK; John Wiley & Sons, New York, NY, USA, 1991.
- [6] M. W. Hirsch and H. Smith, "Monotone dynamical systems," in *Handbook of Differential Equations: Ordinary Differential Equations*, vol. 2, pp. 239–357, Elsevier B. V., Amsterdam, The Netherlands, 2005.
- [7] M. R. S. Kulenović and O. Merino, *Discrete Dynamical Systems and Difference Equations with Mathematica*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2002.
- [8] M. R. S. Kulenović and M. Nurkanović, "Asymptotic behavior of a competitive system of linear fractional difference equations," *Advances in Difference Equations*, vol. 2006, Article ID 19756, 2006.
- [9] A. Brett, M. Garić-Demirović, M. R. S. Kulenović, and M. Nurkanović, "Global behavior of two competitive rational systems of difference equations in the plane," *Communications on Applied Nonlinear Analysis*, vol. 16, no. 3, pp. 1–18, 2009.
- [10] M. Garić-Demirović, M. R. S. Kulenović, and M. Nurkanović, "Global behavior of four competitive rational systems of difference equations in the plane," *Discrete Dynamics in Nature and Society*, vol. 2009, Article ID 153058, 34 pages, 2009.
- [11] H. L. Smith, "Planar competitive and cooperative difference equations," *Journal of Difference Equations and Applications*, vol. 3, no. 5-6, pp. 335–357, 1998.
- [12] H. L. Smith, "The discrete dynamics of monotonically decomposable maps," *Journal of Mathematical Biology*, vol. 53, no. 4, pp. 747–758, 2006.
- [13] A. Brett and M. R. S. Kulenović, "Basins of attraction of equilibrium points of monotone difference equations," *Sarajevo Journal of Mathematics*, vol. 5, pp. 211–233, 2009.
- [14] M. R. S. Kulenović and O. Merino, "Competitive-exclusion versus competitive-coexistence for systems in the plane," *Discrete and Continuous Dynamical Systems B*, vol. 6, no. 5, pp. 1141–1156, 2006.
- [15] M. R. S. Kulenović and O. Merino, "Global bifurcation for discrete competitive systems in the plane," *Discrete and Continuous Dynamical Systems B*, vol. 12, no. 1, pp. 133–149, 2009.
- [16] C. Robinson, *Dynamical Systems: Stability, Symbolic Dynamics, and Chaos*, CRC Press, Boca Raton, Fla, USA, 1995.
- [17] C. A. Clark, M. R. S. Kulenović, and J. F. Selgrade, "On a system of rational difference equations," *Journal of Difference Equations and Applications*, vol. 11, no. 7, pp. 565–580, 2005.
- [18] M. R. S. Kulenović and O. Merino, "Invariant manifolds for competitive discrete systems in the plane," *International Journal of Bifurcation and Chaos*, vol. 20, no. 8, pp. 2471–2486, 2010.
- [19] E. Camouzis, G. Ladas, and L. Wu, "On the global character of the system $x_{n+1} = (\alpha_1 + \gamma_1 y_n)/(x_n)$, $y_{n+1} = (\beta_2 x_n + \gamma_2 y_n)/(B_2 x_n + C_2 y_n)$," *International Journal of Pure and Applied Mathematics*, vol. 53, no. 1, pp. 21–36, 2009.
- [20] Y. S. Huang and P. M. Knopf, "Global convergence properties of first-order homogeneous systems of rational difference equations," *Journal of Difference Equations and Applications*, vol. 18, no. 10, pp. 1683–1707, 2012.
- [21] M. R. S. Kulenović and G. Ladas, *Dynamics of Second Order Rational Difference Equations: With Open Problems and Conjectures*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2001.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

