

2007

Respiration of *Escherichia coli* in the Mouse Intestine

Shari A. Jones

Fatema Z. Chowdhury

Andrew J. Fabich

April Anderson

Darrel M. Schreiner

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/cmb_facpubs

Terms of Use

All rights reserved under copyright.

Citation/Publisher Attribution

Jones, S. A., Chowdhury, F. Z., Fabich, A. J., Anderson, A., Schreiner, D. M., House, A. L., Autieri, S. M.,...Conway, T. (2007). Respiration of *Escherichia coli* in the Mouse Intestine. *Infection and Immunity*, 75(10), 4891-4899. doi: 10.1128/IAI.00484-07

Available at: <http://dx.doi.org/10.1128/IAI.00484-07>

This Article is brought to you for free and open access by the Cell and Molecular Biology at DigitalCommons@URI. It has been accepted for inclusion in Cell and Molecular Biology Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors

Shari A. Jones, Fatema Z. Chowdhury, Andrew J. Fabich, April Anderson, Darrel M. Schreiner, Anetra L. House, Steven M. Autieri, Mary P. Leatham, Jeremy J. Lins, Mathias Jorgensen, Paul S. Cohen, and Tyrrell Conway

Respiration of *Escherichia coli* in the Mouse Intestine[∇]

Shari A. Jones,¹ Fatema Z. Chowdhury,¹ Andrew J. Fabich,¹ April Anderson,¹ Darrel M. Schreiner,¹
Anetra L. House,¹ Steven M. Autieri,² Mary P. Leatham,² Jeremy J. Lins,² Mathias Jorgensen,²
Paul S. Cohen,² and Tyrrell Conway^{1*}

Advanced Center for Genome Technology, University of Oklahoma, Norman, Oklahoma 73019,¹ and Department of Cell and Molecular Biology, University of Rhode Island, Kingston, Rhode Island 02881²

Received 3 April 2007/Returned for modification 1 June 2007/Accepted 30 July 2007

Mammals are aerobes that harbor an intestinal ecosystem dominated by large numbers of anaerobic microorganisms. However, the role of oxygen in the intestinal ecosystem is largely unexplored. We used systematic mutational analysis to determine the role of respiratory metabolism in the streptomycin-treated mouse model of intestinal colonization. Here we provide evidence that aerobic respiration is required for commensal and pathogenic *Escherichia coli* to colonize mice. Our results showed that mutants lacking ATP synthase, which is required for all respiratory energy-conserving metabolism, were eliminated by competition with respiratory-competent wild-type strains. Mutants lacking the high-affinity cytochrome *bd* oxidase, which is used when oxygen tensions are low, also failed to colonize. However, the low-affinity cytochrome *bo₃* oxidase, which is used when oxygen tension is high, was found not to be necessary for colonization. Mutants lacking either nitrate reductase or fumarate reductase also had major colonization defects. The results showed that the entire *E. coli* population was dependent on both microaerobic and anaerobic respiration, consistent with the hypothesis that the *E. coli* niche is alternately microaerobic and anaerobic, rather than static. The results indicate that success of the facultative anaerobes in the intestine depends on their respiratory flexibility. Despite competition for relatively scarce carbon sources, the energy efficiency provided by respiration may contribute to the widespread distribution (i.e., success) of *E. coli* strains as commensal inhabitants of the mammalian intestine.

The intestinal microflora is dominated by diverse anaerobes, providing both a health benefit to the host (15) and a barrier to infection (16, 21). Despite being present in substantially lower numbers, facultative anaerobes, primarily *Escherichia coli* and *Enterococcus faecalis*, are ubiquitous in mammalian intestines (40). While the intestine is commonly thought to be anaerobic (4), the tissues surrounding the lumen are oxygen rich and oxygen diffuses into the intestine at appreciable levels (20). Furthermore, oxygen from swallowed air is present in flatus (29). Oxygen in the intestine apparently has minimal impact on persistence of anaerobes, and it was recently shown that at least one predominant anaerobe, *Bacteroides fragilis*, respire oxygen at low concentrations (5). In contrast to obligate anaerobes, facultative anaerobes (e.g., *E. coli*) grow most rapidly when respiring oxygen and switch to anaerobic respiration in the absence of oxygen or to fermentation in the absence of alternative electron acceptors (17). However, the extent to which facultative anaerobes utilize oxygen to maximize their growth rate in the intestine is not known.

Nutrients consumed for growth of the microflora are thought primarily to be fermentable carbohydrates, the bulk of which are in the form of polysaccharides (37). *E. coli* colonizes the mouse intestine by growing within the polysaccharide-rich mucus layer covering the epithelium but is unable to degrade polysaccharides. Apparently, *E. coli* consumes the mono- and

disaccharides released during degradation of mucosal polysaccharides and dietary fiber (8) by polysaccharide hydrolase enzymes secreted by members of the anaerobic microflora (11) and, perhaps, host colonic epithelial cells (6). Recent studies from our laboratory demonstrate that seven mucus-derived sugars contribute to *E. coli* colonization of the mouse intestine, suggesting that biochemical flexibility is key to its competitiveness in vivo (8). *E. coli* is nearly equally flexible in its respiratory metabolism (17), but nothing is known about the role of bacterial respiration for coupling ATP generation to carbohydrate oxidation in vivo. Thus, it is important to test the hypothesis that respiration confers a competitive advantage to *E. coli* in the intestine.

Enterohemorrhagic *E. coli* (EHEC), has an infectious dose for humans as low as 10 microorganisms and, following ingestion, grows rapidly to a population approaching a billion bacteria per gram of feces (24). Since colonization is the first step in the infection process, it is crucial to understand how EHEC colonizes the intestine because low numbers can survive transport to consumers in foodstuffs such as leafy vegetables, which have caused recent outbreaks in the United States (36). It is not known how EHEC acquires nutrients and generates energy for growth in vivo. While respiration is not a virulence factor per se, our experiments seek to establish the fundamental importance of housekeeping functions, such as energy metabolism, for pathogenesis. Since most mucosal pathogens are facultative anaerobes, these studies of *E. coli* may be extended to include many diseases.

Here we report the results of a systematic mutational analysis designed to identify which respiratory pathways contribute to the ability of commensal and pathogenic *E. coli* to colonize

* Corresponding author. Mailing address: Department of Botany and Microbiology, The University of Oklahoma, 101 David L. Boren Blvd., Norman, OK 73019-0245. Phone: (405) 325-1683. Fax: (405) 325-3442. E-mail: tconway@ou.edu.

[∇] Published ahead of print on 13 August 2007.

TABLE 1. Competitive colonization between respiratory mutants and wild-type *E. coli* strains^a

Respiratory enzyme	Mutant(s)	<i>E. coli</i> colonization (log ₁₀ CFU/g) by strain:			
		EDL933		MG1655	
		Day 1	Day 9	Day 1	Day 9
ATP synthetase	$\Delta(atpA-atpG)$	2.7 ± 0.1	5.6 ± 0.1	5.3 ± 0.5	5.9 ± 0.1
Regulator of aerobic genes	$\Delta arcA$	4.7 ± 0.1	5.6 ± 0.2	3.0 ± 0.2	6.5 ± 0.3
Regulator of anaerobic genes	Δfnr	1.4 ± 0.2	5.6 ± 0.2	1.3 ± 0.1	6.8 ± 0.1
Cytochrome <i>bo</i> oxidase	$\Delta(cyoA-cyoB)$	0.1 ± 0.1	0.4 ± 0.3	0.4 ± 0.1	0.8 ± 0.1
Cytochrome <i>bd</i> oxidase	$\Delta(cydA-cydB)$	2.5 ± 0.1	5.2 ± 0.1	3.8 ± 0.2	5.0 ± 0.1
Cytochrome <i>bd</i> oxidase (assembly)	$\Delta(cydD-cydC)$	2.4 ± 0.1	5.0 ± 0.2	3.3 ± 0.1	6.0 ± 0.2
Nitrate reductase	$\Delta narG$	0.3 ± 0.1	2.0 ± 0.1	0.4 ± 0.2	2.2 ± 0.1
Nitrate reductase	$\Delta narZ$	0.6 ± 0.1	0.1 ± 0.1	0.3 ± 0.1	0.1 ± 0.2
Periplasmic nitrate reductase	$\Delta(napD-napA)$	0.4 ± 0.1	0.1 ± 0.1	0.2 ± 0.1	0.6 ± 0.1
Nitrate reductases	$\Delta narG, \Delta narZ$	0.9 ± 0.1	2.3 ± 0.1	0.1 ± 0.1	2.9 ± 0.2
Nitrate reductases	$\Delta narG, \Delta narZ, \Delta(napD-napA)$	1.3 ± 0.1	3.1 ± 0.1	0.9 ± 0.1	3.8 ± 0.1
Fumarate reductase	$\Delta frdA$	0.8 ± 0.1	3.1 ± 0.2	0.1 ± 0.1	2.8 ± 0.2

^a Mice were fed 10⁵ CFU each of a mutant and its wild-type parent. Mice were transferred to fresh cages every day, and feces no older than 24 h were assayed every other day for 15 days. At each time point, the log₁₀ CFU/gram of feces for the mutant was subtracted from the log₁₀ CFU/gram of feces for the wild type. The average ± standard error of the mean of day 1 and day 9 data from 6 mice is shown. Differences of at least 1 order of magnitude (10-fold) are in boldface type; all values shown in bold are statistically significant ($P < 0.005$; Student's *t* test).

the streptomycin-treated mouse intestine. Our findings lead us to conclude that respiration provides an enormous competitive advantage to *E. coli* in vivo. The results challenge the traditional view that the intestine is strictly anaerobic (4). Instead, we obtained evidence that *E. coli* colonization of the mouse intestine is maximized by the ability to respire oxygen.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The bacterial strains used in this study were derived from *E. coli* MG1655 Str^r (streptomycin resistant), a K-12 strain (33), and *E. coli* EDL933 Str^r, the prototypical O157:H7 strain (31). Cultures were grown at 37°C in Luria-Bertani (LB) medium with gyratory shaking at 250 rpm. Null alleles were constructed by using the allelic replacement method of Datsenko and Wanner (14), as described previously (8), such that target genes were deleted and replaced with kanamycin or chloramphenicol resistance cassettes (used as selectable markers in mouse colonization assays, as described below). The null allele strains are identified in the text by the genes that were deleted; single gene deletions began with the start codon and ended with the stop codon, and multiple gene deletions began with the start codon of the first gene deleted and ended with the stop codon of the last gene deleted. Strains containing multiple mutations were constructed by sequential allelic replacement; the first inserted cassette was removed with FLP recombinase (14), followed by subsequent allelic replacement(s) and removal of the insertion as necessary, leaving the selected marker in the last mutation made. Mutant strains were verified by phenotype analysis and DNA sequencing.

Phenotypic analysis. MOPS [3-(*N*-morpholino) propanesulfonic acid] defined medium was used to grow cultures for growth curves, as described previously (8). Anaerobic cultures were grown in culture tubes filled to the top with N₂-sparged medium, sealed, and incubated in Balch tubes. To test for nitrate or fumarate reductase activity, mutant strains were grown anaerobically overnight in MOPS medium with glycerol (1.6%) as the carbon source and either 50 mM nitrate or fumarate as the electron acceptor. Cell growth was monitored spectrophotometrically by the optical density at 600 nm.

High-performance liquid chromatography analysis. A Dionex DX-500-Microbore system was used with an IonPac AS11 column for anion analysis of mucus. Standards and blanks were analyzed, and standard curves were developed for a 1-mg sample of cecal mucus. Mouse cecal mucus was isolated from the cecum of CD-1 male mice and lyophilized as described previously (50). Regression coefficients for standard curves were calculated and used to demonstrate the linearity of the peak area with respect to concentration.

Mouse colonization experiments. The streptomycin-treated mouse model has been used extensively to study colonization of the mouse large intestine by *E. coli* and *Salmonella enterica* serovar Typhimurium (8, 12, 22, 50). Briefly, three CD-1 male mice, 6 weeks of age, were given drinking water containing streptomycin sulfate (5 g/liter) for 24 h to remove the existing resident facultative microflora

and then starved for food and water for 18 to 24 h. The mice were then fed approximately 10⁵ CFU of both the wild-type and mutant strains in 1 ml of 20% sucrose. The wild-type strains were *E. coli* MG1655 Str^r Nal^r (nalidixic acid resistance) (33) and *E. coli* EDL933 Str^r Nal^r (31); Nal^r was used to distinguish the wild-type (reference strain) from the null allele mutants in fecal plate counts. After the bacterial suspension was ingested, food and streptomycin-water were restored and fecal plate counts were determined at 5 h, 24 h, and on every other day thereafter for 15 days. Fecal samples were homogenized and diluted in 1% tryptone broth and plated on MacConkey agar containing either streptomycin (100 µg/ml) and nalidixic acid (50 µg/ml) to count the wild type or streptomycin and kanamycin (40 µg/ml) or chloramphenicol (30 µg/ml) to count the null allele mutants. Each colonization experiment was repeated on separate occasions, and the plotted values (in figures) represent the average for six mice. The log₁₀ mean number of CFU per gram of feces ± the standard error for each strain in the mice was calculated for each time point. In all experiments, a difference between two strains of ≥10 CFU/g feces was statistically significant (i.e., $P < 0.005$ in Student's *t* test [two tailed with unequal variance]). The limit of detection in fecal plate counts was 10² CFU/g feces. To determine the role of strains during the maintenance stage of colonization, mice were colonized for 10 days with a mutant strain of *E. coli* EDL933, starved for food and streptomycin-water overnight, and fed 10¹⁰ CFU of the wild-type strain *E. coli* EDL933 Str^r Nal^r, after which food and streptomycin-water were replaced.

RESULTS

Colonization assays. The preferred animal model for measuring the relative fitness of two bacterial strains for intestinal colonization is the streptomycin-treated mouse. Streptomycin treatment selectively removes facultative anaerobes while leaving the anaerobic microflora essentially intact; this opens a previously unavailable niche, which can then be colonized by newly introduced microorganisms such as *E. coli* (12, 27, 50). In this model, competing wild-type and mutant strains are fed together to mice and their populations are monitored in fecal plate counts. Previously we showed that colonization involves an initiation stage (5 h to 3 days postfeeding), in which nutrients are not limiting and the population increases from low to high numbers, and a maintenance stage (7 days post-feeding and beyond), in which nutrients are limiting and the population persists at a level correlated with the mutant strain's relative fitness for colonization (8). For this reason, the colonization data shown in Table 1 are given for day 1 (initiation) and day 9 (maintenance) of the 15-day-long experiments. To




FIG. 1. Respiratory mutants exhibited colonization defects in competitive colonization assays. *E. coli* EDL933 $\Delta(atpA-atpG)$ (A) and *E. coli* EDL933 $\Delta(cydA-cydB)$ (C) mutants were eliminated during competition with wild-type *E. coli* EDL933 but were able to colonize when fed alone to mice (B and D, respectively).

compare the bioenergetics of a commensal strain to those of a pathogenic strain, each of the respiratory mutations described in this report was constructed in *E. coli* MG1655 (3), derived from the human isolate *E. coli* K-12, and *E. coli* EDL933 (34), the prototypical strain of *E. coli* O157:H7. Please note that the streptomycin-treated mouse serves as a colonization model for EHEC strains, which do not cause disease in CD-1 mice (49). We found that each of the mutations tested in the commensal and pathogenic strains had a nearly identical impact on colonization. For this reason, colonization curves are shown for *E. coli* EDL933 experiments only.

ATP synthase is necessary for colonization. Since respiratory energy conservation, i.e., ATP generation, requires ATP synthase (32), we tested mutants that lack ATPase for their ability to compete with the wild type in the streptomycin-treated mouse colonization model. The ATPase mutant constructions deleted the *atpA* and *atpG* genes, which encode the F_1 alpha and gamma subunits, respectively, resulting in strains capable only of fermentative energy metabolism (13). In competition with their respective wild-types, *E. coli* EDL933 $\Delta(atpA-atpG)::cat$ and *E. coli* MG1655 $\Delta(atpA-atpG)::cat$ mutants were eliminated from mice within 5 days (Table 1 and Fig. 1A). It is formally possible that mutations inadvertently introduced elsewhere on the genome caused the observed colonization phenotype. However, we would argue that this was not the case, for the following reasons. First, each of the allelic replacements described here and elsewhere (8, 28, 31, 33) was obtained with a frequency that varied less than 1

order of magnitude. Second, half of the mutants tested here and in similar studies (8) had no colonization defects. Third, the results were essentially identical in two different genetic backgrounds (EDL933 and MG1655). The failure of the $\Delta(atpA-atpG)::cat$ mutants to initiate colonization could be due to a general inability to grow in the intestine rather than an inability to compete with the wild type. To distinguish these possibilities, the mutants were fed alone to mice and found to colonize at wild-type levels (Fig. 1B) (data not shown). The ability of the respiratory-defective mutants to colonize alone indicates that fermentation is sufficient for growth of *E. coli* in the mouse intestine, but oxidative phosphorylation is essential for competition with respiratory-competent strains. This finding led us to consider whether one or more modes of respiration are crucial to occupation of intestinal niches formed by electron acceptor availability.

Respiration of oxygen is necessary for colonization. Proof that aerobic respiration is essential for colonization by *E. coli* was obtained by competing mutants lacking the high-affinity cytochrome *bd* oxidase with their respective wild types. The $\Delta(cydA-cydB)::cat$ mutant constructions deleted genes encoding both subunits (I and II) of cytochrome *bd* oxidase, as shown previously (19). The *E. coli* EDL933 $\Delta(cydA-cydB)::cat$ and *E. coli* MG1655 $\Delta(cydA-cydB)::cat$ strains were eliminated by day 11 during competition with their wild-type parents (Table 1 and Fig. 1C). The fitness defect observed for the $\Delta(cydA-cydB)::cat$ strains was not merely a growth defect, since the mutants colonized when fed alone to mice (Fig. 1D) (data not shown). To

confirm the requirement for high-affinity oxygen respiration, $\Delta(\text{cydD-cydC})::\text{cat}$ mutants, which cannot assemble cytochrome *bd* oxidase in the membrane (39), were tested and found to have similar colonization defects (Table 1 and Fig. 2A). The $\Delta(\text{cydD-cydC})::\text{cat}$ mutants were able to colonize when fed to mice alone (Fig. 2B) (data not shown). We note that in addition to CydDC being required for cytochrome *bd* oxidase assembly and activity (39), *cydDC* encodes a glutathione transport system (38), which if relevant might also be required for colonization. Since the $\Delta(\text{cydA-cydB})::\text{cat}$ and $\Delta(\text{cydD-cydC})::\text{cat}$ mutants failed to initiate colonization (i.e., 24 h), it was necessary to determine if they were also defective in the maintenance stage (beyond 7 days). Therefore, mice were precolonized (for 10 days) with *E. coli* EDL933 $\Delta(\text{cydD-cydC})::\text{cat}$ and then challenged with the *E. coli* EDL933 wild type; the mutant was eliminated by the wild type in 7 days, confirming that functional cytochrome *bd* was important for maintenance of colonization (Fig. 2C).

We also examined the importance of the low-affinity cytochrome *bo*₃ oxidase for colonization. The construction of $\Delta(\text{cyoA-cyoB})::\text{cat}$ mutants deleted the genes encoding subunits I and II of the cytochrome *bo*₃ oxidase, as shown previously (2). The results showed that *E. coli* EDL933 $\Delta(\text{cyoA-cyoB})::\text{cat}$ and *E. coli* MG1655 $\Delta(\text{cyoA-cyoB})::\text{cat}$ strains cocolonized with their respective wild-type parents, indicating that respiration of high oxygen levels was not necessary for colonization (Table 1). In summary, the colonization defects of the cytochrome *bd* oxidase mutants challenge the traditional view that the intestine is anaerobic (4). Instead, the results support the hypothesis that a microaerobic niche is critical for both establishing and maintaining *E. coli* in the intestine. Thus, a competitive advantage in vivo is conferred on strains that respire oxygen.

Aerobic respiratory control is necessary for colonization. *E. coli* governs respiratory flexibility via the global regulators ArcA and Fnr. ArcA is a two-component regulator of several hundred genes (30) that responds to the oxidation state of the quinone pool, which is sensed by ArcB (18). Under high oxygen tension, *E. coli* expresses the low-affinity oxidase cytochrome *bo*₃ (encoded by *cyoABCDE*) and the high-affinity oxidase cytochrome *bd* (encoded by *cydAB*) is repressed. Under microaerobic conditions, where oxygen scavenging may play an important role for growth and survival, ArcB phosphorylates ArcA, which represses the *cyoABCDE* operon and activates the *cydAB* and *cydDC* operons (1). Since the experiments described above indicated the importance of cytochrome *bd*, we reasoned that *arcA* mutants would have colonization defects. The construction of *E. coli* EDL933 $\Delta\text{arcA}::\text{cat}$ and *E. coli* MG1655 $\Delta\text{arcA}::\text{kan}$ deleted the gene encoding the response regulator of the ArcAB two-component system, as previously described (23). Although ΔarcA mutants colonized when fed alone (Fig. 3B) (data not shown), they could not compete with their respective wild types and were eliminated from mice within 3 days (Table 1 and Fig. 3A). While it is tempting to speculate that the colonization defect of the ΔarcA mutants resulted solely from failure to induce *cydAB*, the pleiotropic phenotype of the ΔarcA strain makes a number of alternative explanations possible. What is clear from this experiment is that appropriate regulation of aerobic respiratory genes is necessary for *E. coli* to be competitive in vivo.




FIG. 2. Cytochrome *bd* oxidase assembly mutants of *E. coli* EDL933 exhibited colonization defects in competitive colonization assays. *E. coli* EDL933 $\Delta(\text{cydD-cydC})$ was defective in competition with wild-type *E. coli* EDL933 (A) but colonized at wild-type levels when fed alone (B). The $\Delta(\text{cydD-cydC})$ mutant exhibited a maintenance defect when mice were precolonized with *E. coli* EDL933 $\Delta(\text{cydD-cydC})$ and challenged with wild-type *E. coli* EDL933 at day 10 (C).




FIG. 3. Aerobic respiratory and anaerobic global regulatory mutants exhibited colonization defects in competitive colonization assays. *E. coli* EDL933 $\Delta arcA$ was eliminated during competition with wild-type *E. coli* EDL933 (A) but was able to colonize when fed alone (B). *E. coli* EDL933 Δfnr was eliminated during competition with wild-type *E. coli* EDL933 (C) but was able to colonize when fed alone (D).

Anaerobic control is necessary for colonization. Induction of anaerobic processes in *E. coli* is controlled by Fnr, an oxygen-labile transcription factor, which activates transcription of hundreds of genes, including genes that encode respiratory pathways for nitrate and fumarate (25, 35). Since induction of anaerobic respiratory pathways requires Fnr, we constructed $\Delta fnr::kan$ mutants to test the contribution of this global regulator during colonization. The *E. coli* EDL933 $\Delta fnr::kan$ and *E. coli* MG1655 $\Delta fnr::kan$ mutants were fed together with the respective wild types and found to initiate colonization (although not as well as the wild type), and then they were eliminated by day 5 (Table 1 and Fig. 3C). The defect was not an inability to grow in the intestine, since the $\Delta fnr::kan$ mutants colonized when fed alone to mice (Fig. 3D) (data not shown). Since the $\Delta fnr::kan$ mutants exhibit colonization defects in competition with the wild type, this implies that *E. coli* experiences conditions in the intestine that are required for Fnr function (7): i.e., anaerobic or nearly anaerobic conditions. Thus, we conclude that Fnr-dependent genes contribute to colonization success. While these results indicate that appropriate regulation of anaerobic respiration is important for colonization, the specific roles of the alternative pathways were unclear.

Nitrate reductase is necessary for colonization. To determine which anaerobic respiratory pathways were used during colonization, we considered the *in vivo* role of nitrate reduction. Since *E. coli* has three systems for nitrate respiration, it was necessary to consider strains with mutations that eliminated each individually and in combination (17). Mutation of

the primary nitrate reductase was accomplished by deleting *narG* (47) to create *E. coli* EDL933 $\Delta narG::kan$ and *E. coli* MG1655 $\Delta narG::kan$. When $\Delta narG::kan$ strains were fed together with their respective parent strains, they initiated colonization but then declined numerically (2.3 logs; $P < 0.003$) (Table 1 and Fig. 4A). To test the involvement in colonization of the secondary nitrate reductase, mutants were constructed in which *narZ* was deleted, the phenotype of which was described previously (46). Likewise, the involvement of the periplasmic nitrate reductase was tested with a construction that deleted *napD* and *napA*, which encode the assembly protein and large subunit of the reductase, respectively (46). In colonization assays, *E. coli* EDL933 $\Delta narZ::cat$, *E. coli* MG1655 $\Delta narZ::cat$, *E. coli* EDL933 $\Delta(napD-napA)::cat$, and *E. coli* MG1655 $\Delta(napD-napA)::cat$ cocolonized with their wild-type parents, indicating that strains with these individual mutations had no phenotype in the intestine (Table 1).

Although the individual $\Delta narZ::cat$ and $\Delta(napD-napA)::cat$ mutants did not exhibit colonization defects, there is reason to believe these gene systems should be expressed in the wild types under the conditions present in the intestine. First, the intestine contains regions that are microaerobic and others that are anaerobic (20), yet apparently is sufficiently anaerobic overall for Fnr control to be a factor in success of the entire *E. coli* population (Fig. 3). Second, we measured a concentration of 2.62 ± 0.21 mM nitrate in mouse cecal mucus (data not shown), a concentration which is physiologically relevant for controlling expression of the two nitrate-inducible systems (52). Third, *E. coli* cells isolated from the intestine show both




FIG. 4. Anaerobic respiration mutants exhibited colonization defects in competitive colonization assays. *E. coli* EDL933 $\Delta narG$ (A), *E. coli* EDL933 $\Delta narG \Delta narZ$ (B), *E. coli* EDL933 $\Delta narG \Delta narZ \Delta(napD-napA)$ (C), and *E. coli* EDL933 $\Delta frdA$ (D) mutants were defective in competition with wild-type *E. coli* EDL933.

logarithmic and stationary-phase characteristics (26). The *narZYWV* operon that encodes the secondary nitrate reductase is not regulated by anaerobiosis or nitrate but instead is RpoS dependent and induced in stationary phase (9). The *napFD AGHBC* operon that encodes the periplasmic nitrate reductase is maximally induced at 1 mM nitrate and is expressed at one-half of the maximal level at 2.5 mM (52). Thus, it is reasonable to expect that both the periplasmic and secondary nitrate reductases are expressed in the intestine. To investigate the phenotypes of these nitrate reductase mutations in the absence of the primary nitrate reductase, we constructed $\Delta narG \Delta narZ::cat$ and $\Delta narG \Delta narZ \Delta(napD-napA)::cat$ mutants. In colonization assays, *E. coli* EDL933 $\Delta narG \Delta narZ::cat$ and *E. coli* MG1655 $\Delta narG \Delta narZ::cat$ mutants declined 3.2 log units ($P < 0.000006$) relative to the wild-type strain by day 15 (Table 1 and Fig. 4B). Also, the *E. coli* EDL933 $\Delta narG \Delta narZ \Delta(napD-napA)::cat$ and *E. coli* MG1655 $\Delta narG \Delta narZ \Delta(napD-napA)::cat$ mutants declined by 5 log units ($P < 9 \times 10^{-15}$) to populations just above the limit of detection ($< 10^3$ CFU/g feces) by the conclusion of the experiments (Table 1 and Fig. 4D). The additive effect of the sequential nitrate reductase mutations in competition with the wild-type *E. coli* EDL933 in these experiments is statistically significant between the $\Delta narG::kan$ and $\Delta narG \Delta narZ::cat$ mutants ($P < 0.003$) and between the $\Delta narG \Delta narZ::cat$ and $\Delta narG \Delta narZ \Delta(napD-napA)::cat$ mutants ($P < 0.004$). From these results, we conclude the primary nitrate reductase plays the larger role of the three systems in the mouse intestine. In contrast, the $\Delta narZ$ and $\Delta(napD-napA)$ mutations affected colonization only

in the $\Delta narG$ background, suggesting a synergy, rather than mere redundancy, between the primary nitrate reductase and the other two systems. Apparently, conditions in the intestine signal induction of the three nitrate reductase gene systems, which together confer a competitive advantage to *E. coli*.

Fumarate reductase is necessary for colonization. To test the importance of fumarate as an alternative electron acceptor in vivo, we constructed mutants with *frdA*, which are known to inactivate fumarate reductase (43). *E. coli* EDL933 $\Delta frdA::kan$ and *E. coli* MG1655 $\Delta frdA::kan$ mutants were fed together with their respective parent strains and found to initiate colonization but then declined by approximately 4 logs relative to the wild type (Table 1 and Fig. 4D). Thus, fumarate reductase mutants competed poorly, indicating that fumarate is used in the intestine as an alternative electron acceptor. Since fumarate was not detected in intestinal mucus (data not shown), it most likely was generated endogenously, giving rise to succinate as a fermentation product during anaerobiosis (41).

DISCUSSION

Figure 5 shows a model of the respiratory pathways that are critical for successful colonization by both EHEC and commensal *E. coli*. The results indicate that the gut is not strictly anaerobic, because the high-affinity cytochrome *bd* oxidase is required to successfully compete with the wild-type for colonization (Fig. 1 and 2). Also, anaerobic respiration of nitrate and fumarate is essential for *E. coli* in vivo (Fig. 4). We therefore conclude that success of *E. coli* in the gastrointestinal tract




FIG. 5. Model of *E. coli* respiratory pathways. Respiratory oxidases and reductases used by *E. coli* in vivo are shown in black. The oxidase not affecting colonization is shown in gray. The environmental conditions affecting the key regulators of the genes encoding the oxido-reductases are shown. Activation is shown with arrowheads, and repression is shown with diamond heads.

demands respiratory flexibility and use of the best available electron acceptor. The results allow us to deduce the intestinal environment as it is perceived by *E. coli*. Accordingly, the niches defined by mutational analysis of respiratory pathways should correspond with in vivo availability of exogenous electron acceptors (i.e., oxygen and nitrate) but not necessarily fumarate, which is generated endogenously by sugar degradation via central metabolism. These respiratory niches could be open to the entire population, wherein each bacterium simultaneously uses both oxygen and nitrate, or individual cells and microcolonies could each use different electron acceptors. If distinct microaerobic and anaerobic niches were to exist in the intestine, then mutants in the respective respiratory pathways would be maintained at populations corresponding to availability of the cogent electron acceptor when in competition with respiratory-competent wild types. However, since the loss of the high-affinity oxygen, nitrate, or fumarate respiratory pathways leads to the near or complete elimination of the entire population in competition with respiratory-competent strains (Fig. 1C and 4C and D), the data give us reason to think that both aerobic and anaerobic niches are equally crucial. Thus, the behavior of the *E. coli* respiratory mutants implies that the intestinal habitat is at one time microaerobic and at another time anaerobic. Indeed, oxygen tension in the intestine may fluctuate due to dynamic cycles of oxygen diffusion and respiratory consumption by facultative anaerobes.

Cole (10) postulated that the redundancy of respiratory systems and complexities of their regulation ready bacteria for changes in electron acceptor availability: when respiring nitrate, *E. coli* can respond rapidly to the “arrival” of oxygen or cope when nitrate is exhausted. Thus, the physiology, biochemistry, and genetic control of respiratory pathways provide flex-

ibility that is thought to be essential for survival in a changing environment (10). It would appear this strategy is singularly important during animal colonization. We summarize these factors in Fig. 5. For *E. coli* to colonize the intestine, appropriate aerobic respiratory control (ArcA) and anaerobic control (Fnr) are required (Fig. 3). ArcA is most active under microaerobic conditions (i.e., oxygen tension of 2 to 15% of air saturation) (1), and Fnr is most active during transition to anaerobic conditions (i.e., oxygen tension of less than 2% of air saturation) (7). Whole-animal measurements indicated oxygen tensions in the 2 to 7% air saturation range for the mouse colon (20). The measured nitrate concentration in intestinal mucus (2.6 mM [see results]) is near that which results in maximal expression of both nitrate reductase systems (i.e., 1 mM for the periplasmic nitrate reductase and 7 mM for the primary nitrate reductase) (52). Regulation of the periplasmic nitrate reductase genes requires both NarL and NarP (45), while regulation of the primary reductase genes requires NarL only (44). Since NarP exerts its control at low nitrate levels (<4 mM) and NarL control dominates at higher nitrate levels (51), and since both the primary and periplasmic nitrate reductases are necessary for efficient colonization, this suggests that nitrate availability might fluctuate in the intestine. Thus, if the intestinal oxygen tension fluctuates in the anaerobic to microaerobic range and the nitrate concentration fluctuates in the 1 to 7 mM range, then cytochrome *bd* oxidase, the primary nitrate reductase, the periplasmic nitrate reductase, and fumarate reductase all will be expressed in vivo. Indeed, regulation of these gene systems is poised to be most responsive to changes in oxygen and nitrate availability in these concentration ranges (48).

The inference that oxygen availability fluctuates because it is

consumed by bacterial respiration suggests the interesting possibility that facultative anaerobes may make the intestinal environment more anaerobic. Indeed, this conclusion is supported by previous studies of the effect of streptomycin treatment on the mouse anaerobic microflora, which selectively removes facultative anaerobes (i.e., *E. coli*, enterococci, streptococci, and lactobacilli). Following administration of streptomycin, populations of strict anaerobes (e.g., bifidobacteria and clostridia) decreased, while populations of so-called “nanaerobes” were unchanged (22): e.g., *Bacteroides fragilis*, which respire oxygen when available in low concentrations (5). Thus, comparisons of the populations of anaerobes in mice with or without facultative anaerobes present support the hypothesis that oxygen-scavenging facultative anaerobes (e.g., *E. coli*) promote the stability of the predominantly anaerobic microflora, exemplifying how a minor member can have a large impact on an ecosystem.

Despite the apparent competitive advantage gained by oxygen respiration, the *E. coli* population is limited to between 10^8 and 10^9 CFU/g feces: i.e., *E. coli* represents between 1 in 1,000 and 1 in 10,000 bacteria in the intestine. The nutrient-niche hypothesis states that to be successful each species of the intestinal microflora must use at least one carbon source better than all other species (16). Corollary to this hypothesis, the population size of any member of the microflora is determined by the concentration of its preferred nutrient(s). The available concentrations of the seven sugars that contribute to colonization by *E. coli* MG1655 are quite low (12, 37). Since *E. coli* does not secrete polysaccharide-degrading enzymes, its preferred substrates are most likely provided by anaerobes, which degrade mucosal polysaccharides and dietary fiber and are thought to release the breakdown products for use by the host and other microbes (11). These facts lead to the conclusion that *E. coli* maximizes its growth yield by coupling oxidation of low nutrient concentrations to respiration in the intestine. This may be a general strategy of facultative anaerobes, which generally grow well on simple sugars but do not secrete polysaccharide-degrading enzymes. Thus, high-efficiency respiration may ensure the success of facultative anaerobes in the intestine, albeit always in lower numbers, by allowing them to maximize cell yield on scarce resources.

Since most mucosal pathogens are facultative anaerobes, our conclusions may extend to other mucosal pathogens. In support of this idea, *Mycobacterium tuberculosis* genes encoding cytochrome *bd* oxidase and the nitrate transporter were induced during mouse lung infection; a cytochrome *bd* oxidase mutant was attenuated during transition to chronic infection in mice (42). Likewise, *Shigella flexneri* cytochrome *bd* mutants showed decreased intracellular survival and attenuated virulence in mouse infections (53). These examples demonstrate the importance of respiration during infection by particular mucosal pathogens and support the idea that oxygen stimulates infectious disease by providing a competitive advantage for pathogens. Since there appears to be no distinction between enterohemorrhagic and commensal *E. coli* with respect to the respiratory pathways used in vivo, we suggest caution in targeting respiratory metabolism for combating EHEC infections because of potential collateral damage to commensal facultative anaerobes and the resulting instability of the intestinal microbiota (16).

In summary, we have shown that *E. coli* uses both aerobic and anaerobic respiratory pathways during colonization. The results presented in this study support the conclusion that the intestine is microaerobic and that aerobic bacterial respiration in the intestine is essential for competition and therefore successful colonization. Apparently, *E. coli* respire oxygen to optimize its reproduction in animals despite the low availability of its preferred carbon sources, which maximizes its colonization efficiency.

ACKNOWLEDGMENTS

We thank Bruce Roe and David Laux for helpful comments.

This work was supported by grant AI48945 from the National Institutes of Health.

REFERENCES

- Alexeeva, S., K. J. Hellingwerf, and M. J. Teixeira de Mattos. 2003. Requirement of ArcA for redox regulation in *Escherichia coli* under microaerobic but not anaerobic or aerobic conditions. *J. Bacteriol.* **185**:204–209.
- Au, D. C.-T., R. M. Lorence, and R. B. Gennis. 1985. Isolation and characterization of an *Escherichia coli* mutant lacking the cytochrome *o* terminal oxidase. *J. Bacteriol.* **161**:123–127.
- Bachmann, B. J. 1996. Derivations and genotypes of some mutant derivatives of *Escherichia coli* K-12, p. 2460–2488. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed., vol. 2. ASM Press, Washington, DC.
- Backhed, F., R. E. Ley, J. L. Sonnenburg, D. A. Peterson, and J. I. Gordon. 2005. Host-bacterial mutualism in the human intestine. *Science* **307**:1915–1920.
- Baughn, A. D., and M. H. Malamy. 2004. The strict anaerobe *Bacteroides fragilis* grows in and benefits from nanomolar concentrations of oxygen. *Nature* **427**:441–444.
- Beaulieu, J. F., and A. Quaroni. 1991. Clonal analysis of sucrose-isomaltase expression in the human colon adenocarcinoma Caco-2 cells. *Biochem. J.* **280**:599–608.
- Becker, S., G. Holighaus, T. Gabrielyczyk, and G. Unden. 1996. O₂ as the regulatory signal for FNR-dependent gene regulation in *Escherichia coli*. *J. Bacteriol.* **178**:4515–4521.
- Chang, D. E., D. J. Smalley, D. L. Tucker, M. P. Leatham, W. E. Norris, S. J. Stevenson, A. B. Anderson, J. E. Grissom, D. C. Laux, P. S. Cohen, and T. Conway. 2004. Carbon nutrition of *Escherichia coli* in the mouse intestine. *Proc. Natl. Acad. Sci. USA* **101**:7427–7432.
- Chang, L., L. I. Wei, J. P. Audia, R. A. Morton, and H. E. Schellhorn. 1999. Expression of the *Escherichia coli* NRZ nitrate reductase is highly growth phase dependent and is controlled by RpoS, the alternative vegetative sigma factor. *Mol. Microbiol.* **34**:756–766.
- Cole, J. 1996. Nitrate reduction to ammonia by enteric bacteria: redundancy, or a strategy for survival during oxygen starvation? *FEMS Microbiol. Lett.* **136**:1–11.
- Comstock, L. E., and M. J. Coyne. 2003. *Bacteroides thetaiotaomicron*: a dynamic, niche-adapted human symbiont. *Bioessays* **25**:926–929.
- Conway, T., K. A. Kroghfelt, and P. S. Cohen. 29 December 2004, posting date. Chapter 8.3.1.2, The life of commensal *Escherichia coli* in the mammalian intestine. *In* A. Böck, R. Curtiss III, J. B. Kaper, F. C. Neidhardt, T. Nyström, K. E. Rudd, and C. L. Squires (ed.), *EcoSal—Escherichia coli* and *Salmonella*: cellular and molecular biology. ASM Press, Washington, DC.
- Cox, G. B., J. A. Downie, F. Gibson, and J. Radik. 1978. Genetic complementation between two mutant *unc* alleles (*unc* A401 and *unc* D409) affecting the F1 portion of the magnesium ion-stimulated adenosine triphosphatase of *Escherichia coli* K12. *Biochem. J.* **170**:593–598.
- Datsenko, K. A., and B. L. Wanner. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* **97**:6640–6645.
- Eckburg, P. B., E. M. Bik, C. N. Bernstein, E. Purdom, L. Dethlefsen, M. Sargent, S. R. Gill, K. E. Nelson, and D. A. Relman. 2005. Diversity of the human intestinal microbial flora. *Science* **308**:1635–1638.
- Freter, R. 1983. Mechanisms that control the microflora in the large intestine, p. 33–54. *In* D. J. Hentges (ed.), *Human intestinal microflora in health and disease*. Academic Press, Inc., New York, NY.
- Gennis, R. B., and V. Stewart. 1996. Respiration, p. 217–261. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed., vol. 1. ASM Press, Washington, DC.

18. Georgellis, D., O. Kwon, and E. C. Lin. 2001. Quinones as the redox signal for the arc two-component system of bacteria. *Science* **292**:2314–2316.
19. Green, G. N., and R. B. Gennis. 1983. Isolation and characterization of an *Escherichia coli* mutant lacking cytochrome *d* terminal oxidase. *J. Bacteriol.* **154**:1269–1275.
20. He, G., R. A. Shankar, M. Chzhan, A. Samouilov, P. Kuppusamy, and J. L. Zweier. 1999. Noninvasive measurement of anatomic structure and intraluminal oxygenation in the gastrointestinal tract of living mice with spatial and spectral EPR imaging. *Proc. Natl. Acad. Sci. USA* **96**:4586–4591.
21. Hentges, D. J. 1983. Role of the intestinal microflora in host defense against infection, p. 311–331. *In* D. J. Hentges (ed.), *Human intestinal microflora in health and disease*. Academic Press, New York, NY.
22. Hentges, D. J., J. U. Que, S. W. Casey, and A. J. Stein. 1984. The influence of streptomycin on colonization in mice. *Microecol. Theor.* **14**:53–62.
23. Iuchi, S., and E. C. Lin. 1988. *arcA* (*dye*), a global regulatory gene in *Escherichia coli* mediating repression of enzymes in aerobic pathways. *Proc. Natl. Acad. Sci. USA* **85**:1888–1892.
24. Kaper, J. B., J. P. Nataro, and H. L. Mobley. 2004. Pathogenic *Escherichia coli*. *Nat. Rev. Microbiol.* **2**:123–140.
25. Kiley, P. J., and H. Beinert. 1998. Oxygen sensing by the global regulator, FNR: the role of the iron-sulfur cluster. *FEMS Microbiol. Rev.* **22**:341–352.
26. Krogfelt, K. A., L. K. Poulsen, and S. Molin. 1993. Identification of coccoid *Escherichia coli* BJ4 cells in the large intestine of streptomycin-treated mice. *Infect. Immun.* **61**:5029–5034.
27. Laux, D. C., P. S. Cohen, and T. Conway. 2005. Role of the mucus layer in bacterial colonization of the intestine, p. 199–212. *In* J. P. Nataro, P. S. Cohen, H. L. T. Mobley, and J. N. Weiser (ed.), *Colonization of mucosal surfaces*. ASM Press, Washington, DC.
28. Leatham, M. P., S. J. Stevenson, E. J. Gauger, K. A. Krogfelt, J. J. Lins, T. L. Haddock, S. M. Autieri, T. Conway, and P. S. Cohen. 2005. Mouse intestine selects nonmotile *flhDC* mutants of *Escherichia coli* MG1655 with increased colonizing ability and better utilization of carbon sources. *Infect. Immun.* **73**:8039–8049.
29. Levitt, M. D., and J. H. Bond. 1980. Flatulence. *Annu. Rev. Med.* **31**:127–137.
30. Lynch, A. S., and E. C. C. Lin. 1996. Responses to molecular oxygen, p. 1526–1538. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed., vol. 1. ASM Press, Washington, DC.
31. Miranda, R. L., T. Conway, M. P. Leatham, D. E. Chang, W. E. Norris, J. H. Allen, S. J. Stevenson, D. C. Laux, and P. S. Cohen. 2004. Glycolytic and gluconeogenic growth of *Escherichia coli* O157:H7 (EDL933) and *E. coli* K-12 (MG1655) in the mouse intestine. *Infect. Immun.* **72**:1666–1676.
32. Mitchell, P. 1961. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. *Nature* **191**:144–148.
33. Møller, A. K., M. P. Leatham, T. Conway, P. J. M. Nuijten, L. A. M. de Haan, K. A. Krogfelt, and P. S. Cohen. 2003. An *Escherichia coli* MG1655 lipopolysaccharide deep-rough core mutant grows and survives in mouse cecal mucus but fails to colonize the mouse large intestine. *Infect. Immun.* **71**:2142–2152.
34. O'Brien, A. D., A. R. Melton, C. K. Schmitt, M. L. McKee, M. L. Batts, and D. E. Griffin. 1993. Profile of *Escherichia coli* O157:H7 pathogen responsible for hamburger-borne outbreak of hemorrhagic colitis and hemolytic uremic syndrome in Washington. *J. Clin. Microbiol.* **31**:2799–2801.
35. Overton, T. W., L. Griffiths, M. D. Patel, J. L. Hobman, C. W. Penn, J. A. Cole, and C. Constantinidou. 2006. Microarray analysis of gene regulation by oxygen, nitrate, nitrite, FNR, NarL and NarP during anaerobic growth of *Escherichia coli*: new insights into microbial physiology. *Biochem. Soc. Trans.* **34**:104–107.
36. Pearson, H. 2007. The dark side of *E. coli*. *Nature* **445**:8–9.
37. Peekhaus, N., and T. Conway. 1998. What's for dinner? Entner-Doudoroff metabolism in *Escherichia coli*. *J. Bacteriol.* **180**:3495–3502.
38. Pittman, M. S., H. C. Robinson, and R. K. Poole. 2005. A bacterial glutathione transporter (*Escherichia coli* CydDC) exports reductant to the periplasm. *J. Biol. Chem.* **280**:32254–32261.
39. Poole, R. K., F. Gibson, and G. Wu. 1994. The *cydD* gene product, component of a heterodimeric ABC transporter, is required for assembly of periplasmic cytochrome *c* and of cytochrome *bd* in *Escherichia coli*. *FEMS Microbiol. Lett.* **117**:217–223.
40. Savage, D. C. 1977. Microbial ecology of the gastrointestinal tract. *Annu. Rev. Microbiol.* **31**:107–133.
41. Sowers, R. G., and D. P. Clark. 27 July 2004, posting date. Chapter 3.5.3, Fermentative pyruvate and acetyl-coenzyme A metabolism. *In* A. Böck, R. Curtiss III, J. B. Kaper, F. C. Neidhardt, T. Nyström, K. E. Rudd, and C. L. Squires (ed.), *EcoSal—Escherichia coli* and *Salmonella*: cellular and molecular biology. ASM Press, Washington, DC.
42. Shi, L., C. D. Sohaskey, B. D. Kana, S. Dawes, R. J. North, V. Mizrahi, and M. L. Gennaro. 2005. Changes in energy metabolism of *Mycobacterium tuberculosis* in mouse lung and under in vitro conditions affecting aerobic respiration. *Proc. Natl. Acad. Sci. USA* **102**:15629–15634.
43. Spencer, M. E., and J. R. Guest. 1973. Isolation and properties of fumarate reductase mutants of *Escherichia coli*. *J. Bacteriol.* **114**:563–570.
44. Stewart, V. 1993. Nitrate regulation of anaerobic respiratory gene expression in *Escherichia coli*. *Mol. Microbiol.* **9**:425–434.
45. Stewart, V., P. J. Bledsoe, and S. B. Williams. 2003. Dual overlapping promoters control *napF* (periplasmic nitrate reductase) operon expression in *Escherichia coli* K-12. *J. Bacteriol.* **185**:5862–5870.
46. Stewart, V., Y. Lu, and A. J. Darwin. 2002. Periplasmic nitrate reductase (NapABC enzyme) supports anaerobic respiration by *Escherichia coli* K-12. *J. Bacteriol.* **184**:1314–1323.
47. Stewart, V., and C. H. MacGregor. 1982. Nitrate reductase in *Escherichia coli* K-12: involvement of *chlC*, *chlE*, and *chlG* loci. *J. Bacteriol.* **151**:788–799.
48. Tseng, C.-P., J. Albrecht, and R. P. Gunsalus. 1996. Effect of microaerophilic cell growth conditions on expression of the aerobic (*cyoABCDE* and *cydAB*) and anaerobic (*narGHJI*, *frdABCD*, and *dmsABC*) respiratory pathway genes in *Escherichia coli*. *J. Bacteriol.* **178**:1094–1098.
49. Wadolkowski, E. A., J. A. Burris, and A. D. O'Brien. 1990. Mouse model for colonization and disease caused by enterohemorrhagic *Escherichia coli* O157:H7. *Infect. Immun.* **58**:2438–2445.
50. Wadolkowski, E. A., D. C. Laux, and P. S. Cohen. 1988. Colonization of the streptomycin-treated mouse large intestine by a human fecal *Escherichia coli* strain: role of growth in mucus. *Infect. Immun.* **56**:1030–1035.
51. Wang, H., and R. P. Gunsalus. 2003. Coordinate regulation of the *Escherichia coli* formate dehydrogenase *fhnGHI* and *fdhF* genes in response to nitrate, nitrite, and formate: roles for NarL and NarP. *J. Bacteriol.* **185**:5076–5085.
52. Wang, H., C.-P. Tseng, and R. P. Gunsalus. 1999. The *napF* and *narG* nitrate reductase operons in *Escherichia coli* are differentially expressed in response to submicromolar concentrations of nitrate but not nitrite. *J. Bacteriol.* **181**:5303–5308.
53. Way, S. S., S. Sallustio, R. S. Magliozzo, and M. B. Goldberg. 1999. Impact of either elevated or decreased levels of cytochrome *bd* expression on *Shigella flexneri* virulence. *J. Bacteriol.* **181**:1229–1237.