2-27-2015

The Effect of Light Intensity on Prey Detection Behavior in Two Lake Malawi Cichlids, *Aulonocara stuartgranti* and *Tramitichromis* sp.

Margot A.B. Schwalbe

Jacqueline F. Webb

University of Rhode Island, jacqueline_webb@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/bio_facpubs

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Schwalbe MA, Webb JF. (2015). "The Effect of Light Intensity on Prey Detection Behavior in Two Lake Malawi Cichlids, *Aulonocara stuartgranti* and *Tramitichromis* sp." *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology*. Available at: <http://link.springer.com/article/10.1007/s00359-015-0982-y#>

This Article is brought to you for free and open access by the Biological Sciences at DigitalCommons@URI. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

The effect of light intensity on prey detection behavior in two Lake Malawi cichlids,
***Aulonocara stuartgranti* and *Tramitichromis* sp.**

Margot A. B. Schwalbe¹ and Jacqueline F. Webb¹

¹Department of Biological Sciences, University of Rhode Island, 120 Flagg Road, Kingston, RI
02881, USA

Author for Correspondence: Margot A. B. Schwalbe, Department of Biology, Tufts University,
200 Boston Ave, Ste. 4700, Medford, MA USA; margot.schwalbe@tufts.edu,
margot.schwalbe@gmail.com, telephone: (617) 627-0558, fax: (617) 627-0309

Tables: 5

Figures: 7 (2 color)

Abstract

Two Lake Malawi sand-dwelling cichlids (*Aulonocara stuartgranti*, *Tramitichromis* sp.) that have different lateral line phenotypes, but feed on benthic invertebrates, have been shown to use lateral line and/or visual cues to detect prey under light versus dark conditions. The current study examined how ecologically relevant variation in light intensity (0-800 lux) influences detection of prey (mobile, immobile) in each species by analyzing six well-defined behavioral parameters. Both species fed at light intensities ≥ 1 lux; prey type and/or time of day (but not light intensity) predicted all four parameters analyzed with generalized linear mixed models in *A. stuartgranti*, whereas the interaction of light intensity and time of day predicted three of these parameters in *Tramitichromis* sp. Data for all six parameters suggest that the critical light intensity is 1-12 lux for both species, the integration of visual and lateral line input explains differences in the detection of mobile and immobile prey and the behavioral changes that occur at the transition from 1 to 0 lux in *A. stuartgranti*, and that *Tramitichromis* sp. likely uses binocular vision to locate prey. The sensory biology of species that exploit similar food resources will have important implications for the trophic ecology of African cichlid fishes.

Keywords

Vision, lateral line, detection distance, prey detection, sensory ecology

Abbreviations

AICC Akaike information criterion

GLMM Generalized linear mixed model

47	Lx	Lux
48	PAR	Photosynthetically active radiation
49	SL	Standard length
50	TL	Total length

51

52 **Introduction**

53

54 Light in aquatic habitats varies in quality and quantity over time and space (Kirk 2011) and
55 influences the ability of visual predators to detect and capture mobile prey (Vinyard and O'Brien
56 1976; Confer et al. 1978; Lythgoe 1979; Ryer and Olla 1999; Vogel and Beauchamp 1999;
57 Rickel and Genin 2005). Fishes occupying similar habitats may demonstrate variation in
58 visually-mediated prey detection abilities, such as visual thresholds and absorption spectra of
59 visual pigments, which may provide a competitive advantage under particular light conditions
60 (Vogel and Beauchamp 1999; Hofmann et al. 2009). Many fishes are also able to detect prey at
61 low light intensities (e.g., dawn, dusk, at depth, or with increased turbidity), but with reduced
62 capabilities compared to that at higher light intensities. The distance at which free swimming
63 prey are detected dramatically decreases below a certain light intensity ("critical light intensity,"
64 Confer et al. 1978) in salmonids (Dunbrack and Dill 1984; Henderson and Northcote 1985) and
65 some freshwater percomorphs (bluegill, *Lepomis macrochirus*, Vinyard and O'Brien 1976;
66 largemouth bass, *Micropterus salmoides*, Howick and O'Brien 1983; yellow perch, *Perca*
67 *flavescens*, Richmond et al. 2004).

68 Given the importance of multimodal sensory integration in the formulation of behavior,
69 the contributions of the non-visual sensory systems to prey detection (e.g., mechanosensory

lateral line, auditory, olfactory, gustatory, somatosensory/tactile, and in some cases, the electrosensory system; reviewed in Montgomery et al. 2014) must also be considered. Morphological and/or physiological specializations of non-visual sensory systems, including the olfactory system (Parzefall 1993; Montgomery et al. 1999), gustatory system (Atema 1971) and the lateral line system (Janssen 1997; Schwalbe et al. 2012, reviewed in Webb 2014), have been used to predict how these senses provide alternatives to vision for prey detection in light-limited environments. Furthermore, the integration of different combinations of sensory inputs may explain variation in behavior under different environmental conditions (Partridge and Pitcher 1980; Moller 2002; Montgomery et al. 2003; Gardiner and Motta 2012). Several species of fishes have been shown to modulate feeding strategies using a combination of visual and non-visual cues that allow them to feed under a range of light conditions, including darkness (Townsend and Risebrow 1982; Batty et al. 1986; Diehl 1988; Schwalbe et al. 2012).

The mechanosensory lateral line system is known to play important roles in prey detection, as well as in predator avoidance, communication, and navigation around obstacles (Webb et al. 2008; Montgomery et al. 2014). The system demonstrates a great deal of variation, which is defined by the morphology of the cranial and trunk lateral line canals and neuromast receptor organs within them, and the distribution of superficial neuromasts on the skin of the head, trunk and tail (reviewed in Webb 2014). Widened lateral line canals, one of five cranial lateral line canal phenotypes found among bony fishes, has evolved convergently in ~12 teleost families (including deep sea taxa) and appears to be an adaptation for enhanced sensitivity to water flows and prey detection (Denton and Gray 1988, 1989; Montgomery and Coombs 1992; discussed in Schwalbe et al. 2012; reviewed in Webb 2014).

The speciose cichlid fishes of the African Rift Lakes are typically described as visual feeders (Fryer and Iles 1972) and most genera have narrow cranial lateral line canals, but all members of a few genera (e.g., *Alticorpus*, *Aulonocara*, *Aulonocranus*, *Trematocara*, *Trematocranus*, Konings 2007) have widened lateral line canals suggesting the capacity for lateral line mediated prey detection (Konings 1990). Two genera of non-mbuna, haplochromine cichlids in Lake Malawi, *Aulonocara* (widened canals) and *Tramitichromis* (narrow canals; Fig. 1), provide an interesting taxon pair for comparison of prey detection strategies since both feed on benthic invertebrates in the sand, and thus appear to be ecologically similar. Schwalbe et al. (2012) and Schwalbe and Webb (2014) analyzed the behavioral responses of *Aulonocara stuartgranti* and *Tramitichromis* sp. to tethered live and dead prey (=adult brine shrimp, *Artemia* sp.), in experiments carried out under light and dark conditions in which the lateral line system was experimentally inactivated. These studies demonstrated that *A. stuartgranti* uses a combination of inputs to its visual and lateral line systems to detect prey in the light, but depends on its lateral line system to detect prey in the dark. Furthermore, these studies showed that deactivation of the lateral line system of *A. stuartgranti* significantly affected prey detection behavior and revealed that other senses (olfaction, gustation, and somatosensory/tactile) were insufficient to initiate prey detection behavior in the dark. In contrast, *Tramitichromis* sp. did not feed in the dark, and the inactivation of the lateral line system had little effect on prey detection behavior in the presence of light, demonstrating that it is a visual predator.

Aulonocara and *Tramitichromis* species appear to share a food resource (benthic invertebrates in sandy substrates), but occupy different depth ranges (*Aulonocara* species at depths of 5-120 m and *Tramitichromis* species at depths of < 15 m; Fryer and Iles 1972; Konings 1990, 2007) and use different strategies to detect and capture benthic invertebrate prey in the

field. Species of *Aulonocara* swim just above the substrate to sense hydrodynamic flows generated by benthic invertebrates in or on the substrate in the field (Konings 2007). In contrast, species of *Tramitichromis* typically capture invertebrate prey by plunging into the substrate, filling their mouth with sand, and sifting out prey with their gill rakers in the field (= “sand sifting,” Fryer, 1959). This sand sifting behavior appears to be synonymous with the “winnowing” behaviors observed in some surfperches (Laur and Ebeling 1983) and vision likely contributes to the ability to locate patches of high quality food resources where “winnowing” takes place (Holbrook and Schmitt 1984; Schmitt and Holbrook 1984).

This study uses the same methods used in prior studies (Schwalbe et al. 2012; Schwalbe and Webb 2014) to test the hypothesis that variation in light intensity (0-800 lx) will have different effects on the detection of live (mobile) and dead (immobile) benthic invertebrate prey in *A. stuartgranti* and *Tramitichromis* sp.

Materials and methods

Study species

Adult *Aulonocara stuartgranti* and *Tramitichromis* sp. (unidentifiable to species level, J. Stauffer, pers. commun.), is referred to as *Tramitichromis* throughout. These fish were reared in the laboratory from breeding stock originally acquired from commercial suppliers (*A. stuartgranti*: Bluegrass Aquatics, Louisville, KY, USA; *Tramitichromis*: Old World Exotic Fish, Inc., Homestead, FL, USA and Life Fish Direct, Draper, UT, USA). They were housed in small groups by species in 190 L aquaria at $26\pm1^{\circ}\text{C}$ and 1.0 ± 0.2 p.p.t. salinity (using Cichlid Lake Salt,

Seachem Laboratories, Inc., Madison, GA, USA) with standard white fluorescent light on a 12h:12h diurnal cycle and, equipped with appropriate mechanical and biological filtration. Fish were fed daily with cichlid pellets (New Life Spectrum Cichlid Formula; New Life International, Inc., Homestead, FL) and supplemented with live adult brine shrimp. Individual fish were not used in feeding experiments if breeding behavior was observed. Animal care and all experimental procedures followed an approved IACUC protocol.

Light environment in the experimental tank

Light in the experimental tank was provided by two fluorescent light fixtures (Lithonia Lighting, Model GRW 2 14 CSW CO M4, Conyers, GA, USA) fitted with full spectrum bulbs (BlueMax lamps, Full Spectrum Solutions, Jackson, MI, USA) positioned above the tank and within an opaque curtain enclosure. The curtain (black canvas) was suspended from a rectangular plywood frame placed 2 m above the top of the tank in order to exclude ambient light from entering the set-up during all behavioral trials (Fig. 2a). Light intensity was varied by changing the height of the lights above the water surface and using combinations of different neutral density filters covering the lights (Lee Filters, Burbank, CA, USA). Light intensity (in lux [lx], lumen/m², and photosynthetically active radiation [PAR], $\mu\text{mol photons/m}^2/\text{s}$) and color spectrum were measured using a spectrometer (range: 340-1028 nm, Jaz spectrometer, Ocean Optics, Dunedin, FL, USA) connected to a 2 m optical fiber (QP400-2-UV/VIS, Ocean Optics) fitted with a cosine corrector (CC-3, Ocean Optics). Water temperature was monitored during experiments and the fluorescent bulbs did not raise the temperature of the experimental tank.

Light intensities used in this study were based on the following data and calculations. First, light levels present during sunrise/sunset to darkness are known for other freshwater habitats (Harden Jones 1956; Ali 1959) and can range from 1000 lx (early twilight) to 0 lx (new moon, Table 1). Second, few direct measurements of light intensities at different depths in Lake Malawi are available, so the light intensity at specific depths were estimated with the following equation:

$$I_t = I_s \times e^{-\varepsilon \times T}$$

where I_s and I_t are the light intensities at the surface (S) and at depth (T); and ε is the light extinction coefficient. The average light intensity at the surface of Lake Malawi at midday on a clear sunny day is approximately 2000 $\mu\text{mol photons/m}^2/\text{s}$ (~108,000 lx). This photon flux was derived from cloudless surface irradiance for Lake Malawi (Guildford et al. 2000). Using light extinction coefficient of either 0.10 m^{-1} (Patterson et al. 2000), 0.13 m^{-1} (Guildford et al. 2007), or 0.43 m^{-1} (Guildford et al. 2007) depending on location and season, the light intensity at many depths can be estimated under these conditions (Table 1).

Full spectrum bulbs were used because they provide the range of wavelengths that correspond to the range of known absorption peaks of retinal photopigments in species of *Aulonocara* and *Tramitichromis*. For instance, absorption peaks for *A. hueseri* are at 415 nm (violet), 484 nm (blue-green) and 526 nm (green; Jordan et al. 2006) and absorption peaks for *T. intermedius* are at 455 nm (blue), 532 nm (green) and 569 nm (red; Parry et al. 2005). In the experimental tank, full spectrum bulbs generated major and minor light peaks at 404, 435, 487, 545, 587, and 611 nm, and neutral density filters were used to change light intensity did not appreciably change the light spectrum in the experimental tank (Fig. 2b)

Experiments

Behavioral trials and video analysis of six well-defined behavioral parameters (number of prey strikes, detection distance, detection angle, detection-to-strike velocity, swimming phase [glide, pause] at detection and, prey type preference [order of prey strikes]) were carried out as in Schwalbe et al. (2012) and Schwalbe and Webb (2014) with slight modifications. A total of sixty trials were conducted using *A. stuartgranti* (30 trials, $n = 6$ fish, 75-85 mm total length [TL], 4 females, 2 male) and *Tramitichromis* (30 trials, $n = 6$ fish, 75-98 mm TL, 1 female, 5 males) in order to quantify variation in behavioral responses to live (mobile) and dead (immobile) prey (= tethered adult brine shrimp) at five light intensities between 0 and 800 lx.

Trials were conducted in an experimental tank (120 x 75 x 60 cm; 560 L) with 5 cm of sand covering the bottom of the tank. Light intensity and spectral measurements (with ± 0.01 accuracy, measured in lx and PAR) were taken directly above the center of each mesh platform (to which live and dead prey were tethered, see below) before and after each trial, and light intensity and spectrum were found to be consistent at all six platforms and trials (Figs. 2b, c). Each fish was acclimated to a particular light intensity in the experimental tank for at least 30 minutes prior to a trial. The transition between photopic (cone-mediated) and scotopic (rod-mediated) vision occurs at approximately 1 lx, and light-adapted fish may take 30 minutes (and up to 3 hours) to become dark-adapted (Ali 1959). Thus, the 30+ minute light adaptation period was judged to be sufficient to allow the fish's visual system to adjust to the light level for a given trial.

Before each trial, 12 adult brine shrimp (*Artemia* sp.) were tethered in pairs (1 live and 1 dead, freshly frozen) on each of six mesh platforms (10 x 10 cm), which were positioned in a 2 x

3 matrix so that the top of each platform flush with the sand surface. The water filtration system for the experimental tank was then turned off to eliminate acoustic and hydrodynamic noise. At the start of a trial, a fish was released into the experimental arena from behind an opaque divider and feeding behavior was recorded for 30 min using an HD digital video camera (Sony © HDR-CX550V, 30 frames per second) mounted directly above the tank, which provided a dorsal view of the experimental arena. Trials at 1-800 lx were carried out with standard fluorescent room lights on for all but the lowest light levels (1-12 lx). Dark trials (0 lx) were conducted with room lights off, but with infrared illumination (peak = 840 nm, range 800-880 nm; Speco Provideo, IR-200/24, Amityville, NY) to allow video recording of behavior.

Each fish was run through five trials, one trial per day each at a single light intensity, progressing from highest to lowest intensity on subsequent days (e.g., 800, 112, 12, 1, and then 0 lx). Trials were carried out in this order to increase the likelihood that a fish would respond to prey at lower light intensities (especially in the dark, 0 lx), as was suggested by preliminary results. Trials were conducted over four months and the mean time between the first (800 lx) and last (0 lx) trial for a given fish was 11 days (range = 6-19 days).

“Light” trials (1-800 lx) started midday to late afternoon (11:00-17:00) and “dark” trials (0 lx) took place shortly after sunset (19:00-21:00; soon after room lights had automatically shut off; as in Schwalbe et al. 2012; Schwalbe and Webb 2014). Dark trials (0 lx) were not carried out during the day (during the light phase of the lab’s light:dark cycle) in order to avoid the introduction of extraneous light. In addition, it was known that placing fish in low light or darkness during normal daylight hours would disrupt feeding behavior (M.A.B. Schwalbe and A. Mensinger, pers. obs.), and that species that normally feed both in full light during the day and at

night (e.g., during the dark phase of a lab's light:dark cycle) were unresponsive in dark (0 lx) trials carried out during the day.

To assess the number of prey detections that lead to prey strikes, unconsumed prey were counted at the end of each 30-minute trial and strike success was also confirmed in video recordings. Video sequences leading to each prey strike were exported to Premier Pro (Adobe, CS5) for further analysis. Analysis of sequential video frames was used to identify the phase of swimming behavior (thrust, glide, or pause) during which prey detections occurred. Detection distance and detection angle were measured in these images using ImageJ (NIH, v. 1.41o). Detection distance was defined as the distance from the tip of a fish's mouth to the prey, measured in the frame immediately before the fish oriented towards it (e.g. before a turn defining detection). For each prey strike, detection-to-strike velocity was calculated by dividing detection distance by the time interval between detection and initiation of a strike. Detection angle was measured in the same video frame in which detection distance was measured, and was defined as the angle between a line extending anteriorly along midline of the fish (body axis) and a line drawn from the prey to the tip of the fish's mouth.

Statistical analysis

Four of the six behavioral parameters were analyzed using generalized linear mixed models (GLMMs; number of prey strikes, detection distance, detection-to-strike velocity, phase of search behavior during which detections occurred). In addition, a ranking method (Taplin 2007) was used to analyze prey preferences (live versus dead prey) and circular statistics were used to analyze detection angles. All continuous data (e.g. detection distance and detection-to-strike

velocity) were tested for normality (Kolmogorov-Smirnov test) and were $\log_{10}$ transformed to achieve normality (detection distance and detection-to-strike velocity). All statistical tests were considered significant at $P < 0.05$ and values are given as means $\pm$ s.e.m.

Start time (= time of day, 0-24 hr) for trials conducted at the five different light intensities was analyzed with nonparametric tests (e.g. Kruskal-Wallis test and Mann-Whitney U test) to determine whether time of day affected feeding behavior. This analysis showed that the times at which light trials (1-800 lx) and dark trials (0 lx) started did not differ between species (Mann-Whitney U test, $P > 0.05$), but trial start time varied among light intensities in each species (Kruskal-Wallis test, *A. stuartgranti*: $K = 22.804$, $P < 0.001$; *Tramitichromis*: $K = 20.141$, $P < 0.001$). Thus, time of day (=trial start time) was included in all GLMM analyses.

Four-way GLMM analyses (SPSS, IBM, v. 22) were used to test whether species (*A. stuartgranti*, *Tramitichromis*), light intensity (0-800 lx), prey type (live, dead), and/or trial start time (0-24 hr) predict differences in each of four behavioral parameters (number of prey strikes, detection distance, detection-to-strike velocity, and phase of search behavior during which detections occurred). Three-way GLMM analyses were used to further examine whether light intensity, prey type, and/or trial start time predict differences in the four behavioral parameters in each species separately. The selection of random (individual) and fixed effects (species, light intensity, prey type, and trial start time), including repeated measures for the same individual, was addressed in all analyses. Different types of GLMMs were used to account for the different types of data collected in this study (summarized in Table 2) and the most parsimonious model was selected for each behavioral parameter based on the corrected Akaike information criterion (AICC).

The order in which live (mobile) and dead (immobile) prey were struck was analyzed in each species following Taplin (2007). This method assumes that when presented with equal numbers of two or more types of prey, the order in which prey are consumed provides information about prey preference – that prey consumed first are more highly preferred than prey consumed second, third, etc. and the last prey consumed is the least preferred. While differences in handling time, encounter rates, and relative mobility of prey can potentially complicate the results of this sort of analysis (Durham et al. 2012, McWilliam et al. 2013), such variation was minimized in the current study by offering equal numbers of live and dead brine shrimp tethered in the same arrangement to platforms placed in a 2 x 3 matrix in all trials. The null hypothesis for this analysis was that live and dead prey would be consumed randomly during a trial. Videos were analyzed so that each prey consumed was assigned a rank number (first prey consumed=1, second prey consumed=2, etc.), and any remaining prey were assigned an average of the remaining preference scores, and considered “tied for last.” A pair of preference scores for live and dead prey at each light intensity was calculated for each fish. The pairs of scores from all of the fish were considered independent samples and thus grouped by light intensity and species for analysis. A score of 6.5 (based on presentation of six live and six dead prey, 12 total prey in a trial) indicated no preference, a score of <6.5 revealed a preference for that prey type, and a score of >6.5 indicated no preference or that prey type was ignored or avoided. Scores for live and dead prey at each light intensity and for each species, were compared separately using paired *t*-tests (SPSS, IBM, v. 22).

Detection angles were analyzed with circular statistics (Oriana v. 3, Kovach Computing Services, Anglesey, UK). Rayleigh tests were performed for each species to test whether detections of live and dead prey at each of the five light intensities (0-800 lx) occurred at

uniformly or non-uniformly distributed positions around the fishes' body relative to the body axis (e.g. to define the receptive field). Watson's U^2 tests were used to determine if detection angles differed with prey type and with light intensity within and between species.

Results

Aulonocara stuartgranti and *Tramitichromis* sp. actively swam around the experimental tank during trials at all light intensities, including darkness (0 lx), and used a saltatory search strategy (a cyclic sequence of a caudal fin thrust, glide, and pause) while exploring the tank. Of the 360 total prey presented to fish during all 60 trials, *A. stuartgranti* struck at 299 prey (=83%) and *Tramitichromis* struck at 231 prey (=64%; see Figs. 3, 4). Prey were detected by both species during a glide or a pause, but never during a thrust (see Fig. 5).

Four-way GLMM analyses (Table 3) indicated that species alone did not predict differences in any of the four behavioral parameters (number of prey detections, detection distance, detection-to-strike velocity, or swimming phase at prey detection), and that only the interaction of species, light intensity, and prey type had a significant effect only on number of prey detections. The interaction of light intensity and time of day predicted differences in all four behaviors. Light intensity alone predicted differences in all four behaviors, and time of day predicted differences in three behavioral parameters (number of prey detections, detection distance, detection-to-strike velocity), but not in swimming phase at prey detection. Separate three-way GLMM analyses for each species (Table 4, see below) revealed interesting trends that are indicative of species differences in prey detection behavior. Analyses of prey preference (live

vs. dead prey) and prey detection angle, carried out using other statistical methods, also indicated differences in behavior between species, but did not consider time of day.

Feeding behavior of *Aulonocara stuartgranti*

GLMM analyses (Table 4) showed that light intensity did not significantly predict any of the four behavioral parameters in *A. stuartgranti* (number of prey detections, detection distance, detection-to-strike velocity, or swimming phase at prey detection; GLMMs, $P > 0.05$; Table 4). However, time of day predicted the number of prey detections, and the interaction of time of day and prey type predicted both detection distance and detection-to-strike velocity (GLMMs, $P < 0.05$; Table 4). Neither light intensity, time of day, prey type, nor their interactions, predicted swimming phase at prey detection.

An examination of data for each of the behavioral parameters revealed informative trends. *A. stuartgranti* struck at high numbers of both live (mobile) and dead (immobile) prey at light intensities of 1-800 lx (Figs. 3a, 4a). They detected prey during a pause about half of the time, but detected 61% of prey in a pause at the highest light intensity (800 lx; Fig. 5a). Detection distance and detection-to-strike velocity appeared to not vary among light intensities of 12-800 lx, but fish tended to detect live prey from greater distances (mean = 8.7-9.6 cm) than dead prey (6.0-6.9 cm) and to detect live prey at higher detection-to-strike velocities (9.7-10.4 cm/s) than dead prey (6.9-7.4 cm/s; Figs. 4c, e). At 1 lx, fish tended to detect live and dead prey from similar distances (mean = 6.5 and 6.3 cm, respectively; Fig. 4c) and similar detection-to-strike velocities (6.8 cm/s and 7.8 cm/s, respectively; Fig. 4e). Live and dead prey (combined) were detected at non-uniformly distributed positions around the fishes' bodies at light intensities

≥1 lx (Rayleigh test, $P < 0.001$; $\pm 90^\circ$ from body axis) with no differences in the distribution of angles among pairs of light intensities with the exception of the two highest light intensities (112 lx versus 800 lx; Watson's U^2 test, $U^2 = 0.19$, $P < 0.05$; Fig. 6a). Finally, fish tended to prefer live prey at all light intensities, but only demonstrated a statistically significant preference for live prey at 112 lx, but not at 800 lx (Table 5; Fig. 7a), which is not easily explained.

In the dark (0 lx), prey detection behavior of *A. stuartgranti* was different than at light intensities ≥ 1 lx. Fish struck at only 22 prey (=30.6% of total prey presented; Fig. 3a), and tended to detect prey at even shorter distances (Fig. 4c) and at slower detection-to-strike velocities (Fig. 4e) than when at least some light was present. Fish tended to detect more live prey than dead prey (mean of 2.7 and 1.0, respectively; Figs. 3a, 4a), showed a statistically significant preference for live prey (Table 5, Fig. 7a), and detected live prey from more than twice the distance than dead prey (3.2 and 1.4 cm, respectively; Fig. 4c). In addition, detection-to-strike velocity at 0 lx was about one half of that at higher light intensities (~ 3.5 -5 cm/s at 0 lx versus ~ 7 -10 cm/s at ≥ 1 lx), but fish tended to detect live prey at somewhat higher detection-to-strike velocities than dead prey (Fig. 4e). In the dark, 95% of prey were detected during a glide and only a few prey (5%) were detected during a pause (Fig. 5a). Prey (live and dead combined) were detected at positions uniformly distributed around fishes' bodies (Rayleigh test, $P > 0.05$) at a wide range of angles ($\pm 180^\circ$ from body axis, Fig. 6a), but the distribution of detection angles did not differ for live versus dead prey (Watson's U^2 test, $P > 0.05$).

Feeding behavior of *Tramitichromis*

GLMM analyses (Table 4) showed that, in contrast to *A. stuartgranti*, the interaction of light intensity and time of day predicted three of four behavioral parameters (number of prey detections, detection distance, and detection-to-strike velocity). As in *A. stuartgranti*, neither light intensity, time of day, nor prey type, or their interactions, predicted swimming phase at prey detection. Prey type did not predict any of the four behavioral parameters in *Tramitichromis*, and the interaction of light intensity and prey type predicted only detection distance (Table 4).

An examination of trends for each of the behavioral parameters revealed that *Tramitichromis* tended to strike at high numbers of prey (Fig. 3b, 4b), and >60% of prey (live and dead, combined) were detected during a pause at light intensities of 1-800 lx (Fig. 5b). At light intensities of 12-800 lx, fish struck at live and dead prey from similar, long detection distances (means = 9.8-10.1 and 8.5-10.0 cm, respectively) and at high detection-to-strike velocities (9.6-10.5 and 8.7-9.3 cm/s, respectively). In contrast, at 1 lx, fish tended to strike at both live and dead prey at similar, but shorter detection distances (6.9 and 6.3, respectively) and lower detection-to-strike velocities (7.0 and 6.1, respectively; Figs. 4d, f) than at higher light intensities. Both live and dead prey were detected at non-uniform positions around the body (Rayleigh test, $P < 0.001$), which defined a very narrow range of detection angles from the body axis ($\pm 40^\circ$); distributions were the same for live prey and dead prey at light intensities of 1-800 lx (Watson's U^2 , $P > 0.05$; Fig. 6b). Fish tended to prefer live prey at different light intensities, but only showed a statistically significant preference for live prey at the highest light intensity (800 lx; Table 5; Fig. 7b).

Despite being active in the dark (0 lx), *Tramitichromis* only struck at only 3 prey (=4.2% of the 72 prey presented). These strikes are likely to have been the result of random encounters with prey as opposed to being the result of active search and directed strikes.

Discussion

The multiple statistical analyses presented here, and the detailed examination of trends in the detection of live and dead prey at different light intensities in each species indicate that light intensity affects prey detection behavior in different ways in *Aulonocara stuartgranti* and *Tramitichromis*.

Feeding behavior of *Aulonocara stuartgranti* and *Tramitichromis*

Prey type and/or time of day, but not light intensity, were predictors of three of the four behavioral parameters (number of prey detections, detection distance, and detection-to-strike velocity) analyzed using GLMMs in *A. stuartgranti*. The lack of significance for light intensity is consistent with the use of lateral line cues (see also Schwalbe et al. 2012), but also suggests that *A. stuartgranti* may use a light-independent circadian rhythm to interpret time of day. This is consistent with the occurrence of *Aulonocara* species at depths up to 120 m in Lake Malawi where light is limited or absent (Konings 1990, 2007) and in caves where spawning has been reported (Grant et al. 1987), and thus where normal diurnal variation in light intensity may not be a consistent or reliable cue for the regulation of behavior. The significance of prey type as a predictor of detection distance and detection-to-strike velocity is illustrated by apparent differences in numbers of live and dead prey detected at the same light intensities (Fig. 4; see also Schwalbe et al. 2012; Schwalbe and Webb 2014), the tendency to prefer live prey at all light intensities, and the statistically significant preference for live prey in the dark.

In contrast, in *Tramitichromis*, it is the interaction of light intensity and time of day that predict these same three behavioral parameters. The importance of light intensity not surprising because *Tramitichromis* uses visual, but not lateral line cues, for prey detection and does not feed in the dark (Schwalbe and Webb 2014). Furthermore, these two factors are correlated both in the lab where the fish were reared (on a 12:12 hr light/dark cycle) as well as in the relatively shallow waters in their natural habitat in Lake Malawi, which is just 9-17° south of the equator where fish experience 11-13 hours of daylight per day on an annual basis (<http://astro.unl.edu>). Thus, these fish have evolved and are reared in environments where light intensity and time of day are tightly correlated. The independent roles of these two factors in predicting behavior would need to be addressed in additional experiments, which were out of the scope of this study.

Swimming phase (glide, pause) during which prey were detected was predicted neither by light intensity nor by time of day in either species. The ability to detect prey during a glide or pause will affect both the stabilization of the visual field (for vision-mediated detection) and/or the magnitude of environmental and self-generated hydrodynamic noise (for lateral line-mediated detection). *A. stuartgranti* and *Tramitichromis* both detected between 40% and 70% of prey during a pause at light intensities of ≥ 1 lx, suggesting the importance of stabilizing the visual field for prey detection at these light intensities. Prey type (which defines the presence or absence of an additional visual motion stimulus) did not predict swimming phase at detection for *Tramitichromis* ($P < 0.053$), but a larger sample size may have yielded a different statistical outcome. Prey type also did not predict swimming phase at prey detection for *A. stuartgranti*, but the shift to 95% of prey detections during a glide in the dark (where stabilization of visual field is irrelevant), and their preference for live prey (that generate hydrodynamic flows detected in the dark; Schwalbe and Webb 2014), are important indicators of the overall importance of prey type.

A. stuartgranti detected live prey at distances of less than half of a body length and at lower detection-to-strike velocities at a low light intensity (1 lx) and in the dark (0 lx). In the presence of at least some light, lower detection-to-strike velocities would also reduce self-generated hydrodynamic noise (Montgomery et al. 2009), enhancing lateral line-mediated prey detection, which would suggest that fish would tend to detect prey during a pause. However, the high proportion of detections (95%) at relatively low detection-to-strike velocities, while not eliminating self-generated noise, would bring a fish into the vicinity of potential prey that are generating detectable hydrodynamic flows (Schwalbe et al. 2012).

Role of vision and critical light intensities

The importance of vision in *A. stuartgranti* and *Tramitichromis* is further supported by a consideration of critical light intensities and the potential differences in the use of binocular vision. Prey detection at relatively long distances is consistent with vision-mediated prey detection in fishes (Vinyard and O'Brien 1976, Confer et al. 1978, Henderson and Northcote 1985, Mazur and Beauchamp 2003), and at higher light intensities detection of free swimming prey generally occurs at longer distances (Vinyard and O'Brien 1976; Richmond et al. 2004; Bergstrom and Mensinger 2009). In this study, both *A. stuartgranti* and *Tramitichromis* tended to demonstrate the longest detection distances at the highest light intensities, which is thus consistent with vision-mediated prey detection. Detection distances may not increase as light intensity increases further in a given species (Schmidt and O'Brien 1982), but may decrease sharply below a "critical light intensity" (Confer et al. 1978). Trends in behavioral parameters in the current study reveals that the critical light intensity for fish feeding on tethered adult brine

shrimp is between 12 and 1 lx for both *A. stuartgranti* and *Tramitichromis*. This is comparable to the critical light intensities for other freshwater teleosts in studies feeding on free-swimming *Daphnia* (11-50 lx, in bluegill, Vinyard and O'Brien 1976; in lake trout, brook trout, and bluegill, Confer et al. 1978), amphipods (5-25 lx, in round goby, logperch, slimy sculpin, and spoonhead sculpin, Bergstrom and Mensinger 2009), or on small fish (~6-18 lx, in largemouth bass, Howick and O'Brien 1983; lake trout, Vogel and Beauchamp 1999). At low light intensities (below the critical intensity, at 1 lx), the ability of *A. stuartgranti* to detect more prey than *Tramitichromis*, but at comparable distances suggests that *A. stuartgranti* may have superior visual abilities for prey capture at these lower light intensities. This is consistent with their distribution at a wider depth range than *Tramitichromis* and the observation of reproductive behaviors in caves (Grant et al. 1987), but whether *Aulonocara* species possess adaptations for increased sensitivity and/or visual acuity as found in known crepuscular or nocturnal teleosts (reviewed in Warrant 2004; Schmitz and Wainwright 2011) requires further study.

The potential for binocular vision can be revealed by looking at behavioral evidence for differences in the size of visual fields under different light conditions and between species. While visual predators may respond differently to stimuli in different portions of their visual fields (Collin 1989; McComb and Kajiura 2007; Miyazaki et al. 2011), it is detection angle that reflects the overall size of the visual field, which is defined by the size, shape, and position of the eyes (Collin and Shand 2003). *A. stuartgranti* demonstrates a wide range of detection angles at light intensities ≥ 1 lx ($\pm 90^\circ$ from body axis) and an even wider range of angles in darkness (0 lx, $\pm 180^\circ$ from body axis). This shift is correlated with differences in behavioral parameters at 1 lx versus 0 lx, which are interpreted as a shift between primarily vision-mediated prey detection to lateral line-mediated prey detection. Lateral line-mediated detection of prey around the body is

477 enabled by the more sensitive widened cranial lateral line canals that characterize *Aulonocara*
478 species, and by the broad distribution of canal and superficial neuromasts on the skin of the head,
479 trunk and tail, which is typical of cichlids and of most teleosts (reviewed in Webb 2014). In
480 contrast, *Tramitichromis* detected prey at a range of angles ($\pm 40^\circ$ from body axis) that was less
481 than half of that for *A. stuartgranti* ($\pm 90^\circ$ from body axis) at light intensities of 1-800 lx, with
482 one exception (Watson's U^2 test, $P < 0.05$). This suggests that *Tramitichromis*, but likely not *A.*
483 *stuartgranti*, uses binocular vision and depth perception to detect prey at a distance (as
484 demonstrated in other teleosts, Sivak 1978; Blanco-Vives et al. 2011; Miyazaki et al. 2011).
485 Furthermore, *Tramitichromis* tends to swim higher above the substrate than *A. stuartgranti* when
486 searching for prey in the laboratory (Schwalbe and Webb 2014). Coupled with the use of
487 binocular vision, this search strategy could explain the tendency for *Tramitichromis* to detect
488 benthic prey at somewhat longer distances than *A. stuartgranti* (Figs. 4c, d).

489 The movements of the appendages of the live prey used in this study presumably generate
490 a visual motion stimulus, and an enhanced dispersal of an odor plume (not evaluated here), in
491 addition to a hydrodynamic stimulus, which addresses the importance of multimodal integration
492 in the formulation of prey detection behavior. However, prey type predicted detection distance
493 and detection-to-strike velocity only in *A. stuartgranti*, which tended to strike at live prey at
494 longer detection distances and at higher velocities than for dead prey at the same light intensities
495 (12-800 lx). At 1 lx, detection distances were about one body length or less, which is within the
496 effective range of the lateral line system (Coombs 1999). Behavior is consistent with the use of
497 the lateral line system in addition to vision for detection of live prey by *A. stuartgranti* in full
498 light.

In contrast, in *Tramitichromis*, prey type did not predict any of the four behavioral parameters analyzed using GLMMs, although the interaction of prey type and light intensity did predict detection distance. However, the examination of data trends showed that *Tramitichromis* demonstrates comparable values and trends for live and dead prey with reference to number of prey detections, detection distance (despite the significance of its interaction with prey type) and detection-to-strike velocity at light intensities of 1-800 lx. These results also substantiate results of a prior laboratory study (Schwalbe and Webb 2014) that showed that *Tramitichromis* is a visual predator, which is not dependent on the detection of hydrodynamic stimuli generated by live prey. However, the lack of significance of prey type indicates that *Tramitichromis* does not respond to a visual motion stimulus that are likely to have been generated by live (but not dead) prey, which is surprising given the feeding strategies that these fish employ in nature. In the relatively shallow, well-lit waters of Lake Malawi, *Tramitichromis* species typically capture prey by plunging into the substrate, filling their mouth with sand, and sifting out prey with their gill rakers (= sand sifting, Fryer 1959). The sensory basis for the plunge and sift feeding behavior needs to be determined experimentally, but the results of this study suggest that it is a visual stimulus and not an associated motion stimulus generated by live prey that influences where *Tramitichromis* initiates feeding behavior in the field.

The connection between experimental light conditions and light levels in Lake Malawi

As in other lakes, the photic conditions in Lake Malawi are dynamic and many factors influence the light environment, including habitat type, water depth, and proximity to the lake bottom (Sabbah et al. 2011), as well as meteorological events, eutrophication, turbidity, and both diurnal

and seasonal changes in light quality and quantity. In shallow water, full spectrum light is typically present and middle wavelengths transmit best, but shorter and longer wavelengths attenuate rapidly (Dalton et al. 2010). Further, the irradiance spectrum differs between waters overlying sandy and rocky substrates, where light transmission in water above sand is shifted to longer wavelengths compared to that above rocky habitats (Sabbah et al. 2011).

The extent to which species of *Aulonocara* and *Tramitichromis* forage using vision at different depths can be approximated by comparing behavioral data from the current study to estimates of depths at which particular light intensities are predicted to occur in Lake Malawi. The light extinction coefficients in Table 1 are representative of pelagic ($\epsilon = 0.10 \text{ m}^{-1}$, Patterson et al. 2000; $\epsilon = 0.13 \text{ m}^{-1}$, Guildford et al. 2007) and nearshore ($\epsilon = 0.43 \text{ m}^{-1}$, Guildford et al. 2007) habitats in Lake Malawi, but disparities in water clarity between these areas are likely influenced by nutrient loading and sedimentation from deforestation, intense agricultural practices, and erosion in nearshore areas (Bootsma and Jorgensen 2006). Estimations based on low light extinction coefficients (e.g. $\epsilon = 0.10 \text{ m}^{-1}$ or 0.13 m^{-1}) suggest that *Aulonocara* species could visually detect prey at 71 to 92 m ($\geq 12 \text{ lx}$) and with some visual limitations at ~89 to 115 m where light levels are at ~1 lx. Some *Aulonocara* species are found to depths of 120 m (Konings 2007), so they may be able to visually detect prey in these depths at midday when light intensities are highest. Alternatively, when light extinction coefficients are used ($\epsilon = 0.43 \text{ m}^{-1}$), the maximum depths at which *Aulonocara* species could reliably detect prey are greatly reduced (to 21 m and 27 m, respectively). In the lab, *Tramitichromis* was able detect prey at a light intensity of 1 lx, which translates to depths of 89 to 115 m if the light extinction coefficient is low. However, these fish are typically found in shallower waters ($<15 \text{ m}$, Konings 1990, 2007), so the ability of *Tramitichromis* to find prey at 1 lx is more relevant for the potential for feeding

early or late during the day. Given its dependence on vision for prey detection (Schwalbe and Webb 2014), *Tramitichromis* species may be limited to shallow habitats so that the visual detection of prey is not compromised. In contrast, *Aulonocara* species can feed at low light intensities and in the dark, which can explain the wider range of depths at which they occur in Lake Malawi. They may also be crepuscular or nocturnal in habit, which may also facilitate other behaviors (e.g., social interactions) at low light intensities.

Conclusions

A. stuartgranti fed on prey at a range of ecologically relevant light intensities, including darkness, and *Tramitichromis* was also able to feed at low light intensities, but not in darkness. In *A. stuartgranti*, the influence of time of day on several aspects of its behavior suggests that it may use circadian rhythms to regulate behavior in nature where diurnal light cues may not be available (e.g. at greater depth, in caves). The integration of visual and non-visual (e.g., lateral line) sensory modalities can explain the statistically non-significant trends in behavior. Similarly, the dramatic change in behavior from 1 lx to 0 lx is consistent with a transition from primarily vision-mediated to exclusively lateral line-mediated prey detection behavior. In contrast to *A. stuartgranti*, *Tramitichromis* depends on vision-mediated prey detection (Schwalbe and Webb 2014); in this study its behavior was significantly affected by the interaction of light intensity with time of day, but these two factors could not be teased apart. Finally, in an ecological context, the tendency of *Tramitichromis* species to live in shallower, well-lit habitats, in contrast to *Aulonocara* species, which live at a wide range of depths and light environments, suggests that sensory capabilities may allow *Aulonocara* species to escape competition with *Tramitichromis*

species for prey resources, thus facilitating niche differentiation between these taxa. Field observations in Lake Malawi are needed to test this hypothesis, which would provide an important link between the morphology, feeding behavior, and ecology of cichlid fishes.

Acknowledgements

We thank Karen Carleton (University of Maryland) for her expertise in cichlid vision, Stephanie Guildford (University of Minnesota Duluth) for her expertise on light levels in Lake Malawi, and Graham Forrester (University of Rhode Island) for statistical expertise. We also thank Edward Baker (Facilities Manager, RI NSF EPSCoR Marine Life Science Facility), Emily Becker, Brandon Fuller, Christopher Holland, Callie Veelenturf, Rebecca Scott, and Benjamin Sevey for assistance with fish husbandry, Christopher Kenaley (Harvard University) for doing the μ CT scans, and Benjamin Sevey for generating the 3D images in Fig. 1. This work was carried out under an approved University of Rhode Island IACUC protocol (#AN08-11-005). This research was assisted by funds provided by NSF EPSCoR Cooperative Agreement EPS-0554548 and EPS-100405, the College of the Environment and Life Sciences (University of Rhode Island) and was supported by NSF grant IOS 0843307 to JFW.

References

Ali MA (1959) The ocular structure, retinomotor and photobehavioral responses of juvenile pacific salmon. *Can J Zoology* 37:965-996

590 Atema J (1971) Structures and functions of the sense of taste in the catfish (*Ictalurus natalis*).
 591 Brain Behav Evol 4:273-294
 592 Batty RS, Blaxter JHS, Libby DA (1986) Herring (*Clupea harengus*) filter-feeding in the dark.
 593 Mar Biol 91:371-375
 594 Bergstrom MA and Mensinger AF (2009) Interspecific resource competition between the
 595 invasive round goby and three native species: logperch, slimy sculpin, and spoonhead
 596 sculpin. Trans Am Fish Soc 138:1009-1017
 597 Blanco-Vives B, Aliago-Guerrero M, Cañavate JP, García-Mateos G, Martín-Robles AJ,
 598 Herrera-Pérez P, Muñoz-Cueto JA, Sánchez-Vázquez FJ (2011) Metamorphosis induces
 599 a light-dependent switch in Senegalese sole (*Solea senegalensis*) from diurnal to
 600 nocturnal behavior. J Biol Rhythms 27:135-144
 601 Bootsma H, Jorgensen SE (2005) Lake Malawi/Nyasa: Experience and lessons learned brief. pp
 602 259–276. In: companion CD-ROM for: ILEC (2005). Managing lakes and their basins for
 603 sustainable use: A report for lake basin managers and stakeholders. International Lake
 604 Environment Committee Foundation, Kusatsu, Japan
 605 Collin SP (1989) Topographic organization of the ganglion cell layer and intraocular
 606 vascularization in the retinae of two reef teleosts. Vision Res 29:765-775
 607 Collin SP, Shand J (2003) Retinal sampling and the visual field in fishes. In: Collin SP, Marshall
 608 NJ (eds) Sensory processing in aquatic environments. Springer, New York, pp 139-169
 609 Confer JL, Howick GL, Corzette MH, Kramer SL, Fitzgibbon S, Landesberg R (1978) Visual
 610 predation by planktivores. Oikos 31:27-37
 611 Coombs S (1999) Signal detection theory, lateral-line excitation patterns and prey capture
 612 behaviour of mottled sculpin. Anim. Behav. 58: 421–430

613 Dalton BE, Cronin TW, Marshall NJ, Carleton KL (2010) The fish eye view: are cichlids
 614 conspicuous? J Exp Biol 213:2243-2255

615 Denton EJ, Gray JAB (1988) Mechanical factors in the excitation of the lateral lines of fish. In:
 616 Atema J, Fay RR, Popper AN, Tavolga WN (eds) Sensory biology of aquatic animals.
 617 Springer. New York, pp 595-617

618 Denton EJ, Gray JAB (1989) Some observations on the forces acting on neuromasts in fish
 619 lateral line canals. In: Coombs S, Gorner P, Münz H (eds) The mechanosensory lateral
 620 line: neurobiology and evolution. Springer, New York, pp 229-246

621 Diehl S (1988) Foraging efficiency of three freshwater fishes: effects of structural complexity
 622 and light. Oikos 53:207-214

623 Dunbrack RL, Dill LM (1984) Three-dimensional prey reaction field of the juvenile coho salmon
 624 (*Oncorhynchus kisutch*). Can J Fish Aquat Sci 41:1176-1182

625 Durham SR, Dietl GP, Visaggi CC (2012) The mismeasure of behavior: a natural history
 626 revision of prey preference in the banded tulip snail. J Shellfish Res 31:101-109

627 Fryer G (1959) The trophic interrelationships and ecology of some littoral communities of Lake
 628 Nyasa with especial reference to the fishes, and a discussion of the evolution of a group
 629 of rock-frequenting Cichlidae. Proc Zool Soc Lond 132:153-281

630 Fryer G, Iles TD (1972) The cichlid fishes of the great lakes of Africa: their biology and
 631 evolution. Oliver and Boyd, Edinburgh

632 Gardiner JM, Motta PJ (2012) Largemouth bass (*Micropterus salmoides*) switch feeding
 633 modalities in response to sensory deprivation. Zoology 115: 78-83

634 Grant SM, Dieckhoff HW, Mayland HJ, Meyer MK (1987) Ecology of *Aulonocara* Regan, 1922
 635 in Lake Malawi. Cour. Forsch.-Inst. Senckenberg 94:131-139

636 Guildford SJ, Bootsma HA, Fee EJ, Hecky RE, Patterson G (2000) Phytoplankton nutrient status
 637 and mean water column light intensity in Lakes Malawi and Superior. *Aquat Ecosyst*
 638 *Health* 3:35-45
 639 Guildford SJ, Bootsma HA, Taylor WD, Hecky RE (2007) High variability of phytoplankton
 640 photosynthesis in response to environmental forcing in oligotrophic Lake Malawi/Nyasa.
 641 *J Great Lakes Res* 33:170-185
 642 Harden Jones, FR (1956) The behaviour of minnows in relation to light intensity. *J Exp Biol*
 643 33:271-281
 644 Henderson MA, Northcote TG (1985) Visual prey detection and foraging in sympatric cutthroat
 645 trout (*Salmo clarki clarki*) and dolly varden (*Salvelinus malma*). *Can J Fish Aquat Sci*
 646 42:785-790
 647 Hofmann CM, O'Quin KE, Marhsall NJ, Cronin TW, Seehausen O, Carleton KL (2009) The
 648 eyes have it: regulatory and structural changes both underlie cichlid visual pigment
 649 diversity. *PLOS Biol* 7(12):e1000266
 650 Holbrook SJ, Schmitt RJ (1984) Experimental analyses of patch selection by foraging black
 651 surfperch (*Embiotoca jacksoni* Agazzi). *J Exp Mar Biol Ecol* 79:39-64
 652 Howick GL, O'Brien WJ (1983) Piscivorous feeding behavior of largemouth bass: experimental
 653 analysis. *Can J Fish Aquat Sci* 42:785-790
 654 Janssen J (1997) Comparison of response distance to prey via the lateral line in the ruffe and
 655 yellow perch. *J Fish Biol* 51:921-930
 656 Jordan R, Kellogg K, Howe D, Juanes F, Stauffer J, Loew E (2006) Photopigment spectral
 657 absorbance of Lake Malawi cichlids. *J Fish Biol* 68:1291-1299

658 Kirk JTO (2011) Light and photosynthesis in aquatic ecosystems, 3rd edn. Cambridge University
659 Press, New York

660 Konings A (1990) Koning's book of cichlids and other fishes of Lake Malawi. TFH Publications
661 Inc., Neptune City, New Jersey

662 Konings A (2007) Malawi cichlids in their natural habitat, 4th edn. Cichlid Press, El Paso, Texas

663 Laur DR, Ebeling AW (1983) Predator-prey relationships in surfperches. *Env Biol Fish* 8:217-
664 229

665 Lythgoe JN (1979) The ecology of vision. Clarendon Press, Oxford

666 Mazur MM, Beauchamp DA (2003) A comparison of visual prey detection among species of
667 piscivorous salmonids: effects of light and low turbidities. *Environ Biol Fish* 67:397-405

668 McComb DM, Kajiura SM (2008) Visual fields of four batoid fishes: a comparative study. *J Exp*
669 *Biol* 211:482-490

670 McWilliam RA, Minchinton TE, Ayre DJ (2013) Despite prolonged association in closed
671 populations, an intertidal predator does not prefer abundant local prey to novel prey. *Biol*
672 *J Linn Soc* 108:812-820

673 Miyazaki T, Iwami T, Meyer-Rochow VB (2011) The position of the retinal area centralis
674 changes with age in *Champscephalus gunnari* (Channichthyidae), a predatory fish from
675 coastal Antarctic waters. *Polar Biol* 34:1117-1123

676 Moller P (2002) Multimodal sensory integration in weakly electric fish: a behavioral account. *J*
677 *Physiology-Paris* 96: 547-556

678 Montgomery JC, Bleckmann H, Coombs S (2014) Sensory ecology and neuroethology of the
679 lateral line. In: Coombs S, Bleckmann H, Fay RR, Popper AN (eds) The lateral line
680 system. Springer, New York, pp 121-150

681 Montgomery JC, Coombs S (1992) Physiological characterization of lateral line function in the
682 Antarctic fish *Trematomus bernacchii*. Brain Behav Evol 40:209-216

683 Montgomery JC, Diebel C, Halstead MBD, Downer J (1999) Olfactory search tracks in the
684 Antarctic fish *Trematomus bernacchii*. Polar Bio 21:151-154

685 Montgomery JC, McDonald F, Baker CF, Carton AG, Ling N (2003) Sensory integration in the
686 hydrodynamic world of rainbow trout. Roy Soc Lond B Bio 270(2): S195-S197

687 Montgomery JC, Windsor S, Basset D (2009) Behavior and physiology of mechanoreception:
688 separating signal and noise. Integr Zool 4:3-12

689 Parry JWL, Carleton KL, Spady T, Carboo A, Hunt DM, Bowmaker JK (2005) Mix and match
690 color vision: tuning spectral sensitivity by differential opsin gene expression in Lake
691 Malawi cichlids. Curr Biol 15:1734-1739

692 Partridge BL, Pitcher TJ (1980) The sensory basis of fish schools: relative roles of lateral line
693 and vision. J Comp Physiol 135: 315-325

694 Parzefall J (1993) Behavioural ecology of cave-dwelling fishes. In: Pitcher T (ed) Behaviour of
695 teleost fishes, 2nd edn. Chapman & Hall, London, pp 573-608

696 Patterson G, Hecky RE, Fee EJ (2000) Effect of hydrological cycles on planktonic primary
697 productivity in Lake Malawi/Niassa. Adv Ecol Res 31:421-430

698 Richmond HE, Hrabik TR, Mensinger AF (2004) Light intensity, prey detection and foraging
699 mechanisms of age 0 year yellow perch. J Fish Biol 65:195-205

700 Rickel A, Genin A (2005) Twilight transitions in coral reef fish: the input of light-induced
701 changes in foraging behaviour. Anim Behav 70:133-144

702 Ryer CH, Olla BL (1999) Light-induced changes in the prey consumption and behavior of two
703 juvenile planktivorous fish. Mar Ecol Prog Ser 181:41-51

704 Sabbah S, Gray SM, Boss ES, Fraser JM, Zatha R, Hawryshyn CW (2011) The underwater
 705 photic environment of Cape Maclear, Lake Malawi: comparison between rock- and sand-
 706 bottom habitats and implications for cichlid fish vision. *J Exp Biol* 214:487-500
 707 Schmidt D, O'Brien WJ (1982) Planktivorous feeding ecology of arctic grayling (*Thymallus*
 708 *arcticus*). *Can J Fish Aquat Sci* 39:475-482
 709 Schmitt R, Holbrook SJ (1984) Ontogeny of prey selection by black surfperch *Embiotoca*
 710 *jacksoni* (Pisces: Embiotocidae): the roles of fish morphology, foraging behavior, and
 711 patch selection. *Mar Ecol Prog Ser* 63:6-12
 712 Schmitz L, Wainwright PC (2011) Nocturnality constrains morphological and functional
 713 diversity in the eyes of reef fishes. *BMC Evol Biol* 11:338
 714 Schwalbe MAB, Bassett DK, Webb JF (2012) Feeding in the dark: lateral-line-mediated prey
 715 detection in the peacock cichlid *Aulonocara stuartgranti*. *J Exp Biol.* 215:2060-2071
 716 Schwalbe MAB, Webb, JF (2014) Sensory basis for detection of benthic prey in two Lake
 717 Malawi cichlids. *Zoology* 117:112-121
 718 Sivak JG (1978) The functional significance of the aphakic space of the fish eye. *Can J Zool*
 719 56:513-516
 720 Taplin RH (2007) Experimental design and analysis to investigate predator preferences for prey.
 721 *J Exper Mar Biol Ecol* 344:116-122
 722 Townsend CR, Risebrow AJ (1982) The influence of light level on the functional response of a
 723 zooplanktonivorous fish. *Oecologia* 53:293-295
 724 Vinyard GL, O'Brien WJ (1976) Effects of light and turbidity on reaction distance of bluegill
 725 (*Lepomis macrochirus*). *J Fish Res Board Can* 33:2845-2849

726 Vogel JL, Beauchamp DA (1999) Effects of light, prey size, and turbidity on reaction distances
 727 of lake trout (*Salvelinus namaycush*) to salmonid prey. Can J Fish Aquat Sci 56:1293-
 728 1297
 729 Warrant EJ (2004) Vision in the dimmest habitats on Earth. J Comp Physiol A 190:765-789
 730 Webb JF (2014) Morphological diversity, development, and evolution of the mechanosensory
 731 lateral line system. In: Coombs S, Bleckmann H, Fay RR, Popper AN (eds) The lateral
 732 line system. Springer, New York, pp 17-72
 733 Webb JF, Montgomery JC, Mogdans J (2008) Bioacoustics and the lateral line system of fishes.
 734 In: Webb JF, Fay RR, Popper AN (eds) Fish Bioacoustics. Springer, New York, pp 145-
 735 182
 736

Figure Legends

Fig. 1 Lateral line canals and canal neuromasts in *Aulonocara stuartgranti* (widened canals) and *Tramitichromis* (narrow canals) visualized using fluorescent vital staining (4-di-2-ASP, 63 μ M, 5 min; **a**, **c**), μ CT imaging (reconstructed from 14 μ m slices; **b**, **d**, **e**, **g**, **h**), and scanning electron microscopy (**f**). **a** Lateral view of *A. stuartgranti* revealing series of larger infraorbital (IO), mandibular (MD), and preopercular (PO) canal neuromasts and very small superficial neuromasts on the skin (juvenile, 25 mm standard length [SL]). Neuromast number and distribution is the same in *Tramitichromis* **b** μ CT reconstruction of *A. stuartgranti* (adult, 78 mm SL) indicating the location of the supraorbital (SO), IO, MD, and PO canals in dermatocranial bones. **c** Ventral view of the head of *A. stuartgranti* (juvenile, 28 mm SL), revealing canal neuromasts in the MD and PO canals. **d** *A. stuartgranti* (adult, 78 mm SL) and **e** *Tramitichromis* (adult, 79 mm SL) in ventral view. Asterisks (*) denote the locations of the MD and PO canal neuromasts, as visualized in **c**; canal neuromasts are found in floor of the canal, between canal pore positions in the canal roof. Note the much larger pores in *A. stuartgranti* (**d**) than in *Tramitichromis* (**e**). **f** MD canal neuromast in a juvenile *A. stuartgranti*. Ciliary bundles of the sensory hair cells are evident in an elongate sensory strip in the middle of the diamond-shaped neuromast. Double-headed arrow below the sensory strip indicates the axis of physiological sensitivity of the hair cells, as well as the long axis of the canal in which the neuromast is found. Scale bar = 10 μ m. **g** *A. stuartgranti* and **h** *Tramitichromis* in frontal-ventral view with the pores of the SO, IO, MD, and PO canals that are directed ventrally, toward the source of stimuli generated by benthic prey. The pores on the right side of each fish in **g** and **h** have been enhanced to increase their visibility.

Fig. 2 Experimental setup and light conditions used to record feeding behavior of *A. stuartgranti* and *Tramitichromis*. **a** Diagram of experimental tank with front panel of light curtain removed. Light sources originate from two fluorescent light fixtures (full spectrum light) and two infrared (IR) lights. **b** Sample light spectra generated by fluorescent lights in behavioral trials. The y-axis was compressed to illustrate peaks in the visual spectrum (400-700 nm) that occurred in 1-800 lx trials. These peaks were consistent when light intensity was decrease with the addition of neutral density filters among trials (see Table 1). The peak at 840 nm is from two IR lights in 0 and 1 lx trials only. **c** Mean ($\pm$ s.e.m.) light intensities measured before and after trials indicating that light intensity did not differ before and after trials at any of the light intensities used (students *t*-test, $P > 0.05$)

Fig. 3 Total number of prey detections by prey type (white bars = live tethered brine shrimp, grey bars = dead tethered brine shrimp) for **a** *A. stuartgranti* ($n = 6$ fish) and **b** *Tramitichromis* ($n = 6$ fish) at five different light intensities. Maximum number of possible prey detections = 72 for each light intensity

Fig. 4 Three behavioral parameters defining prey detection in *A. stuartgranti* ($n = 6$ fish) and *Tramitichromis* ($n = 6$ fish) at five different light intensities. (**a**, **c**, **e**) Mean ($\pm$ s.e.m.) number of prey detections (maximum 6 live, 6 dead tethered brine shrimp), detection distance, and detection-to-strike velocity, for *A. stuartgranti* feeding on live (—■—) and dead (---□---) prey, and (**b**, **d**, **f**) *Tramitichromis* feeding on live (—●—) and dead (---○---) prey

Fig. 5 Frequency of prey strikes (live and dead prey combined, 12 total prey/trial) during glide or pause phases of swimming at five different light intensities in **a** *A. stuartgranti* ($n = 6$ fish) and **b** *Tramitichromis* ($n = 6$ fish)

Fig. 6 Detection angle for live and dead prey combined (=12 total prey/trial) at five different light intensities for **a** *A. stuartgranti* ($n = 6$ fish) and **b** *Tramitichromis* ($n = 6$ fish). Black lines represent the proportion of prey detections grouped into 20° intervals. Fish snout is at the center and fish is facing 0° (indicated by the grey arrow in the top plot in **a**). The thin line represents the mean angle for all trials. Results for *Tramitichromis* at 0 lx were not included here due to the small number of strikes ($n = 3$ strikes)

Fig. 7 Mean ($\pm$ s.e.m.) prey preference scores (following Taplin 2007) for **a** *A. stuartgranti* ($n = 6$ fish) and **b** *Tramitichromis* ($n = 6$ fish) feeding on six live (white bars) and six dead (gray bars) tethered adult brine shrimp in trials at five different light intensities. Preferences scores were calculated by taking the mean of the rank order in which prey were captured. The dotted line (= 6.5) indicates the mean preference score with no preference for either prey type. Scores <6.5 (below dotted line) indicate a preference. Significantly different preference scores between live and dead prey indicated by an asterisk (*, paired t -test, $P < 0.05$, Table 5)

Table 1 The relationship of measured light intensity (mean lux and PAR, $\pm$ s.e.m. measured immediately after behavioral trials) and predicted depths at which these intensities occur in Lake Malawi. Calculations were based on midday sunlight levels, three light extinction coefficients (0.10 m^{-1} , Patterson et al. 2000; 0.13 m^{-1} , 0.43 m^{-1} , Guildford et al. 2007), and light intensities under natural conditions (Harden Jones 1955, Ali 1959). Light intensities were achieved by varying the height of two fluorescent fixtures (ballasts) and/or covering these fixtures with several neutral density filters

Light Intensity		Light Extinction Coefficient			
Lux	PAR	$\epsilon = 0.10 \text{ m}^{-1}$	$\epsilon = 0.13 \text{ m}^{-1}$	$\epsilon = 0.43 \text{ m}^{-1}$	
Lumen/m ²	$\mu\text{mol photons/m}^2/\text{s}$	Depth (m)	Depth (m)	Depth (m)	Light intensities under natural conditions
800 (800.8 ± 5.4)	11.0 (11.0 ± 0.10)	52	40	12	Very cloudy day
112 (112.4 ± 1.9)	1.5 (1.51 ± 0.03)	72	55	17	Twilight
12 (12.0 ± 0.3)	0.2 (0.16 ± 0.01)	92	71	21	Twilight
1 (1.4 ± 0.1)	0.03 (0.03 ± 0.01)	115	89	27	Full moon/deep twilight
0 (0 ± 0.1)	0 (0.000 ± 0.003)	NA	NA	NA	New moon

Table 2 Determination of GLMM types used to analyze four parameters of feeding behavior at five different light intensities (0-800 lx) in interspecific and intraspecific comparisons

Source	Distribution	Link	Covariance Structure	AICC
4-way GLMMs				
Number of prey strikes	Multinomial	Negative log-log	AR(1)	1,713.0
Detection distance	Normal*	Identity	AR(1)	177.9
Detection-to-strike velocity	Normal*	Identity	AR(1)	-32.4
Swimming phase at prey detection	Binomial	Probit	AR(1)	1932.8
3-way GLMMs				
<i>Aulonocara stuartgranti</i>				
Number of prey strikes	Multinomial	Probit	AR(1)	698.1
Detection distance	Normal*	Identity	AR(1)	44.9
Detection-to-strike velocity	Normal*	Identity	AR(1)	-24.6
Swimming phase at prey detection	Binomial	Probit	AR(1)	1065.9
<i>Tramitichromis</i>				
Number of prey strikes	Multinomial	Negative log-log	AR(1)	955.4
Detection distance	Normal*	Identity	AR(1)	130.6
Detection-to-strike velocity	Normal*	Identity	AR(1)	-7.2
Swimming phase at prey detection	Binomial	Probit	AR(1)	869.1

Note: the table includes information on the error distribution and link function. The first-order auto-regressive process [AR(1)] was used for the covariance structure in all models. The most parsimonious model was selected based on the corrected Akaike information criterion (AICC).

*Data was log10 transformed to achieve normality (normality assessed with Kolmogorov-Smirnov test).

Table 3 Summary of 4-way GLMM statistics for prey detection behavior for two species (*A. stuartgranti*, $n = 6$ fish; *Tramitichromis*, $n = 6$ fish) feeding on two prey types (live, dead) at five light intensities (0-800 lx). Only those factors that are significant for at least one behavioral parameter are listed. See Table 2 for details of GLMMs used

Source	Number of prey detections		Detection distance		Detection-to-strike velocity		Swimming phase at prey detection	
	<i>F</i> (<i>df</i>)	<i>P</i> value	<i>F</i> (<i>df</i>)	<i>P</i> value	<i>F</i> (<i>df</i>)	<i>P</i> value	<i>F</i> (<i>df</i>)	<i>P</i> value
Species (S)		n.s.		n.s.		n.s.		n.s.
Light intensity (L)	14.390 (1,99)	<0.001	9.480 (1,481)	0.002	8.919 (1,511)	0.003	8.276 (1,18)	0.010
Time of day (T)	22.203 (1,99)	<0.001	17.342 (1,512)	<0.001	16.838 (1,513)	<0.001	0.038 (1,129)	0.847
Prey type (P)	4.549 (1,99)	0.035	1.145 (1,503)	0.285	3.876 (1,504)	0.050	3.037 (1,514)	0.082
S × L	8.950 (1,99)	0.004		n.s.		n.s.		n.s.
S × P		n.s.		n.s.		n.s.	4.288 (1,514)	0.039
L × T	14.101 (1,99)	<0.001	10.479 (1,482)	0.001	9.957 (1,512)	0.002	6.044 (1,62)	0.017
S × L × T	10.464 (1,99)	0.002		n.s.		n.s.		n.s.

n.s. = not significant ($P > 0.05$)

P values < 0.05 are shown in bold

Table 4 Summary of 3-way GLMM statistics for prey detection behavior for *A. stuartgranti* ($n = 6$ fish) and *Tramitichromis* ($n = 6$ fish) feeding on two prey types (live, dead) at five different light intensities (0-800 lx). See Table 2 for details of GLMMs used

Model term	<i>Aulonocara stuartgranti</i>		<i>Tramitichromis</i>	
	<i>F</i> (<i>df</i>)	<i>P</i> value	<i>F</i> (<i>df</i>)	<i>P</i> value
Number of prey detections				
Light intensity (L)	0.282 (1,47)	0.598	11.867 (1,47)	0.001
Time of day (T)	11.212 (1,47)	0.002	23.887 (1,47)	<0.001
Prey type (P)	1.649 (1,47)	0.205	0.109 (1,47)	0.743
L $\times$ T	0.293 (1,47)	0.591	12.162 (1,47)	0.001
L $\times$ P	0.845 (1,47)	0.363	0.078 (1,47)	0.780
T $\times$ P	0.968 (1,47)	0.330	0.003 (1,47)	0.956
L $\times$ T $\times$ P	0.675 (1,47)	0.416	0.037 (1,47)	0.847
Detection distance				
Light intensity (L)	2.772 (1,242)	0.097	6.185 (1,217)	0.014
Time of day (T)	26.812 (1,291)	<0.001	5.655 (1,223)	0.018
Prey type (P)	8.220 (1,286)	0.004	0.000 (1,217)	0.986
L $\times$ T	3.408 (1,240)	0.066	6.677 (1,217)	0.010
L $\times$ P	0.002 (1,286)	0.965	4.019 (1,220)	0.046
T $\times$ P	4.604 (1,286)	0.033	0.026 (1,217)	0.872
L $\times$ T $\times$ P	0.004 (1,286)	0.949	4.211 (1,220)	0.041
Detection-to-strike velocity				
Light intensity (L)	3.158 (1,200)	0.077	4.695 (1,223)	0.031
Time of day (T)	16.895 (1,290)	<0.001	8.058 (1,221)	0.005
Prey type (P)	20.107 (1,286)	<0.001	0.012 (1,218)	0.912
L $\times$ T	3.465 (1,198)	0.064	5.495 (1,223)	0.020

L × P	0.385 (1,286)	0.535	2.220 (1,219)	0.138
T × P	14.330 (1,286)	<0.007	0.014 (1,218)	0.907
L × T × P	0.393 (1,286)	0.531	2.300 (1,219)	0.131

Swimming phase at prey detection

Light intensity (L)	0.289 (1,291)	0.592	3.208 (1,223)	0.075
Time of day (T)	1.593 (1,291)	0.208	0.068 (1,223)	0.794
Prey type (P)	0.147 (1,291)	0.701	3.794 (1,223)	0.053
L × T	0.128 (1,291)	0.720	2.873 (1,223)	0.091
L × P	0.220 (1,291)	0.639	0.433 (1,223)	0.511
T × P	0.202 (1,291)	0.653	2.588 (1,223)	0.109
L × T × P	0.211 (1,291)	0.646	0.438 (1,223)	0.509

P values < 0.05 are shown in bold

Table 5 Summary of the paired *t*-tests comparing prey preference scores of live versus dead prey of *A. stuartgranti* and *Tramitichromis* (following Taplin 2007) by light intensity

Light intensity (lx)	<i>T</i> (<i>df</i>)	<i>P</i> value
<i>Aulonocara stuartgranti</i>		
0	-2.853 (5)	0.036
1	-0.618 (5)	0.564
12	-1.395 (5)	0.222
112	-6.102 (5)	0.002
800	-1.892 (5)	0.117
<i>Tramitichromis</i>		
0	-0.797 (5)	0.461
1	-2.396 (5)	0.062
12	-1.379 (5)	0.226
112	-2.441 (5)	0.059
800	-13.647 (5)	<0.001

P values < 0.05 are shown in bold

a **b** **c**

a*Aulonocara stuartgranti* **b***Tramitichromis*

Aulonocara stuartgranti *Tramitichromis*

a*Aulonocara stuartgranti* **b***Tramitichromis*

a*Aulonocara stuartgranti* **b***Tramitichromis*

a **b**