5-2015

Exploring the Planetary Boundary for Chemical Pollution

Miriam L. Diamond

Cynthia A. de Wit

Sverker Molander

Martin Scheringer

Thomas Backhaus

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Miriam L. Diamond, Cynthia A. de Wit, Sverker Molander, Martin Scheringer, Thomas Backhaus, Rainer Lohmann, Rickard Arvidsson, Åke Bergman, Michael Hauschild, Ivan Holoubek, Linn Persson, Noriyuki Suzuki, Marco Vighi, Cornelius Zetzsch. (2015). "Exploring the planetary boundary for chemical pollution." *Environment International*. 78: 8-15. doi: 10.1016/j.envint.2015.02.001
Available at: <http://dx.doi.org/10.1016/j.envint.2015.02.001>

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Authors

Miriam L. Diamond, Cynthia A. de Wit, Sverker Molander, Martin Scheringer, Thomas Backhaus, Rainer Lohmann, Rickard Arvidsson, Åke Bergman, Michael Hauschild, Ivan Holoubek, Linn Persson, Noriyuki Suzuki, Marco Vighi, and Cornelius Zetzsch

Exploring the planetary boundary for chemical pollution

Miriam L. Diamond^{†*}, Cynthia A. de Wit[‡], Sverker Molander,[§] Martin Scheringer,[%] Thomas Backhaus,^{||} Rainer Lohmann,[√] Rickard Arvidsson,[§] Åke Bergman,[⊥] Michael Hauschild,[#] Ivan Holoubek,[¶] Linn Persson,[&] Noriyuki Suzuki,[@] Marco Vighi,[¤] Cornelius Zetzsch^Δ

[†] Department of Earth Sciences, University of Toronto, 22 Russell Street, Toronto, M5S 3B1 Ontario, Canada

[‡] Department of Environmental Science and Analytical Chemistry (ACES), Stockholm University, SE-106 91 Stockholm, Sweden

[§]Environmental Systems Analysis, Department of Energy and Environment, Chalmers University of Technology, SE-412 96 Gothenburg, Sweden

[%]Institute for Chemical and Bioengineering, ETH Zürich, Wolfgang-Pauli-Str. 10, 8093 Zürich, CH-8093, Switzerland, and Leuphana University Lüneburg, D-21335 Lüneburg, Germany

^{||}Department of Biological and Environmental Sciences, University of Gothenburg, Box 100, SE-405 30 Gothenburg, Sweden

[√]Graduate School of Oceanography, University of Rhode Island, South Ferry Road, Narragansett, Rhode Island, 02882, United States

[⊥]Department of Materials and Environmental Chemistry, Stockholm University, SE-106 91 Stockholm, Sweden

[#]Department of Management Engineering, Technical University of Denmark (DTU), Nils Koppels Allé, Building 426 D, DK-2800 Kgs. Lyngby, Denmark

[¶]Research Centre for Toxic Compounds in the Environment (RECETOX), Faculty of Science,
Masaryk University, Kamenice 753/5, 625 00 Brno, Czech Republic

[&]Stockholm Environment Institute, Linnégatan 87D, Box 24218, Stockholm, Sweden

[@]Strategic Risk Management Research Section, Center for Environmental Risk Research,
National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan

[¤]Department of Earth and Environmental Sciences, University of Milano Bicocca, Piazza della
Scienza 1, Milan, 20126 Italy

[^]Forschungsstelle für Atmosphärische Chemie, Dr. Hans-Frisch-Str. 1-3, Universität Bayreuth,
D-954 48 Bayreuth, Germany

ABSTRACT (323 words)

Rockström et al. (2009a, 2009b) have warned that humanity must reduce anthropogenic impacts defined by nine planetary boundaries if “unacceptable global change” is to be avoided. Chemical pollution was identified as one of those boundaries for which continued impacts could erode the resilience of ecosystems and humanity. The central concept of the planetary boundary (or boundaries) for chemical pollution (PBCP or PBCPs) is that the Earth has a finite assimilative capacity for chemical pollution, which includes persistent, as well as readily degradable chemicals released at local to regional scales, which in aggregate threaten ecosystem and human viability. The PBCP allows humanity to explicitly address the increasingly global aspects of chemical pollution throughout a chemical’s life cycle and the need for a global response of internationally coordinated control measures. We submit that sufficient evidence shows stresses on ecosystem and human health at local to global scales, suggesting that conditions are transgressing the safe operating space delimited by a PBCP. As such current local to global pollution control measures are insufficient. However, while the PBCP is an important conceptual step forward, at this point single or multiple PBCPs are challenging to operationalize due to the extremely large number of commercial chemicals or mixtures of chemicals that cause myriad adverse effects to innumerable species and ecosystems, and the complex linkages between emissions, environmental concentrations, exposures and adverse effects. As well, the normative nature of a PBCP presents challenges of negotiating pollution limits amongst societal groups with differing viewpoints. Thus, a combination of approaches is recommended as follows: develop indicators of chemical pollution, for both control and response variables, that will aid in quantifying a PBCP(s) and gauging progress towards reducing chemical pollution, develop new technologies and technical and social approaches to mitigate global chemical

pollution that emphasize a preventative approach, coordinate pollution control and sustainability efforts, and facilitate implementation of multiple (and potentially decentralized) control efforts involving scientists, civil society, government, non-governmental organizations and international bodies.

KEYWORDS: planetary boundary, chemical pollution, chemical emissions, Stockholm Convention, tipping point, global threshold, pollution controls, ecosystem health protection, human health protection, chemical management

1. INTRODUCTION

Rockström et al. (2009a, 2009b) presented nine anthropogenic impacts of global relevance, including climate change, biodiversity loss, anthropogenic changes of the nitrogen and phosphorus cycles, stratospheric ozone depletion, ocean acidification, global freshwater use, changes in land use, atmospheric aerosol loading, and chemical pollution. The authors proposed that humanity may be moving beyond a “safe operating space” as the magnitude of these impacts approach or exceed certain thresholds that represent tipping points of the global system or a natural limit for processes without clear thresholds (so-called “dangerous levels” in the Rockström et al. articles) (Fig. 1). As discussed in detail below, the authors defined a “safe operating space” as those global conditions that allow for continued human development. Rockström et al. (2009a, 2009b) challenged the global scientific community to determine these “non-negotiable” thresholds or natural limits, which are science-based limits of the Earth’s systems, reflecting conditions that are favorable for human life and cultural development, and then to define human-determined boundaries at an appropriate distance from these limits that allow humanity to “avoid unacceptable global change” (Carpenter and Bennett, 2011). A critical

goal of defining the boundaries is to move governance and management away from a piecemeal and sectorial approach, towards an integrated global approach that is necessary to address global phenomena.

For chemical pollution, Rockström et al. (2009a, 2009b) did not define the scope of chemicals considered, natural limits or a planetary boundary, but stated that these remain to be determined. However, they suggested that possible measurable control variables for natural limits could be emissions, concentrations or effects of Persistent Organic Pollutants (POPs), plastics, endocrine disruptors, heavy metals and nuclear wastes. Persson et al. (2013) added to the discussion by suggesting three conditions that must be met simultaneously for chemical pollution to present a global threat. Here we consider a broad range of chemicals including synthetic organic substances and metals, and those intentionally and unintentionally released. We do not consider the nutrients nitrogen and phosphorus that are considered under a separate planetary boundary, or sulfates that can also fall under another planetary boundary (atmospheric aerosol loading).

A large primary literature and numerous reviews document the extent and diversity of chemical pollution and attendant adverse health effects to humans and ecosystems (e.g., UNEP, 2012; AMAP, 2004, 2009; Letcher et al., 2010; WHO and UNEP, 2013; *inter alia*). Indeed, the number of scientific studies providing such evidence fills environmental journals and conference halls. Examples of widespread effects are diminishing populations of wildlife (e.g., Oaks et al., 2004; Tapparo et al., 2012; EFSA, 2013) and increasing burdens of human clinical and

subclinical illness related to environmental toxicants (WHO and UNEP, 2013; Grandjean and Landrigan, 2006; Stillerman et al., 2008). Mounting evidence also indicates that the assessment of individual chemicals is insufficient, as complex mixtures might cause significant toxic effects, even if all individual chemicals are present only at individually non-toxic concentrations, as discussed below. This pattern has been observed repeatedly in a broad range of bioassays at different levels of complexity and for different types of chemicals (see reviews by Kortenkamp et al., 2007, 2009; Kortenkamp, 2008; Backhaus et al., 2010; SCHENIHR et al., 2012). Together, this evidence implies that if emissions of increasing numbers and amounts of chemicals continue at current and anticipated increasing rates (UNEP, 2012), concentrations of such chemicals in many parts of the world, alone or as mixtures, will push the global system beyond the safe operating space. In turn, reaching this point will lead to erosion of vital ecosystems and ecosystem services, and threaten human well-being. Some argue that this point has already been reached (WHO and UNEP, 2013; *inter alia*). Furthermore, the boundary of global chemical pollution cannot be ignored because it is inextricably connected to the other planetary boundaries by the manifold impacts across the life-cycle of chemicals at a global scale, e.g., energy and water use for extraction and manufacturing, land use change that accompanies waste disposal with a potential loss of biodiversity.

This paper explores the definitions and meaning of, and arguments for, a planetary boundary or boundaries for chemical pollution (PBCP). We discuss the many challenges that indicate that defining a boundary or boundaries for chemical pollution is not easily within reach. Our intent here is not to reproduce or re-summarize evidence of widespread adverse effects due to chemical

pollution. Rather, we submit that this evidence points to the need for considering a planetary boundary or more likely *boundaries* for chemical pollution to help humanity remain within the Earth's safe operating space. Thus, the paper closes with recommendations for steps that hopefully will move humanity towards a safe operating space with respect to chemical pollution.

We start the discussion by acknowledging that defining natural limits and a PBCP(s) is challenging for many reasons. In the framework presented by Rockström et al. (2009a, 2009b), defining a PBCP is more difficult than for other planetary boundaries (e.g. for global warming), due to the difficulty of identifying a single or a few measurable control variables. A control variable is defined, according to Rockström et al. (2009a, 2009b), as a measureable parameter that can be related to a specific planetary boundary, e.g., atmospheric CO₂ or temperature for global warming. However, agreeing on one or more control variables for chemical pollution is challenging because chemical pollution is caused by an enormous number of chemicals emitted from innumerable sources and in extremely different amounts in different regions of the world. In the same way, the response variable is difficult to define and measure in a clear-cut way, since chemicals cause a wide variety of adverse effects in a similarly wide variety of species, including humans. The links to the related boundary of biodiversity are evident (Steffen et al. 2015). The critical point is that the Earth's assimilative capacity, or the number and capacities of the sinks capable of degrading or immobilizing anthropogenically-released chemicals, is limited at the global level, even for readily biodegradable chemicals.

2. WHY A PLANETARY BOUNDARY FOR CHEMICAL POLLUTION?

Several policy instruments aimed at controlling chemical pollution have been developed and are in varying degrees of implementation (Table S1). How does a PBCP differ from existing instruments for chemical management and how or why might it be useful rather than redundant? In order to answer these questions we first expand on the concept of planetary boundaries and a “safe operating space” introduced by Rockström et al. (2009a, 2009b) and then move to put a PBCP into the context of existing instruments for chemicals management.

Rockström et al. (2009a, 2009b) identified that several Earth processes and subsystems behave non-linearly, with thresholds that, once crossed, could tip them into new, undesirable states. For these processes, a sharp “tipping point” may exist beyond which the system may transition into a qualitatively different stage, such as much more rapid global warming at CO₂ concentrations above a certain value (Fig. 1a). Examples of Earth systems with such global thresholds or tipping points include the global climate and ocean acidification (e.g., Lenton et al., 2008; Doney et al., 2009; 2014). The planetary boundary can then be set at a level somewhere below the tipping point.

Other processes and subsystems may not have sharp thresholds (Fig. 1b), but their continued erosion or depletion at continental to global scales may cause functional collapse in an increasing number of globally interconnected systems. Here, examples are freshwater use, land use change and loss of biodiversity (May, 1977; Gerten et al., 2013; Baronsky et al., 2012; Brook et al.,

2013). For these, the planetary boundary can be set at a level where the risk of functional collapse is deemed acceptably low. In aggregate, planetary boundaries may thus be defined as a set of critical values for one or several control variables defined by humans to be at a safe distance from such thresholds or dangerous levels (if no threshold is evident) that, if crossed, could lead to abrupt global environmental change. The domain below the boundary can be considered a “safe operating space”.

Figure 1. Illustration of the concept of the planetary boundary (a) for phenomena with a clear tipping point or threshold, where the system moves into a new state, such as CO₂-driven climate change, and (b) without a tipping point, where the system is constantly eroded (modified figure from Rockström et al. (2009a), reprinted with permission of the Stockholm Resilience Center, Stockholm University, Sweden). We suggest that aggregated chemical pollution is illustrated by (b) where there is no clear tipping point.

Although the intention was to define planetary boundaries for systems or processes affecting the Earth at the global scale, Rockström et al. (2009a, 2009b) recognized that many of the identified

boundaries have thresholds that are more evident at local and/or regional scales where disturbance is concentrated or the affected ecosystem is more sensitive. These were identified as “slow processes without known global scale thresholds”. As such, they become a global problem when they occur at many sites at the same time, aggregating to a level that undermines the resilience of ecosystems or that adversely affects human health. In turn, these effects would make it more likely that a threshold with global consequences will be crossed. Examples include biodiversity loss, land use change, global nitrogen and phosphorus biogeochemical cycles, and chemical pollution (Erisman et al., 2013; Hooper et al., 2012; Diaz and Rosenberg, 2008). Slow processes without global thresholds may also exert their effects by affecting other planetary boundaries, for example, chemical pollution of ecosystems linked to biodiversity loss (Voeroesmartty et al., 2010; Lenzen et al., 2012; Steffen et al. 2015).

The distance between the planetary boundary and the threshold or natural limit ideally depends on the uncertainty that surrounds the scientific knowledge about the threshold or natural limit (Fig. 2). If the uncertainty is high, a larger distance between the threshold and the boundary is advisable.

Figure 2. Illustration of where global impacts are located with respect to the safe operating space.

For the planetary boundaries where critical limits were estimated, most of these could be based on one or two specific control variables, such as atmospheric CO₂ concentrations and radiative forcing for climate change. Most of the planetary boundaries that were quantified are preliminary, rough estimates with large uncertainties and for which knowledge gaps were acknowledged.

Although some preliminary boundaries have been proposed, Rockström et al. (2009a, 2009b) pointed out the normative quality of a “safe” distance, as it is based on how societies deal with risk and uncertainty. By normative we mean that decisions on what constitutes a “safe operating space” are societal decisions, supported by scientific evidence. This implies that the diversity of viewpoints held by different societal groups have to be heard in order to come to a decision on what constitutes a safe operating space.

What does the PBCP offer that existing pollution control instruments lack? The planetary boundary concept allows us to explicitly address the *global aspects of chemical pollution*. By recognizing the global nature of chemical pollution, including aggregated local effects or where distance separates emissions from effects, we highlight the need for an integrated global response and acknowledge that pollution control activities of local to national entities alone, are insufficient.

Chemical pollution is a global issue. Several groups of chemicals are distributed around the globe by virtue of their persistence and ability to undergo long-range transport, for example chlorofluorocarbons (CFCs) and persistent organic pollutants (POPs). Others, such as high-production-volume metals that are inherently persistent, are used and emitted globally because of their high production volumes, global trade and widespread use in a broad range of applications. Additionally, the global economy is undergoing chemical “intensification”, as described by the UNEP “Global Chemicals Outlook” analysis (UNEP, 2013). Chemical intensification is due to rapidly increasing global production of chemicals (Wilson and Schwarzman, 2009), to the increasing use of synthetic substances to replace natural materials, and to the use of increasingly complex chemicals in more and more applications. Chemical intensification is predicted to lead to increasing per-capita chemical usage amongst a growing global population (UNEP, 2013).

In addition, chemical product chains, which span the life cycle stages from resource extraction to product manufacturing, use and disposal, are increasing in complexity, often covering several continents and decades of time, and offer new challenges to pollution control. For example, chemical production today can result in future emissions, particularly for chemicals in

infrastructure and goods with long lifetimes. Brunner and Rechberger (2001) have estimated that whereas ~10% of all chemical stocks is contained in waste deposits from primary production and ~10% is contained in land filled waste, ~80% is contained in in-use and “hibernating” stocks. Most documentation of uncontrolled releases concern the two former sources (i.e., 20%) but not the 80% (e.g., Brunner and Rechberger, 2001; Weber et al., 2013; *inter alia*). Examples of the “20%” include long-term emissions from tailings, waste rock piles, nuclear waste repositories, abandoned industrial sites, and numerous landfills in developing countries (Turk et al., 2007; Torres et al., 2013; Weber et al., 2011). One example of long-term emissions from an in-use chemical stock is that of polychlorinated biphenyls (PCBs, listed as a POP under the Stockholm Convention) from equipment that was still in use in Canada in 2006 despite the ban on PCB production nearly 40 years ago (Diamond et al., 2010; Csiszar et al., 2013). Another example is that of CFCs contained in blown building insulation that is subject to uncontrolled releases as the generation of buildings using that foam undergoes renovation or destruction over the next 30 years (Brunner and Rechberger, 2001)

Similar application patterns of chemical technologies and similar uses of chemical products in almost all regions of the world result in widespread chemical releases. Chemical manufacturing and industrial usage are rapidly shifting from Western industrialized countries to developing countries and countries with economies in transition, including BRICS countries (Brazil, Russia, and especially India and China, and most recently South Africa) (UNEP, 2013). New and increasing resource extraction and chemical manufacturing, usage and waste disposal are leading to increased chemical pollution, particularly in jurisdictions with insufficient control mechanisms (Schmidt, 2006; Gottesfeld and Cherry, 2011). Short-lived chemicals are also being released in

many regions at rates that exceed degradation rates and hence environmental assimilative capacities. Examples of such chemicals include pharmaceuticals, high production volume plastics and plasticizers such as bisphenol A and di-ester phthalates, and “D4” and “D5” siloxanes (e.g., WHO and UNEP, 2013; Kolpin et al., 2002; Rosi-Marshall et al., 2013; Peck and Hornbuckle, 2004; Fromme et al., 2002; Fries and Mihajlovic, 2011; Wang et al., 2013).

As pointed out above, the global nature of chemical pollution demands a global response of internationally coordinated control measures, in addition to multiple local, regional and national efforts covering different groups of substances, which are disconnected in time and space. One example of a global governance instrument is the Stockholm Convention on Persistent Organic Pollutants (POPs), which seeks elimination at best, or more broadly, the sound management, of a set of POPs agreed upon through international negotiations (Stockholm Convention, 2008). While achieving many successes (Stockholm Convention, 2012), the Convention is limited to a small number of chemicals or chemical classes (currently 22 are listed, with four more under review), includes numerous exemptions, and has no instrument for sanctions to ensure national implementation. This is not a shortcoming of the Convention because the intention of the Convention is not to address the totality of chemical pollution. As such, the Stockholm Convention is not adequate for challenge presented by developing a PBCP. Similarly, the Montreal Protocol is limited to substances that deplete the stratospheric ozone layer (UNEP 2010-2011) and the Minamata Convention is limited to mercury (UNEP 2015). The Convention on Long-range Transboundary Air Pollution, under the aegis of the United Nations Economic Commission for Europe and to which there are 51 parties, addresses a range of chemical pollutants including metals and POPs (UNECE 2004).

285
286 Another example of a global governance tool is the United Nations Framework Convention on
287 Climate Change where global negotiations and agreements have led to reduction goals for
288 greenhouse gases that are intended to be implemented at national levels (UNFCCC, 2013).
289 International climate negotiations have seen the emergence of control instruments of largely two
290 types. The first is an absolute limit for total CO₂-equivalent emissions (a “cap”) to assure that
291 total global emissions are on target to prevent the global atmospheric CO₂ concentration
292 exceeding an agreed-upon boundary. The second type of control scheme links emissions to
293 activity or intensity such as CO₂-equivalent emissions per unit of electricity generated or per
294 kilometre driven, or to an economic cost resulting in reductions of CO₂-equivalent
295 emissions/capita (Azar and Rodhe, 1997; Ellerman and Sue Wing, 2003). These intensity or
296 efficiency-based emission controls acknowledge the need to reduce greenhouse gas emissions
297 but cannot ensure that global emissions are within the global safe operating space because of
298 population and economic growth that increase the demand for energy services, most of which are
299 based on fossil fuels (IEA, 2014).
300
301 Implicit in the concept of a safe operating space for CO₂ and other greenhouse gases, ocean
302 acidification, nitrogen and phosphorus cycles, and “chemical pollution”, is that there is a finite
303 global assimilative capacity. Here we define assimilative capacity as the ability of an ecosystem
304 to render substances harmless, i.e. avoiding adverse effects. By seeing the problem in this light,
305 it leads us towards exploring the need for a globally coordinated cap for emissions, rather than
306 jurisdiction-specific, intensity-based controls, which may be sufficient in some circumstances but
307 fail to account for cumulative, global effects.

308

309 **3. CHALLENGES OF DEFINING A PLANETARY BOUNDARY FOR** 310 **CHEMICAL POLLUTION**

311 Moving the idea of a PB beyond a conceptual model requires that the impact of anthropogenic
312 stressor(s) on all ecosystems can be described and quantified as a function of a measurable
313 control variable(s) that is (are) related to a measurable response variable(s). For a PBCP, the
314 ultimate effect or response variable (Fig. 1) subject to control is widespread adverse impact(s) to
315 ecological and/or human health caused by exposure to (a) substance(s). Exposure can be
316 identified as the critical control variable since it is the necessary prerequisite for any kind of
317 chemically induced effect or response we want to safeguard against. Ideally, chemical exposure
318 can be used to define a threshold(s) or natural limit(s) that, in turn, can be translated into a global
319 boundary (boundaries) and a safe operating space. As noted above, the boundary (boundaries) is
320 (are) established by humans and is (are) a product of societal demands, needs, value judgments
321 and negotiations. The control variable(s) must also be amenable to translation into possible
322 mitigation or control activities, which in this case would reduce exposure and thus, would
323 maintain human and ecosystem health within the safe operating space, the latter reflected in
324 maintained biodiversity, ecosystem functionality and human health.

325

326 Challenges arise at all stages in the definition process that starts with a control variable(s) and
327 ends with “actionable” activities. First, operationalizing “exposure” as the control variable is
328 difficult because of the high and poorly defined number of chemicals that fall under the umbrella
329 of “chemical pollution”. More than 100 000 substances are in commerce (Egeghy et al., 2012),

including pesticides, biocides and pharmaceuticals, industrial chemicals, building materials and substances in personal care products and cosmetics (e.g., Howard and Muir, 2010, 2011; ECHA, 2013) and very few of them have undergone adequate risk assessment for adverse effects. A recent screening of 95 000 chemicals for persistence (P), bioaccumulation (B) and toxicity (T) properties (REACH criteria) identified 3% or approximately 3000 chemicals as potential PBT chemicals (uncertainty range of 153-12 500 chemicals) (Strempel et al., 2012). Similarly, 93 000 chemicals were screened for P, B and long range transport potential according to the Stockholm Convention criteria, plus T (REACH criteria) resulting in the identification of 510 potential POPs (uncertainty range of 190-1 200 chemicals) (Scheringer et al., 2012). Unintentionally produced substances, such as the combustion by-products polycyclic aromatic hydrocarbons (PAH) and polychlorinated and polybrominated dibenzo-*p*-dioxins and furans (PCDD/F and PBDDs/Fs), are emitted as a consequence of human activity and many emitted chemicals are transformed to a multitude of other chemicals by biological and physical-chemical processes. Whereas some limits have been placed on a few selected chemicals that are highly persistent, bioaccumulative and toxic such as PCDD/F, those with intermediate PBT properties have received insufficient attention (Muir and Howard, 2006; Howard and Muir, 2010; Scheringer et al., 2012). In addition, an enormous number of organisms in a diversity of ecosystems are exposed to chemical pollution (which is invariably a complex chemical mixture) and they will respond in myriad ways. Moreover, chemicals have specific modes of actions and can show very different toxicological potencies. Humans take a specific place among affected organisms. Any approach to establishing a PBCP(s) must include impacts on human health, even if this is in contrast to the framework of Rockström et al. (2009a, 2009b) or which the objects of protection

are biogeochemical systems and ecosystems, e.g., the climate system, the ozone layer, and freshwater.

Second, we acknowledge that boundaries for chemical pollution have been developed at a global scale for selected POPs and mercury, and at local and regional scales for chemicals in foods, water and air (Table S1). However, only a few of these boundaries account for exposure to multiple chemicals simultaneously that can act in an additive fashion. Moving beyond a chemical-by-chemical approach to acknowledge mixture effects is of growing importance if limits are to be protective (e.g., Kortenkamp, 2007; Kortenkamp et al., 2007; Backhaus et al., 2010; Meek et al., 2011; SCHENIHR et al., 2012). An increasing body of evidence suggests that, *de facto*, the existing boundaries are not sufficiently protective for endocrine disrupting chemicals that can cause transgenerational effects (e.g., Baccarelli and Bollati, 2009; Bollati and Baccarelli, 2010; Bouwman et al., 2012; Mani et al., 2012; WHO and UNEP, 2013; *inter alia*). This is not surprising since accepted and validated methods for identifying and testing endocrine disrupting chemicals, particularly after exposure during critical early life stages, are generally lacking or have not yet been implemented in chemicals risk assessment (WHO and UNEP, 2013; *inter alia*).

Third, connecting exposure as the control variable to an “actionable” activity (such as controlling emissions) is difficult because of the diversity of fate and transformation processes at play between an initial emission of a chemical or a chemical mixture and the concentration(s)

373 resulting in exposure and then an adverse effect. Establishing the release-fate-concentration-
374 effect linkage is necessary for other planetary boundaries such as CO₂, stratospheric ozone,
375 phosphorus and nitrogen cycles. Establishing this linkage for chemical pollution is also
376 necessary but it is more challenging because of the large number of chemicals of varying
377 persistence and toxicity that are captured by this boundary.

378
379 Finally, in addition to the scientific challenges of defining a boundary(s), it must be remembered
380 that most of the world's countries do not have the capacity or resources to measure a control
381 variable such as exposure and to implement effective controls such as those listed in Table S1
382 (e.g., Klanova et al., 2009; Adu-Kumi et al., 2012). Furthermore, as noted above, a boundary(s)
383 is normative and as such, a diversity of viewpoints will be held on what constitutes an
384 "acceptable" level of pollution.

385
386 The combination of numerous substances with different use and emission patterns, affecting a
387 multitude of different endpoints in a plethora of exposed species in the vastly different
388 ecosystems of the world, plus consideration of human health, makes the derivation of a single
389 quantitative PBCP or multiple PBCPs a daunting, if not impossible task. However, the situation
390 of increasing chemical production, emissions and adverse effects cannot be allowed to continue
391 unabated. Thus, we believe that the concept of a planetary boundary or boundaries for chemical
392 pollution is a useful framework for global action, but that it needs to be modified to account for
393 these complexities and challenges.

394

395 4. STEPS TOWARD GLOBAL CHEMICALS MANAGEMENT

396

397 Although it may not be possible to establish a single or even multiple PBCP(s) at this time, an
398 increasing body of evidence strongly suggests that we need more effective global chemicals
399 management. What has been accomplished in global chemicals management? Global
400 cooperation amongst nations has, amongst others, resulted in the Stockholm Convention on
401 POPs, the Montreal Protocol on CFCs, the Basel Convention on Control of Transboundary
402 Movements of Hazardous Wastes, and the Rotterdam Convention on Prior Informed Consent
403 Procedure for Certain Hazardous Chemicals and Pesticides in International Trade. These
404 Multilateral Environmental Agreements have come together under the aegis of UNEP. The
405 Stockholm and Montreal agreements strive towards zero-emissions of the listed chemicals. In
406 January 2013, UNEP brokered the Minamata Convention on mercury, the language of which has
407 gained support from 94 signatory countries (UNEP, 2015). The Minamata Convention specifies
408 the banning of production, export and import of a range of mercury-containing products, calls for
409 the drafting of strategies to limit the use of mercury in artisanal and small-scale gold mining, and
410 aims to work towards minimizing mercury emissions from combustion sources such as
411 conventional fossil fuel power plants and cement factories. Like the Stockholm Convention, the
412 Minamata Convention includes the provision to develop a compliance mechanism that will be
413 established through negotiation after the official signing of the Convention.

414

415 These five agreements address priority chemical pollutants at the global scale, reflect the insight
416 that global dilution is not the solution to local or global pollution, and that environmental
417 safeguards are the right of all countries. Well over 100 countries have adopted them (except for
418 the most recent Minamata Convention), which in itself is a great accomplishment. However,
419 these agreements have limitations due to numerous official exemptions and unofficial
420 “loopholes”, they cover only a limited number of chemicals, implementation costs are largely
421 left to individual countries of which many lack such capacity, and sanctions cannot be levied for
422 a lack of compliance. As such, these agreements are not adequate to address the totality of
423 chemical pollution (which was never their intent). Importantly, the fact that these agreements
424 have been enacted is a reflection that humanity has come close to or crossed boundaries for these
425 chemicals. A PBCP provides an overarching conceptual basis to characterize the achievements
426 of these agreements and to accommodate additional necessary controls.

427
428 For chemicals listed by the Stockholm and Minamata Conventions and the Montreal Protocol,
429 the planetary boundary is set at a *de minimus* level (ideally zero emissions but exemptions
430 preclude this). In addition to the zero emissions boundary, several other types of boundaries
431 have been defined during the past decades under many jurisdiction-specific regulations and
432 initiatives spanning local to national scales. As summarized in Table S1, the initiatives, which
433 come from international agencies, Europe, Japan, North America, China, India and Nigeria,
434 include limits to levels of pesticides in groundwater and surface water, levels of priority
435 pollutants in surface waters, and acceptable daily intakes (ADIs) for a wide range of food
436 contaminants. However, as noted above, not all of these agencies are able to monitor for, and
437 enforce compliance.

438

439 Another major global initiative is the Strategic Approach to International Chemicals
440 Management (SAICM), which is also under the aegis of UNEP. The ultimate goal of SAICM is
441 to facilitate activities to ensure that "...chemicals will be produced and used in ways that
442 minimize significant adverse impacts on the environment and human health" (SAICM, 2006).
443 The role of SAICM is advisory by acting as a source of information to governmental and extra-
444 governmental bodies regarding safe chemical management and funding projects to fulfill the aim
445 of the initiative. SAICM is a non-binding agreement with broad participation of countries and
446 other stakeholders such as the chemical industry. In comparison to the five chemical
447 agreements, SAICM is much broader in scope by addressing all agricultural and industrial
448 chemicals from cradle to grave, aiming at overall sound chemicals management. However,
449 SAICM does not have a compliance mechanism.

450

451 To move towards a truly global approach encompassing the aggregated impacts from all
452 anthropogenic chemical pollution, we need to learn from experience and build on successes (and
453 failures). What are the key lessons learned? One lesson learned is that implementation of
454 stringent controls by specific jurisdictions has led to improved local conditions in those
455 jurisdictions. However, increased global trade and the fluidity of global finance have moved
456 more chemical and goods production and waste disposal to locations without stringent controls
457 (e.g., Skelton et al., 2011; Breivik et al., 2011; Sindiku et al., 2014). Thus, one intention of a
458 global boundary is avoiding "pollution free" jurisdictions at the expense of creating "pollution
459 havens" in developing nations (e.g. Gottesfeld, 2013). Examples of developed nations achieving
460 their pollution control goals by shipping waste and waste products to developing nations have

been described elsewhere (Schmidt, 2006; Breivik et al., 2011, 2014; Gioia et al., 2011; Abdullah et al., 2013).

A second lesson learned is that despite the challenges, as scientists we need to avoid calling for more scientific certainty before action is taken as this delays adoption of control measures, which in this case translates to measures that will help stem widespread chemical pollution. Gee and others (Gee, 2006; Gee et al., 2013; Harremoës et al., 2001) have documented examples of where the call for more research to improve risk assessments of chemicals often led to delays in action of up to several decades although early warnings of adverse effects were already apparent (e.g. tobacco smoking and asbestos). Persson et al. (2013) provide a persuasive argument in this regard.

As a result of these considerations, we submit that the PBCP is a useful aspirational framework that allows natural and social scientists, policy makers, industry and civil society to visualize the idea of a safe operating space, see the limited assimilative capacity of the Earth, recognize chemical pollution at a global scale, and see the inadequacy of current control measures to deal with the totality of global chemical pollution. Having said that, we recognize that defining a single or multiple quantitative PBCP(s), or even a single approach for its definition, is not now within reach. Rather, we recommend advancing in multiple directions that involve globally coordinated action in scientific, technical and political domains (e.g., Conklin, 2005; Horn and Weber, 2007). For the scientific domain we propose the following:

1. Explore advancing the concept of, and methods for quantifying a PBCP(s). We advocate making stepwise progress using a few well-known chemicals such as POPs, intermediate PBT chemicals (demonstrated toxicity but not highly persistent), and a few high production volume chemicals with demonstrated toxicity.
2. Continue to identify and develop indicators of global chemical pollution, initially based on proxies for chemical exposure and potency. Information on indicator status should then be used to gauge progress towards staying within the safe operating space for chemical pollution. Useful information to guide this task can be taken from the Drivers, Pressures, States, Impacts, Responses (DPSIR) approach (OECD, 1991; Harremoës, 1998), and suggestions of how this could be accomplished are given in the Supporting information. This proposal builds on the global monitoring networks that have achieved considerable success such as those under the Stockholm Convention (e.g., the Global Atmospheric Passive Sampling network or GAPS (Gawor et al., 2014) and Human milk survey (UNEP et al., 2013)).
3. Conduct research into new technologies and methods that will aid in implementing the goals of the six global chemical agreements (Montreal Protocol; Stockholm, Minamata, Rotterdam, Basel and UNECE LRTAP Conventions) and in lowering production and emissions of non-POP priority chemicals. This research includes methods for identifying and characterizing stocks of chemicals scheduled for elimination, developing technologies for efficient and effective destruction of stockpiles, research into societal and cultural considerations that will maximize the likelihood of policy implementation, etc.
4. Connect activities aimed at chemical pollution control in the context of PBCP to efforts aimed at moving towards sustainable resource use. This should include investigating ways to

chemically “de-intensify” economies, to use “green chemistry” substitutes and non-chemical solutions, and to implement social solutions aimed at reducing resource consumption.

Efforts are underway in this regard, such as the U.S. EPA’s Design for the Environment Program (U.S.EPA, 2014) and the GreenScreen® for Safer Chemicals (Clean Production Action, 2015). These two issues, PBCP and sustainable resource use, are intertwined such that chemical pollution is a manifestation of unsustainable and inefficient resource use.

Thus, efforts directed towards achieving both goals would benefit from coordinated action.

Progressing towards a PBCP(s) will require scientific, political, social and economic strategies.

In the political domain, it will be important to raise more awareness for chemical pollution problems in all parts of the world, and to aid individual countries in implementing existing local and regional boundaries and international agreements. The shift of chemical production from OECD countries primarily to the BRICS countries needs to be complemented by a process that helps to develop chemical regulation and enforcement in these regions to a level comparable or better than that of OECD countries.

To address these needs, organizations at the global level such as WHO and UNEP can be drivers for effective exchange and collaboration amongst the public, environmental NGOs, industry and national government institutions to enable significant pollution control. Civil society and local jurisdictions also have and continue to implement effective pollution controls using a variety of tools. Examples here include the activities of the International POPs Elimination Network

(IPEN), the Pesticides Action Network (PAN), and C40 Cities for “Global Leadership on Climate Change” (C40 Cities, 2013).

In closing, 50 years ago Rachel Carson pointed out for the first time that the extensive use of pesticides is dangerous not only to wildlife, but also to humans. This is still an ongoing concern, emphasized by the recent finding that neonicotinoid pesticides are contributing to the massive collapse of bee populations (Tapparo et al., 2012; Henry et al., 2012; Whitehorn et al., 2012). Now we need to go beyond Rachel Carson’s clarion call about pesticides. Today’s phenomenon of locally to globally distributed chemicals that are causing adverse effects, demands that a wide range of chemical products and uses be restrained and many chemicals in commerce need to be used with much more prudence and precaution. It is time to harness the knowledge, capacity and commitment held by many to see Rachel Carson’s vision moved to a truly global scale.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge financial support by the Swedish Research Council FORMAS and the International Panel on Chemical Pollution, which funded a workshop on this topic.

REFERENCES

- 550 Abdullah, H.M., Mahboob, M.G., Banu, M.R., Seker, D.Z., 2013. Monitoring the drastic growth
551 of ship breaking yards in Sitakunda: a threat to the coastal environment of Bangladesh.
552 Environmental Monitoring & Assessment 185, 3839-3851.
- 553 Adu-Kumi, S., Kares, R., Literak, J., Boruvkova, J., Yeboah, P.O., Carboo, D., Akoto, O.,
554 Darko, G., Osae, S., Klanova, J., 2012. Levels and seasonal variations of organochlorine
555 pesticides in urban and rural background air of southern Ghana. Environmental Science
556 & Pollution Research 19, 1963-1970.
- 557 AMAP (Arctic Monitoring and Assessment Programme), 2004. AMAP Assessment 2002:
558 Persistent Organic Pollutants in the Arctic. Arctic Monitoring and Assessment
559 Programme (AMAP), Oslo, Norway, p. 310.
- 560 AMAP (Arctic Monitoring and Assessment Programme), 2009. AMAP Assessment 2009:
561 Human Health in the Arctic. Arctic Monitoring and Assessment Programme, Oslo,
562 Norway, p. 254.
- 563 Azar, C., Rodhe, H., 1997. Targets for Stabilization of Atmospheric CO₂. Science 276, 1818-
564 1819.
- 565 Baccarelli, A., Bollati, V., 2009. Epigenetics and environmental chemicals. Current Opinion in
566 Pediatrics 21, 243-251.
- 567 Backhaus, T., Blanck, H., Faust, M., 2010. Hazard and Risk Assessment of Chemical Mixtures
568 under REACH–State of the Art, Gaps and Options for Improvement. Swedish Chemicals
569 Agency, Sundbyberg, Sweden.
- 570 Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H., Fortelius, M., Getz,
571 W.M., Harte, J., Hastings, A., Marquet, P.A., Martinez, N.D., Mooers, A., Roopnarine,
572 P., Vermeij, G., Williams, J.W., Gillespie, R., Kitzes, J., Marshall, C., Matzke, N.,

- 573 Mindell, D.P., Revilla, E., Smith, A.B., 2012. Approaching a state shift in Earth's
574 biosphere. *Nature* 486, 52-58.
- 575 Bollati, V., Baccarelli, A., 2010. Environmental epigenetics. *Heredity* 105, 105-112.
- 576 Bouwman, H., Kylin, H., Yive, N., Tatayah, V., Loken, K., Skaare, J.U., Polder, A., 2012. First
577 report of chlorinated and brominated hydrocarbon pollutants in marine bird eggs from an
578 oceanic Indian Ocean island. *Environmental Research* 118, 53-64.
- 579 Breivik, K., Gioia, R., Chakraborty, P., Zhang, G., Jones, K.C., 2011. Are reductions in
580 industrial organic contaminants emissions in rich countries achieved partly by export of
581 toxic wastes? *Environmental Science & Technology* 45, 9154-9160.
- 582 Breivik, K., Armitage, J.M., Wania, F., Jones, K.C., 2014. Tracking the Global Generation and
583 Exports of e-Waste. Do Existing Estimates Add up? *Environmental Science &*
584 *Technology* 48, 8735-8743.
- 585 Brook, B.W., Ellis, E.C., Perring, M.P., Mackay, A.W., Blomqvist, L., 2013. Does the terrestrial
586 biosphere have planetary tipping points? *Trends in Ecology & Evolution* 28, 396-401.
- 587 Brunner, P., Rechberger, H.H., 2001. Anthropogenic metabolism and environmental legacies.
588 Ed: Munn, T. *Encyclopedia of Global Environmental Change*. Wiley, Chichester, UK.
589 pp. 54-72
- 590 C40 Cities, 2013. C40 Cities: Climate Leadership Group. www.c40.org.
- 591 Carpenter, S.R., Bennett, E.M., 2011. Reconsideration of the planetary boundary for phosphorus.
592 *Environmental Research Letters* 6, DOI: 10.1088/1748-9326/6/1/014009
- 593 Clean Production Action. 2015. GreenScreen® for Safer Chemicals.
594 <http://www.greenscreenchemicals.org/>

- 595 Conklin, J., 2005. Dialogue Mapping: Building Shared Understanding of Wicked Problems.
596 Wiley, West Sussex, UK. 266 pp.
- 597 Csiszar, S.A., Daggupaty, S.M., Diamond, M.L., 2013. SO-MUM: a coupled atmospheric
598 transport and multimedia model used to predict intraurban-scale PCB and PBDE
599 emissions and fate. *Environmental Science & Technology* 47, 436-445.
- 600 Diamond, M.L., Melymuk, L., Csiszar, S.A., Robson, M., 2010. Estimation of PCB stocks,
601 emissions, and urban fate: will our policies reduce concentrations and exposure?
602 *Environmental Science & Technology* 44, 2777-2783.
- 603 Diaz, R.J., Rosenberg, R., 2008. Spreading dead zones and consequences for marine ecosystems.
604 *Science* 321, 926-929.
- 605 Doney, S.C., Fabry, V.J., Feely, R.A., Kleypas, J.A., 2009. Ocean acidification: the other CO₂
606 problem. *Annual Review of Marine Science* 1, 169-192.
- 607 Doney, S.C., Bopp, L., Long, M.C., 2014. Historical and future trends in ocean climate and
608 biogeochemistry. *Oceanography* 27, 108-119.
- 609 ECA (European Chemicals Agency), 2013. Pre-registered substances.
610 <http://echa.europa.eu/en/information-on-chemicals/pre-registered-substances>. Accessed
611 May 6, 2013.
- 612 EFSA (European Food Safety Authority), 2013. EFSA identifies risks to bees from
613 neonicotinoids. European Food Safety Authority.
614 <http://www.efsa.europa.eu/en/press/news/130116.htm>
- 615 Egeghy, P.P., Judson, R., Gangwal, S., Mosher, S., Smith, D., Vail, J., Hubal, E.A.C., 2012. The
616 exposure data landscape for manufactured chemicals. *Science of the Total Environment*
617 414, 159-166.

- 618 Ellerman, A.D., Sue Wing, I., 2003. Absolute vs. intensity-based emission caps. *Global Change*
619 *Science and Policy*, MIT. MIT, Cambridge MA, 11 pp.
- 620 Erisman, J.W., Galloway, J.N., Seitzinger, S., Bleeker, A., Dise, N.B., Petrescu, A.M.R., Leach,
621 A.M., de Vries, W., 2013. Consequences of human modification of the global nitrogen
622 cycle. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368.
- 623 Fries, E., Mihajlovic, I., 2011. Pollution of soils with organophosphorus flame retardants and
624 plasticizers. *Journal of Environmental Monitoring* 13, 2692-2694.
- 625 Fromme, H., Kuchler, T., Otto, T., Pilz, K., Muller, J., Wenzel, A., 2002. Occurrence of
626 phthalates and bisphenol A and F in the environment. *Water Research* 36, 1429-1438.
- 627 Gawor, A., Shunthirasingham, C., Hayward, S.J., Lei, Y.D., Gouin, T., Mmereki, B.T.,
628 Masamba, W., Ruepert, C., Castillo, L.E., Shoeib, M., Lee, S.C., Harner, T., Wania, F.,
629 2014. Neutral polyfluoroalkyl substances in the global atmosphere. *Environmental*
630 *Science & Process Impacts* 16, 404-413.
- 631 Gee, D., 2006. Late lessons from early warnings: Toward realism and precaution with endocrine-
632 disrupting substances. *Environmental Health Perspectives* 114, 152-160.
- 633 Gee, D., Grandjean, P., Hansen, S.F., Hove, S.V.D., MacGarvin, M., Martin, J., Nielsen, G.,
634 Quist, D., Stanners, D., 2013. Late Lessons from Early Warnings: Science, Precaution,
635 Innovation. EEA (European Environment Agency), Luxembourg. 760 pp.
- 636 Gerten, D., Hoff, H., Rockström, J., Jaegermeyr, J., Kummu, M., Pastor, A.V., 2013. Towards a
637 revised planetary boundary for consumptive freshwater use: role of environmental flow
638 requirements. *Current Opinion in Environmental Sustainability* 5, 551-558.

- 639 Gioia, R., Eckhardt, S., Breivik, K., Jaward, F.M., Prieto, A., Nizzetto, L., Jones, K.C., 2011.
640 Evidence for major emissions of PCBs in the West African region. *Environmental*
641 *Science & Technology* 45, 1349-1355.
- 642 Gottesfeld, P., 2013. The West's toxic hypocrisy over lead paint. *New Scientist* 218, 26-27.
- 643 Gottesfeld, P., Cherry, C.R., 2011. Lead emissions from solar photovoltaic energy systems in
644 China and India. *Energy Policy* 39, 4939-4946.
- 645 Grandjean, P., Landrigan, P.J., 2006. Developmental neurotoxicity of industrial chemicals.
646 *Lancet* 368, 2167-2178.
- 647 Harremoës, P., 1998. The challenge of managing water and material balances in relation to
648 eutrophication. *Water Science & Technology* 37, 9-17.
- 649 Harremoës, P., Gee, D., MacGarvin, M., Stirling, A., Keys, J., Wynne, B., Vaz, S.G., 2001. Late
650 Lessons from Early Warnings: The Precautionary Principle 1896-2000. EEA (European
651 Environment Agency), Copenhagen. 211 pp.
- 652 Henry, M., Beguin, M., Requier, F., Rollin, O., Odoux, J.F., Aupinel, P., Aptel, J., Tchamitchian,
653 S., Decourtye, A., 2012. A common pesticide decreases foraging success and survival in
654 honey bees. *Science* 336, 348-350.
- 655 Hooper, D.U., Adair, E.C., Cardinale, B.J., Byrnes, J.E.K., Hungate, B.A., Matulich, K.L.,
656 Gonzalez, A., Duffy, J.E., Gamfeldt, L., O'Connor, M.I., 2012. A global synthesis reveals
657 biodiversity loss as a major driver of ecosystem change. *Nature* 486, 105-U129.
- 658 Horn, R.E., Weber, R.P., 2007. New tools for resolving wicked problems: mess mapping and
659 resolution mapping processes. MacroVU, Inc and Stanford University; Strategy Kinetics,
660 LLC, Stanford, CA; Watertown, MA. 31 pp.

- 661 Howard, P.H., Muir, D.C.G., 2010. Identifying new persistent and bioaccumulative organics
662 among chemicals in commerce. *Environmental Science & Technology* 44, 2277-2285.
- 663 Howard, P.H., Muir, D.C.G., 2011. Identifying new persistent and bioaccumulative organics
664 among chemicals in commerce II: pharmaceuticals. *Environmental Science &*
665 *Technology* 45, 6938-6946.
- 666 IEA (International Energy Agency), 2014. *World Energy Outlook 2014*. OECD (Organization
667 for Economic Cooperation and Development), Paris. 748 pp.
- 668 Klanova, J., Cupr, P., Holoubek, I., Boruvkova, J., Pribylova, P., Kares, R., Tomsej, T., Ocelka,
669 T., 2009. Monitoring of persistent organic pollutants in Africa. Part 1: passive air
670 sampling across the continent in 2008. *Journal of Environmental Monitoring* 11, 1952-
671 1963.
- 672 Kolpin, D.W., Furlong, E.T., Meyer, M.T., Thurman, E.M., Zaugg, S.D., Barber, L.B., Buxton,
673 H.T., 2002. Pharmaceuticals, hormones, and other organic wastewater contaminants in
674 US streams, 1999-2000: A national reconnaissance. *Environmental Science &*
675 *Technology* 36, 1202-1211.
- 676 Kortenkamp, A., 2007. Ten years of mixing cocktails: a review of combination effects of
677 endocrine-disrupting chemicals. *Environmental Health Perspectives* 115, 98-105.
- 678 Kortenkamp, A., 2008. Low dose mixture effects of endocrine disruptors: implications for risk
679 assessment and epidemiology. *International Journal of Andrology* 31, 233-237.
- 680 Kortenkamp, A., Faust, M., Scholze, M., Backhaus, T., 2007. Low-level exposure to multiple
681 chemicals: reason for human health concerns? *Environmental Health Perspectives* 115,
682 106-114.

- 683 Kortenkamp, A., Backhaus, T., Faust, M., 2009. State of the Art Report on Mixture Toxicity.
684 Commission of the European Union (Directorate General for the Environment), London.
685 391 pp.
- 686 Lenton, T.M., Held, H., Kriegler, E., Hall, J.W., Lucht, W., Rahmstorf, S., Schellnhuber, H.J.,
687 2008. Tipping elements in the Earth's climate system. Proceedings of the National
688 Academy of Sciences 105, 1786-1793.
- 689 Lenzen, M., Moran, D., Kanemoto, K., Foran, B., Lobefaro, L., Geschke, A., 2012. International
690 trade drives biodiversity threats in developing nations. Nature 486, 109-112.
- 691 Letcher, R.J., Bustnes, J.O., Dietz, R., Jenssen, B.M., Jorgensen, E.H., Sonne, C., Verreault, J.,
692 Vijayan, M.M., Gabrielsen, G.W., 2010. Exposure and effects assessment of persistent
693 organohalogen contaminants in arctic wildlife and fish. Science of the Total Environment
694 408, 2995-3043.
- 695 Mani, M., Markandya, A., Sagar, A., Strukova, E., 2012. An Analysis of Physical and Monetary
696 Losses of Environmental Health and Natural Resources in India. The World Bank, South
697 Asia Region, Disaster Risk Management and Climate Change, Washington DC. 41 pp.
- 698 May, R.M., 1977. Thresholds and breakpoints in ecosystems with a multiplicity of stable states.
699 Nature 269, 471-477.
- 700 Meek, M.E., Boobis, A.R., Crofton, K.M., Heinemeyer, G., Van Raaij, M., Vickers, C., 2011.
701 Risk Assessment of Combined Exposure to Multiple Chemicals: A WHO/IPCS
702 framework. Regulatory Toxicology & Pharmacology 60, S1-S14.
- 703 Muir, D.C.G., Howard, P.H., 2006. Are there other persistent organic pollutants? A challenge for
704 environmental chemists. Environmental Science & Technology 40, 7157-7166.

- 705 OECD (Organization for Economic Cooperation and Development), 1991. Core Set of Indicators
706 for Environmental Performance Reviews, a Synthesis Report by the Group on the State of
707 the Environment. OECD, Paris.
- 708 Oaks, J.L., Gilbert, M., Virani, M.Z., Watson, R.T., Meteyer, C.U., Rideout, B.A., Shivaprasad,
709 H.L., Ahmed, S., Chaudhry, M.J.I., Arshad, M., Mahmood, S., Ali, A., Khan, A.A., 2004.
710 Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature* 427,
711 630-633.
- 712 Peck, A.M., Hornbuckle, K.C., 2004. Synthetic musk fragrances in Lake Michigan.
713 *Environmental Science & Technology* 38, 367-372.
- 714 Persson, L.M., Breitholtz, M., Cousins, I.T., de Wit, C.A., MacLeod, M., McLachlan, M.S.,
715 2013. Confronting unknown planetary boundary threats from chemical pollution.
716 *Environmental Science & Technology* 47, 12619-12622.
- 717 Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S., Lambin, E., Lenton, T.M.,
718 Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van
719 der Leeuw, S., Rodhe, H., Sorlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark,
720 M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B., Liverman, D.,
721 Richardson, K., Crutzen, P., Foley, J., 2009. Planetary boundaries: exploring the safe
722 operating space for humanity. *Ecology and Society* 14. [online] URL:
723 <http://www.ecologyandsociety.org/vol14/iss2/art32/>
- 724 Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S., Lambin, E.F., Lenton, T.M.,
725 Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van
726 der Leeuw, S., Rodhe, H., Sorlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark,
727 M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B., Liverman, D.,

- 728 Richardson, K., Crutzen, P., Foley, J.A., 2009. A safe operating space for humanity.
729 Nature 461, 472-475.
- 730 Rosi-Marshall, E.J., Kincaid, D.W., Bechtold, H.A., Royer, T.V., Rojas, M., Kelly, J.J., 2013.
731 Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial
732 communities in stream biofilms. Ecological Applications 23, 583-593.
- 733 SAICM (The Strategic Approach to International Chemicals Management), 2006. Texts and
734 Resolutions of the International Conference on Chemicals Management in 2006. United
735 Nations Environment Programme and the World Health Organization, Geneva,
736 Switzerland. 125 pp.
- 737 Scheringer, M., Stempel, S., Hukari, S., Ng, C.A., Blepp, M., Hungerbühler, K., 2012. How
738 many persistent organic pollutants should we expect? Atmospheric Pollution Research 3,
739 383-391.
- 740 Schmidt, C.W., 2006. Unfair trade - E-waste in Africa. Environmental Health Perspectives 114,
741 A232-A235.
- 742 SCHENIHR (Scientific Committee on Emerging and Newly Identified Health Risks), SCCS
743 (Scientific Committee on Consumer Safety), SCHER (Scientific Committee on Health
744 and Environmental Risks), 2012. Addressing the New Challenges for Risk Assessment.
745 European Commission, DG Health & Consumers, Directorate D: Public Health Systems
746 and Products, Unit D3 – Risk Assessment, Brussels. 157 pp.
- 747 Sindiku, O., Babayemi, J., Osibanjo, O., Schlummer, M., Schleup, M., Watson, A., Weber, R.
748 2014. Polybrominated diphenyl ethers listed as Stockholm Convention POPs, other
749 brominated flame retardants and heavy metals in e-waste polymers in Nigeria.
750 Environmental Science & Pollution Research DOI: 10.1007/s11356-014-3266-0.

- 751 Skelton, A., Guan, D., Peters, G.P., Crawford-Brown, D., 2011. Mapping Flows of Embodied
752 Emissions in the Global Production System. *Environmental Science & Technology* 45,
753 10516-10523.
- 754 Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R.,
755 Carpenter, S.R., de Vries, W., de Wit, C., Folke, C., Gerten, D., Heinke, J., Mace, G.M.,
756 Persson, L.M., Ramanathan, V., Reyers, B., Sörlin, S. 2015. Planetary boundaries:
757 guiding human development on a changing planet. *Science* DOI:
758 10.1126/science.1259855.
- 759 Stillerman, K.P., Mattison, D.R., Giudice, L.C., Woodruff, T.J., 2008. Environmental exposures
760 and adverse pregnancy outcomes: a review of the science. *Reproductive Sciences* 15,
761 631-650.
- 762 Stockholm Convention, 2008. Convention text.
763 http://ozone.unep.org/new_site/en/Treaties/treaties_decisions-hb.php?sec_id=5
- 764 Stockholm Convention, 2012. Success Stories Stockholm Convention 2001-2011. UNEP (United
765 Nations Environment Programme), Geneva. 169 pp.
- 766 Strempel, S., Scheringer, M., Ng, C.A., Hungerbühler, K., 2012. Screening for PBT chemicals
767 among the “existing” and “new” chemicals of the EU. *Environmental Science &*
768 *Technology* 46, 5680–5687.
- 769 Tapparo, A., Marton, D., Giorio, C., Zanella, A., Solda, L., Marzaro, M., Vivan, L., Girolami,
770 V., 2012. Assessment of the environmental exposure of honeybees to particulate matter
771 containing neonicotinoid insecticides coming from corn coated seeds. *Environmental*
772 *Science & Technology* 46, 2592-2599.

- 773 Torres, J.P.M., Froes-Asmus, C.I.R., Weber, R., Vijgen, J.M.H., 2013. HCH contamination from
774 former pesticide production in Brazil- A challenge for the Stockholm Convention
775 implementation. *Environmental Science & Pollution Research* 20, 1951-1957.
- 776 Turk, M., Jaksic, J., Miloradov, M.V., Klanova, J., 2007. Post-war levels of persistent organic
777 pollutants (POPs) in air from Serbia determined by active and passive sampling methods.
778 *Environmental Chemistry Letters* 5, 109-113.
- 779 UNECE (United Nations Economic Commission for Europe), 2004. Handbook for the 1979
780 Convention on Long-Range Transboundary Air Pollution and its Protocols. New York
781 and Geneva. United Nations Publication, ISBN 92-1-116895-3.
- 782 UNEP (United Nations Environment Programme), 2010-2011. The Montreal Protocol on
783 Substances that Deplete the Ozone Layer. Ozone Secretariat.
784 http://ozone.unep.org/new_site/en/Treaties/treaties_decisions-hb.php?sec_id=5
- 785 UNEP (United Nations Environment Programme), 2012. Global Environmental Outlook 5:
786 Environment for the future we want. Ed. UNEP, Nairobi.
- 787 UNEP (United Nations Environment Programme), 2013. GCO Global Chemicals Outlook:
788 Towards sound management of chemicals. Ed. United Nations Environment Programme,
789 Nairobi, Kenya.
- 790 UNEP (United Nations Environment Programme), 2015. Minamata Convention Agreed by
791 Nations. <http://www.mercuryconvention.org/>
- 792 UNEP (United Nations Environment Programme), WHO (World Health Organisation), CVUA
793 (The State Institute for Chemical and Veterinary Analysis of Food), MTM Research
794 Centre, 2013. Human exposure to POPs across the globe: POPs levels and human health

- 795 implications. Results of the WHO/UNEP human milk survey. WHO; UNEP; Stockholm
796 Convention, Geneva; Nairobi; Châtelaine.
- 797 U.S. EPA (United States Environmental Protection Agency), 2014. Design for the Environment.
798 <http://www.epa.gov/dfe/>
- 799 UNFCCC, 2013. United Nations Framework Convention on Climate Change.
800 <http://unfccc.int/2860.php>.
- 801 Voeroesmart, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P.,
802 Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R., Davies, P.M., 2010. Global
803 threats to human water security and river biodiversity. *Nature* 467, 555-561.
- 804 Wang, D.G., Norwood, W., Alae, M., Byer, J.D., Brimble, S., 2013. Review of recent advances
805 in research on the toxicity, detection, occurrence and fate of cyclic volatile methyl
806 siloxanes in the environment. *Chemosphere* 93, 711-725.
- 807 Weber, R., Watson, A., Forter, M., Oliaei, F., 2011. Persistent organic pollutants and landfills - a
808 review of past experiences and future challenges. *Waste Management & Research* 29,
809 107-121.
- 810 Weber, R., Aliyeva, G., Vijgen, J., 2013. The need for an integrated approach to the global
811 challenge of POPs management. *Environmental Science & Pollution Research* 20, 1901-
812 1906.
- 813 Whitehorn, P.R., O'Connor, S., Wackers, F.L., Goulson, D., 2012. Neonicotinoid pesticide
814 reduces bumble bee colony growth and queen production. *Science* 336, 351-352.
- 815 WHO (World Health Organisation), UNEP (United Nations Environment Programme), 2013.
816 State of the science of endocrine disrupting chemicals 2012. United Nations Environment
817 Programme and the World Health Organization, Geneva, Switzerland, 260 pp.

818 Wilson, M.P., Schwarzman, M.R., 2009. Toward a new U.S. chemicals policy: rebuilding the
819 foundation to advance new science, green chemistry, and environmental health.
820 Environmental Health Perspectives 117, 1202-1209.