

2019

Maternal–Infant Supplementation with Small-Quantity Lipid-Based Nutrient Supplements Does Not Affect Child Blood Pressure at 4–6 Y in Ghana: Follow-up of a Randomized Trial

Sika M. Kumordzie

Seth Adu-Afarwuah

Rebecca R. Young

Brietta M. Oaks

University of Rhode Island, boaks@uri.edu

Solace M. Tamakloe

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/nfs_facpubs

Citation/Publisher Attribution

Kumordzie, S. M., Adu-Afarwuah, S., Young, R. R., Oaks, B. M., Tamakloe, S. M., Ocansey, M. E.,...Dewey, K. G. (2019). Maternal–Infant Supplementation with Small-Quantity Lipid-Based Nutrient Supplements Does Not Affect Child Blood Pressure at 4–6 Y in Ghana: Follow-up of a Randomized Trial. *The Journal of Nutrition*, 149(3), 522-531. doi: 10.1093/jn/nxy285

Available at: <https://doi.org/10.1093/jn/nxy285>

This Article is brought to you for free and open access by the Nutrition and Food Sciences at DigitalCommons@URI. It has been accepted for inclusion in Nutrition and Food Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Maternal–Infant Supplementation with Small-Quantity Lipid-Based Nutrient Supplements Does Not Affect Child Blood Pressure at 4–6 Y in Ghana: Follow-up of a Randomized Trial

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Authors

Sika M. Kumordzie, Seth Adu-Afarwuah, Rebecca R. Young, Brietta M. Oaks, Solace M. Tamakloe, Maku E. Ocansey, Harriet Okronipa, Elizabeth L. Prado, and Kathryn G. Dewey

Maternal–Infant Supplementation with Small-Quantity Lipid-Based Nutrient Supplements Does Not Affect Child Blood Pressure at 4–6 Y in Ghana: Follow-up of a Randomized Trial

Sika M Kumordzie,¹ Seth Adu-Afarwuah,² Rebecca R Young,¹ Brietta M Oaks,^{1,3} Solace M Tamakloe,² Maku E Ocansey,¹ Harriet Okronipa,¹ Elizabeth L Prado,¹ and Kathryn G Dewey¹

¹Program in International and Community Nutrition, Department of Nutrition, University of California Davis, Davis, CA; ²Department of Nutrition and Food Science, University of Ghana, Legon, Ghana; and ³Department of Nutrition and Food Sciences, University of Rhode Island, Kingston, RI

ABSTRACT

Background: In the International Lipid-Based Nutrient Supplements (iLiNS)-DYAD-Ghana trial, prenatal small-quantity lipid-based nutrient supplements (LNSs) had a positive effect on birth weight. Birth weight may be inversely related to blood pressure (BP) later in life.

Objectives: We examined the effect of the intervention on BP at 4–6 y of age, and maternal and child factors related to BP.

Methods: The iLiNS-DYAD-Ghana study was a partially double-blind, randomized controlled trial which assigned women ($n = 1320$) ≤ 20 weeks of gestation to daily supplementation with: 1) iron and folic acid during pregnancy and 200 mg Ca for 6 mo postpartum, 2) multiple micronutrients during pregnancy and postpartum, or 3) LNSs during pregnancy and postpartum plus LNSs for infants from 6 to 18 mo of age. At 4–6 y of age ($n = 858$, 70% of live births), we compared BP, a secondary outcome, between non-LNS and LNS groups and examined whether BP was related to several factors including maternal BP, child weight-for-age z score (WAZ), and physical activity.

Results: Non-LNS and LNS groups did not differ in systolic (99.2 ± 0.4 compared with 98.5 ± 0.6 mm Hg; $P = 0.317$) or diastolic (60.1 ± 0.3 compared with 60.0 ± 0.4 mm Hg; $P = 0.805$) BP, or prevalence of high BP (systolic or diastolic BP ≥ 90 th percentile of the US National Heart, Lung, and Blood Institute reference: 31% compared with 28%; $P = 0.251$). BP at 4–6 y of age was positively related to birth weight; this relation was largely mediated through concurrent WAZ in a path model. Concurrent WAZ and maternal BP were the factors most strongly related to child BP.

Conclusions: Despite greater birth weight in the LNS group, there was no intervention group difference in BP at 4–6 y. In this preschool population at high risk of adult hypertension based on BP at 4–6 y, high maternal BP and child WAZ were key factors related to BP. This trial was registered at clinicaltrials.gov as NCT00970866. *J Nutr* 2019;149:522–531.

Keywords: supplementation, blood pressure, lipid-based nutrient supplements, Ghanaian children; prenatal; infant nutrition

Introduction

The maternal intrauterine environment plays a key role in fetal development and conditions the offspring for risk of certain metabolic diseases later in life (1). Because fetal growth is strongly influenced by nutrient and oxygen supply in utero, prenatal nutrition is considered a key programming stimulus (2). There is considerable evidence for an inverse relation between birth weight and subsequent blood pressure (BP) later in life (3), most of which comes from animal research

and human cohort studies (4). Low birth weight is associated with a reduction in nephron number in humans (5, 6), which may increase susceptibility to renal injury and thereby affect BP. Besides birth weight, elevated BP and adiposity in childhood are independent predictors of adult hypertension and cardiovascular disease (7).

In children, hypertension is defined as systolic or diastolic BP ≥ 95 th percentile and elevated BP as values ≥ 90 th percentile and < 95 th percentile (8). In a recent meta-analysis, the prevalence of hypertension and elevated BP among 2- to 19-y-old African

children was 5.5% and 12.7%, respectively (9). Results from a meta-analysis and systematic review indicate that BP tracks from childhood into adulthood (10) and 1 study reported that even short periods of hypertension in childhood may increase the risk of hypertension as an adult (11). The prevalence of adult hypertension (defined as systolic BP \geq 140 mm Hg or diastolic BP \geq 90 mm Hg) in Ghana is relatively high, ranging between 19.3% and 54.6% among adults (12), which parallels the general trend towards chronic disease in a country undergoing the nutrition transition. The double burden of undernutrition in early life and subsequent exposure to risk factors for overweight puts young children at high risk of later development of chronic diseases. Interventions that can improve nutrition in early life may thus help to prevent these consequences.

The International Lipid-Based Nutrient Supplements (iLiNS)-DYAD trial in Ghana evaluated the effects of maternal and child supplementation with small-quantity lipid-based nutrient supplements (LNSs) on birth outcomes and child growth. As reported previously, LNS supplementation had a positive effect on birth weight (13) and weight and length by 18 mo of age (14). The current study reports results from a follow-up of the children from the iLiNS-DYAD trial at 4–6 y of age. The first objective of this analysis was to determine the effect of LNS supplementation on systolic BP (SBP) and diastolic BP (DBP). Because birth weight has been inversely related to BP in later life, we hypothesized that children in the LNS group would have lower BP than children in the control groups. The second objective was to examine the maternal and child factors related to BP in this cohort of preschool children, including maternal BMI and BP, and child anthropometric indexes, dietary variables, and physical activity.

Methods

Location and study design of the main trial

The iLiNS-DYAD-Ghana trial (NCT00970866) was conducted in the Yilo and Lower Manya Krobo districts of the Eastern Region of Ghana between December 2009 and March 2014. Details of the study are published elsewhere (13). Briefly, the study was a partially double-blind, randomized controlled trial that enrolled 1320 women 18 y or older at \leq 20 weeks of gestation attending antenatal clinics in 4 main health facilities in the study area. The women were randomly assigned to 1 of 3 supplementation groups: 1) daily iron and folic acid capsules during pregnancy, 200 mg/d calcium tablets (placebo) during the first 6 mo postpartum, and no infant supplementation; 2) daily multiple micronutrient capsules (1–2 times the RDA of 18 vitamins and minerals) during pregnancy and the first 6 mo postpartum and no infant supplementation; or 3) daily 20-g (118-kcal) LNSs during pregnancy and the first 6 mo postpartum followed by infant LNS supplementation from 6 to 18 mo of age. The maternal LNSs had the same micronutrient content as the multiple micronutrient supplement, plus calcium, magnesium, phosphorus, potassium, and macronutrients

(essential fatty acids and a small amount of protein), whereas the infant LNSs had the same macronutrients and 22 micronutrients based on infant Recommended Nutrient Intakes (15). Women and children were followed \leq 6 mo postpartum and 18 mo of age, respectively. Growth status of children was measured at birth, 6, 12, and 18 mo.

Follow-up when children were 4–6 y of age

Participants in the follow-up study (also NCT00970866) were the children born to the pregnant women who were randomly assigned into the trial. In the main trial, when a mother delivered twins, one of the twins was randomly selected as the study child. All children who were alive at the time of the follow-up study were potentially eligible to participate. At follow-up, after excluding misdiagnosed pregnancies ($n = 5$), miscarriages and stillbirths ($n = 66$), and child deaths before the end of the main trial ($n = 27$), 1222 children were potentially eligible to participate. Details of the study profile are shown in Figure 1.

The follow-up study protocols were approved by the Institutional Review Board of University of California Davis; the Ethics Committee for the College of Basic and Applied Sciences at the University of Ghana; and the Ghana Health Service Ethical Review Committee. Written informed consent was given by children's primary caregivers before data collection.

Data collection procedures.

Participants were contacted using the last known address and contact number provided. Field staff called to set up an appointment and visited the caregiver to obtain informed consent for the follow-up activities. After consent had been given, participants were scheduled for the data collection visits, the last of which was the laboratory visit, which took place in a central location. BP measurements for mothers and children were done in duplicate by a trained nurse and her assistant using automated BP monitors (Riester ri-champion N, for children; Omron BP742N, for mothers) with age-appropriate arm cuffs. Measurements were done while the child was sitting with the child's arm at their chest level on a table and feet resting flat on a footstool. For mothers, measurements were done following the same procedure as for children except mothers had their feet on the floor. Participants rested (sitting) for \geq 5 min before measurements were taken.

Anthropometric measurements were taken according to WHO standard procedures (16) by trained anthropometrists. Height was measured using a stadiometer (Seca 217) to the nearest 0.1 cm, weight using a scale (Seca 875) to the nearest 50 g. All measurements were taken in duplicate and in triplicate if the first 2 measurements differed by a predefined amount: 0.1 kg for weight and 0.5 cm for height.

Data on child food and beverage preferences and consumption, with an emphasis on sweet foods and beverages, were collected by administering a questionnaire to the caregiver (17). The questionnaire asked about the number of times different food and beverage items (or food and beverage groups) were consumed by the child in the week preceding the interview. The reported numbers of times for each of the items were summed up to obtain a score for total frequency of consumption of sweet foods and beverages.

Anthropometry, BP measurements, and recall of child food and beverage consumption were targeted for the full sample. Physical activity was to be estimated in a random subsample of 630 ($n = 420$ and 210, non-LNS and LNS, respectively) children. Of this number, 376 children were fitted with a device and physical activity was successfully estimated for 353 children ($n = 224$ and 129, non-LNS and LNS, respectively). Every child in the subsample was fitted with a single accelerometer (Actigraph GT3X) for 1 wk to estimate physical activity. Accelerometers were fitted to an elastic belt and fastened to the child's right hip by trained data collectors. Caregivers were instructed to let the child wear the accelerometer continuously day and night for the 1-wk period unless they experienced discomfort. Activity was stored in 60-s intervals or epochs. Physical activity data were analyzed using ActiLife data analysis software version 6.13.1 to generate mean vector magnitude accelerometer counts per minute per participant.

All data collection staff were blinded to group assignments. The data analyst remained blinded until all decisions regarding outliers for our first objective had been made.

Supported by a Bill & Melinda Gates Foundation grant (# OPP49817) to the University of California, Davis.

Author disclosures: SMK, SA-A, RRY, BMO, SMT, MEO, HO, ELP, and KGD, no conflicts of interest.

The findings and conclusions contained within this work are those of the authors and do not necessarily reflect the positions or policies of the Bill & Melinda Gates Foundation.

Supplemental Tables 1–4 are available from the "Supplementary data" link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/jn/>.

Address correspondence to SMK (e-mail: smkumordzie@ucdavis.edu).

Abbreviations used: BP, blood pressure; DBP, diastolic blood pressure; iLiNS, International Lipid-Based Nutrient Supplements; LNS, lipid-based nutrient supplement; MAP, mean arterial pressure; PP, pulse pressure; SBP, systolic blood pressure; WAZ, weight-for-age z score.




FIGURE 1 Study profile. IFA, iron and folic acid; LNS, lipid-based nutrient supplement; MMN, multiple micronutrients.

Sample size and data analysis.

For these analyses, our sample size was the 858 children who had BP data at the 4–6 y follow-up. With this sample size we were powered to detect a difference of 1.07 mm Hg in systolic BP (an effect size of 0.11, based on the SD of 9.87) between the 2 groups (non-LNS and LNS) at 80% power.

For our first objective, to determine the effect of the intervention on BP (a secondary objective of the study), a statistical analysis plan was posted on the iLiNS Project website (www.ilins.org) before data

analysis. The primary outcomes for this analysis were SBP and DBP and secondary outcomes were SBP and DBP z scores standardized based on gender, age, and height, as well as high BP defined as SBP or DBP ≥ 90 th percentile of the reference population. BP percentiles and z scores were calculated using the equations given in the National Heart, Lung, and Blood Institute Fourth Report on the Diagnosis, Evaluation, and Treatment of High Blood Pressure in Children and Adolescents (8). Analysis was carried out based on the intention-to-treat principle using SAS version 9.4 (SAS Institute) and all tests




FIGURE 2 Conceptual model of maternal and child factors related to child blood pressure. WAZ, weight-for-age z score.

were 2-sided and considered significant at the 5% level. Because our hypothesis was based on comparing the LNS group with the 2 non-LNS groups, the primary analysis for the first objective was a 2-group comparison. However, a sensitivity analysis was performed comparing all 3 groups. The effect of the intervention on continuous outcomes was examined using ANCOVA. Logistic regression was used for analysis of categorical outcomes. For all analyses, we controlled for child age at follow-up (minimally adjusted models). For fully adjusted models, several prespecified covariates were considered for inclusion in addition to child age: child sex, and maternal characteristics at enrollment in the main trial (gestational age, nulliparity, BMI, and household asset score). These additional covariates were included only if they were associated with an outcome at a 10% level of significance in bivariate analysis. Maternal SBP, DBP, and BMI at enrollment in the main trial, nulliparity, and child sex were tested as effect modifiers, and if the interaction term was significant (P -interaction < 0.05) we conducted further analysis by stratifying the intervention groups on categories of the effect modifier. In addition, we conducted a per-protocol analysis by limiting the analysis to mothers who self-reported $\geq 80\%$ adherence to the supplements during the pregnancy period.

For the second objective, to identify the factors related to BP in this cohort of children, the analysis was based on the conceptual model in [Figure 2](#). The outcomes were SBP, DBP, mean arterial pressure (MAP), and pulse pressure (PP) calculated as follows: $MAP = DBP + [0.333 \times (SBP - DBP)]$ and $PP = SBP - DBP$ (18). We included MAP and PP because they may be helpful in interpreting the relative role of both arterial stiffness and vascular resistance in contributing to cardiovascular disease risk (19). We examined the relation of each of these outcomes to several maternal (maternal prepregnancy BMI, maternal BP at enrollment and follow-up) and child [birth WAZ (weight-for-age z score); postnatal weight gain (0–6 mo, i.e., difference between weight at 6 mo and birth weight); concurrent WAZ, percentage body fat, and BMI; sweet food and beverage intake at follow-up; physical activity at follow-up] factors. Estimated prepregnancy BMI was calculated from estimated prepregnancy weight (based on polynomial regression with gestational age, gestational age squared, and gestational age cubed as predictors) and height at enrollment (14). All variables were standardized before analysis of the data. For each

model, the outcome and the factor were examined in an independent regression model adjusting for child age in minimally adjusted models and including in fully adjusted models other covariates (household asset score at enrollment, maternal age, education, and nulliparity at enrollment, child sex, and intervention group) if these covariates were associated with the outcome at a 10% level of significance in bivariate analysis. Household asset score was constructed based on ownership of a set of assets (radio, television, refrigerator, and stove), lighting source, drinking water supply, sanitation facilities, and flooring materials, developed into an index (with a mean of 0 and SD of 1) using principal components analysis (20). For each model, the value of the covariate for observations that had high leverage (> 0.02) was truncated to the 2.5th or 97.5th percentile. We performed a sensitivity analysis with and without truncation of the observations and indicated where appropriate whether truncation made a difference to the results. Normality of the residuals was tested with the Shapiro–Wilk test and the linearity assumption was evaluated by plotting a residual against fit scatterplot. We also examined how the factors related to each other. First, for each path in the model, factor–outcome or factor–factor, we removed any path that was nonsignificant ($P < 0.10$). Second, for variables that were retained, we checked for mediation between independent variables and the outcome according to the model in [Figure 2](#), using the binary mediation command in Stata version 15.0 (StataCorp) and we included any significant mediation pathways in the overall path model. The mediators tested were child WAZ and BMI z score at 4–6 y. Finally, for variables and pathways that were retained in the first 2 steps, the overall model coefficients were estimated using the SEM command in Stata specifying standardized coefficients and the maximum likelihood for missing values method to account for missing data. The overall path was modeled for only 1 outcome: child SBP.

Results

Of the 1222 children eligible to participate in the follow-up study, we enrolled 662 and 352 in the non-LNS and LNS groups, respectively, and measured BP for 557 in the non-LNS and 301

TABLE 1 Background characteristics of women and children in the iLiNS-DYAD-Ghana trial included in the follow-up¹

	Non-LNS (<i>n</i> = 557)	LNS (<i>n</i> = 301)
Maternal characteristics at baseline		
Age, y	26.7 ± 5.4	26.7 ± 5.5
Gestational age at enrolment, wk	16.0 ± 3.2	16.1 ± 3.4
Formal education, y	7.6 ± 3.4	7.6 ± 3.7
Married or cohabiting, % (<i>n</i> / <i>N</i>)	93.5 (521/557)	92.7 (279/301)
Asset score ²	0.07 ± 0.94	−0.11 ± 0.97
Nulliparous, % (<i>n</i> / <i>N</i>)	32.1 (179/557)	33.6 (101/301)
Weight, kg	61.5 ± 12.0	62.9 ± 12.4
Height, cm	158.8 ± 5.8	159.3 ± 5.4
Prepregnancy BMI, ³ kg/m ²	24.4 ± 4.5	24.8 ± 4.5
Overweight (BMI ≥ 25), % (<i>n</i> / <i>N</i>)	30.5 (166/545)	34.3 (102/297)
Systolic blood pressure, mm Hg	110.1 ± 10.8	110.7 ± 13.9
Diastolic blood pressure, mm Hg	62.3 ± 8.1	63.1 ± 9.3
Child characteristics		
Birth weight, g	2962.5 ± 425.9	3019.9 ± 415.9
Current age of child, y	5.0 ± 0.6	5.1 ± 0.6
Sex of child, % male	46.8	50.2
Weight at 4–6 y, kg	16.4 ± 2.1	16.7 ± 2.3
Height at 4–6 y, cm	106.1 ± 5.4	107.0 ± 5.9
Height-for-age z score at 4–6 y	−0.30 ± 0.91	−0.25 ± 0.99
BMI z score at 4–6 y	−0.57 ± 0.80	−0.55 ± 0.82
Fat mass at 4–6 y, %	15.3 ± 4.6	15.2 ± 5.0
Fat-free mass at 4–6 y, %	84.9 ± 5.5	84.8 ± 5.1

¹Values are means ± SDs unless otherwise stated. iLiNS, international lipid-based nutrient supplements; LNS, lipid-based nutrient supplement group; non-LNS, IFA (iron and folic acid group) + MMN (multiple micronutrients group).

²Household asset score was constructed based on ownership of a set of assets and access to certain amenities using principal components analysis.

³Estimated prepregnancy BMI was calculated from estimated prepregnancy weight (based on polynomial regression with gestational age, gestational age squared, and gestational age cubed as predictors) and height at enrollment.

in the LNS group (70% of live births from the main trial; 84.6% of re-enrolled children) by the end of the period for follow-up data collection (January to December, 2016). The reasons for nonparticipation were inability to trace the mother (*n* = 96), consent refusal (*n* = 43), child death after the main trial ended (*n* = 5), no longer residing in the study area (*n* = 64), and loss to follow-up after enrollment in the follow-up (*n* = 156). Details of the study flow are presented in [Figure 1](#). There was a trend for a difference in proportion lost to follow-up between the groups (non-LNS: 36.7%; LNS: 31.6%, *P* = 0.07).

Maternal characteristics at enrollment into the main trial for women in the current analysis are shown in [Table 1](#). More than 30% of the women were overweight (31% in the non-LNS group and 34% in the LNS group, *P* = 0.035) and >90% were married or cohabiting. The non-LNS group had a higher household assets score than the LNS group (0.07 compared with −0.11, *P* = 0.009) whereas the LNS women were heavier (61.5

compared with 62.9 kg, *P* = 0.035). About half the children were male and mean ± SD age was 5.0 ± 0.6 y. There were no significant differences in maternal enrollment characteristics between those included in this analysis (*n* = 858) and those lost to follow-up ([Supplemental Table 1](#)).

Mean ± SD SBP and DBP of children at 4–6 y were 99.0 ± 9.9 and 60.1 ± 6.7 mm Hg, respectively, and prevalence of high BP was 30.4%. Continuous and categorical BP outcomes adjusted for age are shown in [Tables 2](#) and [3](#), respectively. There were no significant differences between the intervention groups for any of the BP outcomes. The aforementioned results did not change after adjusting for additional prespecified covariates. The sensitivity analysis revealed no significant differences in the 3-group comparisons for any of the continuous ([Supplemental Table 2](#)) or categorical ([Supplemental Table 3](#)) outcomes.

Tests for interaction showed no significant (*P* < 0.05) interactions of intervention group with any of the following

TABLE 2 Comparison of continuous BP measurements of children in the iLiNS-DYAD-Ghana trial follow-up at 4–6 y¹

	Non-LNS (<i>n</i> = 557)	LNS (<i>n</i> = 300)	<i>P</i> value	Difference in mean (95% CI)
SBP, mm Hg	99.2 ± 0.4	98.5 ± 0.6	0.317	0.71 (−0.68, 2.09)
SBP z score	0.67 ± 0.04	0.59 ± 0.05	0.199	0.09 (−0.05, 0.22)
DBP, mm Hg	60.1 ± 0.3	60.0 ± 0.4	0.805	0.12 (−0.83, 1.07)
DBP z score	0.69 ± 0.03	0.69 ± 0.04	0.891	0.01 (−0.08, 0.09)

¹Values represent means ± SEs and differences in means (95% CIs). Values in the table are adjusted for child age. Results are based on ANCOVA (SAS PROC GLIMMIX). BP, blood pressure; DBP, diastolic blood pressure; iLiNS, international lipid-based nutrient supplements; LNS, lipid-based nutrient supplements group; non-LNS, IFA (iron and folic acid group) + MMN (multiple micronutrients group); SBP, systolic blood pressure.

TABLE 3 Comparison of categorical BP outcomes of children in the iLiNS-DYAD-Ghana trial follow-up at 4–6 y¹

	Non-LNS (<i>n</i> = 536)	LNS (<i>n</i> = 289)	<i>P</i> value	OR (95% CI)
Prehypertensive/hypertensive (SBP ≥ 90th percentile)				
Prevalence, %	26.3	22.2	0.146	0.78 (0.55, 1.09)
Prehypertensive/hypertensive (DBP ≥ 90th percentile)				
Prevalence, %	14.4	14.2	0.999	1.00 (0.66, 1.51)
Prehypertensive/hypertensive (SBP or DBP ≥ 90th percentile)				
Prevalence, %	31.7	28.0	0.251	0.83 (0.61, 1.14)

¹Values are the percentage of participants whose response was “yes” for the outcome in question and OR (95% CI) obtained by comparing the groups. Reference = non-LNS group for all outcomes. Values in the table are adjusted for child age. Results are based on logistic regression (SAS PROC GLIMMIX). BP, blood pressure; DBP, diastolic blood pressure; iLiNS, international lipid-based nutrient supplements; LNS, lipid-based nutrient supplements group; non-LNS, IFA (iron and folic acid group) + MMN (multiple micronutrients group); SBP, systolic blood pressure.

variables at baseline: maternal prepregnancy BMI, maternal SBP or DBP, nulliparity, and child sex (data not shown). When the analysis was restricted to mothers who reported ≥80% adherence during the pregnancy period (*n* = 316 in the non-LNS group and *n* = 134 in the LNS group), we observed a marginally significantly higher SBP in the non-LNS group than in the LNS group (99.6 ± 0.6 compared with 97.7 ± 0.9 mm Hg; *P* = 0.06), a borderline significant difference in SBP *z* score (0.70 ± 0.05 compared with 0.51 ± 0.08; *P* = 0.049), and a marginally significant difference in the prevalence of high BP (34.9% compared with 26.2%; *P* = 0.08). The results remained the same in the adjusted analysis (data not shown). There was no difference in any of the outcomes for the sensitivity analyses among high adherers (data not shown).

Summary statistics for the variables used in the analyses for objective 2 are presented in **Supplemental Table 4**. WAZ at 4–6 y was positively associated with sweet food and beverage intake [β (SE) = 0.07 (0.03), *P* = 0.018], WAZ at birth was inversely associated with postnatal weight gain [β (SE) = –0.07

(0.04), *P* = 0.07], and WAZ and BMI at 4–6 y were positively associated [β (SE) = 0.65 (0.02), *P* < 0.0001]. The relation between BMI at 4–6 y and sweet food and beverage intake was nonsignificant [β (SE) = 0.02 (0.03), *P* = 0.53].

Table 4 shows the associations between each independent factor and each outcome, adjusting only for child age. SBP, DBP, PP, and MAP were all positively associated (*P* < 0.10) with postnatal weight gain, WAZ at 4–6 y, BMI *z* scores at 4–6 y, and maternal SBP and DBP at follow-up. In addition, child SBP and MAP were positively associated with WAZ at birth and sweet food and beverage intake, whereas child DBP and MAP were positively associated with maternal SBP and DBP at enrollment.

For the path analysis, we modeled the overall path only for SBP because of the 4 outcomes, the β -coefficients for the associations with the predictors were highest for SBP and MAP but SBP is measured more commonly than MAP. We also included maternal BP at follow-up and not enrollment because the enrollment measures were not significantly related to child SBP. Maternal SBP at enrollment and follow-up were,

TABLE 4 Associations between outcomes (child SBP, DBP, PP, and MAP) and each factor in the iLiNS-DYAD-Ghana trial follow-up at 4–6 y¹

Factors	SBP		DBP		PP		MAP	
	β (SE)	<i>P</i> value						
WAZ at birth	0.08 (0.04)	0.026	0.06 (0.04)	0.132	0.04 (0.04)	0.223	0.08 (0.04)	0.035
Postnatal weight gain (0–6 mo) ²	0.10 (0.04)	0.004	0.06 (0.04)	0.085	0.08 (0.04)	0.035	0.09 (0.04)	0.012
WAZ at 4–6 y	0.29 (0.04)	<0.0001	0.23 (0.04)	<0.0001	0.16 (0.04)	<0.0001	0.29 (0.04)	<0.0001
BMI <i>z</i> score at 4–6 y	0.21 (0.04)	<0.0001	0.17 (0.04)	0.0002	0.12 (0.04)	0.006	0.20 (0.04)	<0.0001
Percentage fat mass at 4–6 y	–0.04 (0.04)	0.260	0.02 (0.04)	0.638	–0.05 (0.04)	0.145	–0.01 (0.04)	0.846
Physical activity ³ at 4–6 y	0.04 (0.06)	0.465	0.05 (0.06)	0.345	0.01 (0.06)	0.802	0.06 (0.06)	0.331
Sweet food and drink intake	0.06 (0.03)	0.062	0.04 (0.03)	0.206	0.04 (0.03)	0.282	0.06 (0.03)	0.084
Maternal SBP at enrollment	0.05 (0.03)	0.124	0.08 (0.03)	0.016	–0.01 (0.03)	0.881	0.07 (0.03)	0.034
Maternal SBP at follow-up	0.13 (0.03)	0.0002	0.16 (0.03)	<0.0001	0.03 (0.03)	0.316	0.16 (0.03)	<0.0001
Maternal DBP at enrollment	0.05 (0.04)	0.130	0.13 (0.03)	0.0002	–0.04 (0.03)	0.233	0.10 (0.10)	0.002
Maternal DBP at follow-up	0.12 (0.03)	0.0008	0.16 (0.03)	<0.0001	0.02 (0.03)	0.593	0.15 (0.03)	<0.0001
Maternal prepregnancy ⁴ BMI	0.04 (0.03)	0.260	0.05 (0.03)	0.184	0.01 (0.03)	0.882	0.05 (0.03)	0.185

¹Values in the table are coefficients (SEs) from separate models of each independent factor with each outcome, adjusted only for child age. Coefficients are standardized. Results are based on regression (SAS PROC REG). DBP, diastolic blood pressure; iLiNS, international lipid-based nutrient supplements; MAP, mean arterial pressure; PP, pulse pressure; SBP, systolic blood pressure; WAZ, weight-for-age *z* score.

²Postnatal weight gain = weight at 6 mo – birth weight.

³Physical activity was estimated as vector magnitude count per minute.

⁴Estimated prepregnancy BMI was calculated from estimated prepregnancy weight (based on polynomial regression with gestational age, gestational age squared, and gestational age cubed as predictors) and height at enrollment.




FIGURE 3 Final path model of the maternal and child factors related to child systolic blood pressure at 4–6 y. Numbers in the model are standardized β -coefficients. ** $P > 0.0001$ but $P < 0.001$; *** $P < 0.0001$. SBP, systolic blood pressure; WAZ, weight-for-age z score.

however, correlated with each other [correlation coefficient (r) = 0.47; $P < 0.0001$]. Child BMI z score and WAZ at 4–6 y were highly correlated ($r > 0.65$; $P < 0.0001$) but the β -coefficient for the association of child BMI z score with SBP was lower than that for WAZ, so BMI z score was dropped from the final model. In the final path model (Figure 3), WAZ at 4–6 y and maternal concurrent SBP remained positively associated with child SBP. Birth WAZ and postnatal weight gain were not significantly associated with child SBP in the path model even though both factors were significantly associated with child SBP in a regression analysis controlling only for child age. Maternal prepregnancy BMI was significantly associated with both child WAZ at follow-up (4–6 y) and maternal SBP at follow-up.

Birth WAZ, postnatal weight gain, and sweet food and beverage intake were also positively associated with child WAZ at 4–6 y. WAZ at 4–6 y was a significant mediator of the relations between birth WAZ and child SBP and between postnatal weight gain and child SBP. Figure 4A illustrates the relation between birth WAZ and child SBP mediated by WAZ at 4–6 y. Birth WAZ was positively associated with WAZ at 4–6 y in a fully adjusted model. Child SBP was positively associated with WAZ at 4–6 y but not with birth WAZ. In Figure 4B we examine WAZ at 4–6 y as a mediator between postnatal weight gain and child SBP. Postnatal weight gain was positively associated with WAZ at 4–6 y whereas SBP at 4–6 y was inversely associated with postnatal weight gain but positively associated with WAZ at 4–6 y. Birth WAZ was a significant mediator of the relation between maternal prepregnancy BMI and WAZ at 4–6 y (Figure 4C). However, the direct association between maternal prepregnancy BMI and WAZ at 4–6 y also remained significant.

Discussion

Because increased birth weight has been associated with lower BP later in life (21–30), and the provision of LNSs during pregnancy increased birth weight in the iLiNS-DYAD-Ghana

trial (13), we hypothesized that children in the LNS group would have lower BP at 4–6 y than those in the non-LNS group. The primary results did not support this hypothesis: we observed no effect of LNSs provided to mothers prenatally and in the first 6 mo of lactation and to children from 6 to 18 mo of age on child BP in this follow-up of the iLiNS-DYAD-Ghana cohort. However, in the per-protocol analysis among high adherers ($\geq 80\%$ adherence during pregnancy), we




FIGURE 4 Path diagrams representing mediation analysis within the final path model. (A) Mediation between birth WAZ and SBP at 4–6 y by WAZ at 4–6 y. Parameter estimates of mediation: 0.08 (0.02)***; proportion mediated 97.2%. (B) Mediation between postnatal weight gain (0–6 mo) and SBP at 4–6 y by WAZ at 4–6 y. Parameter estimates of mediation: 0.13 (0.03)***; proportion mediated 75.6%. (C) Mediation between maternal prepregnancy BMI and WAZ at 4–6 y by birth WAZ. Parameter estimates of mediation: 0.07 (0.02)***; proportion mediated 25.9%. Numbers in the model were obtained from mediation analysis in STATA software version 15.0 and represent parameter estimates (SEs). *** $P < 0.0001$. SBP, systolic blood pressure; WAZ, weight-for-age z score.

observed lower SBP z score ($P < 0.05$), SBP, and prevalence of high BP ($P < 0.10$) in the LNS group. There was a high overall prevalence of high BP (30%) among the children. Child BP at 4–6 y was significantly associated with concurrently measured maternal SBP and child WAZ. Birth weight and postnatal weight gain (0–6 mo of age) were both related indirectly to child SBP via their positive associations with child WAZ at 4–6 y.

Strengths of our study include the large sample size and the blinding of field staff (study nurses and anthropometrists) to group assignment. A potential limitation is that loss to follow-up was somewhat higher in the non-LNS group. However, there were no differences in characteristics between the sample included in this analysis and those lost to follow-up, suggesting that our conclusions are generalizable to the full cohort. Another limitation is that the non-LNS group had a lower household asset score and a higher percentage of maternal overweight than the LNS group. However, we adjusted for asset score, maternal prepregnancy BMI, and other prespecified covariates in our analysis. A further limitation is that we did not measure certain dietary factors associated with BP, such as sodium intake. Although we did not measure BP by the recommended auscultation procedure (manual BP cuff and stethoscope), we used a validated oscillometric device (automated BP cuff). We also used appropriately sized cuffs (child and adult) for the measurement of BP and allowed participants to rest for ≥ 5 min before measurements were taken to prevent artificially elevated values.

Our primary results are consistent with those of a previous trial providing both prenatal and postnatal nutrition supplementation that included a long-term follow-up: the INCAP Longitudinal Study in Guatemala (31). The investigators found no independent effect of the supplements (Atole compared with Fresco, intervention compared with control supplement) on SBP or DBP in young adulthood. Results of other follow-up studies, in which the intervention included only prenatal supplementation (with food or micronutrients) (32–38), have been mixed. Two trials in Nepal examined the effect of maternal micronutrient supplementation on offspring BP (33, 37, 38). One showed lower SBP but not DBP in the multiple micronutrient group than in the control group when the children were 2.5 y of age (38), but no difference at 8.5 y (33). The other trial showed no difference in BP when children were ~ 7.5 y of age (37). Four other trials involved prenatal food supplementation (with or without micronutrient supplements) (34, 36) or prenatal calcium supplementation (32, 35); 3 indicated no effect on child BP, but 1 of the calcium trials (32) showed lower SBP and lower risk of hypertension in the calcium group than in the placebo group. Although we cannot directly compare the results of these trials to ours because of differences in study design, all except the latter trial agree with our results showing no main effect of prenatal or postnatal supplementation on later BP. It is important to note that the increase in birth weight observed in the LNS group in the main iLiNS-DYAD-Ghana trial was < 100 g, whereas the magnitude of the association between birth weight and later BP reported in observational studies was a decrease of 1–4 mm Hg in BP for every 1-kg increase in birth weight (25, 26, 30). Thus, the increase in birth weight resulting from prenatal nutrition interventions may not be large enough to lead to a detectable difference in BP later in life.

The prevalence of high BP was 30% in this cohort, which is similar to the prevalence observed in a Spanish population of 4- to 6-y-old children: 27.5% in boys and 30.6% in

girls (18). In a study of 5- to 8-y-old children in Brazil, the investigators used 2 different definitions for prevalence of prehypertension/hypertension, based on elevated SBP (35%) or elevated DBP (5%) (39). When we considered prehypertension/hypertension due only to elevated SBP, prevalence was 25% (data not shown) in our cohort, which is lower than in Brazil (39), but when based only on elevated DBP, prevalence was 14%, higher than the 5% rate reported in Brazil.

We did not observe an inverse relation between birth weight and SBP in this cohort as has been reported in several prospective cohort studies (21, 22, 25, 26, 28–30) and a systematic review (3). Rather, we found a positive relation between birth WAZ and BP at 4–6 y in regression analysis controlling only for child age. Similar results have been reported for an Aboriginal community where low birth weight individuals had lower weight, height, BMI, cholesterol concentration, and BP at 20–38 y compared to normal birth weight individuals (40) and in the Pelotas, Brazil birth cohort where SBP and DBP at 11 y were positively associated with birth weight and length (41). However, in the Pelotas cohort, further adjustment for several covariates and current BMI resulted in an inverse association of birth weight with SBP. When we adjusted for WAZ or BMI at 4–6 y the relation between birth WAZ and child SBP did not become inverse. Thus, our results do not support the hypothesis that low birth weight is a risk factor for hypertension later in life. The mean birth weight in our study was 3.0 kg, comparable to the mean birth weight of ≥ 3 kg in most of the studies (22, 26, 28–30) reporting an inverse association between birth weight and BP, except for 2 studies with lower mean birth weights (21, 25). However, in our cohort the prevalence of low birth weight was low and at 4–6 y the prevalence of overweight was low, which may explain why we did not observe an inverse association between birth weight and BP. In addition, although the prevalence of overweight in our cohort was low, the prevalence of elevated BP was high, suggesting that in this population factors other than body size may also be critical determinants of BP.

We observed a positive association between postnatal weight gain (0–6 mo) and all BP outcomes at 4–6 y in regression analysis controlling only for age, similar to the findings of the Manchester Children's Growth and Vascular Health study (42), which showed that change in weight from birth to 3 mo of age in children of South Asian and Caucasian origin was positively associated with SBP at 12 mo of age. A similar result was also observed in a prospective cohort in Amsterdam, in which faster relative weight gain after the first month was associated with higher SBP and DBP at 5 y of age (43).

We found that child WAZ at 4–6 y of age largely mediated the associations of both birth WAZ and postnatal weight gain with BP at 4–6 y, suggesting that current size is a more important determinant than body size or growth in infancy. In addition to WAZ, BMI z score at 4–6 y was positively associated with BP in a regression analysis controlling only for age, but the association with BMI z score was not significant when WAZ was taken into account. Other studies have also reported a positive association between BMI and BP in children (28, 44, 45). In children and adolescents, the normal range of BP depends on body size and age (8). We did not observe a relation between percentage body fat and BP in this cohort of children, although BMI was associated with BP, suggesting that the association is driven by overall weight. In the Amsterdam Born Child and their Development study, which examined ethnic differences in the

relations of children's BP to BMI, waist-to-height ratio, and fat mass index, the strongest association was observed for BMI, and the associations of BP with BMI and fat mass index differed by ethnicity (46).

We also found a positive association between maternal BP (at both baseline and follow-up) and child BP, although the association was stronger with maternal BP at follow-up. In the Generation R prospective cohort study in the Netherlands (47), maternal BP at all time points in pregnancy (early, mid-, and late pregnancy) was positively associated with childhood BP at 6 y, with the strongest association observed with early pregnancy maternal SBP. A similar result was observed in the Avon Longitudinal Study of Parents and Children in the United Kingdom, in which higher maternal BP in early pregnancy was associated with higher child BP at 7 y (48). The study investigators concluded that the observed association may be driven by shared genetic or environmental factors.

In the final path model accounting for all factors, there was no direct relation between SBP and either birth WAZ or postnatal weight gain. These 2 early-life factors operated through current WAZ, which was the strongest factor directly related to BP at preschool age. Postnatal weight gain was a stronger predictor of current WAZ than was birth WAZ. Sweet food and drink consumption at 4–6 y and maternal prepregnancy BMI were also positively related to current WAZ, and maternal prepregnancy BMI was also related to maternal BP. We did not observe a direct relation between maternal prepregnancy BMI and child SBP but we observed a significant relation between WAZ at age 4–6 y and child SBP. A recent systematic review (49) suggested that the relation between maternal prepregnancy body weight and offspring BP may be an indirect effect mediated through offspring anthropometric status, as we observed. To further understand the relation between maternal prepregnancy BMI and WAZ at age 4–6 y, we tested birth WAZ as a mediator of the relation. Birth WAZ was a significant mediator but the direct relation between maternal prepregnancy BMI and WAZ at 4–6 y also remained significant, indicating the relation between maternal prepregnancy BMI and WAZ at 4–6 y is not solely driven by WAZ at birth.

In conclusion, we did not find a difference in BP between the intervention groups at the 4–6 y follow-up in the iLiNS-DYAD-Ghana cohort, which is consistent with findings from previous trials. Our results indicate that this preschool population is at high risk of hypertension in later life. Child weight and maternal BP were directly and significantly related to child BP. Given that BP tends to remain consistent from childhood to adulthood, further follow-up of this population would be informative. These results suggest the need to evaluate the environmental and lifestyle factors that increase the risk of noncommunicable diseases in this population, in order to design effective intervention strategies.

Acknowledgments

We thank Richard Azumah and Ebenezer Adjetey for data management, and Hannah Enniful and Emelia Asamoah for the blood pressure measurements. We are grateful to Drs. Christine P Stewart and Lars F Berglund for their insightful comments and guidance while serving as dissertation committee members for SMK. We would also like to acknowledge Dr. Anna Lartey who supervised the iLiNS-DYAD-Ghana trial. The authors' responsibilities were as follows—SMK, SA-A, RRY, BMO, and KGD: designed the study; SMK, SA-A, SMT, and MEO: conducted the research; RRY: was the study statistician; RRY and SMK: performed the statistical analysis; RRY, ELP,

and KGD: advised on the analysis; SMK and KGD: wrote the manuscript; SA-A, RRY, SMT, MEO, HO, ELP, and BMO: reviewed the draft manuscript; and all authors: had primary responsibility for the final content and read and approved the final manuscript.

References

1. Kereliuk SM, Brawerman GM, Dolinsky VW. Maternal macronutrient consumption and the developmental origins of metabolic disease in the offspring. *Int J Mol Sci* 2017;18(7):1451.
2. Adair LS, Kuzawa CW, Borja J. Maternal energy stores and diet composition during pregnancy program adolescent blood pressure. *Circulation* 2001;104(9):1034–9.
3. Huxley RR, Shiell AW, Law CM. The role of size at birth and postnatal catch-up growth in determining systolic blood pressure: a systematic review of the literature. *J Hypertens* 2000;18(7):815–31.
4. Adair L, Dahly D. Developmental determinants of blood pressure in adults. *Annu Rev Nutr* 2005;25:407–34.
5. Nuyt AM, Alexander BT. Developmental programming and hypertension. *Curr Opin Nephrol Hypertens* 2009;18(2):144–52.
6. Hoy WE, Hughson MD, Bertram JF, Douglas-Denton R, Amann K. Nephron number, hypertension, renal disease, and renal failure. *J Am Soc Nephrol* 2005;16(9):2557–64.
7. Sun SS, Liang R, Huang TT, Daniels SR, Arslanian S, Liu K, Grave GD, Siervogel RM. Childhood obesity predicts adult metabolic syndrome: the Fels Longitudinal Study. *J Pediatr* 2008;152(2):191–200.
8. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004;114(2 Suppl 4th Report):555–76.
9. Noubiap JJ, Essouma M, Bigna JJ, Jingi AM, Aminde LN, Nansseu JR. Prevalence of elevated blood pressure in children and adolescents in Africa: a systematic review and meta-analysis. *Lancet Public Health* 2017;2(8):e375–e86.
10. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. *Circulation* 2008;117(25):3171–80.
11. Blumfield M, Nowson C, Hure A, Smith R, Simpson S, Raubenheimer D, MacDonald-Wicks L, Collins C. Lower protein-to-carbohydrate ratio in maternal diet is associated with higher childhood systolic blood pressure up to age four years. *Nutrients* 2015;7(5):3078.
12. Addo J, Agyemang C, Smeeth L, de-Graft Aikins A, Edusei AK, Ogedegbe O. A review of population-based studies on hypertension in Ghana. *Ghana Med J* 2012;46(2 Suppl):4–11.
13. Adu-Afarwuah S, Lartey A, Okronipa H, Ashorn P, Zeilani M, Pearson JM, Arimond M, Vosti S, Dewey KG. Lipid-based nutrient supplement increases the birth size of infants of primiparous women in Ghana. *Am J Clin Nutr* 2015;101:834–46.
14. Adu-Afarwuah S, Lartey A, Okronipa H, Ashorn P, Pearson JM, Arimond M, Ashorn U, Zeilani M, Vosti S, Dewey KG. Small-quantity, lipid-based nutrient supplements provided to women during pregnancy and 6 mo postpartum and to their infants from 6 mo of age increase the mean attained length of 18-mo-old children in semi-urban Ghana: a randomized controlled trial. *Am J Clin Nutr* 2016;104(3):797–808.
15. Arimond M, Zeilani M, Jungjohann S, Brown KH, Ashorn P, Allen LH, Dewey KG. Considerations in developing lipid-based nutrient supplements for prevention of undernutrition: experience from the International Lipid-Based Nutrient Supplements (iLiNS) Project. *Matern Child Nutr* 2015;11(Suppl 4):31–61.
16. WHO Multicentre Growth Reference Study Group. WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr Suppl* 2006;450:76–85.
17. Okronipa H, Arimond M, Young RR, Arnold CD, Adu-Afarwuah S, Tamakloe SM, Bentil HJ, Ocansey ME, Kumordzie SM, Oaks BM, et al. Exposure to a slightly sweet lipid-based nutrient supplement during early life does not increase the preference for or consumption of sweet foods and beverages by 4–6-y-old Ghanaian preschool children: follow-up of a randomized controlled trial. *J Nutr* 2019;149(3):532–41.

18. Martín-Espinosa N, Díez-Fernández A, Sánchez-López M, Rivero-Merino I, Lucas-De La Cruz L, Solera-Martínez M, Martínez-Vizcaíno V. Prevalence of high blood pressure and association with obesity in Spanish schoolchildren aged 4–6 years old. *PLoS One* 2017;12(1):e0170926.
19. Franklin SS, Lopez VA, Wong ND, Mitchell GF, Larson MG, Vasan RS, Levy D. Single versus combined blood pressure components and risk for cardiovascular disease: the Framingham Heart Study. *Circulation* 2009;119(2):243–50.
20. Vyas S, Kumaranayake L. Constructing socio-economic status indices: how to use principal components analysis. *Health Policy Plan* 2006;21(6):459–68.
21. Walker SP, Gaskin P, Powell CA, Bennett FI, Forrester TE, Grantham-McGregor S. The effects of birth weight and postnatal linear growth retardation on blood pressure at age 11–12 years. *J Epidemiol Community Health* 2001;55(6):394–8.
22. Taine M, Stengel B, Forhan A, Carles S, Botton J, Charles MA, Heude B. Rapid early growth may modulate the association between birth weight and blood pressure at 5 years in the EDEN cohort study. *Hypertension (Dallas, Tex: 1979)* 2016;68(4):859–65.
23. Steinthorsdottir SD, Eliasdottir SB, Indridason OS, Palsson R, Edvardsson VO. The relationship between birth weight and blood pressure in childhood: a population-based study. *Am J Hypertens* 2013;26(1):76–82.
24. Martínez-Aguayo A, Aglony M, Bancalari R, Avalos C, Bolte L, Garcia H, Loureiro C, Carvajal C, Campino C, Inostroza A, et al. Birth weight is inversely associated with blood pressure and serum aldosterone and cortisol levels in children. *Clin Endocrinol (Oxf)* 2012;76(5):713–8.
25. Loos RJJ, Fagard R, Beunen G, Derom C, Vlietinck R. Birth weight and blood pressure in young adults: a prospective twin study. *Circulation* 2001;104(14):1633–8.
26. Gunnarsdottir I, Birgisdottir BE, Benediktsson R, Gudnason V, Thorsdottir I. Relationship between size at birth and hypertension in a genetically homogeneous population of high birth weight. *J Hypertens* 2002;20(4):623–8.
27. Roseboom TJ, van der Meulen JH, van Montfrans GA, Ravelli AC, Osmond C, Barker DJ, Bleker OP. Maternal nutrition during gestation and blood pressure in later life. *J Hypertens* 2001;19(1):29–34.
28. Cruickshank JK, Mzayek F, Liu L, Kieleyka L, Sherwin R, Webber LS, Srinivasan SR, Berenson GS. Origins of the “black/white” difference in blood pressure: roles of birth weight, postnatal growth, early blood pressure, and adolescent body size: the Bogalusa heart study. *Circulation* 2005;111(15):1932–7.
29. Hulst AV, Barnett TA, Paradis G, Roy-Gagnon MH, Gomez-Lopez L, Henderson M. Birth weight, postnatal weight gain, and childhood adiposity in relation to lipid profile and blood pressure during early adolescence. *J Am Heart Assoc* 2017;6(8):e006302.
30. Law CM, Shiell AW, Newsome CA, Syddall HE, Shinebourne EA, Fayers PM, Martyn CN, de Swiet M. Fetal, infant, and childhood growth and adult blood pressure: a longitudinal study from birth to 22 years of age. *Circulation* 2002;105(9):1088–92.
31. Webb AL, Conlisk AJ, Barnhart HX, Martorell R, Grajeda R, Stein AD. Maternal and childhood nutrition and later blood pressure levels in young Guatemalan adults. *Int J Epidemiol* 2005;34(4):898–904.
32. Belizan JM, Villar J, Bergel E, del Pino A, Di Fulvio S, Galliano SV, Kattan C. Long-term effect of calcium supplementation during pregnancy on the blood pressure of offspring: follow up of a randomised controlled trial. *BMJ* 1997;315(7103):281–5.
33. Devakumar D, Chaube SS, Wells JC, Saville NM, Ayres JG, Manandhar DS, Costello A, Osrin D. Effect of antenatal multiple micronutrient supplementation on anthropometry and blood pressure in mid-childhood in Nepal: follow-up of a double-blind randomised controlled trial. *Lancet Glob Health* 2014;2(11):e654–63.
34. Hawkesworth S, Prentice AM, Fulford AJ, Moore SE. Maternal protein-energy supplementation does not affect adolescent blood pressure in The Gambia. *Int J Epidemiol* 2009;38(1):119–27.
35. Hawkesworth S, Sawo Y, Fulford AJ, Goldberg GR, Jarjou LM, Prentice A, Moore SE. Effect of maternal calcium supplementation on offspring blood pressure in 5- to 10-y-old rural Gambian children. *Am J Clin Nutr* 2010;92(4):741–7.
36. Hawkesworth S, Wagatsuma Y, Kahn AI, Hawlader MD, Fulford AJ, Arifeen SE, Persson LA, Moore SE. Combined food and micronutrient supplements during pregnancy have limited impact on child blood pressure and kidney function in rural Bangladesh. *J Nutr* 2013;143(5):728–34.
37. Stewart CP, Christian P, Schulze KJ, Leclercq SC, West KP, Jr, Khatri SK. Antenatal micronutrient supplementation reduces metabolic syndrome in 6- to 8-year-old children in rural Nepal. *J Nutr* 2009;139(8):1575–81.
38. Vaidya A, Saville N, Shrestha BP, de L Costello AM, Manandhar DS, Osrin D. Effects of antenatal multiple micronutrient supplementation on children’s weight and size at 2 years of age in Nepal: follow-up of a double-blind randomised controlled trial. *Lancet* 2008;371(9611):492–9.
39. Rondo PH, Pereira JA, Lemos JO. High sensitivity C-reactive protein concentrations, birthweight and cardiovascular risk markers in Brazilian children. *Eur J Clin Nutr* 2013;67(6):664–9.
40. Hoy WE, Rees M, Kile E, Mathews JD, Wang Z. A new dimension to the Barker hypothesis: low birthweight and susceptibility to renal disease. *Kidney Int* 1999;56(3):1072–7.
41. Menezes AM, Hallal PC, Horta BL, Araújo CL, de Fátima Vieira M, Neutzling M, Barros FC, Victora CG. Size at birth and blood pressure in early adolescence: a prospective birth cohort study. *Am J Epidemiol* 2007;165(6):611–6.
42. Bansal N, Ayoola OO, Gemmell I, Vyas A, Koudsi A, Oldroyd J, Clayton PE, Cruickshank JK. Effects of early growth on blood pressure of infants of British European and South Asian origin at one year of age: the Manchester Children’s Growth and Vascular Health Study. *J Hypertens* 2008;26(3):412–8.
43. de Beer M, Vrijkotte TG, Fall CH, van Eijsden M, Osmond C, Gemke RJ. Associations of infant feeding and timing of weight gain and linear growth during early life with childhood blood pressure: findings from a prospective population based cohort study. *PLoS One* 2016;11(11):e0166281.
44. Edstedt Bonamy AK, Mohlert LA, Hallberg J, Liuba P, Fellman V, Domellof M, Norman M. Blood pressure in 6-year-old children born extremely preterm. *J Am Heart Assoc* 2017;6(8):e005858.
45. Falaschetti E, Hingorani AD, Jones A, Charakida M, Finer N, Whincup P, Lawlor DA, Davey Smith G, Sattar N, Deanfield JE. Adiposity and cardiovascular risk factors in a large contemporary population of prepubertal children. *Eur Heart J* 2010;31(24):3063–72.
46. de Hoog MLA, van Eijsden M, Stronks K, Gemke RJ, Vrijkotte TG. Association between body size and blood pressure in children from different ethnic origins. *Cardiovasc Diabetol* 2012;11(1):136.
47. Miliku K, Bergen NE, Bakker H, Hofman A, Steegers EAP, Gaillard R, Jaddoe VVW. Associations of maternal and paternal blood pressure patterns and hypertensive disorders during pregnancy with childhood blood pressure. *J Am Heart Assoc* 2016;5(10):e003884.
48. Staley JR, Bradley J, Silverwood RJ, Howe LD, Tilling K, Lawlor DA, Macdonald-Wallis C. Associations of blood pressure in pregnancy with offspring blood pressure trajectories during childhood and adolescence: findings from a prospective study. *J Am Heart Assoc* 2015;4(5):e001422.
49. Ludwig-Walz H, Schmidt M, Gunther ALB, Kroke A. Maternal prepregnancy BMI or weight and offspring’s blood pressure: systematic review. *Matern Child Nutr* 2018;14(2):e12561.