

4-24-2018

Beyond the Biomarker: Understanding the Diverse Roles of Human Epididymis Protein 4 in the Pathogenesis of Epithelial Ovarian Cancer

Nicole E. James
University of Rhode Island

Clinton Chichester
University of Rhode Island, chichester@uri.edu

Jennifer R. Ribeiro

Follow this and additional works at: https://digitalcommons.uri.edu/bps_facpubs

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Citation/Publisher Attribution

James NE, Chichester C and Ribeiro JR (2018) Beyond the Biomarker: Understanding the Diverse Roles of Human Epididymis Protein 4 in the Pathogenesis of Epithelial Ovarian Cancer. *Front. Oncol.* 8:124.

<https://doi.org/10.3389/fonc.2018.00124>

Available at: <https://doi.org/10.3389/fonc.2018.00124>

This Article is brought to you for free and open access by the Biomedical and Pharmaceutical Sciences at DigitalCommons@URI. It has been accepted for inclusion in Biomedical and Pharmaceutical Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.



Beyond the Biomarker: Understanding the Diverse Roles of Human Epididymis Protein 4 in the Pathogenesis of Epithelial Ovarian Cancer

Nicole E. James^{1,2}, Clinton Chichester² and Jennifer R. Ribeiro^{1*}

¹Division of Gynecologic Oncology, Program in Women's Oncology, Department of Obstetrics and Gynecology, Women and Infants Hospital, Providence, RI, United States, ²Department of Biomedical and Pharmaceutical Sciences, University of Rhode Island, Kingston, RI, United States

OPEN ACCESS

Edited by:

Viive Maarika Howell,
University of Sydney, Australia

Reviewed by:

Raghav Sundar,
National University Health
System, Singapore
Dayanidhi Raman,
University of Toledo,
United States

*Correspondence:

Jennifer R. Ribeiro
jribeiro@wihri.org

Specialty section:

This article was submitted to
Women's Cancer,
a section of the journal
Frontiers in Oncology

Received: 15 February 2018

Accepted: 05 April 2018

Published: 24 April 2018

Citation:

James NE, Chichester C and
Ribeiro JR (2018) Beyond the
Biomarker: Understanding the
Diverse Roles of Human Epididymis
Protein 4 in the Pathogenesis of
Epithelial Ovarian Cancer.
Front. Oncol. 8:124.
doi: 10.3389/fonc.2018.00124

Human epididymis protein 4 (HE4) is an important clinical biomarker used for the detection of epithelial ovarian cancer (EOC). While much is known about the predictive power of HE4 clinically, less has been reported regarding its molecular role in the progression of EOC. A deeper understanding of HE4's mechanistic functions may help contribute to the development of novel targeted therapies. Thus far, it has been difficult to recommend HE4 as a therapeutic target owing to the fact that its role in the progression of EOC has not been extensively evaluated. This review summarizes what is collectively known about HE4 signaling and how it functions to promote tumorigenesis, chemoresistance, and metastasis in EOC, with the goal of providing valuable insights that will have the potential to aid in the development of new HE4-targeted therapies.

Keywords: human epididymis protein 4, epithelial ovarian cancer, tumorigenesis, chemoresistance, metastasis

INTRODUCTION

Approximately 22,280 new cases of epithelial ovarian cancer (EOC) are diagnosed each year, resulting in 14,240 deaths annually in the United States (1). The 5-year survival rate for stage III ovarian cancer is only 39% (1). These dire statistics are due to the fact that the disease is frequently detected at an advanced stage, which drastically impacts overall patient survival. Initially, many patients respond well to first-line therapy that includes cytoreduction surgery and platinum-based treatment. However, many patients experience a chemoresistant recurrence within the first 2 years following treatment (2). Therefore, there is an urgent need for tools to aid in the early diagnosis of ovarian cancer when the disease is fundamentally curable, as well as improved treatment options for later stage disease.

Human epididymis protein 4 (HE4) is a secretory protein that is member of the whey acidic protein domain family, bearing a conserved motif found in a number a protease inhibitors (3). HE4 was initially suggested to be involved in the innate immune defense of multiple epithelia and has also been found to function in epithelial host defense (4). In ovarian tissue, HE4 is highly over-expressed in EOC compared normal tissue (5, 6). Clinically, HE4 has been identified as a novel therapeutic biomarker for EOC and has also proven useful in detection of recurrent disease (7) Serum HE4 level predicts EOC with equal sensitivity to the established biomarker CA125 and is less likely to be elevated in benign disease (5). A multicenter study led by our institution established the

FDA-approved Risk of Ovarian Malignancy Algorithm (ROMA), which combines menopausal status and serum levels of both HE4 and CA125 to detect and monitor EOC. ROMA demonstrates improved sensitivity and specificity over the Risk of Malignancy Index that uses CA125 alone as a serum based biomarker (6). Recently, it has been reported that HE4 can be detected in EOC patient urine, indicating the possibility that it may be utilized as a non-invasive biomarker (8).

While HE4 has been well studied in the clinical setting, less is known regarding its specific molecular and biological roles in EOC. Several studies have investigated its effect on gene expression in EOC cells, as well as on events associated with aggressive disease. This review will summarize HE4's effect on cell proliferation and tumor growth; invasion, migration, and adhesion; chemoresistance; and steroid biosynthesis (**Figure 1**). Each section will detail associated pathways and factors that are reported to be involved in these HE4-mediated effects, with the goal of revealing common themes in signaling pathways affected by HE4 and exposing gaps in our knowledge of HE4 molecular and biological functions.

CELL PROLIFERATION AND TUMOR GROWTH

Within the past 5 years, a handful of *in vitro* and *in vivo* studies have begun to examine HE4's role in proliferation and tumor growth in EOC. A study by Wang et al. examined the role of HE4 in cell proliferation and found that cells treated with recombinant HE4 formed a statistically greater number of colonies compared with control treated cells (9). Furthermore, cells stimulated with recombinant HE4 exhibited greater cell viability compared with respective controls. In another study by Zhu et al. (10), proliferation rate in two different HE4-overexpressing cell lines was significantly higher than in the control cells. Likewise, Zhu et al. (11) and Lee et al. (12) determined that when HE4 was ablated *via* shRNA, cell proliferation decreased accordingly. Kong et al. report conflicting results, stating that HE4 inhibits proliferation in ovarian cells (13); however, no other studies support these claims, necessitating further explanation to understand the implications of their results.

Several *in vitro* studies suggest that HE4 promotes proliferation through its involvement in cell cycle regulation (11). Silencing of HE4 causes G0/G1 cell cycle arrest and blocks the transition from the G1 to the S phase of the cell cycle. Conversely, when cells are stimulated with recombinant HE4, the number of cells in the G2/M phase is increased, while the number of cells in the G0/G1 phase is reduced (9). These results indicate that HE4 may mediate the cell cycle by promoting the G0/G1 transition. In addition, *in vivo* tumorigenicity studies using HE4 knockdown clones revealed a marked inhibition in the growth of ovarian tumors in nude mice (14), while injection of HE4-overexpressing cells led to more aggressive tumor growth and an overall higher tumor volume compared with controls (10, 15). Taken together, results from numerous *in vitro* and *in vivo* studies provide compelling evidence that HE4 plays a role in cell proliferation and the promotion of tumorigenesis. A full list of factors associated with

HE4-mediated cell proliferation and tumor growth can be found in **Table 1A** and is outlined in greater detail below.

Associated Pathways and Factors—Cell Proliferation and Tumor Growth

Human epididymis protein 4 has been connected to several oncogenic signaling cascades that play key roles in ovarian cancer progression, including the PI3K/AKT pathway, HIF1 α , and ERK/mitogen-activated protein kinase (MAPK) signaling. Evidence of HE4's effect on activation of each of these pathways is discussed below.

Protein Kinase B Signaling

AKT has been established as a strong promoter of tumorigenesis, and the PI3K/AKT pathway is one of the most commonly hyperactivated pathways in many types of human cancers (16). Its diverse signaling regulates proliferation, growth, survival, motility, angiogenesis, and glucose metabolism (17). HE4-overexpressing OVCAR3 ovarian cancer cells were found to have a marked increase in activation of protein kinase B (AKT) compared with control cells, while HE4 knockdown in OVCAR3 cells reduced AKT activation (12). Moreover, it was found that HE4-overexpressing SKOV3 clones had naturally higher gene levels of *AKT3* compared with the null-vector control (18), bolstering the claim that HE4 affects the PI3K/AKT pathway.

Hypoxia-Inducible Factor-1 Alpha (HIF1 α)

Adaptation of malignant cells to hypoxic conditions is a key step in the promotion of tumorigenesis and angiogenesis (19–21), a process that is regulated by the transcription factor HIF1 α . Co-immunoprecipitation revealed an interaction between HIF1 α and HE4 in HE4-overexpressing SKOV3 xenografts. There was also strong colocalization of HE4 and HIF1 α in SKOV3 ovarian xenograft tissue. In addition, when SKOV3 cells were treated with HIF1 α siRNA or 2-methoxyestradiol (a HIF1 α inhibitor), there was a marked decrease in HE4 protein levels (15). It is important to note that 2-methoxyestradiol is not a specific HIF1 α inhibitor as it primarily causes the depolymerization of microtubules, which in turn prevents HIF1 α expression (22). Thus, the specificity of the effect of HIF1 α inhibition on HE4 levels may require further investigation. Although the exact mechanism and significance of the HE4-HIF1 α interaction is not understood, this evidence suggests that HE4 could play a role in regulating HIF1 α functions in angiogenesis.

MAPK Signaling

The MAPK pathway is composed of a family of conserved kinases that mediate essential cellular processes such as migration, growth, proliferation, differentiation, and apoptosis (23). The extracellular signal-regulated kinase (ERK) pathway is the best characterized of all MAPK pathways and is deregulated in approximately one-third of all cancers. Several studies have shown activation of ERK in response to HE4 treatment or overexpression, or suppression of ERK phosphorylation in response to HE4 knockdown (11, 12, 18). Using microarray analysis, Zhu et al. determined that seven genes involved in the MAPK pathway (CHUK, GADD45A, IL1A, RPS6KA1, HSPA1B, DUSP1,



FIGURE 1 | Graphical representation of clinical, *in vivo* and *in vitro* studies completed relating to HE4 and EOC, as well as associated pathways and mechanisms.

TABLE 1 | Summary of factors associated with human epididymis protein 4 (HE4) in epithelial ovarian cancer (EOC).

Gene symbol	Description	Association
(A) Cell proliferation and tumor growth		
<i>AKT</i>	Protein kinase B	<ul style="list-style-type: none"> Upregulated in overexpressing OVCAR3 HE4 cell lines (10) Decreased in response to HE4 knockdown in OVCAR3 cells (10) Upregulated in HE4 SKOV3 clones compared with vector control (16)
<i>HIF1α</i>	Hypoxia-inducible factor-1 alpha	<ul style="list-style-type: none"> Co-immunoprecipitation with HE4 SKOV3 xenograft tissue (13) Colocalization with HE4 in SKOV3 xenograft tissue (13) Treatment with 2-methoxyestradiol leads to marked decrease in HE4 (13)
<i>ERK</i>	Extracellular signal-regulated kinase	<ul style="list-style-type: none"> Decrease in p-ERK when HE4 was silenced in SKOV3 cells (9) Corresponding decrease and increase in ERK when HE4 was downregulated and overexpressed in OVCAR3 cells (10) Increase in p-ERK in SKOV3 and OVCAR8 cells with recombinant HE4 treatment (16)
<i>CHUK</i> <i>GADD45A</i>	Conserved helix-loop-helix ubiquitous kinase Growth arrest and DNA-damage-inducible protein GADD45 alpha	<ul style="list-style-type: none"> Upregulation when HE4 was induced in ES-2 cells (8)
<i>IL1A</i> <i>RPS6KA1</i> <i>HSPA1B</i> <i>DUSP1</i> <i>JUND</i>	Interleukin-1 alpha Ribosomal protein S6 kinase alpha 1 Heat shock 70 kDa protein 1B, Dual specificity protein phosphatase 1 Transcription factor JunD	
<i>EGF/EGFR</i>	Epidermal growth factor/epidermal growth factor receptor	<ul style="list-style-type: none"> Co-immunoprecipitation with HE4 SKOV3 xenograft tissue (13) Colocalization with HE4 in SKOV3 xenograft tissue (13) HE4 overexpressed in OVCAR8 cells when stimulated with recombinant protein (13) HE4 increased when inhibited by IRESSA (13)
<i>VEGF</i> <i>INS</i>	Vascular endothelial growth factor Insulin	<ul style="list-style-type: none"> HE4 overexpressed in OVCAR8 cells when stimulated with recombinant protein (13)
(B) Invasion, migration, and adhesion		
<i>MMP-9</i> <i>MMP-2</i> <i>CTSB</i>	Matrix metalloproteinase 9 Matrix metalloproteinase 9 Cathepsin B	<ul style="list-style-type: none"> Downregulated when HE4 is silenced in ovarian cell lines (9)
<i>IL1A</i>	Interleukin-1 alpha	<ul style="list-style-type: none"> Microarray results reveal correlation with HE4 levels (8, 71) One microarray reports an inverse correlation with HE4 (16)
<i>ITGβ5</i>	Integrin β 5	<ul style="list-style-type: none"> Differentially regulated by HE4 in ES-2 and CaOV3 cells (8) Correlation with HE4 in paraffin embedded ovarian human tissue (8)
<i>SDC1</i> <i>COL1A1</i> <i>DAG1</i>	Syndecan 1 Collagen type 1 alpha 1 Dystroglycan 1	<ul style="list-style-type: none"> Differentially regulated in response to HE4 (71)
<i>LAMB3</i> <i>LAMC2</i> <i>GREM1</i> <i>TNC</i> <i>SERPIND2</i>	Laminin- β 3 Laminin- γ 2 Gremlin 1 Tenascin C Serine peptidase inhibitor member 2	<ul style="list-style-type: none"> Increased expression when stimulated with recombinant HE4 in OVCAR8 cells (57)
<i>LAMA3</i>	Laminin 332 Lewis y antigen	<ul style="list-style-type: none"> Increased in response to HE4 stimulation of <i>LAMB3</i> and <i>LAMC2</i> (57) Colocalized with HE4 in human ovarian tissue (98) Immunohistochemistry stained found correlative staining with HE4 (98) Overexpression promoted HE4-mediated invasion and metastasis in <i>in vitro</i> cell lines (99) Knockdown promoted a decrease in invasion and metastatic properties of HE4 (99)
<i>SERPIN D1</i>	Heparin cofactor II	<ul style="list-style-type: none"> Upregulated in HE4-overexpressing clones and downregulated in knockout in <i>in vitro</i> lines (71) Spearman analysis revealed positive correlation with HE4 in human EOC tissue immunohistochemistry staining (71) Poor patient prognosis when levels upregulated in combination with HE4 (71)
<i>ANXA2</i>	Annexin II	<ul style="list-style-type: none"> Mass spectrometry and co-immunoprecipitation identify as strong interacting partner of HE4 (58) Gene levels co-dependent with HE4 (58) Higher along with HE4 in EOC patients with lymph node metastasis than those without (58)

(Continued)

TABLE 1 | Continued

Gene symbol	Description	Association
<i>LAMB2</i>	Laminin subunit beta-2	• Gene levels decreased in presence of HE4 knockdown cell line (58)
<i>MKN2</i>	MAP kinase-interacting serine/threonine-protein kinase 2	
(C) Chemoresistance		
<i>EGR1</i>	Early growth response protein 1	• Suppressed in overexpressing HE4 clones (16)
<i>p38</i>	p38 mitogen-activated protein kinase	• Activated in NV cells treated with cisplatin and suppressed in overexpressing HE4 clones (16)
<i>BCL2</i>	B-cell lymphoma 2	• Increased in response to recombinant HE4 <i>in vitro</i> (7)
<i>BAX</i>	bcl-2-like protein 4	• Decreased in response to recombinant HE4 <i>in vitro</i> (7)
<i>MAPT</i> <i>SEPT3</i>	Microtubule-associated protein tau Septin 3	• Upregulated in HE4-overexpressing clones (16)
<i>TUBB</i>	β -Tubulin	• Increased in response to recombinant HE4 <i>in vitro</i> (16)
<i>ERK</i> <i>AKT</i>	Extracellular signal-regulated kinase Protein kinase B	• Knockdown with HE4 lead to a reduction in cell growth and resensitization to cisplatin and paclitaxel (10)
(D) Steroid biosynthesis		
<i>FOXA2</i>	Forkhead box protein A2	• Differentially expressed in -overexpressing clones and knockouts (71)
<i>SQLE</i> <i>DHCR7</i> <i>NSDHL</i>	Squalene monoxygenase Dehydrocholesterol reductase Sterol-4-alpha-carboxylate-3 dehydrogenase	• Differentially expressed in HE4-overexpressing clones and knockouts (71)
<i>5-MC</i>	5-Methylcytosine	• Downregulated in HE4-overexpressing clones compared with wild-type SKOV3 cells and null vector (131)
<i>ESR1</i>	Estrogen/estrogen receptor	• Abolished in HE4-overexpressing clones (131) • When stimulated in H08910 cells HE4 increased in gene and protein levels. Effect not observed in SKOV3 cells (131) • Increase expression in HE4 SKOV3 knockdown (131)

and JUND) were differentially regulated in response to HE4 overexpression in ES-2 cells (10).

Activation of the MAPK/ERK pathway occurs through EGF binding of its membrane bound receptor, EGFR (24). Using co-immunoprecipitation studies in SKOV3 cells, Moore et al. found that HE4 interacts with EGFR, with a greater degree of immunoprecipitation seen in HE4-overexpressing clones than wild-type cells (15). Furthermore, ovarian xenograft tissue showed colocalization of HE4 and EGFR. In addition, when SKOV3 and OVCAR8 cells were stimulated with growth factors EGF, VEGF, and Insulin, nuclear localization of HE4 was significantly increased. Finally, when EGF was repressed by the small molecule inhibitor Iressa, relative intensity of HE4 staining was decreased in ovarian cancer cell lines. Collectively, these results provide several layers of evidence that HE4 is tied to growth factor signaling and the MAPK/ERK pathway, although further research is needed to elucidate the precise mechanisms involved.

HE4's Role in Proliferation in Other Cancers

Human epididymis protein 4 has been investigated as a putative biomarker in endometrial (25–39), lung (40–52), breast (53, 54), pancreatic (55, 56), and gastric cancer (57). While the majority of these studies examine the value of HE4 as a clinical biomarker for detecting and monitoring disease, one study investigated the molecular mechanisms of HE4 in pancreatic and endometrial cancer. Lu et al. stimulated both pancreatic and endometrial cancer cell lines with recombinant HE4 and found that cell viability,

cell growth, and DNA synthesis was increased prominently in both cancer types (56). They also report that HE4 upregulates gene expression of proliferating cell nuclear antigen (PCNA) and downregulates p21 in both cancer cell lines in a dose dependent manner. PCNA, which is expressed in the late G1/S phase of the cell cycle, is required for DNA repair, replication, cell proliferation, and cell cycle progression (58), while p21 is an important effector of tumor suppressor pathways by promoting cell cycle arrest. Specifically, p21 is able to facilitate p53-dependent G1 growth arrest (59). Therefore, results from this study highlight HE4's role in proliferation in both pancreatic and endometrial cancer and lend support to similar evidence from studies published on EOC.

INVASION, MIGRATION, AND ADHESION

Several studies have associated HE4 with metastatic properties, including invasion, migration, and adhesion of ovarian cancer cells. Lu et al. found that adhesion to a fibronectin substrate was twofold greater in SKOV3 cells overexpressing HE4 than in mock cells. In addition, a transwell migration assay demonstrated that the HE4-overexpressing clones had a 1.8-fold greater migration capacity than mock transfected cells. By contrast, immunofluorescence analysis showed that HE4 knockout clones displayed inhibited cell-spreading ability in a statistically significant fashion compared with respective controls. Furthermore, cell invasion, proliferation, and migration were significantly decreased in these clones (14). In agreement with this study, Ribeiro et al.

also found that OVCAR8 ovarian cells treated with recombinant HE4 exhibited 2.07-fold greater invasion capacity and 1.29-fold greater adhesion to a fibronectin matrix compared with untreated controls. Interestingly, there was no change in adhesion to collagen I, IV, laminin I, and fibrinogen matrices, suggesting that HE4 has a specific effect on fibronectin adhesion. Haptotaxis toward a fibronectin substrate also was increased in the ovarian cancer cells treated with recombinant HE4 by 1.72-fold (60).

Zhu et al. used wound healing and transwell invasion assays to show that HE4-overexpressing ES-2 and CaOV3 cells possess enhanced cell migration and invasion capacities. In addition, *in vivo* tail vein injection of HE4-overexpressing ES-2 cells into nude mice resulted in significantly more metastatic lung nodules than mock transfected cells (10). Using the same ovarian cancer cell lines, Zhuang et al. report the importance of HE4 interaction with annexin II (ANXA2) to promote invasion and migration *in vitro* and metastasis *in vivo* (61). Finally, Zou et al. found that knockdown of HE4 in SKOV3.ip1 cells inhibited migration and invasion (62). Taken together, these studies strongly suggest that HE4 plays a prominent role in the promotion of ovarian cancer metastasis. A full list of factors associated with HE4-mediated invasion, migration, and adhesion can be found in **Table 1B** and is outlined in greater detail below.

Associated Pathways and Factors—Invasion, Migration, and Adhesion

Human epididymis protein 4 appears to interact with numerous molecular pathways that promote metastasis in ovarian cancer. However, it is still not entirely known how HE4 affects signaling pathways and gene expression signatures to promote invasion, migration, and adhesion of ovarian cancer cells. Following is a summary of HE4-mediated molecular pathways that are involved in metastatic events in EOC.

Matrix Metalloproteinases (MMPs)

Human epididymis protein 4 has been associated with MMPs MMP-9 and MMP-2, and Cathepsin B. MMPs are a family of zinc-dependent endopeptidases that are vital for the remodeling of the extracellular matrix (63). They are expressed in almost all types of cancers and are responsible for stimulating angiogenesis, tumor growth, and metastasis (64, 65). Cathepsin B is a lysosomal cysteine protease that has been linked to cancer progression (66), specifically in signaling pathways related to angiogenesis (67). In addition, it can promote MMP activity by degrading MMP inhibitors (68). Interestingly, silencing of HE4 in ovarian cancer cells led to a decrease in protein levels of MMP-9, MMP-2, and Cathepsin B, suggesting these factors may be involved in HE4-mediated tumor promoting effects (11).

Interleukin-1 Alpha (IL1A)

Interleukin-1 alpha is a pro-inflammatory cytokine that is involved in angiogenesis and metastasis. IL1A can directly stimulate the synthesis of VEGF (69) and fibroblastic pro matrix metalloproteinase I (70, 71). IL1A causes resistance to EGFR inhibitors in both colon and head and neck cancers (72, 73). IL1A was also found to be differentially expressed in three separate microarray studies involving HE4. In two microarrays, IL1A

levels positively associated with HE4 levels (10, 74), while in one study their levels were inversely associated (18). While there may be some ambiguity as to how HE4 and IL1A are mechanistically linked, the consistent connection between IL1A with HE4 merits further investigation.

Extracellular Matrix Proteins

Integrins are a family of transmembrane proteins that are vital to ECM adhesion and play important roles in wound healing as well as the pathogenesis of cancer (75–77). Integrin $\beta 5$ (ITG $\beta 5$) gene expression was differentially regulated by HE4 in ES-2 and CaOV3 cells, which was confirmed by positive correlation of ITG $\beta 5$ and HE4 staining in paraffin embedded ovarian tissue samples (10). This finding suggests that integrin signaling is one mechanism by which HE4 can promote increased adhesion of ovarian cancer cells. However, further research is needed to clarify the mechanisms involved.

In addition to ITG $\beta 5$, three other genes related to ECM modeling—syndecan 1 (SDC1), collagen type 1 alpha 1 (COL1A1), and dystroglycan 1 (DAG1)—were more highly expressed in cells overexpressing HE4 and were downregulated in cells with HE4 knockdown (10). SDC1, also known as CD138, is an essential cell surface adhesion molecule that is responsible for maintaining cell morphology and interactions within the surrounding micro-environment (78). Loss of SDC1 in cancer cells is associated with reduced ECM adhesion and enhanced invasion and cell motility (79). Another ECM gene found to be affected by HE4 expression levels, COL1A1, is a crucial component of the ECM as it supports cartilage, bone, and tendon tissues in the body and also functions to maintain the rigidity and elasticity of tissues (80, 81). COL1A1 plays an important role in cancer, since tumor cells that express COL1A1 are able to dissociate from their surrounding stromal components, which is essential for tumor growth (81). The final ECM gene found to be affected by HE4 is DAG1, which is a cell adhesion molecule that plays a key role basement membrane assembly (82), muscle integrity (83), and the maintenance of basolateral cell adhesion in numerous epithelial tissues (84). Loss of DAG1 is associated with cancer progression (85). Taken together, these results show that HE4 is strongly interconnected with ECM related proteins, specifically those involved in the ITG $\beta 5$ signaling pathway.

Our lab has also determined that HE4 regulates several components of the extracellular matrix (60). We performed microarray analyses comparing untreated OVCAR8 wild-type cells to recombinant HE4 treated cells, and OVCAR8 cells overexpressing HE4 to null-vector control cells. Serpin peptidase inhibitor, member 2 (SERPINB2), gremlin 1 (GREM1), laminin- $\beta 3$ (LAMB3), laminin- $\gamma 2$ (LAMC2), fibroblast growth factor 5 (FGF5), and tenascin C (TNC) were all found to be significantly upregulated upon treatment with recombinant HE4. These genes encode for extracellular matrix proteins that promote cell migration and adhesion (60). Specifically, we found that HE4 upregulates LAMC2 and LAMB3 proteins in a time-dependent manner, and this increase of both factors in turn leads to an increase in laminin-332 levels (60). Laminin-332, a heterotrimer composed of LAMC2, LAMB3, and LAMA2, is an important component of the basement membrane in epithelial tissue. Abnormal increases

in its levels have been shown to promote increased invasion in cancers (86). Further evidence suggested involvement of the FAK pathway in these events. In addition, activation of matriptase, a serine protease responsible for cleaving laminin-332 in its β chain and regulating its effects on metastatic properties, increased upon *in vitro* exposure to recombinant HE4 (60). This study provides compelling evidence that HE4 is involved in basement membrane invasion and adhesion.

Lewis y Antigen

Human epididymis protein 4 undergoes glycosylation before it is secreted by ovarian cells (87), prompting Zhuang et al. to examine the relationship between HE4 glycosylation status and metastatic properties. Lewis y antigen is a glycosyl antigen that is overexpressed in ovarian cancer and has been associated with chemoresistance and poor prognosis (88–97). They determined that Lewis y antigen was present in HE4 from benign and malignant ovarian tissues, *in vitro* cancer cells, and culture medium. HE4 from ovarian cancer samples contained higher levels of Lewis y antigen than HE4 from benign tissues, and their expression co-localized in ovarian cancer tissue (98). Furthermore, when Lewis y antigen was over expressed, it promoted HE4-mediated invasion and metastasis in ovarian cancer cell lines. Conversely, when Lewis y antigen was blocked, the invasive and metastatic properties of HE4 were significantly decreased (99). Interestingly, overexpression of Lewis y antigen increased tyrosine phosphorylation of EGFR and HER/neu, which promoted cell proliferation through the PI3K/Akt and Raf/MEK/MAPK pathways (100). Thus, it appears that Lewis y antigen and HE4 affect similar signaling pathways that promote tumor growth and malignancy (101). Taken together, these results show that Lewis y antigen could be a potential therapeutic target to decrease HE4 function in the treatment of EOC.

Heparin Cofactor II (HCII)

SERPIND1 encodes for the protein HCII, which is a serum glycoprotein and protease inhibitor (102). A study in non-small cell lung cancer (NSCLC) showed that HCII promotes cell motility, invasion, and filopodium dynamics through the PI3K/AKT pathway. High HCII expression in NSCLC tissue correlated to an increased recurrence rate and shorter overall survival (103). Furthermore, its levels were upregulated in metastatic brain cell lines compared with non-metastatic parental lines, suggesting an involvement of SERPIND1 in metastatic functions (104). Results from a microarray study by Zhu et al. showed that SERPIND1 was upregulated in HE4-overexpressing cells and conversely downregulated in HE4 knockdown cells. These results were validated *via* qPCR and immunohistochemistry. In addition, they found that 37/50 ovarian cancer samples showed positive expression of both SERPIND1 and HE4, and Spearman correlation analysis confirmed that HE4 and SERPIND1 were positively correlated. Finally, Kaplan–Meier analysis revealed that patients with high levels of HE4 and SERPIND1 had a worse prognosis (74). While these data strongly suggest a connection between HE4 and SERPIND1, which may be related to their roles in promoting ovarian cancer metastasis, further study of the association between these two proteins is required.

Annexin II

Annexin II is a calcium-dependent, phospholipid-binding protein that is overexpressed in a variety of cancers and is involved in angiogenesis, proliferation, apoptosis, cell migration, invasion, and adhesion (105). High levels of Annexin II activate MAPK signaling, which in turn promotes tumor proliferation (106), invasion (107), and metastasis (108). Zhuang et al. employed mass spectrometry and co-immunoprecipitation to identify Annexin II (ANXA2) as a strong HE4 interacting partner (61). This binding promoted invasion and metastasis in ES-2 and CaOV3 ovarian cancer cells. HE4 and ANXA2 gene expression levels were found to be co-dependent, and examination of EOC tissue revealed that both HE4 and Annexin II levels were increased in malignant phenotypes compared with benign and normal ovarian tissues. Both proteins were also more highly expressed in tissues from patients with lymph node metastases than those without. Downregulation of HE4 was found to decrease expression of MKNK2 (MAP kinase-interacting serine/threonine-protein kinase 2) and LAMB2 (laminin, beta-2), two factors associated with MAPK and focal adhesion signaling pathways. When HE4 protein was supplemented, this effect was reversed. Collectively, these results show that HE4 interaction with Annexin II to activate MAPK and focal adhesion signaling is one mechanism by which HE4 may promote ovarian cancer metastasis.

CHEMORESISTANCE

Several studies show that HE4 is associated with chemoresistance clinically. The addition of HE4 serum levels in the ROMA score better predicts platinum resistance in patients than CA125 alone (15). Angioli et al. found that HE4 was able to predict chemotherapy response in EOC patients undergoing first-line therapy (109). In addition, higher levels of serum HE4 are reported in women who are resistant to first-line chemotherapy (110). Finally, higher HE4 levels inversely correlate with clinical outcome (111), optimal cytoreduction (112), progression free survival (113), and overall survival (15, 113). While the mechanism underlying HE4's contribution to chemoresistance has not been established fully, a few studies have begun to delineate HE4's role in this process. A full list of factors associated with HE4-mediated chemoresistance can be found in **Table 1C** and is outlined in detail below.

Associated Pathways and Factors— Chemoresistance Antiapoptotic Gene Expression

A study performed in our lab by Ribeiro et al. determined that HE4 overexpression promotes collateral chemoresistance to both cisplatin and paclitaxel in SKOV3 and OVCAR8 cells (18). Conversely, CRISPR/Cas9 mediated knockdown of HE4 in SKOV3 cells overexpressing HE4 partially reversed their chemoresistance. Microarray analysis revealed suppression of cisplatin-induced early growth response 1 (*EGRI*) gene expression in HE4-overexpressing SKOV3 cells compared with null vector-transfected cells (18). *EGRI* is a transcription factor that regulates apoptosis, proliferation, and differentiation through regulating expression of genes such as p53, BCL2, PTEN, IGF2,

PDGF, VEGF, TGF β 1, and TNF (114, 115). *EGR1* expression is influenced by MAPK signaling, including phospho-ERK and phospho-p38 (115). Ribeiro et al. found that p38 was strongly activated in SKOV3 null vector-transfected cells treated with cisplatin, while its activation was suppressed in HE4-overexpressing clones (18), suggesting that HE4-mediated chemoresistance may involve MAPK signaling.

Similarly, a study by Wang et al. showed that HE4 represses carboplatin-induced apoptosis *in vitro*. Recombinant HE4 caused an increase in expression of antiapoptotic protein B-cell lymphoma 2 (BCL-2) and a decrease in expression of pro-apoptotic Bax (Bcl-2 associated X protein) in SKOV3 cells treated with carboplatin (9). This decrease in the Bax/Bcl-2 ratio, in addition to the suppression of *EGR1* when HE4 is overexpressed, may contribute to the overall decrease in pro-apoptotic factors that leads to chemoresistance in EOC.

Microtubule Stabilization

Microtubule-associated protein tau, which has been associated with paclitaxel resistance in ovarian (116), breast (117), and gastric cancer (118), was upregulated in SKOV3 cells overexpressing HE4 compared with null-vector cells (18). In addition, HE4-overexpressing cells were found to express significantly higher levels of SEPT3 (Septin 3) mRNA compared with null-vector controls (18). Septins are a family of conserved GTP binding proteins that are associated with microtubules and actin filaments and have an important role in cytoskeletal organization (119). Furthermore, recombinant HE4 treatment of SKOV3 cells increased β -tubulin levels, indicating that HE4 might promote microtubule stability, leading to paclitaxel resistance.

Kinase Signaling Pathways

Human epididymis protein 4 knockdown has also been shown to lead to a reduction in cell growth and the resensitization of ovarian cancer cells to both cisplatin and paclitaxel (12). Lee et al. found that this effect was due to corresponding decreases of ERK and AKT in HE4 knockouts. Activation of these pathways suppresses apoptotic signaling in tumors, suggesting that HE4's regulation of these pathways may be an important mechanism of chemoresistance (120).

STEROID BIOSYNTHESIS

Evidence suggests an association between sex steroids and EOC pathogenesis, which is explained by processes that take place during the menstrual cycle. The ovarian surface epithelium (OSE) plays a critical role in ovulation and postovulatory wound repair. During the menstrual cycle, the OSE proliferates during the pro-estrus/estrus transition. After, ovulation the proliferation rate decreases (121). It is hypothesized that when the OSE is repeatedly exposed to high doses of luteinizing hormone and follicle stimulating hormone during the menstrual cycle, this can promote cell proliferation and increase the likelihood of tumor growth over time (121). Furthermore, epidemiological data have suggested that ovarian cancer progression, pathogenesis, and etiology are highly dependent on the activity of estrogens (121), and numerous experimental studies have demonstrated

the promotive effect of estrogens on ovarian tumors in mice and human EOC cell lines (122). However, activation of diverse oncogenic pathways in EOC may lead to the eventual downregulation of ER α levels and the overall decrease in ER α related signaling in ovarian cancers, rendering them resistant to anti-estrogen therapies (122). Some evidence exists that HE4 may be involved in this process by regulating steroid signaling in EOC. A full list of factors associated with HE4-mediated steroid biosynthesis can be found in **Table 1D** and is outlined in detail below.

Steroid Biosynthesis Gene Expression

Two separate microarray pathway analyses identified steroid biosynthesis as a pathway affected by HE4 (10, 74). Important genes that were differentially expressed between HE4-overexpressing and HE4 knockdown cell lines were Forkhead box protein A2 (FOXA2) (74), squalene monoxygenase (SQLE), 7-dehydrocholesterol reductase (DHCR7), 24-dehydrocholesterol (DHCR24), and sterol-4-alpha-carboxylate-3-dehydrogenase (NSDHL) (10). FOXA2, a transcription factor required for normal metabolism (123), promotes cell proliferation, maintains cancer stem cells, and is associated with a higher rate of relapse in triple-negative breast cancer (124).

Another gene differentially regulated by HE4, SQLE, is an enzyme required in the later stages of cholesterol synthesis (125). Out of 22 cancer types, SQLE copy number-driven gene expression was highest in breast, ovarian and colorectal cancer (125). Also affected by HE4 levels was DHCR7, one of the terminal enzymes involved in the production of cholesterol from 7-dehydrocholesterol (7DHC). DHCR7 was found to be an important regulatory determinate between cholesterol and vitamin synthesis, as cholesterol is able to accelerate the proteasomal degradation of DHCR7, which can result in the accumulation of 7DHC and an increased production of vitamin D (126). DHCR24, which was also affected by modulation of HE4 levels, is another enzyme in the cholesterol biosynthesis pathway (127). It interacts physically and functionally with DHCR7 (128) and has a number of different cellular functions including anti-inflammatory and antiapoptotic functions, as well as regulation of oxidative stress and cell differentiation (129). DHCR24 has also been proposed to be involved in tumor progression, as its deregulation has been linked to prostate, ovarian, and urothelial carcinomas (127).

Finally, NSDHL is also involved in cholesterol biosynthesis and produces metabolites that are essential in the conversion of squalene to cholesterol (130). Interestingly, NSLD1 was found to have a role in the control of signaling, vesicular trafficking, and degradation of EGFR and its dimerization partners ERBB2 and ERBB3. A study by Sukhanova et al. showed that NSLD1 knockout *in vivo* leads to a reduction in EGFR activation (131). The results from these microarrays show that modulating HE4 levels results in differential expression of several genes involved in steroid biosynthesis—especially cholesterol—suggesting that HE4 may affect tumor metabolism and ultimately contribute to tumorigenesis.

Estrogen Signaling

In support of the above described pathway analyses, two other studies have shown that HE4 interacts with steroid signaling,

specifically estrogen signaling. Lokich et al. showed that ER α expression was reduced in HE4-overexpressing SKOV3 cells, resulting in increased resistance to tamoxifen and fulvestrant compared with wild-type cells (132). 5-Methylcytosine (5-MC), a methylated form of the DNA base cytosine, is one of the most prominently identified epigenetic modifications, and can cause suppression of ER α gene expression. Deregulation of DNA methylation can result in abnormal gene expression and tumorigenesis (133, 134). Lokich et al. found that 5-MC was readily detected in SKOV3 wild-type and null-vector cells but not in HE4-overexpressing clones, suggesting that HE4 overexpression may have an effect on epigenetic modifications (132). However, methylation of the ER α gene was not specifically examined in this study. It is unclear whether HE4 overexpression would promote increased methylation at the ER α promoter region (even with the presence of global demethylation), which would be expected given the reported suppression of ER α in this study.

Interestingly, Chen et al. reported that when HO8910 ovarian cancer cells were stimulated with estradiol (E2), there was an increase in the expression of HE4 at the mRNA and protein level. This effect was not observed in estrogen-insensitive SKOV3 cells; however, when HE4 was knocked down in SKOV3 cells, their proliferative response to estrogen was restored (135). Collectively with the results shown by Lokich et al, this study suggests that HE4 works to suppress estrogen signaling in ovarian cancer cells, which can contribute to resistance to anti-estrogen therapies. Conversely, it appears that estradiol promotes HE4 expression in estrogen-responsive cells, which could indicate a role for HE4 in the initial tumor promoting effects of estrogen. Further clarification of the effect of HE4 on estrogen signaling may be useful in improving implementation of anti-estrogen based therapies.

CONCLUSION

Ovarian cancer is an extremely deadly disease owing to the fact that patients are typically diagnosed at a late stage. Initially, patients respond well to frontline platinum therapy; however, a majority of tumors recur, and the initial chemosensitivity eventually gives way to a broad chemoresistance (136). Available detection methods have improved in recent years with the discovery of HE4 as a diagnostic and prognostic biomarker. However, there

has yet to be a breakthrough targeted therapy to combat EOC. While PARP inhibitors are used in the maintenance setting for all patients, this therapy has most significantly benefited BRCA-positive patients, who comprise only 20–25% of patients (137, 138). In addition, inhibitors of immune checkpoints, such as programmed death ligand-1 have demonstrated modest benefit in clinical trials for ovarian cancer (139). Therefore, there is still a crucial need for novel targeted EOC treatments.

Although HE4 is well established as a clinical biomarker for ovarian cancer, it has been largely understudied for its therapeutic targeting potential. However, ongoing research continues to support that HE4 is profoundly involved in the pathogenesis of EOC. The individual studies mentioned in this review provide evidence that HE4 promotes EOC progression through pathways associated with cell proliferation, tumor growth, metastasis, chemoresistance, and steroid biosynthesis. These pathways, along with specific genes that have been shown to be associated with HE4, are summarized in **Table 1**. This compilation of HE4 regulated factors and pathways will serve as a starting point for scientists to further elucidate specific mechanisms by which HE4 ultimately drives tumorigenesis. In addition, a comprehensive summary of clinical, *in vivo*, and *in vitro* studies related to each facet of EOC progression and HE4 can be seen in **Figure 1**. This diagram highlights the progress that has been made to establish HE4 as an attractive therapeutic target, while simultaneously denoting areas of research that are still lacking. The results discussed here suggest that inhibition of HE4 *via* a neutralizing antibody or small molecule inhibitor could provide viable treatment options for patients in dire need of more effective therapies.

AUTHOR CONTRIBUTIONS

NJ, CC, and JR contributed conceptually to this review. All the authors reviewed and approved final manuscript.

ACKNOWLEDGMENTS

The authors would like to thank Paul DiSilvestro MD, Richard Moore, MD, and Rakesh Singh, PhD for their support and collaboration.

REFERENCES

1. Cancer facts fig. 2016. *Am Cancer Soc* (2016) 1:1–66.
2. Westin SN, Herzog TJ, Coleman RL. Investigational agents in development for the treatment of ovarian cancer. *Invest New Drugs* (2013) 31:213–29. doi:10.1007/s10637-012-9837-3
3. Bingle L, Singleton V, Bingle CD. The putative ovarian tumour marker gene HE4 (WFDC2), is expressed in normal tissues and undergoes complex alternative splicing to yield multiple protein isoforms. *Oncogene* (2002) 21:2768–73. doi:10.1038/sj.onc.1205363
4. Bingle L, Cross SS, High AS, Wallace WA, Rassl D, Yuan G, et al. WFDC2 (HE4): a potential role in the innate immunity of the oral cavity and respiratory tract and the development of adenocarcinomas of the lung. *Respir Res* (2006) 7:61. doi:10.1186/1465-9921-7-61
5. Heliström I, Raycraft J, Hayden-Ledbetter M, Ledbetter JA, Schummer M, McIntosh M, et al. The HE4 (WFDC2) protein is a biomarker for ovarian carcinoma. *Cancer Res* (2003) 63:3695–700.
6. Moore RG, McMeekin DS, Brown AK, DiSilvestro P, Miller MC, Allard WJ, et al. A novel multiple marker bioassay utilizing HE4 and CA125 for the prediction of ovarian cancer in patients with a pelvic mass. *Gynecol Oncol* (2009) 112:40–6. doi:10.1016/j.ygyno.2008.08.031
7. Piovano E, Attamante L, Macchi C, Cavallero C, Romagnolo C, Maggino T, et al. The role of HE4 in ovarian cancer follow-up: a review. *Int J Gynecol Cancer* (2014) 24:1359–65. doi:10.1097/IGC.0000000000000218
8. Sandow JJ, Rainczuk A, Infusini G, Makanji M, Bilandzic M, Wilson AL, et al. Discovery and validation of novel protein biomarkers in ovarian cancer patient urine. *Proteomics Clin Appl* (2018) 9:1700135. doi:10.1002/prca.201700135
9. Wang H, Zhu L, Gao J, Hu Z, Lin B. Promotive role of recombinant HE4 protein in proliferation and carboplatin resistance in ovarian cancer cells. *Oncol Rep* (2015) 33:403–12. doi:10.3892/or.2014.3549
10. Zhu L, Zhuang H, Wang H, Tan M, Schwab CL, Deng L, et al. Overexpression of HE4 (human epididymis protein 4) enhances proliferation, invasion and metastasis of ovarian cancer. *Oncotarget* (2015) 7:729–44. Available

- from: <http://www.impactjournals.com/oncotarget/index.php?journal=oncotarget&page=article&op=view&path%5B%5D=6327&path%5B%5D=21088> (Accessed: December 14, 2017).
11. Zhu YF, Gao GL, Tang SB, Zhang ZD, Huang QS. Effect of WFDC 2 silencing on the proliferation, motility and invasion of human serous ovarian cancer cells in vitro. *Asian Pac J Trop Med* (2013) 6:265–72. doi:10.1016/S1995-7645(13)60055-3
 12. Lee S, Choi S, Lee Y, Chung D, Hong S, Park N. Role of human epididymis protein 4 in chemoresistance and prognosis of epithelial ovarian cancer. *J Obstet Gynaecol Res* (2017) 43:220–7. doi:10.1111/jog.13181
 13. Kong X, Chang X, Cheng H, Ma R, Ye X, Cui H. Human epididymis protein 4 inhibits proliferation of human ovarian cancer cells via the mitogen-activated protein kinase and phosphoinositide 3-kinase/AKT pathways. *Int J Gynecol Cancer* (2014) 24:427–36. doi:10.1097/IGC.0000000000000078
 14. Lu R, Sun X, Xiao R, Zhou L, Gao X, Guo L. Human epididymis protein 4 (HE4) plays a key role in ovarian cancer cell adhesion and motility. *Biochem Biophys Res Commun* (2012) 419:274–80. doi:10.1016/j.bbrc.2012.02.008
 15. Moore RG, Hill EK, Horan T, Yano N, Kim K, MacLaughlan S, et al. HE4 (WFDC2) gene overexpression promotes ovarian tumor growth. *Sci Rep* (2014) 4:3574. doi:10.1038/srep03574
 16. Crowell JA, Steele VE, Fay JR. Targeting the AKT protein kinase for cancer chemoprevention. *Mol Cancer Ther* (2007) 6:2139–48. doi:10.1158/1535-7163.MCT-07-0120
 17. Testa JR, Tschlis PN. AKT signaling in normal and malignant cells. *Oncogene* (2005) 24:7391–3. doi:10.1038/sj.onc.1209100
 18. Ribeiro JR, Schorl C, Yano N, Romano N, Kim KK, Singh RK, et al. HE4 promotes collateral resistance to cisplatin and paclitaxel in ovarian cancer cells. *J Ovarian Res* (2016) 9:28. doi:10.1186/s13048-016-0240-0
 19. Maxwell PH, Wiesener MS, Chang GW, Clifford SC, Vaux EC, Cockman ME, et al. The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* (1999) 399:271–5. doi:10.1038/20459
 20. Dang CV, Semenza GL. Oncogenic alterations of metabolism. *Trends Biochem Sci* (1999) 24:68–72. doi:10.1016/S0968-0004(98)01344-9
 21. Zhong H, De Marzo AM, Laughner E, Lim M, Hilton DA, Zagzag D, et al. Overexpression of hypoxia-inducible factor 1 α in common human cancers and their metastases. *Cancer Res* (1999) 59:5830–5.
 22. Diaz-Gonzalez JA, Russell J, Rouzaut A, Gil-Bazo I, Montuenga L. Targeting hypoxia and angiogenesis through HIF-1 α inhibition. *Cancer Biol Ther* (2005) 4:1055–62. doi:10.4161/cbt.4.10.2195
 23. Dhillon AS, Hagan S, Rath O, Kolch W. MAP kinase signalling pathways in cancer. *Oncogene* (2007) 26:3279–90. doi:10.1038/sj.onc.1210421
 24. Schulze WX, Deng L, Mann M. Phosphotyrosine interactome of the ErbB-receptor kinase family. *Mol Syst Biol* (2005) 1:E1–13. doi:10.1038/msb4100012
 25. Bian J, Sun X, Li B, Ming L. Clinical significance of serum HE4, CA125, CA724, and CA19-9 in patients with endometrial cancer. *Technol Cancer Res Treat* (2016) 16:435–9. doi:10.1177/1533034616666644
 26. Gasiorowska E, Magnowska M, Izyczka N, Warchol W, Nowak-Markwitz E. The role of HE4 in differentiating benign and malignant endometrial pathology. *Ginekol Pol* (2016) 87:260–4. doi:10.17772/gp/62356
 27. Li J, Chen H, Mariani A, Chen D, Klatt E, Podratz K, et al. HE4 (WFDC2) promotes tumor growth in endometrial cancer cell lines. *Int J Mol Sci* (2013) 14:6026–43. doi:10.3390/ijms14036026
 28. Angioli R, Capriglione S, Scaletta G, Aloisi A, Miranda A, De Cicco Nardone C, et al. The role of HE4 in endometrial cancer recurrence: how to choose the optimal follow-up program. *Tumour Biol* (2016) 37:4973–8. doi:10.1007/s13277-015-4324-z
 29. Minář L, Klabeňšová I, Jandáková E. [The importance of HE4 in differential diagnosis of endometrial cancer]. *Ces Gynecol* (2015) 80:256–63.
 30. Brennan DJ, Hackethal A, Mann KP, Mutz-Dehbalalaie I, Fiegl H, Marth C, et al. Serum HE4 detects recurrent endometrial cancer in patients undergoing routine clinical surveillance. *BMC Cancer* (2015) 15:33. doi:10.1186/s12885-015-1028-0
 31. Capriglione S, Plotti F, Miranda A, Ricciardi R, Scaletta G, Aloisi A, et al. Utility of tumor marker HE4 as prognostic factor in endometrial cancer: a single-center controlled study. *Tumour Biol* (2015) 36:4156. doi:10.1007/s13277-015-3049-3
 32. Antonsen SL, Høgdall E, Christensen IJ, Lydolph M, Tabor A, Loft Jakobsen A, et al. HE4 and CA125 levels in the preoperative assessment of endometrial cancer patients: a prospective multicenter study (ENDOMET). *Acta Obstet Gynecol Scand* (2013) 92:1313–22. doi:10.1111/aogs.12235
 33. Presl J, Novotny Z, Topolcan O, Vlasak P, Kucera R, Fushova R, et al. CA125 and HE4 levels in a Czech female population diagnosed with endometrial cancer in preoperative management. *Anticancer Res* (2014) 34:327–31.
 34. Brennan DJ, Hackethal A, Metcalf AM, Coward J, Ferguson K, Oehler MK, et al. Serum HE4 as a prognostic marker in endometrial cancer – a population based study. *Gynecol Oncol* (2014) 132:159–65. doi:10.1016/j.ygyno.2013.10.036
 35. Saarelainen SK, Peltonen N, Lehtimäki T, Perheentupa A, Vuento MH, Mäenpää JU. Predictive value of serum human epididymis protein 4 and cancer antigen 125 concentrations in endometrial carcinoma. *Am J Obstet Gynecol* (2013) 209:142.e1–6. doi:10.1016/j.ajog.2013.04.014
 36. Prueksaritanond N, Cheanpracha P, Yanaranop M. Association of serum HE4 with primary tumor diameter and depth of myometrial invasion in endometrial cancer patients at Rajavithi hospital. *Asian Pac J Cancer Prev* (2016) 17:1489–92. doi:10.7314/APJCP.2016.17.3.1489
 37. Kalogera E, Scholler N, Powless C, Weaver A, Drapkin R, Li J, et al. Correlation of serum HE4 with tumor size and myometrial invasion in endometrial cancer. *Gynecol Oncol* (2012) 124:270–5. doi:10.1016/j.ygyno.2011.10.025
 38. Zamani N, Modares Gilani M, Zamani F, Zamani MH. Utility of pelvic MRI and tumor markers HE4 and CA125 to predict depth of myometrial invasion and cervical involvement in endometrial cancer. *J Family Reprod Health* (2015) 9:177–83.
 39. Bie Y, Zhang Z. Diagnostic value of serum HE4 in endometrial cancer: a meta-analysis. *World J Surg Oncol* (2014) 12:169. doi:10.1186/1477-7819-12-169
 40. Yoon HI, Kwon OR, Kang KN, Shin YS, Shin HS, Yeon EH, et al. Diagnostic value combining tumor and inflammatory markers in lung cancer. *J Cancer Prev* (2016) 21:187–93. doi:10.15430/JCP.2016.21.4.302
 41. Zeng Q, Liu M, Zhou N, Liu L, Song X. Serum human epididymis protein 4 (HE4) may be a better tumor marker in early lung cancer. *Clin Chim Acta* (2016) 455:102–6. doi:10.1016/j.cca.2016.02.002
 42. Tang QF, Zhou ZW, Ji HB, Pan WH, Sun MZ. Value of serum marker HE4 in pulmonary carcinoma diagnosis. *Int J Clin Exp Med* (2015) 8:19014–21.
 43. Li BT, Lou E, Hsu M, Yu HA, Naidoo J, Zauderer MG, et al. Serum biomarkers associated with clinical outcomes fail to predict brain metastases in patients with stage IV non-small cell lung cancers. *PLoS One* (2016) 11(1):e0146063. doi:10.1371/journal.pone.0146063
 44. Lan WG, Hao YZ, Xu DH, Wang P, Zhou YL, Ma LB. Serum human epididymis protein 4 is associated with the treatment response of concurrent chemoradiotherapy and prognosis in patients with locally advanced non-small cell lung cancer. *Clin Transl Oncol* (2016) 18:375–80. doi:10.1007/s12094-015-1375-y
 45. Lamy P-J, Plassot C, Pujol J-L. Serum HE4: an independent prognostic factor in non-small cell lung cancer. *PLoS One* (2015) 10:e0128836. doi:10.1371/journal.pone.0128836
 46. Ma Q, Wang Q, Zhong DS. Advances of human epididymis protein 4 in lung cancer. *Chin J Lung Cancer* (2015) 18:184–6. doi:10.3779/j.issn.1009-3413.2015.03.10
 47. Jiang Y, Wang C, Lv B, Ma G, Wang L. Expression level of serum human epididymis 4 and its prognostic significance in human non-small cell lung cancer. *Int J Clin Exp Med* (2014) 7:5568–72.
 48. Lou E, Johnson M, Sima C, Gonzalez-Espinoza R, Fleisher M, Kris MG, et al. Serum biomarkers for assessing histology and outcomes in patients with metastatic lung cancer. *Cancer Biomark* (2014) 14:207–14. doi:10.3233/CBM-140399
 49. Yamashita S, Tokuiishi K, Moroga T, Yamamoto S, Ohbo K, Miyahara S, et al. Serum level of HE4 is closely associated with pulmonary adenocarcinoma progression. *Tumour Biol* (2012) 33:2365–70. doi:10.1007/s13277-012-0499-8
 50. Yamashita SI, Tokuiishi K, Hashimoto T, Moroga T, Kamei M, Ono K, et al. Prognostic significance of HE4 expression in pulmonary adenocarcinoma. *Tumour Biol* (2011) 32:265–71. doi:10.1007/s13277-010-0118-5
 51. Iwahori K, Suzuki H, Kishi Y, Fujii Y, Uehara R, Okamoto N, et al. Serum HE4 as a diagnostic and prognostic marker for lung cancer. *Tumour Biol* (2012) 33:1141–9. doi:10.1007/s13277-012-0356-9

52. Tokuishi K, Yamashita SI, Ohbo K, Kawahara K. Splice variant HE4-V3 expression is associated with favorable prognosis in pulmonary adenocarcinoma. *Tumour Biol* (2012) 33:103–9. doi:10.1007/s13277-011-0252-8
53. Gündüz UR, Gunaldi M, Isiksacan N, Gündüz S, Okuturlar Y, Kocoglu H. A new marker for breast cancer diagnosis, human epididymis protein 4: a preliminary study. *Mol Clin Oncol* (2016) 5:355–60. doi:10.3892/mco.2016.919
54. Durur-Karakaya A, Durur-Subasi I, Karaman A, Akcay MN, Palabiyik SS, Erdemchi B, et al. The use of breast magnetic resonance imaging parameters to identify possible signaling pathways of a serum biomarker, HE4. *J Comput Assist Tomogr* (2016) 40:436–41. doi:10.1097/RCT.0000000000000390
55. Huang T, Jiang SW, Qin L, Senkowski C, Lyle C, Terry K, et al. Expression and diagnostic value of HE4 in pancreatic adenocarcinoma. *Int J Mol Sci* (2015) 16:2956–70. doi:10.3390/ijms16022956
56. Lu Q, Chen H, Senkowski C, Wang J, Wang X, Brower S, et al. Recombinant HE4 protein promotes proliferation of pancreatic and endometrial cancer cell lines. *Oncol Rep* (2015) 35. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26497244> (Accessed: December 15, 2017).
57. Guo YD, Wang JH, Lu H, Li XN, Song WW, Zhang XD, et al. The human epididymis protein 4 acts as a prognostic factor and promotes progression of gastric cancer. *Tumour Biol* (2015) 36:2457–64. doi:10.1007/s13277-014-2858-0
58. Thomas H, Nasim MM, Sarraf CE, Alison MR, Love S, Lambert HE, et al. Proliferating cell nuclear antigen (PCNA) immunostaining – a prognostic factor in ovarian cancer? *Br J Cancer* (1995) 71:357–62. doi:10.1038/bjc.1995.72
59. Abbas T, Dutta A. p21 in cancer: intricate networks and multiple activities. *Nat Rev Cancer* (2009) 9:400–14. doi:10.1038/nrc2657
60. Ribeiro JR, Gaudet HM, Kahn M, Schorl C, James NE, Oliver MT, et al. Human epididymis protein 4 promotes events associated with metastatic ovarian cancer via regulation of the extracellular matrix. *Front Oncol* (2018) 7:332. doi:10.3389/fonc.2017.00332
61. Zhuang H, Tan M, Liu J, Hu Z, Liu D, Gao J, et al. Human epididymis protein 4 in association with Annexin II promotes invasion and metastasis of ovarian cancer cells. *Mol Cancer* (2014) 13:1–14. doi:10.1186/1476-4598-13-243
62. Zou S-L, Chang X-H, Ye X, Cheng H-Y, Cheng Y-X, Tang Z-J, et al. Effect of human epididymis protein 4 gene silencing on the malignant phenotype in ovarian cancer. *Chin Med J (Engl)* (2011) 124:3133–40.
63. Cathcart J, Pulkoski-Gross A, Cao J. Targeting matrix metalloproteinases in cancer: bringing new life to old ideas. *Genes Dis* (2015) 2:26–34. doi:10.1016/j.gendis.2014.12.002
64. Bhowmick NA, Neilson EG, Moses HL. Stromal fibroblasts in cancer initiation and progression. *Nature* (2004) 432:332–7. doi:10.1038/nature03096
65. Kalluri R, Zeisberg M. Fibroblasts in cancer. *Nat Rev Cancer* (2006) 6:392–401. doi:10.1038/nrc1877
66. Aggarwal N, Sloane BF, Cathepsin B: multiple roles in cancer. *Proteomics Clin Appl* (2014) 8:427–37. doi:10.1002/prca.201300105
67. Lakka SS, Gondi CS, Rao JS. Proteases and glioma angiogenesis. *Brain Pathol* (2005) 15:327–41. doi:10.1111/j.1750-3639.2005.tb00118.x
68. Gondi CS, Rao JS. Cathepsin B as a cancer target. *Expert Opin Ther Targets* (2013) 17:281–91. doi:10.1517/14728222.2013.740461
69. Dabkeviciene D, Sasnauskiene A, Leman E, Kvietkauskaitė R, Daugelaviciene N, Stankevicius V, et al. MTHPC-mediated photodynamic treatment up-regulates the cytokines VEGF and IL-1alpha. *Photochem Photobiol* (2012) 88:432–9. doi:10.1111/j.1751-1097.2011.01062.x
70. Ma J, Sawai H, Matsuo Y, Ochi N, Yasuda A, Takahashi H, et al. Interleukin-1 α enhances angiogenesis and is associated with liver metastatic potential in human gastric cancer cell lines. *J Surg Res* (2008) 148:197–204. doi:10.1016/j.jss.2007.08.014
71. Löffek S, Zigrino P, Angel P, Anwald B, Krieg T, Mauch C. High invasive melanoma cells induce matrix metalloproteinase-1 synthesis in fibroblasts by interleukin-1alpha and basic fibroblast growth factor-mediated mechanisms. *J Invest Dermatol* (2005) 124:638–43. doi:10.1111/j.0022-202X.2005.23629.x
72. Gelfo V, Rodia MT, Pucci M, Dall'Orta M, Santi S, Solmi R, et al. A module of inflammatory cytokines defines resistance of colorectal cancer to EGFR inhibitors. *Oncotarget* (2016) 7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27708224> (Accessed: January 3, 2018).
73. Stanam A, Gibson-Corley KN, Love-Homan L, Ihejirika N, Simons AL. Interleukin-1 blockade overcomes erlotinib resistance in head and neck squamous cell carcinoma. *Oncotarget* (2016) 7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27738319> (Accessed: January 3, 2018).
74. Zhu L, Guo Q, Jin S, Feng H, Zhuang H, Liu C, et al. Analysis of the gene expression profile in response to human epididymis protein 4 in epithelial ovarian cancer cells. *Oncol Rep* (2016) 36:1592–604. doi:10.3892/or.2016.4926
75. Giancotti FG, Ruoslahti E. Integrin signaling. *Science* (1999) 285:1028–33. doi:10.1126/science.285.5430.1028
76. Duperret EK, Dahal A, Ridky T. Focal-adhesion-independent integrin- α regulation of FAK and c-Myc is necessary for 3D skin formation and tumor invasion. *J Cell Sci* (2015) 128:3997–4013. doi:10.1242/jcs.175539
77. Desgrosellier JS, Cheresch DA. Integrins in cancer: biological implications and therapeutic opportunities. *Nat Rev Cancer* (2010) 10:9–22. doi:10.1038/nrc2748
78. Akl MR, Nagpal P, Ayoub N, Prabhu S, Gilksman M, Tai B, et al. Molecular and clinical profiles of syndecan-1 in solid and hematological cancer for prognosis and precision medicine. *Oncotarget* (2015) 6:28693–715. doi:10.18632/oncotarget.4981
79. Teng YHF, Aquino RS, Park PW. Molecular functions of syndecan-1 in disease. *Matrix Biol* (2012) 31:3–16. doi:10.1016/j.matbio.2011.10.001
80. Bou-Gharios G, Ponticos M, Rajkumar V, Abraham D. Extra-cellular matrix in vascular networks. *Cell Prolif* (2004) 37:207–20. doi:10.1111/j.1365-2184.2004.00306.x
81. Karsenty G, Park RW. Regulation of type I collagen genes expression. *Int Rev Immunol* (1995) 12:177–85. doi:10.3109/08830189509056711
82. Matthew G, Mitchell A, Down JM, Jacobs LA, Hamdy FC, Eaton C, et al. Nuclear targeting of dystroglycan promotes the expression of androgen regulated transcription factors in prostate cancer. *Sci Rep* (2013) 3:2792. doi:10.1038/srep02792
83. Ervasti JM, Ohlendieck K, Kahl SD, Gaver MG, Campbell KP. Deficiency of glycoprotein component of the dystrophin complex in dystrophic muscle deficiency of a glycoprotein component of the dystrophin complex in dystrophic muscle. *Nature* (1990) 345:315–9. doi:10.1038/345315a0
84. Durbeej M, Larsson E, Ibraghimov-Beskrovnya O, Roberds SL, Campbell KP, Ekblom P. Non-muscle alpha-dystroglycan is involved in epithelial development. *J Cell Biol* (1995) 130:79–91. doi:10.1083/jcb.130.1.79
85. Cross SS, Lippitt J, Mitchell A, Hollingsbury F, Balasubramanian SP, Reed MWR, et al. Expression of beta-dystroglycan is reduced or absent in many human carcinomas. *Histopathology* (2008) 53:561–6. doi:10.1111/j.1365-2559.2008.03157.x
86. Tsuruta D, Kobayashi H, Imanishi H, Sugawara K, Ishii M, Jones JCR. Laminin-332-integrin interaction: a target for cancer therapy? *Curr Med Chem* (2008) 15:1968–75. doi:10.2174/092986708785132834
87. Drapkin R, Von Horsten HH, Lin Y, Mok SC, Crum CP, Welch WR, et al. Human epididymis protein 4 (HE4) is a secreted glycoprotein that is overexpressed by serous and endometrioid ovarian carcinomas. *Cancer Res* (2005) 65:2162–9. doi:10.1158/0008-5472.CAN-04-3924
88. Yin BW, Finstad CL, Kitamura K, Federici MG, Welshinger M, Kudryashov V, et al. Serological and immunochemical analysis of Lewis y (Ley) blood group antigen expression in epithelial ovarian cancer. *Int J Cancer* (1996) 65:406–12. doi:10.1002/(SICI)1097-0215(19960208)65:4<406::AID-IJC2>3.0.CO;2-0
89. Li Q, Liu S, Lin B, Yan L, Wang Y. Expression and correlation of Lewis y antigen and integrins α 5 and β 1 in ovarian serous and mucinous carcinoma. *Int J Gynecol Cancer* (2010) 20:1482–9.
90. Sabbatini PJ, Kudryashov V, Ragupath G, Danishefsky SJ, Livingston PO, Bornmann W, et al. Immunization of ovarian cancer patients with a synthetic Lewis(y)-protein conjugate vaccine: a phase I trial. *Int J Cancer* (2000) 87:79–85. doi:10.1002/1097-0215(20000701)87:1<79::AID-IJC12>3.0.CO;2-L
91. Smaletz O, Diz MD, do Carmo CC, Sabbaga J, Cunha-Junior GF, Azevedo S, et al. A phase II trial with anti-Lewis-Y monoclonal antibody (hu3S193) for the treatment of platinum resistant/refractory ovarian, fallopian tube and primary peritoneal carcinoma. *Gynecol Oncol* (2015) 138:272–7. doi:10.1016/j.ygyno.2015.05.023

92. Gao J, Hu Z, Liu J, Liu D, Wang Y, Cai M, et al. Expression of CD147 and Lewis y antigen in ovarian cancer and their relationship to drug resistance. *Med Oncol* (2014) 31:920. doi:10.1007/s12032-014-0920-9
93. Gao J, Hu Z, Liu D, Liu J, Liu C, Hou R, et al. Expression of Lewis y antigen and integrin α v, β 3 in ovarian cancer and their relationship with chemotherapeutic drug resistance. *J Exp Clin Cancer Res* (2013) 32: 36. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3699420&tool=pmcentrez&rendertype=abstract> (Accessed: January 6, 2018).
94. Zhang D, Gao J, Zhu L, Hu Z, Hou R, Liu S, et al. Chemoresistance is associated with MUC1 and lewis y antigen expression in ovarian epithelial cancers. *Int J Mol Sci* (2013) 14:11024–33. doi:10.3390/ijms140611024
95. Hu Z, Gao J, Zhang D, Liu Q, Yan L, Gao L, et al. High expression of Lewis y antigen and CD44 is correlated with resistance to chemotherapy in epithelial ovarian cancers. *PLoS One* (2013) 8:e57250. doi:10.1371/journal.pone.0057250
96. Hu Z, Gao S, Gao J, Hou R, Liu C, Liu J, et al. Elevated levels of Lewis Y and integrin α 5 β 1 correlate with chemotherapeutic drug resistance in epithelial ovarian carcinoma. *Int J Mol Sci* (2012) 13:15588–600. doi:10.3390/ijms131215588
97. Zhang F, Liu J, Lin B, Liu Q, Zhao Y, Zhu L, et al. Increase in docetaxel-resistance of ovarian carcinoma-derived RMG-1 cells with enhanced expression of Lewis Y antigen. *Int J Mol Sci* (2011) 12:7323–34. doi:10.3390/ijms12117323
98. Zhuang H, Gao J, Hu Z, Liu J, Liu D, Lin B. Co-expression of Lewis y antigen with human epididymis protein 4 in ovarian epithelial carcinoma. *PLoS One* (2013) 8:e68994. doi:10.1371/journal.pone.0068994
99. Zhuang H, Hu Z, Tan M, Zhu L, Liu J, Liu D, et al. Overexpression of Lewis y antigen promotes human epididymis protein 4-mediated invasion and metastasis of ovarian cancer cells. *Biochimie* (2014) 105:91–8. doi:10.1016/j.biochi.2014.06.022
100. Liu J-J, Lin B, Hao Y-Y, Li F-F, Liu D-W, Qi Y, et al. Lewis(y) antigen stimulates the growth of ovarian cancer cells via regulation of the epidermal growth factor receptor pathway. *Oncol Rep* (2010) 23:833–41.
101. Burotto M, Chiou VL, Lee JM, Kohn EC. The MAPK pathway across different malignancies: a new perspective. *Cancer* (2014) 120:3446–56. doi:10.1002/cncr.28864
102. Tollefsen DM. Heparin cofactor II modulates the response to vascular injury. *Arterioscler Thromb Vasc Biol* (2007) 27:454–60. doi:10.1161/01.ATV.0000256471.22437.88
103. Liao W-Y, Ho C-C, Hou H-H, Hsu T-H, Tsai M-F, Chen K-Y, et al. Heparin co-factor II enhances cell motility and promotes metastasis in non-small cell lung cancer. *J Pathol* (2015) 235:50–64. doi:10.1002/path.4421
104. Valiente M, Obenauf AC, Jin X, Chen Q, Zhang XHF, Lee DJ, et al. Serpins promote cancer cell survival and vascular co-option in brain metastasis. *Cell* (2014) 156:1002–16. doi:10.1016/j.cell.2014.01.040
105. Lokman NA, Ween MP, Oehler MK, Ricciardelli C. The role of annexin A2 in tumorigenesis and cancer progression. *Cancer Microenviron* (2011) 4:199–208. doi:10.1007/s12307-011-0064-9
106. Ortiz-Zapater E, Peiró S, Roda O, Corominas JM, Aguilar S, Ampurdanó C, et al. Tissue plasminogen activator induces pancreatic cancer cell proliferation by a non-catalytic mechanism that requires extracellular signal-regulated kinase 1/2 activation through epidermal growth factor receptor and annexin A2. *Am J Pathol* (2007) 170:1573–84. doi:10.2353/ajpath.2007.060850
107. Delys L, Detours V, Franc B, Thomas G, Bogdanova T, Tronko M, et al. Gene expression and the biological phenotype of papillary thyroid carcinomas. *Oncogene* (2007) 26:7894–903. doi:10.1038/sj.onc.1210588
108. Zhang Y, Zhou Z-H, Bugge TH, Wahl LM. Urokinase-type plasminogen activator stimulation of monocyte matrix metalloproteinase-1 production is mediated by plasmin-dependent signaling through annexin A2 and inhibited by inactive plasmin. *J Immunol* (2007) 179:3297–304. doi:10.4049/jimmunol.179.5.3297
109. Angioli R, Capriglione S, Aloisi A, Guzzo F, Luvero D, Miranda A, et al. Can HE4 predict platinum response during first-line chemotherapy in ovarian cancer? *Tumour Biol* (2014) 35:7009–15. doi:10.1007/s13277-014-1836-x
110. Chudecka-Glaz AM, Cymbaluk-Płoska AA, Menkiszak JL, Sompolska-Rzechuła AM, Tołoczko-Grabarek AI, Rzepka-Górska IA. Serum HE4, CA125, YKL-40, bcl-2, cathepsin-L and prediction optimal debulking surgery, response to chemotherapy in ovarian cancer. *J Ovarian Res* (2014) 7:62. doi:10.1186/1757-2215-7-62
111. Gonzalez VD, Samusik N, Chen TJ, Savig ES, Aghaeepour N, Quigley DA, et al. Commonly occurring cell subsets in high-grade serous ovarian tumors identified by single-cell mass cytometry. *Cell Rep* (2018) 22:1875–88. doi:10.1016/j.celrep.2018.01.053
112. Angioli R, Plotti F, Capriglione S, Aloisi A, Montera R, Luvero D, et al. Can the preoperative HE4 level predict optimal cytoreduction in patients with advanced ovarian carcinoma? *Gynecol Oncol* (2013) 128:579–83. doi:10.1016/j.ygyno.2012.11.040
113. Yuan C, Li R, Yan S, Kong B. Prognostic value of HE4 in patients with ovarian cancer. *Clin Chem Lab Med* (2018). doi:10.1515/cclm-2017-1176
114. Zhao DY, Jacobs KM, Hallahan DE, Thotala D. Silencing Egr1 attenuates radiation-induced apoptosis in normal tissues while killing cancer cells and delaying tumor growth. *Mol Cancer Ther* (2015) 15:35–7163. doi:10.1158/1535-7163.MCT-14-1051
115. Manente AG, Pinton G, Taviani D, Lopez-Rodas G, Brunelli E, Moro L. Coordinated sumoylation and ubiquitination modulate EGF induced EGR1 expression and stability. *PLoS One* (2011) 6:e25676. doi:10.1371/journal.pone.0025676
116. Gurler H, Yu Y, Choi J, Kajdacsy-Balla AA, Barbolina MV. Three-dimensional collagen type I matrix up-regulates nuclear isoforms of the microtubule associated protein tau implicated in resistance to paclitaxel therapy in ovarian carcinoma. *Int J Mol Sci* (2015) 16:3419–33. doi:10.3390/ijms16023419
117. Wang K, Deng QT, Liao N, Zhang GC, Liu YH, Xu FP, et al. Tau expression correlated with breast cancer sensitivity to taxanes-based neoadjuvant chemotherapy. *Tumour Biol* (2013) 34:33–8. doi:10.1007/s13277-012-0507-z
118. Wu H, Huang M, Lu M, Zhu W, Shu Y, Cao P, et al. Regulation of microtubule-associated protein tau (MAPT) by miR-34c-5p determines the chemosensitivity of gastric cancer to paclitaxel. *Cancer Chemother Pharmacol* (2013) 71:1159–71. doi:10.1007/s00280-013-2108-y
119. Russell SEH, Hall PA. Do septins have a role in cancer? *Br J Cancer* (2005) 93:499–503. doi:10.1038/sj.bjc.6602753
120. Cantley LC. The phosphoinositide 3-kinase pathway. *Science* (2002) 296:1655–7. doi:10.1126/science.296.5573.1655
121. Mungenast F, Thalhammer T. Estrogen biosynthesis and action in ovarian cancer. *Front Endocrinol* (2014) 5. doi:10.3389/fendo.2014.00192
122. Ribeiro JR, Freiman RN. Estrogen signaling crosstalk: implications for endocrine resistance in ovarian cancer. *J Steroid Biochem Mol Biol* (2014) 143:160–73. doi:10.1016/j.jsbmb.2014.02.010
123. Wolfrum C, Shih DQ, Kuwajima S, Norris AW, Kahn CR, Stoffel M. Role of Foxa-2 in adipocyte metabolism and differentiation. *J Clin Invest* (2003) 112:345–56. doi:10.1172/JCI18698
124. Perez-Balaguer A, Ortiz-Martinez F, Garcia-Martinez A, Pomares-Navarro C, Lerma E, Peiró G. FOXA2 mRNA expression is associated with relapse in patients with triple-negative/basal-like breast carcinoma. *Breast Cancer Res Treat* (2015) 153:465–74. doi:10.1007/s10549-015-3553-6
125. Brown DF, Caffa I, Cirmena G, Piras D, Garuti A, Gallo M, et al. Squalene epoxidase is a bona fide oncogene by amplification with clinical relevance in breast cancer. *Sci Rep* (2016) 6:19435. doi:10.1038/srep19435
126. Prabhu AV, Luu W, Sharpe LJ, Brown AJ. Cholesterol-mediated degradation of 7-dehydrocholesterol reductase switches the balance from cholesterol to vitamin D synthesis. *J Biol Chem* (2016) 291:8363–76. doi:10.1074/jbc.M115.699546
127. Dai M, Zhu XL, Liu F, Xu QY, Jiang SH, Yang XM, et al. Cholesterol synthetase DHCR24 induced by insulin aggravates cancer invasion and progesterone resistance in endometrial carcinoma. *Sci Rep* (2017) 7:41404. doi:10.1038/srep41404
128. Luu W, Hart-Smith G, Sharpe LJ, Brown AJ. The terminal enzymes of cholesterol synthesis, DHCR24 and DHCR7, interact physically and functionally. *J Lipid Res* (2015) 56:888–97. doi:10.1194/jlr.M056986
129. Zerenturk EJ, Sharpe LJ, Ikonen E, Brown AJ. Desmosterol and DHCR24: unexpected new directions for a terminal step in cholesterol synthesis. *Prog Lipid Res* (2013) 52:666–80. doi:10.1016/j.plipres.2013.09.002
130. Gabitova L, Restifo D, Gorin A, Manocha K, Handorf E, Yang DH, et al. Endogenous sterol metabolites regulate growth of EGFR/KRAS-dependent tumors via LXR. *Cell Rep* (2015) 12:1927–38. doi:10.1016/j.celrep.2015.08.023

131. Sukhanova A, Gorin A, Serebriiskii IG, Gabitova L, Zheng H, Restifo D, et al. Targeting C4-demethylating genes in the cholesterol pathway sensitizes cancer cells to EGF receptor inhibitors via increased EGF receptor degradation. *Cancer Discov* (2013) 3:96–112. doi:10.1158/2159-8290.CD-12-0031
132. Lokich E, Singh RK, Han A, Romano N, Yano N, Kim K, et al. HE4 expression is associated with hormonal elements and mediated by importin-dependent nuclear translocation. *Sci Rep* (2014) 4:5500. doi:10.1038/srep05500
133. Robertson KD. DNA methylation and human disease. *Nat Rev Genet* (2005) 6:597–610. doi:10.1038/nrg1655
134. Bird A. DNA methylation patterns and epigenetic memory. *Genes Dev* (2002) 16:6–21. doi:10.1101/gad.947102
135. Chen Y, Wang S, Liu T, Wu Y, Li J, Li M. WAP four-disulfide core domain protein 2 gene (WFDC2) is a target of estrogen in ovarian cancer cells. *J Ovarian Res* (2016) 9:10. doi:10.1186/s13048-015-0210-y
136. Cottrill H, Cason S, Caroen S, Oronsky B, Donaldson E. A recurrent platinum refractory ovarian cancer patient with a partial response after RRx-001 resensitization to platinum doublet. *J Investig Med High Impact Case Rep* (2018) 6:2324709618760080. doi:10.1177/2324709618760080
137. Liu G, Yang D, Sun Y, Shmulevich I, Xue F, Sood AK, et al. Differing clinical impact of BRCA1 and BRCA2 mutations in serous ovarian cancer. *Pharmacogenomics* (2012) 13:1523–35. doi:10.2217/pgs.12.137
138. Earl H, Molica S, Rutkowski P. Spotlight on landmark oncology trials: the latest evidence and novel trial designs. *BMC Med* (2017) 15:111. doi:10.1186/s12916-017-0884-7
139. Gaillard SL, Secord AA, Monk B. The role of immune checkpoint inhibition in the treatment of ovarian cancer. *Gynecol Oncol Res Pract* (2016) 3:11. doi:10.1186/s40661-016-0033-6

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 James, Chichester and Ribeiro. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.