

2017

Inappropriate prescribing in outpatient healthcare: an evaluation of respiratory infection visits among veterans in teaching versus non-teaching primary care clinics

Diane M. Parente
University of Rhode Island

T. Timbrook
University of Rhode Island

Aisling R. Caffrey
University of Rhode Island, aisling_caffrey@uri.edu

Kerry L. LaPlante
University of Rhode Island, kerrylaplante@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/php_facpubs

Citation/Publisher Attribution

Parente, D. M., Timbrook, T. T., Caffrey, A. R., & LaPlante, K. L. (2017). Inappropriate prescribing in outpatient healthcare: an evaluation of respiratory infection visits among veterans in teaching versus non-teaching primary care clinics. *Journal Antimicrobial Resistance & Infection Control* 2017 6:33. doi: 10.1186/s13756-017-0190-3
Available at: <http://dx.doi.org/10.1186/s13756-017-0190-3>

This Letter to the Editor is brought to you by the University of Rhode Island. It has been accepted for inclusion in Pharmacy Practice and Clinical Research Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Inappropriate prescribing in outpatient healthcare: an evaluation of respiratory infection visits among veterans in teaching versus non-teaching primary care clinics

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

LETTER TO THE EDITOR

Open Access



Inappropriate prescribing in outpatient healthcare: an evaluation of respiratory infection visits among veterans in teaching versus non-teaching primary care clinics

Diane M. Parente^{1,2}, Tristan T. Timbrook^{1,2}, Aisling R. Caffrey^{1,2,3} and Kerry L. LaPlante^{1,2,3,4*}

Abstract

A recent study led by the Centers for Disease Control and Prevention (CDC) revealed at least 30% of antibiotic prescriptions in the outpatient setting were inappropriate. In this study of all ages, among adult patients, results were similar to the overall population, with the majority of inappropriate prescribing relating to respiratory infections. We applied the same methodology to investigate rates of antibiotic prescribing for respiratory tract infections in outpatient primary care clinics at the Providence Veterans Affairs Medical Center. The results of our evaluation reflected comparable rates of inappropriate prescribing, but when stratified by teaching versus non-teaching primary care clinics, inappropriate prescribing was significantly higher in non-teaching clinics (17.6% vs 44.0%, $p < .0001$). Respiratory infection visits in non-teaching outpatient clinics may be a pragmatic target for antimicrobial stewardship programs.

Keywords: Antibiotic use, Prescribing, Outpatient, Antimicrobial stewardship

Dear Editor

We applaud Fleming-Dutra et al. on their recent analysis, which evaluated the rate of inappropriate outpatient antibiotic prescribing across several age groups and infectious diseases [1]. They found at least 30% of outpatient antibiotics prescribed were inappropriate in the United States. Inappropriate prescribing among adults, excluding children, was slightly higher at 33%, with the majority of inappropriate prescription related to acute diagnoses of respiratory conditions including sinusitis, bronchitis, and pharyngitis.

Misuse of antibiotics has been associated with increased morbidity due to adverse drug reactions, increased healthcare costs, and antibiotic resistant bacteria [2]. The Centers for Disease Control and Prevention and the White House, have declared the issue of antibiotic

resistant bacteria a threat to the health of the public. As time and resources for antimicrobial stewardship programs (ASPs) are limited, outpatient ASPs need to evaluate which interventions will ensure appropriate prescribing [3]. Currently, data on implementation and metrics of ASP initiatives in the outpatient setting are limited [4], including Veteran clinics [5]. To characterize possible opportunities for an ASP initiative, we performed an evaluation of inappropriate antibiotic prescribing among patients with respiratory tract infections based on the high rate of inappropriate prescribing with these diagnoses described by Fleming-Dutra et al.

Our Veterans Affairs (VA) Medical Center is an academic institution that provides inpatient and outpatient healthcare services to Veterans. The outpatient primary care department consists of 28 prescribers providing care during approximately 34,000 visits per year. Within the primary care department, there are two teaching clinics led by 18 medical residents under the supervision of seven attendings and six non-teaching clinics which consist of two to three primary care prescribers per clinic. Over the time period of this study there were 21

* Correspondence: kerrylaplante@uri.edu

¹Rhode Island Infectious Diseases Research Program, Providence Veterans Affairs Medical Center, 830 Chalkstone Avenue, Providence, RI 02908, USA

²Department of Pharmacy Practice, College of Pharmacy, University of Rhode Island, 7 Greenhouse Road, Kingston, RI 02881, USA

Full list of author information is available at the end of the article

Table 1 Rates of prescribing during respiratory infection visits

	Fleming-Dutra et al. ^a	VA clinics total	VA teaching clinics	VA non-teaching clinics
Sinusitis	1206/1748 (69.0)	294/357 (82.4)	38/63 (60.3)	256/294 (87.1)
Pharyngitis	824/1172 (70.3)	123/244 (50.4)	12/53 (22.6)	111/191 (58.1)
Bronchitis	748/1014 (73.8)	431/696 (61.9)	55/168 (32.7)	376/528 (71.2)
Pneumonia	324/478 (67.8)	67/233 (28.8)	15/40 (37.5)	52/193 (26.9)
Total	3102/4412 (70.3)	915/1530(59.8)	120/324 (37.0)	795/1206 (65.9)

All data as no. (%)

VA veterans affairs

^aUnweighted sample data among age ≥ 20

primary care prescribers overall in the non-teaching clinics. As part of our well-established ASP [6–8], we evaluated outpatient primary care oral antibiotic prescribing by diagnosis from March 2013 to February 2015 in Veterans who were 18 years and older. Similar to Flemming-Durta et al., we identified primary care office visits with infection diagnosis codes (ICD-9) of interest from electronic medical records. Antibiotic prescription issue dates were matched to infection diagnosis visit dates and the proportion of oral antibiotics prescribed for each infection diagnosis visit was then calculated. We calculated our rates of prescribing (Table 1), and inappropriate prescribing (Table 2) using methodology described by Flemming-Durta et al. [1] We then compared our data with that from the Flemming-Durta et al. study.

Prescribing was lower among VA clinic visits (59.8%) compared to non-federal office-based physician visits (70.3%, $p < .0001$). However, when stratified by VA teaching versus VA non-teaching clinics, teaching clinics had markedly less prescribing (37.0% vs 65.9%, $p < .0001$), and non-teaching clinics had a rate similar to that of non-federal office visits (70.3%).

Rates of inappropriate prescribing among patients with respiratory infections were similar among non-federal office-based physician visits and VA clinic visits at 34.2% and 38.4%, respectively. However, VA teaching clinics had significantly lower rates of inappropriate prescribing compared with VA non-teaching clinics (17.6% vs 44.0%, $p < .0001$) and non-federal office visits (17.6% vs 34.2%,

$p < .0001$). Our study had several limitations. As with any observational study, misclassification bias may have occurred due to diagnosis coding errors. Since we did not manually review patient electronic medical records and our reports provided limited information, we were not able to separate initial encounters from follow-up visits. It is not uncommon for a patient to be diagnosed and prescribed an antibiotic in the emergency room or hospitalized and upon discharge, follow up with their primary care provider. In addition, we were not able to observe whether patient's had other co-morbidities that may have influenced the patient's probability of getting an antibiotic or compare the patients among the different clinic types.

Nonetheless, our data offer insights into inappropriate outpatient prescribing among government clinics as those data were not available from the data source used in the Fleming-Dutra et al. study. While similar rates of inappropriate prescribing for acute respiratory tract infections have been described in two other VA health systems' outpatient clinics [9], our findings are unique in suggesting that outpatient ASPs may benefit from focusing their efforts in non-teaching clinics, at least among VA clinics. In conclusion, while VA outpatient rates of inappropriate prescribing in respiratory infections were similar to those reported at non-federal office visits, inappropriate prescribing rates among VA teaching clinics were much lower than non-federal office visits. Non-teaching clinics therefore represent an important area of opportunity for ASPs.

Table 2 Inappropriate prescribing^a during respiratory infection visits

	Fleming-Dutra et al. ^b	VA clinics total	VA teaching clinics	VA non-teaching clinics
Sinusitis	150/1748 (8.6)	78/357 (21.8)	0/63 (0)	78/294 (69.1)
Pharyngitis	613/1172 (52.3)	79/244 (32.4)	2/53 (3.8)	77/191 (40.3)
Bronchitis	748/1014 (73.8)	431/696 (61.9)	55/168 (32.7)	376/528 (71.2)
Pneumonia	0/478 (0)	0/233 (0)	0/40 (0)	0/193 (0)
Total	1511/4412 (34.2)	588/1530(38.4)	57/324 (17.6)	531/1206 (44.0)

All data as no. (%)

VA veterans affairs

^aFleming-Dutra et al. inappropriate prescribing definitions^bUnweighted sample data among age ≥ 20

Abbreviations

ASPs: Antimicrobial stewardship programs; VA: Veterans Affairs

Acknowledgements

Not applicable

Funding

The views expressed are those of the authors and do not necessarily reflect the position or policy of the United States Department of Veterans Affairs. Work supported, in part, by Dept. of VA.

Availability of data and materials

Not applicable

Authors' contributions

DMP, TTT, ARC, and KLL obtained and analyzed the data. DMP and TTT drafted the manuscript while critical revision was provided by ARC and KLL. All authors read and approved the final draft of the manuscript.

Competing interests

Diane M. Parente and Tristan T. Timbrook have no conflicts of interest to disclose. Aisling R. Caffrey has received research funding from Pfizer and Merck (Cubist). Kerry L. LaPlante has received research funding or acted as an advisor, or consultant for, Allergan, Cembra, Merck (Cubist), Davol/BARD, The Medicines Company, Ocean Spray, and Pfizer Inc.

Consent for publication

Not applicable

Ethics approval and consent to participate

A waiver of informed consent was obtained from the Providence Veterans Affairs Medical Center's Institutional Review Board for this study.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Rhode Island Infectious Diseases Research Program, Providence Veterans Affairs Medical Center, 830 Chalkstone Avenue, Providence, RI 02908, USA. ²Department of Pharmacy Practice, College of Pharmacy, University of Rhode Island, 7 Greenhouse Road, Kingston, RI 02881, USA. ³Center of Innovation in Long Term Services and Supports, Providence Veterans Affairs Medical Center, 830 Chalkstone Avenue, Providence, RI 02908, USA. ⁴Infectious Diseases Division, Warren Alpert Medical School of Brown University, 222 Richmond Street, Providence, RI 02903, USA.

Received: 2 November 2016 Accepted: 17 March 2017

Published online: 29 March 2017

References

1. Fleming-Dutra KE, Hersh AL, Shapiro DJ, et al. Prevalence of inappropriate antibiotic prescriptions among US ambulatory care visits, 2010–2011. *JAMA*. 2016;315:1864–73.
2. Centers for Disease Control and Prevention. Antibiotic Resistance Threats in the United States. CDC, 2013. <http://www.cdc.gov/drugresistance/pdf/ar-threats-2013-508.pdf>. Accessed 15 Sept 2016.
3. Barlam TF, Cosgrove SE, Abbo LM, et al. Implementing an antibiotic stewardship program. *Clin Infect Dis*. 2016;62:e51–77.
4. National Quality Forum. Antibiotic Stewardship Playbook. NQF, 2016. http://www.qualityforum.org/Publications/2016/05/Antibiotic_Stewardship_Playbook.aspx. Accessed 15 Sept 2016.
5. Drekonja DM, Filice GA, Greer N, et al. Antimicrobial stewardship in outpatient settings: a systematic review. *Infect Control Hosp Epidemiol*. 2015;36:142–52.
6. Morrill HJ, Caffrey AR, Gaitanis MM, LaPlante KL. Impact of a prospective audit and feedback antimicrobial stewardship program at a veterans affairs medical center: a six-point assessment. *PLoS ONE*. 2016;11:e0150795.
7. Morrill HJ, Gaitanis MM, LaPlante KL. Antimicrobial stewardship program prompts increased and earlier infectious diseases consultation. *Antimicrob Resist Infect Control*. 2014;3:12.

8. Morton JB, Curzake DJ, Morrill HJ, Parente DM, Gaitanis MM, LaPlante KL. Verbal communication with providers improves acceptance of antimicrobial stewardship interventions. *Infect Control Hosp Epidemiol*. 2016;37:740–2.
9. Rattinger GB, Mullins CD, Zuckerman IH, et al. A sustainable strategy to prevent misuse of antibiotics for acute respiratory infections. *PLoS ONE*. 2012;7:e51147.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

