

University of Rhode Island

DigitalCommons@URI

Ocean Engineering Faculty Publications

Ocean Engineering

2020

A NUMERICAL MODEL FOR THE EFFICIENT SIMULATION OF MULTIPLE LANDSLIDE-TSUNAMI SCENARIOS

Verdiana Iorio

Giorgio Bellotti

Claudia Cecioni

Stephan Grilli

University of Rhode Island, grilli@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/oce_facpubs

Citation/Publisher Attribution

Iorio, V., Bellotti, G., Cecioni, C., & Grilli, S. (2020). A Numerical Model for the Efficient Simulation of Multiple Landslide-Tsunami Scenarios. *Coastal Engineering*, 36. <https://doi.org/10.9753/icce.v36v.currents.20>

Available at: <https://doi.org/10.9753/icce.v36v.currents.20>

This Article is brought to you for free and open access by the Ocean Engineering at DigitalCommons@URI. It has been accepted for inclusion in Ocean Engineering Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

A NUMERICAL MODEL FOR THE EFFICIENT SIMULATION OF MULTIPLE LANDSLIDE-TSUNAMI SCENARIOS

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

A NUMERICAL MODEL FOR THE EFFICIENT SIMULATION OF MULTIPLE LANDSLIDE-TSUNAMI SCENARIOS

Verdiana Iorio, Roma Tre University, verdiana.iorio@uniroma3.it
Giorgio Bellotti, Roma Tre University, giorgio.bellotti@uniroma3.it
Claudia Cecioni, Roma Tre University, claudia.cecioni@uniroma3.it
Stephan Grilli, University of Rhode Island, grilli@uri.edu

INTRODUCTION

Submarine landslides can pose serious tsunami hazard to coastal communities, occurring frequently near the coast itself. The properties of the tsunami and the consequent inundation depend on many factors, such as the geometry, the rheology and the kinematic of the landslide and the local bathymetry. However, when evaluating the risk related to landslide tsunamis, it is very difficult to accurately predict all of the above mentioned parameters. It is therefore useful to carry out many simulations of tsunami generation and propagation, with reference to different landslide scenarios, in order to deal with such uncertainties (see for example the probabilistic approach by Grilli et al. 2009). Accurate computations of landslide tsunami generation, propagation, and inundation, however, is computationally expensive, thus limiting the possible maximum number of scenarios. To partially overcome this difficulty, in the present research, a numerical model is proposed that can efficiently compute a large number of tsunami simulations triggered by different landslides. The main goal is to provide a numerical tool that can be used in a Monte Carlo approach framework. Following the study by Ward (2001), we propose a methodology taking advantage of the linear superposition of elementary tsunami solutions.

THE NUMERICAL MODEL

The proposed method uses as wave solver the elliptic version of the mild-slope equation (MSE) with a source term representing the seafloor motion (see for example Cecioni & Bellotti, 2010). The area where the landslide can occur is split into small rectangles, as shown in Figure 1 for the test case described in the following. In each of these, a pseudo Green's function is calculated, representing the tsunami triggered by a bottom uplift of unitary magnitude over each small area. Once the actual source term is specified in terms of bottom motion for a generic landslide scenario, the tsunami solution can be quickly built by linearly superposing the effects of the elementary Green's function multiplied by the actual magnitude of the source term in each rectangle. As many preliminary computations must be carried out in order to calculate all the pseudo Green's functions, the definition of the size of the small rectangles is a crucial step in order to make the computational times acceptable.

TEST CASE

Our proposed modeling methodology was tested by reproducing the experimental study by Enet & Grilli (2007), which considers tsunamis generated by a rigid landslide in a three-dimensional tank. First, the experiments were reproduced using the MSE. Due to the fact that the employed model is linear and depth-integrated, the agreement between the numerical and

the experimental results was found to be much better in the far-field than in the near-field. Then, a parametric analysis was carried out by varying the size of the unit source rectangles. As expected, it was found that, when using relatively large rectangles, the accuracy of the solution deteriorates with respect to the benchmark.

Guidance for the selection of the rectangle size with respect to the landslide dimensions will be provided based on results obtained. The superposition method will be further tested by simulating the physical model experiments by Di Risio et al. (2009), which simulate the propagation of tsunamis generated by landslides occurring at the coast of a conical island. The method will then be applied to a real world case.




Figure 1 - Enet & Grilli (2007) test. Gray dots: computational grid. Red line: position of the landslide at the beginning of the test. Small black rectangles: unit source areas.

REFERENCES

- Cecioni, Bellotti (2010). Modeling tsunamis generated by submerged landslides using depth integrated equations, *Applied Ocean Research*, vol. 32, pp. 343-350.
- Enet, Grilli (2007). Experimental study of tsunami generation by three-dimensional rigid underwater landslides, *Journal of waterway, port, coastal and ocean engineering*, ASCE, vol. 133(6), pp. 442-454.
- Grilli, Taylor, Baxter, Maretzki (2009). A probabilistic approach for determining submarine landslide tsunami hazard along the upper east coast of the United States, *Marine Geology*, vol. 264(1-2), pp. 74-97.
- Ward (2001): Landslide tsunamis, *Journal of Geophysical Research: Solid Earth*, vol. 106(B6), pp. 11201-11215.
- Di Risio, De Girolamo, Bellotti, Panizzo, Aristodemo, Molfetta, Petrillo (2009): Landslide-generated tsunamis runup at the coast of a conical island: New physical model experiments, *Journal of Geophysical Research*, vol. 114, pp. 1-16.