2014

Homeoviscous response of *Clostridium pasteurianum* to butanol toxicity during glycerol fermentation

Keerthi P. Venkataramanan

Yogi Kurniawan
University of Rhode Island

Judy J. Boatman

Cassandra H. Haynes

Katherine A. Taconi

See next page for additional authors
Follow this and additional works at: https://digitalcommons.uri.edu/cmb_facpubs

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Venkataramanan, K. P., Kurniawan, Y., Boatman, J. J., Haynes, C. H., Taconi, K. A., Martin, L. M., Bothun, G. D., & Scholz, C. (2014). Homeoviscous response of *Clostridium pasteurianum* to butanol toxicity during glycerol fermentation. *Journal of Biotechnology*, 179, 8-14. doi: 10.1016/j.jbiotec.2014.03.017
Available at: <https://doi.org/10.1016/j.jbiotec.2014.03.017>

This Article is brought to you for free and open access by the Cell and Molecular Biology at DigitalCommons@URI. It has been accepted for inclusion in Cell and Molecular Biology Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Authors

Keerthi P. Venkataramanan, Yogi Kurniawan, Judy J. Boatman, Cassandra H. Haynes, Katherine A. Taconi, Lenore M. Martin, Geoffrey D. Bothun, and Carmen Scholz

Differential Homeoviscous Response of *Clostridium Pasteurianum* by Membrane Composition and Structural Adaptations to Butanol Toxicity

Keerthi P. Venkataramanan^{a,#}, Yogi Kurniawan^{b,#}, Judy J. Boatman^c, Casandra H. Haynes^c,
Katherine A. Taconi^c, Lenore Martin^d, Geoffrey D. Bothun^{b,*} and Carmen Scholz^{c,*}

^aBiotechnology Science and Engineering Program, University of Alabama in Huntsville, 301 Sparkman Dr NW,
MSB 333, Huntsville, AL 35899. USA.

^bDepartment of Chemical Engineering, University of Rhode Island, 16 Greenhouse Rd, 205 Crawford Hall,
Kingston, RI 02881.

^cDepartment of Chemical and Materials Engineering, University of Alabama in Huntsville, 301 Sparkman Dr NW,
EB 130, Huntsville, AL 35899. USA.

^dDepartment of Cell and Molecular Biology, University of Rhode Island, 120 Flagg Rd, 393 CBLS, Kingston, RI
02881. martin@uri.edu.

^eDepartment of Chemistry, University of Alabama in Huntsville, 301 Sparkman Dr NW, MSB 333, Huntsville, AL
35899. USA. scholz@uah.edu. Tel: (+1) 256 824 6188; Fax: (+1) 256 824 6349

#-authors KPV and YK contributed equally and are co-first authors.

*authors GB and CS are co-corresponding authors

Abstract:

Clostridium pasteurianum has been shown to ferment glycerol into butanol at higher yields than when sugars are used as the carbon source. *C. pasteurianum*'s potential to use biodiesel-derived crude glycerol as the carbon source has been gaining importance in the recent past. This study investigated the homeoviscous response of *C. pasteurianum* during butanol stress. *C. pasteurianum*'s lipid composition of the plasma membrane during butanol challenge

was analyzed. *C. pasteurianum* was found to exert two different homeoviscous responses by altering the composition of the lipid membranes in an attempt to counteract the butanol toxicity. Addition of exogenous butanol to a fermentation, when *C. pasteurianum* produced endogenous butanol, led to an increase in the ratio of saturated to unsaturated fatty acids. On the other hand, addition of exogenous butanol to fermentation, when *C. pasteurianum* did not produce any endogenous butanol, led to a decrease in the ratio of saturated to unsaturated fatty acids. This differential response for exogenous butanol during the presence and absence of an active butanol biosynthesis indicates that *C. pasteurianum* is a versatile micro-organism that has the potential to be engineered as an industrial butanol producer using crude glycerol, a promising low cost feedstock for butanol production.

Keywords

Clostridium pasteurianum; Glycerol; Fermentation; Butanol; Lipid composition; Homeoviscous response

Introduction

Clostridium pasteurianum, an anaerobic spore-forming firmicute, ferments glycerol as the sole substrate, resulting in a mixture of butanol, ethanol, and 1,3-propanediol (PDO) along with acetate and butyrate (1-4). *C. pasteurianum* is of particular interest for butanol production, as it has also been shown to ferment biodiesel-derived crude glycerol as the sole substrate (3, 4).

Butanol is toxic to cells, as it partitions into the cell membrane and affects both the structural and functional integrity of the cell. The extent of solvent toxicity correlates to the log

44 P value. Solvents with a log P value less than 4 partitions into the lipid membrane bilayer and
45 are considered extremely toxic. Butanol has a log P value of 0.8 and is considered to be one of
46 the most toxic solvents (5). When *Clostridia* are exposed to solvents, the solvents exhibit a
47 fluidizing effect on the phospholipid bilayer, which causes the organism to alter the lipid
48 composition of the bilayer. This response of bacteria to tolerate toxic solvents by altering the
49 composition of the lipid bilayer is known as homeoviscous response (membrane viscosity is
50 proportional fluidity). To compensate for the fluidizing effects of butanol, *Clostridia* increase
51 the ratio of saturated to unsaturated fatty acids (SFA/UFA) in the lipid membrane, thereby
52 reducing the fluidity of the membrane and increasing butanol tolerance (6-16). The fluidity of
53 the lipid membrane is directly proportional to the amount of saturated fatty acids in the tail of the
54 lipid bilayer (11). Hence, the bacteria that tolerate more butanol have a much higher SFA/UFA
55 ratio in the lipid bilayer (11). This has been observed to be an essential biophysical process of
56 tolerating butanol in various butanol producing organisms from the genus *Clostridium* (6-8, 10,
57 12, 13). It has been established that lipid composition and distribution on cell membrane play an
58 essential role not only in maintaining membrane stability, curvature and membrane fluidity but
59 also in modulating protein function and insertion on the membranes.(17-19) Membrane fluidity
60 is essential for maintaining the proper distribution and diffusion of embedded proteins in
61 membrane.(20)

62 Membrane lipid composition changes in response to alcohol toxicity by increasing the
63 distribution of unsaturated fatty acids observed in *Lactobacillus heterohiochii*, *Lactobacillus*
64 *homohiochii*, *Escherichia coli*, and *Saccharomyces cereeisiae*.(21-23) However, an opposite
65 effect (decreasing in the amount of saturated fatty acids) in response to alcohol toxicity is also

observed in several other microorganisms; *Clostridium acetobutylicum* (10) and *Bacillus subtilis*. Timmons et al has studied a comparison of wild-type and ethanol-adapted (EA) *Clostridium thermocellum*.(24) EA cells preserve the optimum level of fluidity in response to ethanol toxicity with increasing the fatty acids chain lengths of lipid tails and decreasing the unsaturation index in the cell membrane; resulting in higher membrane rigidity (10).

Butanol has been shown to affect the membrane by increasing fluidity and hence reducing lipid ordering(10, 25-27). Also, butanol's toxic effects lead to the formation of interdigitated phases and phase separation(25, 28). Overall, butanol can compromise cellular function of the membrane by altering cell fission, fusion, budding, vesicle formation and cell signaling (28, 29). The passive and active transport of substrates and products is also affected, along with the structure and function of integral membrane proteins. This can hinder the ability of the cell to maintain an internal pH and inhibits membrane-bound ATPases and the uptake of glucose (if present), which subsequently inhibits energy generation (9). Membrane bound ATPases are one such examples, which maintain a transmembrane pH for ATP generation. Butanol inhibits the ATPases and reduces the transmembrane pH resulting in lower ATP formation (30).

To our knowledge, the surface mechanical properties of butanol-tolerant cell membrane producing solvents have not yet been examined. In this study, the membrane extracts of *Clostridia Pasteurianum* exposed to exogenous addition of different butanol concentrations were used to investigate membrane composition, membrane phase behavior, and membrane fluidization.

88 2 **Materials and Methods**

89 2.1 **Materials**

90 All chemicals were purchased from Fisher Scientific and Sigma-Aldrich. Deuterated
91 chloroform (CDCl_3) was obtained from Cambridge Isotope Laboratory. Synthetic lipids, 1,2-
92 Dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC, >99% purity) and 1,2-dioleoyl-*sn*-glycero-3-
93 phosphocholine (DOPC, >99% purity) were obtained from Avanti Polar Lipids, Inc.

94 2.2 **Bacterial Strain**

95 The bacterial strain *C. pasteurianum* ATCC 6013 was purchased from American Type
96 Culture Collection and the glycerol stock was maintained as described earlier [8].

97 2.3 **Effect of n-Butanol**

98 The effect of n-butanol (n-butanol was used throughout all experiments) on the bacterial growth
99 and the stability and change in composition of the membrane were studied by adding butanol to
100 the media containing either glucose or glycerol as the sole carbon source. For investigating the
101 effect of butanol, exogenous butanol was added to the media varied from 0 to 1% (w/v) (0 gL^{-1}
102 to 10 gL^{-1}) containing either glycerol (25 gL^{-1}) or glucose (50 gL^{-1}) in Biebl media [6] were used
103 to study the effect of exogenous butanol during an active butanol biosynthesis on glycerol and its
104 absence when grown in glucose, as *C. pasteurianum* does not produce butanol when growing on
105 glucose. The cells were allowed to grow in the presence of butanol for 24 hours after which the
106 membrane was extracted. All experiments were conducted with 10% (v/v) inoculum, pre-grown
107 in RCM.

108

109 2.4 **Extraction of the Cell Membrane**

Cell membranes were extracted using the modified protocol of Bligh and Dyer using dichloromethane/methanol mixtures [29]. The cells were harvested (0.5 mL cell suspension) by centrifugation at 13000 rpm for 15 minutes and the pellets were resuspended in 0.5 mL of sterile 1.0% (w/v) NaCl in a 10 mL glass sample tube with PTFE lined caps. To the resuspended pellets, 2 mL of dichloromethane/methanol mixture (1:2 v/v) was added and shaken vigorously for 15 minutes. It was followed by a 2 hour incubation at RT, followed by centrifugation at 2500 rpm and the supernatant (S1) was collected in a fresh tube. The pellet was again resuspended in 0.5 mL 1.0% NaCl and 2 mL of a dichloromethane/methanol mixture (2:1 v/v) was added to the resuspended pellet and shaken vigorously for 15 minutes. Following a 2 hour incubation, the samples were centrifuged at 2500 rpm and the supernatant (S2) was collected. Supernatants S1 and S2 were combined and 1 mL of dichloromethane and 1 mL of sterile 1.0% NaCl were added. The top phase (aqueous) was removed and the bottom (organic) phase was retained. The solvent was evaporated under a gentle nitrogen stream. Once a dry film was obtained, the headspace was flushed with nitrogen, capped tightly, and stored for further analysis.

2.5 NMR Analysis

The dry film of the membrane was dissolved in CDCl_3 for ^1H -NMR analysis. Synthetic lipids, DPPC and DOPC were used as standards for NMR analysis. Various ratios of DPPC and DOPC (1:0, 3:1, 1:1, 1:3 and 0:1) in a total lipid concentration of 10 mM were used for calibration (Supplementary Material and Figure S1). The synthetic lipids were dried and dissolved in CDCl_3 . All ^1H -NMR spectra were recorded on a VarianTM Unity Inova 500 (500 MHz) spectrometer equipped with a 5 mm triple resonance inverse detectable probe. The

percentage of unsaturation in the membrane lipid samples was calculated by multiplying the ratio of the intensity of the peak corresponding to the olefinic hydrogen (at 5.31 ppm) to total intensity of all the peaks from the lipid sample by 100 (Eq 1) [30].

$$\%unsaturation = \frac{Intensity_{olefinic}}{Intensity_{total}} \times 100 \quad (\text{Eq 1})$$

2.6 Liposome Preparation and Fluorescence Anisotropy

The dry film of the reconstituted cell membrane and 0.1 mM 1,6-diphenyl 1,3,5-hexatriene (DPH) dissolved in chloroform were mixed and co-evaporated under a gentle stream of nitrogen until a dry reconstituted cell membrane/DPH film remained. Vacuum was used (~ 60 min) to remove residual solvent from the film. The film was then hydrated with distilled water and maintained at 50°C in a water bath for 1 hour before shaking. The reconstituted membrane/DPH film was suspended as multilamellar liposomes by vigorously shaking for approximately 1 hour. The liposomes were sonicated before use for 60 min at 50°C, which has been previously shown to yield unilamellar liposomes. DPPC liposomes at 10 mM lipid generated by the same procedure were used as control. Fluorescence anisotropy (Perkin Elmer LS 55) and melting temperature were measured using the L-format configuration with DPH as the hydrophobic bilayer probe from 25 to 50°C at a rate of 1°C/min under continuous mixing as described previously (25).

2.5 GC/MS Analysis

The lipid samples were methylated for GC/MS analysis. The dried lipid samples were

153 saponified using 1 mL of 3N sodium hydroxide at 90°C for an hour and then cooled to RT. The
154 excess sodium hydroxide was neutralized with 1.8 mL of 3.6 N hydrochloric acid at 90 °C for 10
155 minutes and cooled to RT. The free fatty acids were extracted using 1 mL of hexane and diethyl
156 ether (1:1 v/v). The organic phase was then separated into a clean and dry round bottom flask.
157 The fatty acid hydrolyzates were dried using a rotoevaporator and stored under Argon in
158 desiccators.

159 The dried fatty acid hydrolyzates were derivatized using a 5 mL borontrifluoride-
160 methanol complex at 60 °C for 5 minutes and then cooled to RT. To the cooled solution, 1 mL
161 water and 1 mL hexane were added and the container shaken multiple times to ensure the
162 transfer of esters into the non-polar solvent. The upper organic layer was removed and
163 transferred into an Erlenmeyer flask containing 5 g of anhydrous sodium sulfate. The flask was
164 incubated at RT to dry overnight. The sodium sulfate was filtered and the hexane solution was
165 transferred into a clean round bottom flask and dried in a rotoevaporator and the samples were
166 stored under Argon. For GC-MS analysis, samples were dissolved in 250 µL of
167 dichloromethane.

168 Fatty acid methyl esters (FAME) derivatives were analyzed by a Shimadzu GC-MS QP
169 2010 system using a SHR5xLB silica capillary column (30m× 0.25mm ID, composed of 100%
170 dimethyl polysiloxane). Manufacturers protocol were followed for FAME analysis. The
171 compounds were analyzed with GC-MS solution and identified by comparison with the data in
172 the NIST libraries. The distribution of FAMEs of *C. Pasteurianum* was determined by dividing
173 area counts for each FAME species by the total FAME area, as defined below:

$$\% \text{ FAME} = \frac{\text{Area}_{\text{FAME}}}{\sum \text{FAME}_{\text{area}}} \times 100 \quad (\text{Eq 2})$$

2.6 Surface pressure-area (Π -A) isotherms

Π -A isotherms were conducted in a Langmuir trough (model 102M, Nima technology Ltd, UK) with a deposition area of 70 cm² at 25°C, 37°C and at 50°C. A Wilhelmy plate was used to measure surface pressure with an accuracy of $\pm 1 \mu\text{N/m}$ connected to an electronic microbalance. Isotherms were monitored by NIMA TR 7.4 .vi software. The external water bath system was used to control the sub phase temperature. Monolayers were obtained by spreading diluted solutions of dry lipid sample in chloroform at the air/water interface using a 5 μl microsyringe.

The chloroform was allowed to evaporate and the lipid films to spread at air/water for 15 min. The system was the equilibrated for 15 min before being compressed at a speed of 10 cm²/min.

2.7 Elastic modulus / area compressibility modulus

Mechanical properties of the monolayer films were determined by the compressibility modulus. The elastic modulus (C_s^{-1}) / area compressibility modulus is the reciprocal of the compressibility (C_s). (31) The elastic modulus corresponds to the elasticity of the Langmuir films under the compression force. C_s^{-1} values were defined as:

$$C_s^{-1} = -A \left(\frac{\partial \pi}{\partial A} \right)_T \quad (\text{Eq 3})$$

where, A is the area per molecule at the particular surface pressure and Π is the corresponding surface pressure. The maximum compressibility or elasticity ($C_{s\text{Max}}^{-1}$) provides information on the onset of the plastic region, and the maximum packing condition of monolayer.

Results and Discussion

Clostridial species produce various metabolites in the form of acids and solvents, and have to modify the composition of their lipid membranes to tolerate the toxic effects of the produced metabolites. The fundamental studies to identify the effect of the toxic metabolites on the lipid composition of during the homeoviscous response can be investigated using the tools of chemical biology. This study is focused on investigating the effect of butanol in *C. pasteurianum* leading to the tolerance response involving changes in the lipid membrane composition using tools to analyze the physical and structural compositions of the membranes along with the membrane phase behavior.

Effect of Exogenous Butanol

The homeoviscous response of *C. pasteurianum* to the addition of exogenous butanol was studied under two conditions, where the first condition involved the addition of exogenous butanol during the production of butanol by *C. pasteurianum* during glycerol fermentation (EB1), while the second condition involved the addition of exogenous butanol while no butanol was produced, during the fermentation of glucose (EB2). *C. pasteurianum* cultures do not undergo solventogenesis during the fermentation of glucose, as the fermentation is predominantly only in the acidogenic phase, resulting in butyric acid as the major fermentation

211 product with no butanol formation.

212 Butanol was added at concentrations of 0 g/L (control), 2.5 g/L (0.0335 M), 5 g/L (0.067
213 M), 7.5 g/L (0.105 M) and 10 g/L (0.134 M), respectively, once the cells reached mid
214 exponential phase. The membranes were extracted after 24 hours of butanol exposure and
215 analyzed using ^1H -NMR, fluorescent anisotropy and GC-MS . The ^1H -NMR and fluorescent
216 anisotropy results are summarized in Figure 1, while Figure 2 summarizes the data from the GC-
217 MS analysis on the fatty acid composition of the membranes.

218 The addition of butanol during glycerol fermentation resulted in a homeoviscous
219 response due to a decrease in the degree of unsaturation in the membrane as determined by ^1H -
220 NMR (Figure 1) of the extracted lipid of cells exposed to different butanol concentrations. The
221 fluorescent anisotropy of the reconstituted *C. pasteurianum* cell membranes supports the results
222 obtained from ^1H -NMR. The fluorescence anisotropy data at 37°C shows a decrease in
223 anisotropy with increase in the butanol dose. The anisotropy, $\langle r \rangle$, of the unexposed control was
224 found to be 0.120 and $\langle r \rangle$ was found to reduce as the butanol dose increased in the media. The
225 anisotropy of the membrane had a relatively constant value with a minor increase at higher
226 concentrations of butanol at 7.5 g/L and 10 g/L. The fluorescence anisotropy of DPH in the
227 membranes is inversely related to the fluidity of the reconstituted lipid membranes. The decrease
228 in the anisotropy indicates the fluidizing effect of butanol on the membrane lipids. The
229 stabilization of the anisotropy at higher butanol concentrations can be explained as the response
230 of the bacteria to tolerate the fluidizing effects of butanol. The overlap of the anisotropy data
231 with the ^1H -NMR data substantiates the existence of a dominant homeoviscous response leading
232 to a compositional change in the lipid membranes. This can be further confirmed by analyzing

233 the fatty acid composition of the membrane.

234 The data from GC-MS on the fatty acid composition (Figure 2) of the membranes
235 corroborates the presence of the homeoviscous response to butanol. Furthermore, there was an
236 increase in the percentage of higher carbon saturated fatty acids (C₁₉ to C₂₂), which are almost
237 completely absent in the control samples with no external butanol present. A similar trend was
238 also observed in the disappearance or reduction in the shorter chain fatty acids with an increase
239 in the dose of butanol stress (Figure 2). The shorter length fatty acids C₁₀, were completely
240 absent in the cells exposed to higher butanol concentrations, while the other shorter fatty acids
241 from C₁₁ to C₁₆ were found to decrease proportionately with increase in butanol dose. Similarly,
242 the longer fatty acids, C₁₉ to C₂₂, were not observed in the control which was not exposed to
243 butanol and had a proportionate increase in the percentage with corresponding increase in the
244 butanol dosage. The compositional analysis through GC-MS not only supports the results from
245 NMR and fluorescence anisotropy, but also substantiates that *C. pasteurianum* exerts a
246 homeoviscous response to butanol in EB1 condition by two major changes in the fatty acid
247 composition to counteract the fluidity of the toxic butanol. First, it increases the percentage of
248 longer chain fatty acids at the expense of shorter chain fatty acids and secondly by increasing the
249 ratio of saturated to unsaturated fatty acids.

250 The effect of butanol on the lipid composition and the fluidity of the lipid membrane has
251 also been reported for *C. acetobutylicum* (6, 7, 10, 12). *C. acetobutylicum* produces butanol by
252 fermenting glucose and the effect of butanol challenge was studied during the growth of *C.*
253 *acetobutylicum* in glucose (butanol producing media) (6, 7, 10, 12). *C. acetobutylicum* tolerates
254 butanol through a homeoviscous response that predominantly involves an increase of saturated

fatty acids at the expense of unsaturated fatty acids in the lipid membrane (6, 7, 10, 12). Lepage et al reported the composition of the *C. acetobutylicum*'s fatty acid composition in the lipid membrane, which consisted of fatty acids from C₁₂ to C₁₉ (10). The ratio of the unsaturated to saturated fatty acids was found to be close to 1 without any butanol exposure but the ratio was reduced to 0.87 and 0.77 respectively with an exposure to 4 g/L and 8 g/L butanol respectively, due to the reduction in unsaturation and an increase in saturated fatty acids in the membrane lipids, driven by a homeoviscous response.

As a control, the change in the degree of unsaturation and anisotropy of the lipid membrane during the production of endogenous butanol was studied and compared (Supplementary material and Figure S2 and S3). The addition of butanol during glycerol fermentation results in a similar result as observed during the endogenous butanol production. The only change was observed in the anisotropy data which accounts for the fluidizing effect of the exogenous butanol on the membrane. The difference observed in the anisotropy data is consistent with the previously reported fluidizing effect of butanol observed through fluorescence anisotropy of synthetic lipids and reconstituted membrane lipids (8, 25).

The Differential Response to Butanol Toxicity

The effect of butanol during EB1 showed a conventional homeoviscous response by increasing the fatty acid length and the ratio of saturated to unsaturated fatty acids. To further investigate the sole effect of butanol toxicity the cells grown in glucose were exposed to butanol (EB2). Initially, the EB2 experiment was conducted to match the same butanol stress dosage from 0 g/L to 10 g/L but was also performed at higher concentrations of butanol of up to 20 g/L.

As explained earlier, the extracted membranes were studied using ^1H -NMR and fluorescent anisotropy. Figure 3 summarizes the change in the fluidity of the membrane in terms of anisotropy and the degree of unsaturation in the membranes of *C. pasteurianum* grown on glucose. The ^1H -NMR results indicated an increase in the percentage of unsaturation in the fatty acid tails of the lipid membranes extracted from the cells stressed with exogenous butanol. This result is completely contrasting to the results obtained earlier during butanol stress on glycerol fermentation. The degree of unsaturation in the lipid membranes increased proportionately with increasing concentration of butanol in the media. For an exogenous butanol concentration of 5 g/L, a small drop in the percentage of unsaturation in the fatty acid tail was observed when compared to the control with no external butanol. The data obtained from fluorescent anisotropy of the reconstituted membrane, using DPH as the probe, supported the data obtained from ^1H -NMR (Figure 3). The anisotropy, $\langle r \rangle$, measured at 37°C decreased gradually with an increase in the concentration of butanol in the media.

The lipid membranes extracted from the cells were also analyzed by GC-MS to determine the constituent fatty acids. Figure 3 summarizes the results from GC-MS analysis of the cells stressed with exogenous butanol. Thirteen different fatty acids were identified using GC-MS, of which 11 were saturated. The data on the composition of the fatty acids from Figure 4 shows not only an increase in the degree of unsaturation in the fatty acids, but also an increase in the percentage of fatty acid length ($\geq \text{C}_{16}$) with an increase in the concentration of exogenous butanol.

The degree of unsaturation was found to increase (Figure 3 and 4) in the presence of exogenous butanol, but without endogenous butanol production (cells grown on glucose). An

298 increase in unsaturation coincided with an increase in membrane fluidity (Figure 4). This is in
299 contrast with the response observed for EB1(Figure 2).

300 It has been shown that an increase in the percentage of saturated lipids in the model
301 liposomes (model membranes from DPPC and DOPC) results in an increase in the anisotropy of
302 the liposomes (25). Furthermore, the addition of butanol to model membranes comprised of
303 DPPC, DOPC or mixture of the two have shown a decrease in anisotropy due to the fluidizing
304 effects of butanol on the lipid membrane (25, 29). Butanol fluidizes membranes by reducing
305 inter lipid interactions and the surface tension within the membrane. Increase in the degree of
306 unsaturation in lipid membranes has been shown to augment the fluidizing effects of butanol
307 (25). Hence, the results obtained from the addition of exogenous butanol to the cells of *C.*
308 *pasteurianum* in EB2 contradict the results obtained during EB1. This led to the question of
309 whether other non-lipid entities could be involved in the homeoviscous adaptation of the
310 membrane in EB2.

311 The fatty acid composition of the cells also varied considerably when the cells are grown
312 on glycerol and glucose . The cells grown on glucose and exposed to exogenous butanol show
313 neither a decrease in shorter (C₆ to C₁₀) fatty acids nor an increase in longer (C₁₉ and greater)
314 fatty acids (Figure 4). The fatty acid composition of the cells are also different under the two
315 condition, glucose and glycerol fermentation. The fatty acid chain length is as low as 6 carbons
316 during glucose fermentation, while the the lowest fatty acid in glycerol fermentation is 10.
317 Similar distinction is also observed in the maximum chain length as well for the two carbon
318 sources. Moreover, the percentage of unsaturated fatty acids increased with increasing
319 concentrations of exogenous butanol during EB2 (Figure 4).

320 The reduction in shorter chain fatty acids and an increase in longer chain fatty acids in *C.*
321 *acetobutylicum* resulted in a decrease of membrane fluidity (6, 10). It was also observed in *C.*
322 *acetobutylicum* that challenging the cells with butanol resulted in the formation of longer chain
323 fatty acids at the expense of shorter chain fatty acids (6, 10).

324 The increase in unsaturation (exogenous butanol on glucose) observed with all three
325 different analytical methods cannot be explained by a homeoviscous adaptation, as it led to an
326 increase in the fluidity of the membrane consistent with the toxic effect of butanol (Figure 3 and
327 4). However, an increase in unsaturation in the lipid membrane of cells exposed to solvent stress
328 has been reported for *E. coli* during ethanol stress and for *C. butyricum* during 1,3-PDO stress
329 (14, 32). Dombek and Ingram reported that the plasma membrane became more rigid during
330 ethanol challenge experiments, but the extracted lipid membrane exhibited higher fluidity in
331 comparison to the control cells that were unexposed to ethanol. This shows that the rigidity of
332 the membrane is not only dependent on the ratio of saturated to unsaturated fatty acids, or the
333 presence of shorter or longer fatty acids, but is also dependent on the lipid to protein ratio in the
334 membrane (34). Membrane proteins can rigidify the membrane despite irrespective of the lipid
335 composition. Hence, an increase in protein:lipid ratio can account for the net increase in the
336 rigidity of the membrane and compensates for the increase in fluidity as a result of the increased
337 unsaturation of the lipid tails (Figure 3).

338 During the formation of butanol (EB1), the bacteria must be synthesizing butanol efflux
339 pumps in the membrane that serve as butanol transporters to the extracellular environment (33).
340 Dunlop *et al.* have shown that cloning and expressing of efflux pumps from different
341 microorganisms for various solvents resulted in an increase of solvent tolerance. But,

overexpression of butanol and iso-butanol efflux pumps did not improve butanol tolerance leading to a conclusion that the toleration of butanol is a much more complex phenomenon (33). There should also be a correlation between the transcriptional regulation of the genes involved in butanol production and the genes responsible for butanol tolerance through homeoviscous adaptation. If there was no correlation, the homeoviscous adaptation of *C. pasteurianum* must be similar during both, endogenous production (from glycerol) and exogenous addition (growth on glucose).

***II-A* isotherms of reconstituted cell membrane (RCM) monolayers**

To gain more understanding of the interaction within lipid-lipid and lipid-protein, it is essential to first characterize the phase behavior of the RCM monolayers spread at the air/water interface. The *II-A* isotherms of RCM is plotted in Figure 5. Three different temperatures (25, 37, and 50 °C) and sample extract at different conditions (without butanol and with 10 g/l butanol) have been examined. Without exogenous butanol in media at 25°C, the experimental isotherms exhibited a liquid expanded (LE) phase and a collapse phase. The trends in *II-A* isotherms were slightly different with those of unsaturated phospholipids(34) and to *E. coli* lipid extract(17). *II* of RCM monolayers was much lower than the unsaturated lipids and the *E. coli* lipid extract and lift off area was not observable on RCM monolayers. The collapse of the RCM monolayers took place at ~25 mN/m indicating that the monolayers had low viscosity due to strong lipid interaction and a plausible displacement of relaxation membrane protein. At 37°C RCM monolayers isotherms existed in single *LE* phase and collapse phase disappeared that could be attributed to the increase of disordered lipid tails and disordered organic membrane protein lowering lipid-protein interactions in membrane monolayers. With increasing temperature up to

50 °C, Π - A isotherms occurred at lower pressure. The appearance of the lift-off area was able to observe. A plateau region on Π - A isotherms indicates that the G - LE phase transition occurred. Increases temperature lowered hydrogen bonding between lipid head groups and hydrophobic interactions between lipid tails and extended the relaxation phenomena in membrane proteins.

In addition, the length of plateau region could be associated with the ratio of unsaturated/saturated lipid (U/S). A short plateau region displayed as a high ratio of saturated lipid. The addition of 10 g/l butanol in media had markedly different effects on the Π - A isotherms (Figure 5) and the maximum elastic modulus, C_{Max}^{-1} (Figure 6). With temperature changes (25, 37, and 50 °C) Π - A isotherms of RCM monolayer existed in the LE phase and occurred at higher surface pressure than without butanol present in media. This reveals that the content of *hydrophobic proteins* (Supplementary material and Figure S4) adsorbed *at air/water interface* had increased creating a higher Π - A and an increased C_{Max}^{-1} (more rigid). The C_{Max}^{-1} decreased proportionally with increasing temperature due to disordering structure of lipid and expanded relaxation phenomena.

Based on our interpretation data, the effect of protein can be interpreted as shown in Figure 7. At low surface pressure ($\Pi < \Pi_{Collapse}^{protein}$) 25°C, hydrophobic membrane protein from cell grew in media without butanol exhibited relaxation phenomena within lipid membrane monolayers and water molecules were between the hydrophilic head groups of lipids. As a result, repulsive contribution at polar headgroup was high, and attractive van der Waals interactions between tails of lipids and interfacial interaction at head/tail of tails were low promoting a weak

384 molecular packing. At high surface pressure, $\Pi > \Pi_{\text{Collapse}}^{\text{protein}}$ external pressure compressed the
385 protein network until it failed and displaced from the interface, forming collapse protein
386 multilayers in the aqueous bulk phase near to the interface; thus, the collapse phase appeared. In
387 the case of the reconstituted cell membranes extracted from cells grew in media with butanol, the
388 content of hydrophobic membrane protein was increased resulted in longer relaxation
389 phenomena. Therefore, the collapse pressure appeared higher. It is clear that homeoviscous *C.*
390 *pasteurianum* producing solvents altered the lipid membrane composition to modulate protein
391 function and insertion on the membranes in determining membrane integrity and activity.

392

393 4. Conclusion

394 This study is the first report of the analysis of the homeoviscous adaptation to butanol by
395 *C. pasteurianum* and this study is also the first study that quantifies lipid composition in *C.*
396 *pasteurianum*. The analysis of the extracted membranes from cells exposed to various
397 concentration of butanol in two different media, butanol producing medium containing glycerol
398 and non-butanol producing medium containing glucose, resulted in two completely different
399 responses with respect to the changes in the composition of the lipids. During butanol
400 production, when stressed with exogenous butanol, *C. pasteurianum* responds by increasing the
401 ratio of the saturated to unsaturated fatty acids in the membrane. This is also the response found
402 in other butanol producing species of *Clostridia*, along with the response exhibited by other
403 microbes towards tolerating toxic organic compounds (14-16, 35). *C. pasteurianum* exhibits a
404 different response when grown on glucose (no butanol produced), but challenged with exogenous

butanol, which can be explained on the hypothesis of altering the ratio of the protein to lipids in the membrane (Figure 7 and 8). This hypothesis was further confirmed using by *II-A* isotherms and the presence of hydrophobic membrane proteins in the lipid membrane of *C. pasteurianum* during EB2.

At 25°C *II-A* isotherms of the membrane monolayer were observed in the *LE* phase and the collapse phase. The collapse phase on membrane monolayers was observed because of a greater attractive interaction of saturated lipids and a displacement of lipid monolayers on collapse protein network at the interface. At 50°C *II-A* isotherms membrane monolayer exhibited a *G-LE* phase transition and *LE* phase. The *G-LE* phase transition appeared due to a low interaction of lipid-lipid and lipid-protein associating with disordering protein and lipid structure. This study implies the effect of unsaturated lipids and proteins on membrane monolayers and helps our understanding of the mechanism of butanol tolerant membrane by *C. pasteurianum* could be utilized to develop cultures in higher concentrations of butanol leading to more cost-effective and more efficient on butanol production.

The result from this study substantiates the assumption that a correlation exists between the modes of homeoviscous response, which are in turn, dependent on the activation of butanol production pathway. Furthermore, the existence of two different homeoviscous adaptations to butanol challenge in *C. pasteurianum*, demonstrates the potential of this organism to be studied further in terms of proteomics, functional genomics and metabolic engineering for the development of an industrial strain. The unavailability of the genome sequence and the proteome data must be addressed to explore butanol production and tolerance in *C. pasteurianum*. In the meantime, methods to use genomic data from closely related species can

427 be explored to establish a platform that can be used to perform transcriptional analysis of *C.*
428 *pasteurianum* during butanol stress.

429 Acknowledgements

430 This project was supported by NSF grant (CBET-0966846). We sincerely thank Dr.
431 Vinka Craver, University of Rhode Island, for the GC-MS.

432 References

- 433 1. Biebl, H. (2001) Fermentation of glycerol by *Clostridium pasteurianum*--batch and
434 continuous culture studies, *Journal of industrial microbiology & biotechnology* 27, 18-
435 26.
- 436 2. Dabrock, B., Bahl, H., and Gottschalk, G. (1992) Parameters Affecting Solvent
437 Production by *Clostridium pasteurianum*, *Applied and environmental microbiology* 58,
438 1233-1239.
- 439 3. Taconi, K. A., Venkataramanan, K. P., and Johnson, D. T. (2009) Growth and solvent
440 production by *Clostridium pasteurianum* ATCC® 6013™ utilizing biodiesel-derived
441 crude glycerol as the sole carbon source, *Environ. Prog. Sustainable Energy* 28, 100-110.
- 442 4. Venkataramanan, K. P., Boatman, J. J., Kurniawan, Y., Taconi, K. A., Bothun, G. D., and
443 Scholz, C. (2012) Impact of impurities in biodiesel-derived crude glycerol on the
444 fermentation by *Clostridium pasteurianum* ATCC 6013, *Applied microbiology and*
445 *biotechnology* 93, 1325-1335.
- 446 5. Sardessai, Y., and Bhosle, S. (2002) Tolerance of bacteria to organic solvents, *Research in*
447 *microbiology* 153, 263-268.
- 448 6. Baer, S. H., Blaschek, H. P., and Smith, T. L. (1987) Effect of Butanol Challenge and
449 Temperature on Lipid-Composition and Membrane Fluidity of Butanol-Tolerant
450 *Clostridium-Acetobutylicum*, *Applied and environmental microbiology* 53, 2854-2861.
- 451 7. Baer, S. H., Bryant, D. L., and Blaschek, H. P. (1989) Electron-Spin Resonance Analysis
452 of the Effect of Butanol on the Membrane Fluidity of Intact-Cells of *Clostridium-*
453 *Acetobutylicum*, *Applied and environmental microbiology* 55, 2729-2731.
- 454 8. Baut, F., Fick, M., Viriot, M. L., Andre, J. C., and Engasser, J. M. (1994) Investigation of
455 Acetone-Butanol-Ethanol Fermentation by Fluorescence, *Applied microbiology and*

- 456 *biotechnology* 41, 551-555.
- 457 9. Bowles, L. K., and Ellefson, W. L. (1985) Effects of Butanol on Clostridium-
458 Acetobutylicum, *Applied and environmental microbiology* 50, 1165-1170.
- 459 10. Lepage, C., Fayolle, F., Hermann, M., and Vandecasteele, J. P. (1987) Changes in
460 Membrane Lipid-Composition of Clostridium-Acetobutylicum during Acetone Butanol
461 Fermentation - Effects of Solvents, Growth Temperature and Ph, *Journal of general*
462 *microbiology* 133, 103-110.
- 463 11. Liu, S. Q., and Qureshi, N. (2009) How microbes tolerate ethanol and butanol, *New*
464 *biotechnology* 26, 117-121.
- 465 12. Vollherbst-Schneck, K., Sands, J. A., and Montenecourt, B. S. (1984) Effect of butanol on
466 lipid composition and fluidity of Clostridium acetobutylicum ATCC 824, *Applied and*
467 *environmental microbiology* 47, 193-194.
- 468 13. Isar, J., and Rangaswamy, V. (2012) Improved n-butanol production by solvent tolerant
469 Clostridium beijerinckii, *Biomass Bioenerg* 37, 9-15.
- 470 14. Heipieper, H. J., Weber, F. J., Sikkema, J., Keweloh, H., and Debont, J. A. M. (1994)
471 Mechanisms of Resistance of Whole Cells to Toxic Organic-Solvents, *Trends in*
472 *biotechnology* 12, 409-415.
- 473 15. Lăzăroaie, M. Mechanisms Involved In Organic Solvent Resistance in Gram-Negative
474 Bacteria.
- 475 16. Ramos, J. L., Duque, E., Gallegos, M.-T., Godoy, P., Ramos-González, M. I., Rojas, A.,
476 Terán, W., and Segura, A. (2002) MECHANISMS OF SOLVENT TOLERANCE IN
477 GRAM-NEGATIVE BACTERIA, *Annual Review of Microbiology* 56, 743-768.
- 478 17. Lopez-Montero, I., Arriaga, L. R., Monroy, F., Rivas, G., Tarazona, P., and Velez, M.
479 (2008) High fluidity and soft elasticity of the inner membrane of Escherichia coli
480 revealed by the surface rheology of model Langmuir monolayers, *Langmuir* 24, 4065-
481 4076.
- 482 18. Lingwood, D., Kaiser, H. J., Levental, I., and Simons, K. (2009) Lipid rafts as functional
483 heterogeneity in cell membranes, *Biochem Soc Trans* 37, 955-960.
- 484 19. Simons, K., and Ikonen, E. (1997) Functional rafts in cell membranes, *Nature* 387, 569-
485 572.
- 486 20. Lenaz, G. (1987) Lipid fluidity and membrane protein dynamics, *Biosci Rep* 7, 823-837.
- 487 21. Beaven, M. J., Charpentier, C., and Rose, A. H. (1982) Production and Tolerance of
488 Ethanol in Relation to Phospholipid Fatty-Acyl Composition in Saccharomyces-

- 489 Cerevisiae Ncyc 431, *Journal of General Microbiology* 128, 1447-1455.
- 490 22. Ingram, L. O. (1982) Regulation of Fatty-Acid Composition in Escherichia-Coli - a
491 Proposed Common Mechanism for Changes Induced by Ethanol, Chaotropic Agents, and
492 a Reduction of Growth Temperature, *Journal of Bacteriology* 149, 166-172.
- 493 23. Uchida, K. (1974) Occurrence of Saturated and Mono-Unsaturated Fatty-Acids with
494 Unusually Long Chains (C20-C30) in Lactobacillus Heterohiochii, an Alcoholophilic
495 Bacterium, *Biochimica Et Biophysica Acta* 348, 86-93.
- 496 24. Timmons, M. D., Knutson, B. L., Nokes, S. E., Strobel, H. J., and Lynn, B. C. (2009)
497 Analysis of composition and structure of Clostridium thermocellum membranes from
498 wild-type and ethanol-adapted strains, *Applied Microbiology and Biotechnology* 82, 929-
499 939.
- 500 25. Kurniawan, Y., Venkataramanan, K. P., Scholz, C., and Bothun, G. D. (2012) n-Butanol
501 partitioning and phase behavior in DPPC/DOPC membranes, *The journal of physical*
502 *chemistry. B* 116, 5919-5924.
- 503 26. Aguilar, L. F., Sotomayor, C. P., and Lissi, E. A. (1996) Main phase transition depression
504 by incorporation of alkanols in DPPC vesicles in the gel state: Influence of the solute
505 topology, *Colloid Surface A* 108, 287-293.
- 506 27. Iiyama, S., Toko, K., Murata, T., Ichinose, H., Suezaki, Y., Kamaya, H., Ueda, I., and
507 Yamafuji, K. (1992) Cutoff Effect of N-Alkanols in an Excitable Model Membrane
508 Composed of Dioleoyl Phosphate, *Biophysical chemistry* 45, 91-100.
- 509 28. Ceve, G. (1982) Water and Membranes - the Interdependence of Their Physicochemical
510 Properties in the Case of Phospholipid-Bilayers, *Stud Biophys* 91, 45-52.
- 511 29. Lobbecke, L., and Ceve, G. (1995) Effects of Short-Chain Alcohols on the Phase-
512 Behavior and Interdigitation of Phosphatidylcholine Bilayer-Membranes, *Bba-*
513 *Biomembranes* 1237, 59-69.
- 514 30. Papoutsakis, E. T. (2008) Engineering solventogenic clostridia, *Current opinion in*
515 *biotechnology* 19, 420-429.
- 516 31. Duncan, S. L., and Larson, R. G. (2008) Comparing experimental and simulated pressure-
517 area isotherms for DPPC, *Biophys J* 94, 2965-2986.
- 518 32. Dombek, K. M., and Ingram, L. (1984) Effects of ethanol on the Escherichia coli plasma
519 membrane, *Journal of bacteriology* 157, 233-239.
- 520 33. Dunlop, M. J., Dossani, Z. Y., Szmids, H. L., Chu, H. C., Lee, T. S., Keasling, J. D., Hadi,
521 M. Z., and Mukhopadhyay, A. (2011) Engineering microbial biofuel tolerance and export
522 using efflux pumps, *Molecular systems biology* 7.

- 523 34. Lucero, A., Nino, M. R. R., Gunning, A. P., Morris, V. J., Wilde, P. J., and Patino, J. M. R.
524 (2008) Effect of hydrocarbon chain and pH on structural and topographical characteristics
525 of phospholipid monolayers, *Journal of Physical Chemistry B* 112, 7651-7661.
- 526 35. Segura, A., Duque, E., Mosqueda, G., Ramos, J. L., and Junker, F. (2002) Multiple
527 responses of Gram-negative bacteria to organic solvents, *Environmental microbiology* 1,
528 191-198.
- 529
- 530