

2020

Long-term exposure to PM_{2.5} and incidence of disability in activities of daily living among oldest old

Yuebin Lv

Jinhui Zhou

Virginia Byers Kraus

Tiantian Li

Jeremy A. Sarnat

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/hdf_facpubs

Citation/Publisher Attribution

Lv, Y., Zhou, J., Kraus, V. B., Li, T., Sarnat, J. A., Wang, J., Liu, Y., Chen, H., Brasher, M. S., Mao, C., Zeng, Y., Zheng, T., & Shi, X. (2020). Long-term exposure to PM_{2.5} and incidence of disability in activities of daily living among oldest old. *Environmental pollution (Barking, Essex : 1987)*, 259, 113910. <https://doi.org/10.1016/j.envpol.2020.113910>

Available at: <https://doi.org/10.1016/j.envpol.2020.113910>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Human Development and Family Science Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Long-term exposure to PM_{2.5} and incidence of disability in activities of daily living among oldest old

Authors

Yuebin Lv, Jinhui Zhou, Virginia Byers Kraus, Tiantian Li, Jeremy A. Sarnat, Jiaonan Wang, Yang Liu, Huashuai Chen, Melanie S. Brasher, Cheng Mao, Yi Zeng, and Xiaoming Shi

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).



HHS Public Access

Author manuscript

Environ Pollut. Author manuscript; available in PMC 2021 April 01.

Published in final edited form as:

Environ Pollut. 2020 April ; 259: 113910. doi:10.1016/j.envpol.2020.113910.

Long-term exposure to PM_{2.5} and incidence of disability in activities of daily living among oldest old

Yuebin Lv^a, Jinhui Zhou^a, Virginia Byers Kraus^b, Tiantian Li^a, Jeremy A Sarnat^c, Jiaonan Wang^d, Yang Liu^e, Huashuai Chen^f, Melanie Sereny Brasher^g, Chen Mao^h, Yi Zeng^{f,i}, Tongzhang Zheng^j, Xiaoming Shi^{a,d,*}

^aNational Institute of Environmental Health, Chinese Center for Disease Control and Prevention, Beijing, 100021, China

^bDuke Molecular Physiology Institute and Division of Rheumatology, Department of Medicine, Duke University School of Medicine, Durham, NC 27710, USA

^cRollins School of Public Health, Emory University, Atlanta, GA 30322, USA

^dCenter for Global Health, School of Public Health, Nanjing Medical University, Nanjing, Jiangsu, China

^eDepartment of Environmental and Occupational Health, Emory University, Atlanta, GA 30322, USA

^fCenter for the study of Aging and Human Development and the Geriatric Division of School of Medicine, Duke University, Durham, NC 27708, USA

^gDepartment of Sociology and Anthropology; Department of Human Development and Family Studies, University of Rhode Island, Kingston, RI 02881, USA

* **Correspondence to:** Xiaoming Shi, PhD, National Institute of Environmental Health, Chinese Center for Disease Control and Prevention, #7 Panjiayuan Nanli, Chaoyang, Beijing 100021, China; shixm@chinacdc.cn; Telephone: (+86) 1050930101; Fax: (+86) 1058900247.

Author statement

Conceptualization: Yuebin Lv, Xiaoming Shi

Data curation: Yuebin Lv, Xiaoming Shi, Jinhui Zhou

Formal analysis: Yuebin Lv, Xiaoming Shi

Funding acquisition: Xiaoming Shi, Yi Zeng

Investigation: Yuebin Lv, Jinhui Zhou, Virginia Byers Kraus, Huashuai Chen, Chen Mao, Yi Zeng, Xiaoming Shi

Methodology: Yuebin Lv, Xiaoming Shi

Project administration: Yuebin Lv, Jinhui Zhou, Yi Zeng, Xiaoming Shi

Resources: Xiaoming Shi

Software: Yuebin Lv, Xiaoming Shi

Supervision: Xiaoming Shi

Validation: Yuebin Lv, Xiaoming Shi, Tiantian Li, Jeremy A Sarnat, Jiaonan Wang, aYang Liu, Melanie Sereny Brashe

Visualization: Yuebin Lv, Jinhui Zhou, Jiaonan Wang

Roles/Writing - original draft: Yuebin Lv, Xiaoming Shi

Writing - review & editing: All authors

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declarations of Interest

None.

^hDivision of Epidemiology, School of Public Health, Southern Medical University, Guangzhou, 510515, China

ⁱCenter for Study of Healthy Aging and Development Studies, Peking University, Beijing, 100871, China

^jEpidemiology Department, Brown University, Providence, RI 02912, USA


Abstract

Currently the Chinese government has adopted World Health Organization interim target-1 values as the national ambient air quality standards values. However, the population-based evidence was insufficient, especially for the oldest old (aged 80+). We evaluated the association of fine particulate matters (PM_{2.5}) exposure and incidence of disability in activities of daily living (ADL) in 15 453 oldest old in 886 counties/cities in China from 2002 to 2014 using Cox model with penalized splines and competing risk models to evaluate the linear or non-linear association. After adjusting for potential confounders, a J-shaped association existed between PM_{2.5} exposure with a threshold concentration of 33 µg/m³, and incident disability in ADL. Above this threshold, the risk magnitude significantly increased with increase of PM_{2.5} concentrations; compared to 33 µg/m³, the hazard ratio ranged from 1.03 (1.00-1.06) at 40 µg/m³ to 2.25 (1.54-3.29) at 110 µg/m³. The risk magnitude was not significantly changed below this threshold. Each 10 µg/m³ increase in PM_{2.5} exposure corresponded to a 7.7% increase in the risk of disability in ADL (hazard ratio 1.077, 95% CI 1.051-1.104). Men, smokers, and participants with cognitive impairment might be more vulnerable to PM_{2.5} exposure. The study provided limited population-based evidence for the oldest old and detected a threshold of 33 µg/m³, and supported that reduction to current World Health Organization interim target-1 value (35 µg/m³) and Chinese national ambient air quality standards (35 µg/m³) or lower may be associated with lower risk of disability in ADL.

Capsule:

A J-shaped association existed between long-term PM_{2.5} exposure with a threshold concentration of 33 µg/m³ and incident disability in ADL.

Graphical Abstract



Keywords

PM_{2.5}; disability; air pollution; oldest old; cohort study

1. Introduction

Worldwide, aging-related physical disability has become a critical health concern with the rapidly growing number of older adults, who comprise the largest group of people with disabilities¹. In China, the population aged 60 years and older with physical disability is predicted to reach 64.02 million by 2020, and 140 million by 2050^{2, 3}. Regarded as a key indicator of overall health for older adults, disability represents a person's ability to perform usual tasks and activities, which are essential for self-care and independent living⁴. Elevated exposure to air pollution, especially over a long-term duration, is associated with various adverse health problems, such as increased risks for cardiovascular and respiratory diseases^{5, 6}. Given that physical disability is a common functional consequence of these subclinical pathologic processes and co-morbid chronic diseases, it is likely that exposure to air pollution also influences physical disability in older age.

Evidence is growing for an association between ambient air pollution exposure and physical functioning of older adults^{7–12}. Despite this, several knowledge gaps are identified in the prior literature: (1) prior studies mainly focused on older adults younger than age 80 years, although it is likely that adults within an older cohort may be extremely vulnerable to the negative effects of air pollution exposure given functional organ decline¹³. (2) previous studies have mainly focused on either a composite air pollution index (API) or nitrogen oxides (NO_x) but have not assessed fine matter with a diameter <2.5 μm (PM_{2.5}) exposures, which are generally believed to be a primary driver of air pollution related adverse health. (3) Prior studies have been mostly conducted in developed countries where population exposures to PM_{2.5} are low, such as Europe and North America. China, the largest developing country with a relatively high level of exposure to air pollution¹⁴, is at a different stage of the epidemiologic transition, which means the conclusions drawn from developed countries may not reflect the situation in China. (4) In part, due to the lack of more accurate exposure data, prior studies have been conducted examining primarily community-level exposures partly. For the current analysis, we were able to access monitoring data throughout China, using satellite remote sensing combined with a nationwide networking of PM_{2.5} monitoring¹⁵. (5) Prior studies were mainly cross-sectional study in design, or only accounting for acute effects of short-term exposures, while disability in activities of daily living (ADL) would be expected to relate to chronic, progressive effects of long-term exposure to air pollution¹⁶. (6) Prior studies used different measures of disability; and an internationally accepted measure was not used, such as Katz index of independence in ADL¹⁷.

Currently, the Chinese government has adopted the World Health Organization (WHO) air quality interim target-1 values of 35 μg/m³ as the National Ambient Air Quality Standards (NAAQS) values^{15, 18, 19}; these are expected to be updated when sufficient evidence from epidemiologic research conducted in Chinese populations becomes available. The concurrent rapid population aging and environmental degradation in China²⁰ make it more urgent to study the role of air pollution in disability in ADL for oldest old (80+). With an attempt to contribute to the currently limited evidence, in this study, we analyzed data from a prospective cohort of the oldest old scattered across 866 community units (cities or counties) in 23 provinces of China, by exposure estimates derived from satellite observations; and

determined disability in ADL by means of the Katz index and examined linear and non-linear dose-response relationships to explore the exposure limits most appropriate for the over 80 age group.

2. Materials and Methods

2.1 Study group

The study subjects were participants of the Chinese Longitudinal Healthy Longevity Survey (CLHLS). The study was chosen probabilistically to represent 85% of the total population of Chinese oldest old^{12,20} and a national range of exposures to PM_{9.5} in ambient air (Figure 1). All participants underwent triennial assessments during which they completed questionnaires and underwent evaluation of ADL²¹. In 2002-2014, we enrolled 15 453 participants aged 80 and older (mean age, 92.3±7.3 years), with available geographic topography information, and independent ADL at baseline, including 8141 octogenarians (aged 80-89), 4472 nonagenarians (aged 90-99), and 2840 centenarians (aged 100 and older). Among 15453 participants, 7387, 3177, 4393, 496 participants were enrolled in 2002, 2005, 2008 and 2011 survey respectively; and 2326, 2441, 2836, and 1142 participants died before the incidence of disability in ADL in 2002, 2005, 2008 and 2011 survey respectively. Cohort members were followed from the date of recruitment until the date of disability in ADL, death, emigration, or 1 September 2014 (whichever came first). All procedures performed in studies were approved by the biomedical ethics committee of Peking University (IRB00001052-13074). Written informed consents were obtained from all participants included in the study by themselves or their proxy respondents.

2.2 Exposure assessment

At baseline, we obtained geographic topography information from the community questionnaire of the CLHLS. Data on estimates of PM_{2.5} came from the Atmospheric Composition Analysis Group from using satellite-, simulation- and monitor-based information²². Annual average residential PM_{2.5} exposure with 0.01° spatial resolution (1.1×1.1 km) at the participants' home addresses was estimated according to the year of survey, from satellite observations by combining Aerosol Optical Depth (AOD) retrievals from multiple satellite products (MISR, MODIS Dark Target, MODIS and SeaWiFS Deep Blue, and MODIS MAIAC)²². The resultant PM_{2.5} estimates were highly consistent (R²=0.81) with out-of-sample cross-validated PM_{2.5} concentrations from monitors²². Nearly all of the locations (99.3%) were sufficiently complete and correct to allow geocoding to an exact location; exposures were not updated given that only very few (2.1%) participants moved to another address. The correlation matrix of estimated PM_{2.5} showed that the exposures from the year of 2002 to 2014 were highly correlated, the correlation coefficient ranged from 0.86 to 0.98 (Table 1), which suggested that the baseline PM_{2.5} may be a good estimation of the exposure from recruitment to event incidence. PM_{2.5} in the recruitment year of 2002, 2005, 2008 and 2011 was therefore used in the analysis. The improved ambient air pollution estimates from satellite monitoring may have lower exposure misclassification bias than the exposure estimates that relied only on fixed-location monitoring station.

2.3 Outcome assessment

The internationally accepted Katz index of independence in ADL was used to assess disability¹⁷. Disability was identified if a participant needed help in any areas of the six daily tasks of dressing, eating, bathing, continence, toileting and cleaning themselves afterward, and indoor movement; a participant was considered independent if no help was required with any ADL. This resulted in scores ranging from 0 to 6; higher scores indicated better ADL. Participants who scored 5 or less were identified as disability in ADL; scored 6 were identified as normal ADL. The Katz index has been used in a variety of epidemiologic studies of aging, and has well-established reliability and validity²³, more details on Katz index were described in Supplementary 1.

2.4 Statistical analysis

To explore the linear or non-linear associations of PM_{2.5} with disability in ADL, Cox models with penalized splines were conducted²⁴. Based on the corrected Akaike Information Criterion (AIC), 3 degrees of freedom for PM_{2.5} (coded as a continuous variable) were selected for the association analyses. Cox models with penalized splines showed a statistically significant J-shaped association for PM_{2.5} with disability in ADL. Participants with 33 µg/m³ had the minimum risk of disability in ADL based on the estimated parameters of penalized splines (J-shaped exposure-response function) after adjustment for the confounders. The hazard ratios (HRs) and 95% confidence intervals (CI) were evaluated for every 10 µg/m³ interval of PM_{2.5} concentrations for 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 and 110 µg/m³ in Cox models with penalized splines adjusting for the confounders, with reference to 33 µg/m³. A number of potential confounding factors were measured, including age (continuous), sex, residence, current marital status, living pattern, educational level, occupation, source of income, parity, playing cards, smoking status, alcohol consumption, physical activity, hypertension, heart disease, cerebrovascular disease, diabetes mellitus, respiratory disease and cognitive impairment at baseline, based on the existing literature²⁵, and included in the final model specification. Living pattern has been defined as “living with family members” or “living alone or at nursing home”. Educational level was classified as illiterate and literate since most Chinese adults aged 80 and older didn't receive any formal education. Occupation was categorized as farmer or non-farmer. Source of income was categorized as pension or other. Parity was the number of children and used as continuous variable. Smoking status was defined into three categories: current, former, and never. Alcohol drinking status was defined into three categories: current, former, and never. Data on the prevalence of chronic diseases such as heart disease, cerebrovascular disease, diabetes mellitus, and respiratory disease were collected. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) at baseline enrollment were measured to determine the presence of hypertension (SBP 140 mmHg, DBP 90 mmHg). Cognitive impairment was defined as the score of mini-mental status examination lower than 24. In the final analytic database, although less than 0.7% of individuals had missing data for any individual covariates, multiple imputation methods were used to impute missing values of covariates.

To correctly estimate the marginal probability of disability in ADL in the presence of mortality, competing risk models were constructed to estimate risk of disability in ADL for each 10 µg/m³ increase in PM_{2.5} adjusted for the above confounders. Using Cox models

with penalized splines, we evaluated potential modifiers of an effect between PM_{2.5} and disability in ADL such as sex, smoking status, residence, cognitive function, subgroup analyses in men and women, smokers and non-smokers, urban participants and rural participants, participants with cognitive impairment and normal cognition. Additionally, interactions between those modifiers and PM_{2.5} for disability in ADL were tested.

To assess the robustness of the observed associations, a number of sensitivity analyses were conducted. (1) The predicted PM_{2.5} concentrations were averaged at each of her/his residential locations from recruitment to event incidence, characterizing the historic long-term exposures. (2) Models were repeated for the participants who did not move to another address during the follow-up survey, even though address changes were only observed in very few of participants. (3) Crude models and models adjusted only for sex and age were also constructed, for adjusted variables may be plausible intermediates on the causal pathway. (4) Analyses were conducted excluding the participants lost to follow-up. (5) Models were conducted with a relatively “healthy” designation, by excluding individuals diagnosed with the comorbidities (heart disease, cerebrovascular disease, diabetes mellitus, and respiratory disease).

A P-value threshold of 0.05 was used to assess statistical significance. All analyses were carried out using SAS version 9.4 (SAS Inc., Cary, NC, USA) and R 3.4.2 (R foundation for Statistical Computing, Vienna, Austria).

3. Results

3.1 Study participants

In total, the analysis included data from 15 453 individuals (Table 2). The mean age of the participants was 92.3±7.3 years; 43.9% were men and 18.0% lived in urban areas at baseline. Annual average residential PM_{2.5} exposure at baseline ranged from 8.2 µg/m³ to µg/m³ (Figure 1). The mean annual PM_{2.5} exposure was as follows: 50.2 µg/m³ overall; 49.5 µg/m³ for rural residents; 53.4µg/m³ for urban residents; 49.9 µg/m³ for participants with independent ADL; and 51.4 µg/m³ for participants with disability in ADL. The mean (standard deviation, SD) of the follow-up time for included participants was 4.6 (3.1) years. Over a follow-up period of 71 396 person-years, 3373 (21.8%) incident cases of disability in ADL were ascertained.

3.2 Association between PM_{2.5} and disability in ADL

Using a Cox model with penalized splines adjusted for potential confounders, a J-shaped association was observed between PM_{2.5} exposure and disability in ADL with a threshold concentration of 33 µg/m³. Above this threshold, the risk magnitude significantly increased with increase of PM_{2.5} concentrations; not significantly changed below this threshold. Compared to 33 µg/m³, the hazard ratios for different PM_{2.5} thresholds of exposure of 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 and 110 µg/m³ were 1.09(0.86-1.39), 1.02(0.89-1.17), 0.99 (0.93-1.06), 1.03 (1.00-1.06), 1.11 (1.09-1.14), 1.23 (1.20-1.27), 1.38 (1.31-1.45), 1.54 (1.41-1.71), 1.75 (1.48-2.07), 1.98 (1.53-2.58) and 2.25 (1.54-3.29) respectively (Figure 2). The value at which the curve becomes significantly non-zero was 40 µg/m³. The competing

risk models showed that each 10 $\mu\text{g}/\text{m}^3$ increase in $\text{PM}_{2.5}$ exposure corresponded to a 7.7% increase in the risk of disability in ADL (HR 1.077 [1.051-1.104]).

3.3 Sub-group and sensitivity analyses

Significant interactions of sex ($P=0.012$), smoking status ($P=0.008$), and cognitive function ($P<0.001$) with $\text{PM}_{2.5}$ for disability in ADL were observed; there were no significant interactions of residence location with $\text{PM}_{2.5}$. Men, smokers, and participants with cognitive impairment might be more vulnerable to the effects of long-term $\text{PM}_{2.5}$ exposure (Figure 3). The associations of $\text{PM}_{2.5}$ and disability in ADL were unchanged or slightly strengthened in the various sensitivity analyses that included the following: lagging 1-, 2-, 3-, 4-years exposure period exposure (Supplementary Table S1).; averaging exposures from baseline to the event, excluding participants with address changes, only adjusting for sex and age, excluding participants lost to follow-up, and excluding participants with comorbidities (Supplementary Table S2).

4. Discussion

In this prospective cohort study of community-based oldest old, higher risk of disability in ADL incidence occurred among those with individual, higher long-term exposure to $\text{PM}_{2.5}$. In addition, men, smokers, and participants with cognitive impairment might be more vulnerable to this effect. Our study adds to the current limited evidence of an adverse effect of $\text{PM}_{2.5}$ on physical functioning in oldest old; to our knowledge, this is the first cohort study to demonstrate the association for this specific age group. The major contribution of our study is the identification of the threshold value (33 $\mu\text{g}/\text{m}^3$) of $\text{PM}_{2.5}$ for disability in ADL incidence in a large subsample of oldest-old living in 866 highly diverse counties and cities in a nationwide representative study. The strengths of this study also included standard measurements of disability, exposure estimation at an individual level, extensive information regarding important confounding factors, accounting for possible bias from selective attrition and use of Cox models with penalized splines to estimate the shape of the dose-response relationship.

4.1 Previous studies

Our findings are generally consistent with the limited prior epidemiological evidence focusing on older adults. In the cross-sectional study of the 2002 wave of the CLHLS, a higher API at the county or city level was associated with a 25% higher risk of disability in ADL among Chinese older adults aged 65+⁷. The study of the CLHLS 2002-2005 wave also revealed a decrease in 3-yr self-reported health expectancies with increasing API at the county or city level⁸. In a cohort of 1762 Dutch older adults aged 55 to 85, long-term exposure to air pollution, such as NO_2 , PM_{10} , and $\text{PM}_{\text{coarse}}$, adversely affected performance-based and self-reported physical functioning⁹. A recent study from Chicago using NO_x as an indicator of traffic-related air pollution identified a significant association between long-term air pollution exposure and progression in physical disability among older adults¹⁰. An early study reported an association of short-term exposure to $\text{PM}_{2.5}$ with increased disability days¹¹. A cross-sectional study among more than 45 000 adults (mean age: 56 years) from China, Ghana, India, Mexico, Russia, and South Africa showed that

exposure to ambient PM_{2.5} might be one risk factor for disability with women and older adults as the vulnerable population¹².

Unlike the majority of the prior cross-sectional studies, as a prospective cohort study, the present study makes it possible to identify a more confident relationship between PM_{2.5} exposure and disability in ADL. Our ability to estimate exposures at an individual level provides unique and valuable data to complement those prior studies at a community-based level. Notably, the hazard ratio for oldest old in our study was significantly higher than in other studies conducted in adults or younger elderly in the general population⁷⁻¹². The oldest old generally have worse health conditions, which may diminish their ability to adapt to air pollution exposure. In consideration of the unique characteristics for those aged 80+, the extent to which our results might generalize to other age groups are unclear.

4.2 Explanations

There are known or postulated biological mechanisms that support the plausibility of an association between ambient PM_{2.5} exposure and disability. Inhaled fine particles may enter by direct translocation through the olfactory bulb, then could elicit inflammatory responses and oxidative stress²⁶, which have also been one pathological pathway of the occurrence of disability¹⁰. Higher air quality promotes outdoor physical activities, which may protect against risk for subsequent disability in ADL and enhance recovery after the onset of disability in ADL²³. It is also possible that chronic exposure to PM_{2.5} could result in damage in methylation of DNA repair genes²⁷, which may be linked to disability. Because disability and health status were likely to be results of PM_{2.5} exposure rather than causes of it, and because of our use of a prospective cohort study design, reverse causality is not likely to be a big concern in the present study. Additionally, selection bias was partly examined by a number of sensitivity and subgroup analyses.

The association between PM_{2.5} and disability in ADL has appeared to be stronger in men, has also been reported previously²⁸. Specifically, we hypothesize that this may be attributable more time spent outdoors, higher level of physical activity in daily life required more air inhalation for men, led to higher accumulation of toxic chemicals from the air than women. Notably, participants with cognitive impairment were not as adept at handling these added stressors and, therefore, have a higher risk of disability in ADL as a result of the exposure²¹. Not surprisingly, it was found that the adverse effects of PM_{2.5} exposure were greater in smokers, considering the combined effects of air pollution and smoking on health²⁹.

In a large epidemiologic study, personal air monitoring devices is impractical, especially for measuring long-term exposures. By contrast, GIS-based spatiotemporal models offered the ability to estimate individual annual average exposures at baseline and averaged exposures from baseline to the event for the 15 453 participants, which has been validated and applied in previous studies²². The exposure measurements were based on participants' residential locations, and thus limited by the assumption that participants spent most of their time at home; an assumption we believe to be reasonable for the oldest old in this study.

Policy implications—The study found that the oldest old may be more vulnerable to the negative effects of long-term air pollution exposure. It suggested detailed measurements targeting on different populations including the general adults, younger elderly and the oldest old may be needed to identify the high-risk population and give their priority to health protection in the rural and urban regions. Additionally, in the published integrated exposure-response model for long-term exposure of PM_{2.5} from global burden of diseases, the concentration range of 50-100 µg/m³ were estimated by second-hand smoke due to the gap of epidemiological evidence on high ambient PM_{2.5} exposure and health outcome from developing countries³⁰. The developing countries, was similar to the PM_{2.5} exposure and at a similar different stage of the epidemiologic transition to China. Future studies are needed to investigate whether the conclusions drawn from in China can be transferred to other developing countries with similar characteristics. The results will provide important policy implications to policy-makers and service providers in the disciplines of health and environmental protection, especially in the developing countries such as China and India³¹.

Increasing awareness of the harmful effects of air pollution has led to the development of guidelines to prevent excess exposure to these toxicants. For better air quality management to protect public health, the Chinese NAAQS were revised by adopting the interim target values in air quality guidelines in 2012, in which concentration limits for annual averages of PM_{2.5} were 35 µg/m³.^{18,19} However, the formulation of air quality standards requires integrated assessments of risks of air pollution attributable to disease burden and feasibility of environmental management³². Our results support that reduction to current WHO air quality interim target-1 and Chinese national ambient air quality standards (35 µg/m³) may be associated with lower risk of disability in ADL. It highlight more air pollution health researcher were needed to explore the benefits of continued reductions for cleaner air (i.e., 10ug/m³ for PM_{2.5} annual average) in setting up health risk management-based air quality standards in China..

4.3 Limitations

Several limitations merit attention. First, our study provides only indirect evidence for an effect of PM_{2.5} exposure since air pollution is a complex mixture in which PM_{2.5} is one component among many. Second, only outdoor exposures to PM_{2.5} were estimated based on residential location, insufficient adjustment of indoor air pollution (such as household solid fuel use), second-hand smoking and other air pollutants, which may lead to potential confounding³², may comprise a limitation in this study. Third, in this study, more accurate estimation of PM_{2.5}, such as the PM_{2.5} exposure at actually date of survey, were not available, even though lagging 1-, 2-, 3-, 4-years exposure period exposure were matched. Fourth, it is possible that there is information bias in the design of the present study, as a result that disability in ADL could be occurred at any time during a 2- to 3-year interval of follow-up, the exact date of the event was not collected. Fifth, potential reversibility of the incident disability in ADL when performing Cox models may weaken the robustness of the present finding, although the probability of reversible physical disability was relatively low for this population of oldest old with an average age of 92.3±7.3 years.

5. Conclusions

We observed a clear J-shaped dose-response relationship of long-term exposure to PM_{2.5} with corresponding disability in ADL incidence among Chinese oldest old; the risk magnitude increased with increments of PM_{2.5} concentrations. This association was detected at a threshold of 33 µg/m³, which was basically consistent with current Chinese NAAQS, demonstrating that current annual averages for PM_{2.5} in NAAQS can be sufficient for mitigating excess risk of disability in ADL from long-term PM_{2.5} exposures for those aged 80+. Therefore, air pollution reduction may be a means for reducing the public health burden of disability and increasing the quality of life of the older adults-particularly aged 80+, men, smokers and people with cognitive impairment. However, the generality of our findings should be further confirmed in other studies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to all participating people and their families. We acknowledge all staffs and postgraduates for their great contributions to the program.

Funding

This work was supported by National Natural Sciences Foundation of China (81273160, 81573247, and 71490732), the U.S. National Institute on Aging (2P01AG031719, 5P30AG028716) and United Nations Fund for Population Activities.

Role of the funding source

The financial supporters had no involvement in study design, data collection, analysis and interpretation of data, writing of the report, and decision to submit the article.

References:

1. GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; 390: 1211–1259. [PubMed: 28919117]
2. China aging problem national commission, China ministry of civil affairs, China ministry of finance. The fourth survey on the living conditions of urban and rural elderly in China. 2016.
3. Qian J, Chen Y, Rao K, et al. Trends analysis and prediction on disability of China mainland aging population and policy suggestion. *Chin J Health Stat* 2012; 29: 6–9.
4. Artaud F, Dugravot A, Sabia S, et al. Unhealthy behaviours and disability in older adults: three-City Dijon cohort study. *BMJ* 2013; 347: f4240. [PubMed: 23881930]
5. Kaufman JD, Adar SD, Barr RG; et al. Association between air pollution and coronary artery calcification within six metropolitan areas in the USA (the Multi-Ethnic Study of Atherosclerosis and Air Pollution): a longitudinal cohort study. *Lancet* 2016; 388: 696–704. [PubMed: 27233746]
6. Guan WJ, Zheng XY, Chung KF, et al. Impact of air pollution on the burden of chronic respiratory diseases in China: time for urgent action. *Lancet* 2016; 388: 1939–1951. [PubMed: 27751401]
7. Zeng Y, Gu D, Purser J, et al. Associations of environmental factors with elderly health and mortality in China. *Am J Public Health* 2010; 100:298–305. [PubMed: 20019314]

8. Wen M, Gu D. Air Pollution Shortens Life Expectancy and Health Expectancy for Older Adults: The Case of China. *J Gerontol A Biol Sci Med Sci* 2012; 67:1219–1229. [PubMed: 22518820]
9. de Zwart F, Brunekreef B, Timmermans E, et al. Air Pollution and Performance-Based Physical Functioning in Dutch Older Adults. *Environ Health Perspect* 2018; 126:017009. [PubMed: 29364820]
10. Jennifer W, Kaufman JD, Szpiro AA, et al. Exposure to traffic-related air pollution in relation to progression in physical disability among older adults. *Environ Health Perspect*; 2016, 124:1000–1008. [PubMed: 27022889]
11. Stieb D M, Smith-Doiron M, Brook J R, et al. Air pollution and disability days in Toronto: results from the national population health survey. *Environ Res* 2002; 89:210–219. [PubMed: 12176005]
12. Lin H, Guo Y, Yang Z, et al. Exposure to ambient PM_{2.5}, associated with overall and domain-specific disability among adults in six low- and middle-income countries. *Environ Int* 2017; 104:69–75. [PubMed: 28453972]
13. Zeng Y, Feng Q, Hesketh T, et al. Survival, disabilities in activities of daily living, and physical and cognitive functioning among the oldest-old in China: a cohort study. *Lancet* 2017; 389:1619–1629. [PubMed: 28285816]
14. Yin P, He Q Fan M, et al. Particulate air pollution and mortality in 38 of China's largest cities: time series analysis. *BMJ* 2017; 356: j667. [PubMed: 28292780]
15. World Health Organization. WHO | WHO Global Urban Ambient Air Pollution Database (update 2016). 2016 WHO http://www.who.int/phe/health_topics/outdoorair/databases/cities/en/
16. Tak E, Kuiper R, Chorus A, et al. Prevention of onset and progression of basic ADL disability by physical activity in community dwelling older adults: a meta-analysis. *Ageing Res Rev* 2013; 12: 329–338. [PubMed: 23063488]
17. Katz S, Ford AB, Moskowitz RW, et al. The index of ADL: A standardized measure of biological and psychosocial function. *JAMA* 1963; 185:914–919. [PubMed: 14044222]
18. MEP (Ministry of Environmental Protection of China). Ambient Air Quality Standards. (Document GB 3095-2012). Beijing: Ministry of Environmental Protection of the People's Republic of China 2012.
19. World Health Organization. WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide: global update 2005: summary of risk assessment. Geneva: World Health Organization, 2006.
20. Zhang J, Mauzerall DL, Zhu T, et al. Environmental health in China: progress towards clean air and safe water. *Lancet* 2010; 375: 1110–1119. [PubMed: 20346817]
21. Lv YB, Gao X, Yin ZX, et al. Revisiting the association of blood pressure with mortality in the oldest old: a community-based longitudinal prospective study in China. *BMJ* 2018; 361: k2158. [PubMed: 29871897]
22. van Donkelaar A, Martin RV, Brauer M, et al. Global Estimates of Fine Particulate Matter using a Combined Geophysical-Statistical Method with Information from Satellites, Models, and Monitors. *Environ Sci Technol* 2016; 50: 3762–3772. [PubMed: 26953851]
23. Yang M, Ding X, Dong B. The measurement of disability in the elderly: a systematic review of self-reported questionnaires. *J Am Med Dir Assoc* 2014; 15: 150.e1–9.
24. Eisen EA, Agalliu I, Thurston SW, et al. Smoothing in occupational cohort studies: an illustration based on penalised splines. *Occup Environ Med* 2004; 61:854–860. [PubMed: 15377772]
25. Gill TM, Guralnik JM, Pahor M, et al. Effect of structured physical activity on overall burden and transitions between states of major mobility disability in older persons. *Ann Intern Med* 2016; 165:833–840. [PubMed: 27669457]
26. Tzivian L, Winkler A, Dlugaj M, et al. Effect of long-term outdoor air pollution and noise on cognitive and psychological functions in adults. *Int J Hyg Environ Health* 2015; 218:1–11. [PubMed: 25242804]
27. Neven KY, Saenen ND, Tarantini L, et al. Placental promoter methylation of DNA repair genes and prenatal exposure to particulate air pollution: an ENVIRONAGE cohort study. *Lancet Planet Health* 2018; 2: e174–e183. [PubMed: 29615218]

28. Liu T, Zhang Y H, Xu Y J, et al. The effects of dust-haze on mortality are modified by seasons and individual characteristics in Guangzhou, China. *Environ Pollut* 2014; 187:116–123. [PubMed: 24477104]
29. Lin H, Guo Y, Kowal P, et al. Exposure to air pollution and tobacco smoking and their combined effects on depression in six low- and middle-income countries. *Br J Psychiatry* 2017; 211:157–162. [PubMed: 28798061]
30. Lelieveld J, Klingmüller K, Pozzer A, et al. Cardiovascular disease burden from ambient air pollution in Europe reassessed using novel hazard ratio functions. *Eur Heart J*, 2019, 40: 1590–1596. [PubMed: 30860255]
31. Cohen AJ, Brauer M, Burnett R, et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet*, 2017, 389: 1907–1918. [PubMed: 28408086]
32. Kan HD. A review of standard value of fine particulate matter (PM_{2.5}) ruled by National Ambient Air Quality Standards (GB3095-2012) in China. *Chin J Prev Med* 2012; 46:396–398.
33. GBD 2013 Risk Factors Collaborators, Forouzanfar MH, Alexander L, et al. Global, regional, and national comparative risk assessment of 79 behavioral, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015; 386:2287–2323. [PubMed: 26364544]

Highlights

- The dose-response relationship between long-term PM_{2.5} exposure and ADL was J-shaped
- The identification of a threshold of 33 µg/m³ for PM_{2.5} and ADL disability incidence
- The first cohort study to demonstrate effect of PM_{2.5} on ADL for the oldest old
- Air pollution reduction may reduce the public health burden of disability in the elderly.




Figure 1.

Annual average residential PM_{2.5} exposure with 0.01° spatial resolution (1.1x1.1 km) in 23 provinces of China at baseline for 15,453 participants

Annual average residential PM_{2.5} exposure at baseline ranged from 8.2 µg/m³ to 115.1 µg/m³, the mean annual PM_{2.5} exposure was 50.2 µg/m³ overall.




Figure 2. Association of PM_{2.5} and disability in ADL in Cox models with penalized splines among Chinese oldest old (33 µg/m³ as reference).^a
^a Adjusting for age, sex, residence, current marital status, living pattern, educational level, occupation, source of income, parity, playing cards, smoking status, alcohol consumption, physical activity, hypertension, heart disease, cerebrovascular disease, diabetes mellitus, respiratory disease and cognitive impairment at baseline.




Figure 3.

Subgroup analyses: association of PM_{2.5} and disability in ADL in Cox models with penalized splines among Chinese oldest old (33 µg/m³ as reference).

^a Adjusting for age, sex, residence, current marital status, living pattern, educational level, occupation, source of income, parity, playing cards, smoking status, alcohol consumption, physical activity, hypertension, heart disease, cerebrovascular disease, diabetes mellitus, respiratory disease and cognitive impairment at baseline

Table 1.The correlation matrix of PM_{2.5} exposure from the year of 2002 to 2014 (N = 453) [†]

PM _{2.5}	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
2002	1.00	-	-	-	-	-	-	-	-	-	-	-	-
2003	0.94	1.00	-	-	-	-	-	-	-	-	-	-	-
2004	0.94	0.89	1.00	-	-	-	-	-	-	-	-	-	-
2005	0.96	0.92	0.96	1.00	-	-	-	-	-	-	-	-	-
2006	0.92	0.93	0.89	0.94	1.00	-	-	-	-	-	-	-	-
2007	0.94	0.96	0.89	0.93	0.94	1.00	-	-	-	-	-	-	-
2008	0.95	0.96	0.9	0.92	0.92	0.97	1.00	-	-	-	-	-	-
2009	0.94	0.94	0.88	0.9	0.9	0.95	0.98	1.00	-	-	-	-	-
2010	0.92	0.96	0.86	0.93	0.93	0.95	0.94	0.92	1.00	-	-	-	-
2011	0.95	0.94	0.88	0.93	0.93	0.96	0.96	0.96	0.96	1.00	-	-	-
2012	0.93	0.93	0.9	0.95	0.96	0.96	0.96	0.93	0.94	0.96	1.00	-	-
2013	0.93	0.96	0.84	0.91	0.93	0.95	0.95	0.95	0.96	0.98	0.95	1.00	-
2014	0.93	0.96	0.88	0.91	0.91	0.96	0.97	0.96	0.95	0.96	0.95	0.96	1.00

[†] All the *P*value of the correlation coefficients were <0.001.

Table 2.

Baseline characteristics of the study participants by disability in ADL status (N = 15,453).

Characteristics	All	Independent ADL	Disability in ADL	P value
Number of participants *	15453 (100)	12080 (78.2)	3373 (21.8)	
Age (year) [#]	92.3±7.3	91.9±7.2	93.8±7.4	<0.001
Sex				<0.001
Men *	6790 (43.9)	5507 (45.6)	1283 (38.0)	
Women *	8663 (56.1)	6573 (54.4)	2090 (62.0)	
Residence				<0.001
Urban *	2781 (18.0)	2016 (16.7)	765 (22.7)	
Rural *	12672 (82.0)	10064 (83.3)	2608 (77.3)	
Marital status				0.022
In marriage *	4083 (26.4)	3140 (26.0)	943 (28.0)	
Not in marriage *	11370 (73.6)	8940 (74.0)	2430 (72.0)	
Education				0.009
Illiteracy *	10420 (67.4)	8083 (66.9)	2337 (69.3)	
Literacy *	5033 (32.6)	3997 (33.1)	1036 (30.7)	
Living pattern				<0.001
Live with family members *	12325 (79.8)	9544 (79.0)	2781 (82.4)	
Live alone or at nursing home *	3128 (20.2)	2536 (21.0)	592 (17.6)	
Tobacco smoking				<0.001
Current *	10454 (67.7)	8077 (66.9)	2377 (70.5)	
Former *	2929 (19.0)	2372 (19.6)	557 (16.5)	
Never *	2070 (13.4)	1631 (13.5)	439 (13.0)	
Alcohol drinking				<0.001
Current *	10630 (68.8)	8193 (67.8)	2437 (72.3)	
Former *	3364 (21.8)	2718 (22.5)	646 (19.2)	
Never *	1459 (9.4)	1169 (9.7)	290 (8.6)	
Regular exercise	5329 (34.5)	4090 (33.9)	1239 (36.7)	0.002
Hypertension *	7827 (51.3)	5992 (49.6)	1835 (54.4)	
Heart disease *	937 (6.1)	670 (5.5)	267 (7.9)	<0.001
Diabetes *	178 (1.2)	130 (1.1)	48 (1.4)	0.095
Cerebrovascular disease *	415 (2.7)	292 (2.4)	123 (3.6)	<0.001
Respiratory disease *	1615 (10.5)	1321 (10.9)	294 (8.7)	<0.001
Cognitive impairment *	3488 (22.6)	2751 (22.8)	737 (21.8)	0.257
PM _{2.5} (ug/m ³) [#]	50.2±13.4	49.9±13.2	51.4±14.3	<0.001

ADL: activities of daily living; PM_{2.5}: fine particulate matter with diameter <2.5 μm

*. Indicating categories variables displayed as sample size (%)

#. Indicating continuous variables displayed as mean±SD.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript