1980

The Spin-Wave Continuum of the $S = 1/2$ Linear Heisenberg Antiferromagnet

Gerhard Müller

University of Rhode Island, gmuller@uri.edu

Harry Thomas

Hans Beck

Follow this and additional works at: https://digitalcommons.uri.edu/phys_facpubs

Terms of Use

All rights reserved under copyright.

Citation/Publisher Attribution

Müller G., Thomas H., Beck H. (1980) The Spin-Wave Continuum of The $S=1/2$ Linear Heisenberg Antiferromagnet. In: Riste T. (eds) Ordering in Strongly Fluctuating Condensed Matter Systems. NATO Advanced Study Institutes Series (Series B: Physics), vol 50. Springer, Boston, MA.
Available at: http://dx.doi.org/10.1007/978-1-4684-3626-6_12

This Article is brought to you for free and open access by the Physics at DigitalCommons@URI. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

The Spin-Wave Continuum of the $S = 1/2$ Linear Heisenberg Antiferromagnet

Gerhard Müller, ¹ Harry Thomas ¹ and Hans Beck ²

¹ Institut für Physik, Universität Basel, CH-4056 Basel, Switzerland

² Institut de Physique de l'Université, CH-2000 Neuchâtel, Switzerland

In the $S = 1/2$ linear Heisenberg antiferromagnet (HB AF)

$$H = J \sum_{i=1}^N \vec{S}_i \cdot \vec{S}_{i+1} - h \sum_{i=1}^N S_i^z \quad (1)$$

– although investigated by various theoretical approaches – many important questions concerning the statics and the dynamics have remained open. Recent low-temperature neutron- scattering experiments on $\text{CuCl}_2 \cdot 2\text{N}(\text{C}_5\text{H}_5)$ (CPC) , which is a good realization of an $S = 1/2$ HB AF chain, provided new important information on the dynamics of the system, such as lineshapes and the behaviour in a magnetic field [1]. The important quantity for direct comparison with experiments of this kind is the dynamic spin-correlation function in (q, ω) -space. It is the Fourier transform of $\langle S^z(l, t) S^z(l', 0) \rangle$, and for $T = 0$ it can be written as

$$G_{zz}(q, \omega) = \sum_{\lambda} M_{\lambda} \delta(\omega + E_0 E_{\lambda}), \quad M_{\lambda} = 2\pi |\langle 0 | S^z(q) | \lambda \rangle|^2. \quad (2)$$

In a recent publication [2] an approximate analytical expression for G_{zz} at $T = 0$ and $h = 0$ was obtained by using finite-chain calculations together with the results of two other theoretical approaches. It represents the dominant contribution to $G_{zz}(q, \omega)$ originating from a spin-wave continuum (SWC) bounded between the dispersion branches $E_1(q) = (\pi J/2) |\sin q|$ and $E_2(q) = \pi J |\sin(q/2)|$:

$$G_{zz}^{\text{SWC}}(q, \omega) = 2 \{ \omega^2 - [E_1(q)]^2 \}^{-1/2} \theta(\omega - E_1(q)) \theta(E_2(q) - \omega). \quad (3)$$

G_{zz} increases strongly towards the lower bound $E_1(q)$, which is the desCloizeaux-Pearson spin-wave energy. Result (3) is in good agreement with experimental data for CPC concerning excitation energy, lineshape and integrated intensity [1,2].

In this note we give further arguments supporting (3), which demonstrates the usefulness of finite-chain calculations for properties of the infinite system. Although we cannot give a rigorous derivation of (3), we conjecture that it represents the SWC quantitatively. We have identified those excitations of the finite system which have dominant spectral weight with a special class of eigenstates in the Bethe formalism [3], and we show that these states form, in the thermodynamic limit, a continuum exactly between the two branches $E_1(q)$ and $E_2(q)$. The Bethe Ansatz for the exact eigenfunctions consists of a linear combination $\psi = \sum a(n_1, \dots, n_r) \phi(n_1, \dots, n_r)$ of local basis vectors with reversed spins at lattice sites $n_1, \dots, n_r$ with coefficients of the form

$$a(n_1, \dots, n_r) = \sum_p \exp \left(i \sum_j k_{p_j} n_j + \frac{1}{2} i \sum_{j < l} \psi_{p_j p_l} \right), \quad (4)$$

where the summation $\sum_p$ extends over all permutations of the integers $1, \dots, r$ and p_j is the image of j under the p th permutation. The k_j and the ψ_{ij} obey the coupled equations:

$$2 \cot \frac{\psi_{jl}}{2} = \cot \frac{k_j}{2} - \cot \frac{k_l}{2}, \quad Nk_j = 2\pi\lambda_j + \sum_{l \neq j} \psi_{jl}. \quad (5)$$

The integers λ_j are confined to $1 \leq \lambda_j \leq N-1$, and each choice of a set $\{\lambda_j\}$ (being subject to additional restrictions) determines an eigenstate of the system. Having solved the above equations for k_j , it is straightforward to calculate wave number and energy of the corresponding eigenstate,

$$q = \sum_{j=1}^r k_j = \frac{2\pi}{N} \sum_{j=1}^r \lambda_j, \quad E = - \sum_{j=1}^r (1 - \cos k_j). \quad (6)$$

The ground state, which is a singlet (for even N), corresponds to the $N/2$ integers $\lambda_j = 1, 3, 5, \dots, (N-1)$. Des Cloizeaux and Pearson [4] found the lowest excited states to be given by

$$\begin{aligned} 1, 3, \dots, (N-2n-1), (N-2n+2), \dots, (N-2) \quad q > 0 \\ 2, 4, \dots, (2n-2), (2n+1), \dots, (N-1) \quad q < 0 \end{aligned} \quad (7)$$

($q = 2\pi n/N$) and calculated their energies. The result is the famous DC-P spin-wave branch $E_1(q)$. By generalization of their method we have found the sets $\{\lambda_j\}$ for all SWC states. To the highest branch $E_2(q)$, in particular, belong the sets (always for even N):

$$\begin{aligned} 1, 3, \dots, (N-n-2), (N-n+2), \dots, (N-1) \quad n \text{ odd} \\ 1, 3, \dots, (N-n-3), (N-n), (N-n+3), \dots, (N-1) \quad n \text{ even} \end{aligned} \quad (8a)$$

for $q > 0$ and

$$\begin{aligned} 1, 3, \dots, (n-2), (n+2), \dots, (N-1) \quad n \text{ odd} \\ 1, 3, \dots, (n-3), n, (n+3), \dots, (N-1) \quad n \text{ even} \end{aligned} \quad (8b)$$

for $q < 0$. Using these numbers we can calculate (in the thermodynamic limit) the energies of all the excitations of the two-parameter SWC ($q > 0$ for convenience):

$$E_b(q) = \pi J \left| \sin \frac{q}{2} \cos \left(\frac{q}{2} - \frac{q_b}{2} \right) \right|, \quad (9)$$

where q ($0 \leq q \leq \pi$) is the wave number of the excitation (now with respect to that of the ground state) and q_b ($0 \leq q_b \leq q$) labels the different dispersion branches within the continuum. The lowest branch $E_1(q)$ has $q_b = 0$ and the highest one has $q_b = q$ yielding $E_2(q)$. Furthermore (9) immediately provides the density of states in the SWC

$$D(q, \omega) = \frac{N}{2\pi} \left\{ [E_2(q)]^2 - \omega^2 \right\}^{-1/2} \quad (10)$$

According to (2) $G_{zz}^{\text{SWC}}(q, \omega)$ is the product of the density of states $D(q, \omega)$ and a spectral weight defined by the squared matrix elements between the ground state and the SWC excitations: $M(q, \omega) \equiv |\langle 0, 0 | S^z(q) | \omega, q \rangle|^2$, yielding

$$M(q, \omega) = \frac{4\pi}{N} \sqrt{\frac{[E_2(q)]^2 - \omega^2}{\omega^2 - [E_1(q)]^2}}. \quad (11)$$

Comparison of (11) with finite-chain matrix elements shows good agreement.

This approach to the dynamics of the $S = 1/2$ HB AF at $T = 0$ can be extended to the $h \neq 0$ case. From finite-chain calculations we have determined the excitations contributing significantly

to $G_{zz}(q, \omega)$. Again we have identified this class of excitations unambiguously with a certain class of eigenstates in the Bethe formalism. The calculations to solve eq's (5) for these states are in progress. Preliminary approximate results show that $G_{zz}(q, \omega)$ is dominated by two partly overlapping continua of excitations. Fig. 1 shows the boundaries of these continua for a special value of h . Again, the spectral weight of $G_{zz}(q, \omega)$ increases strongly as the frequency is lowered towards the lower boundary of each continuum. Further we find that $G_{xx}(q, \omega)$ looks for $h \neq 0$ qualitatively different from $G_{zz}(q, \omega)$. In particular, the lowest branch is inverted with respect to the axis $q = \pi/2$. Therefore we expect appropriate neutron scattering experiments to show spectra which are more complex than for $h = 0$ (having at least two dominant peaks), and which strongly depend on the relative weight of G_{xx} and G_{zz} in the scans under consideration. More details will be published elsewhere.

Figure 1. The two continua of excitations dominating $G_{zz}(q, \omega)$ at $T = 0$ and $h = \frac{1}{2}h_{\text{crit}}$. In each continuum the spectral weight increases strongly towards the corresponding lower boundary. The lowest boundary corresponds approximately to the spin-wave frequency obtained numerically by Ishimura and Shiba [5] and to the approximate analytical result by Pytte [6]. The special wave number q_m depends only on the magnetization. It is equal to π at $h = 0$ and decreases as h increases, reaching zero at the critical field h_{crit} .

Acknowledgments

We have used a modified cmpj.sty style file.

References

1. I.U. Heilmann, G. Shirane, Y. Endoh, R.J. Birgeneau, and S.L. Holt, to be published.
2. G. Müller, H. Beck, and J.C. Bonner, to be published.
3. H. Bethe, Z. Phys. **71**, 205, (1931).
4. J. Des Cloizeaux and J.J. Pearson, Phys. Rev. **128**, 2131, (1962).
5. N. Ishimura and H. Shiba, Prog. Theor. Phys. **57**, 1862, (1977).
6. E. Pytte, Phys. Rev. B **10**, 685, (1974).