1981

Quantum Effects in the Dynamics of the One-Dimensional Planar Antiferromagnet

Gerhard Müller

University of Rhode Island, gmuller@uri.edu

Harry Thomas

Marcos W. Puga

Hans Beck

Follow this and additional works at: https://digitalcommons.uri.edu/phys_facpubs

Terms of Use

All rights reserved under copyright.

Citation/Publisher Attribution

Gerhard Müller, Harry Thomas, Marcos W. Puga and Hans Beck. *Quantum effects in the dynamics of the one-dimensional planar antiferromagnet*, in *Physics in one dimension*, J. Bernasconi and T. Schneider (Eds.), Springer Series in Solid State Sciences (Springer, Berlin-Heidelberg-New York, 1981), pp. 165-168. Available at: http://dx.doi.org/10.1007/978-3-642-81592-8_17

This Article is brought to you for free and open access by the Physics at DigitalCommons@URI. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Quantum Effects in the Dynamics of the One-Dimensional Planar Antiferromagnet

Gerhard Müller,¹ Harry Thomas,¹ Marcos W. Puga,² and Hans Beck²

¹ Institut für Physik, Universität Basel, CH-4056 Basel, Switzerland

² Institut de Physique de l'Université, CH-2000 Neuchâtel, Switzerland

In a recent publication [1] we found that the $T = 0$ dynamics of the 1D $s = \frac{1}{2}$ Heisenberg antiferromagnet (HB AF) with nearest-neighbor exchange interaction J is almost completely dominated by a particular continuum of excitations (called spin-wave continuum, SWC) bounded by the dispersion branches $\epsilon_1(q) = (\pi J/2) \sin q$ and $\epsilon_2(q) = \pi J \sin(q/2)$. It differs markedly from the classical 1D HB AF where the spectral weight is concentrated on a single branch of spin-waves. The result for the structure function $S_{\mu\mu}(q, \omega) \equiv \langle S_i^\mu S_{i'}^\mu \rangle_{q, \omega}$, which is a special case of Eq. (6), is in good agreement with low- T neutron scattering data on CPC [2] concerning excitation energies, lineshapes and integrated intensity.

In this paper we outline an extension to the planar $s = \frac{1}{2}$ AF with Hamiltonian

$$H = \sum_{l=1}^N \left\{ J_{\perp} (S_l^x S_{l+1}^x + S_l^y S_{l+1}^y) + J_z S_l^z S_{l+1}^z \right\}, \quad J_{\perp} \equiv J > 0, \quad 0 < J_z < J. \quad (1)$$

By using finite-chain calculations on systems with 4 to 10 spins, we again find a special class of excitations forming a two-parameter continuum (SWC) in (q, ω) -space which dominates $S_{\mu\mu}(q, \omega)$ at $T = 0$. Further, we can identify each one of these excitations with a member of the ‘class C’ of non-bound states defined in the framework of Bethe’s formalism. By use of des Cloizeaux and Gaudin’s method [3], we obtain the SWC energies

$$\omega_m(q) = \frac{\pi J \sin \theta}{\theta} \sin \frac{q}{2} \cos \left(\frac{q}{2} - \frac{q_m}{2} \right); \quad q \geq 0, \quad 0 \leq q_m \leq q, \quad (2)$$

where $\theta = \arccos(J_z/J_{\perp})$, and q_m labels the different dispersion branches of the SWC. In particular, $q_m = 0$ and $q_m = q$ yield the lower and the upper boundary, respectively,

$$\epsilon_1(q) = \frac{\pi J \sin \theta}{2\theta} \sin q, \quad \epsilon_2(q) = \frac{\pi J \sin \theta}{\theta} \sin \frac{q}{2}. \quad (3)$$

From (2) we deduce the density of states in the SWC

$$D(q, \omega) = \frac{N}{2\pi} \left\{ \epsilon_2^2(q) - \omega^2 \right\}^{-1/2}; \quad \epsilon_1(q) \leq \omega \leq \epsilon_2(q). \quad (4)$$

In analogy to our result for the HB AF ($J_z = J_{\perp}$) and to exact results for the XY AF ($J_z = 0$) [1], we conjecture that $S_{zz}(q, \omega)$ can be written as a product $M(q, \omega) \cdot D(q, \omega)$ with

$$M(q, \omega) = \text{const} \cdot \left\{ \frac{\epsilon_2^2(q) - \omega^2}{\omega^2 - \epsilon_1^2(q)} \right\}^{\alpha} \quad (5)$$

representing the squared matrix elements $|\langle G | S^z(q) | \text{SWC} \rangle|^2$ between the non-degenerate ground state and the SWC excitations. The exponent α can be related to exponents of the Baxter model

[4] yielding $\alpha(\theta) = (\pi/2 - \theta)/(\pi - \theta)$. Thus we are led to the analytic expression

$$S_{zz}(q, \omega) = \frac{2A}{B(1 - \alpha, 1/2 + \alpha)} \frac{\Theta(\omega - \epsilon_1(q))\Theta(\epsilon_2(q) - \omega)}{(\omega^2 - \epsilon_1^2(q))^\alpha (\epsilon_2^2(q) - \omega^2)^{1/2 - \alpha}}, \quad (6)$$

where the Beta function serves as a convenient normalizing factor, and the constant A will be determined below. In the limit $J_z/J_\perp = 0$ ($\theta = \pi/2$), this result is exact [1].

Fig. 1 shows $S_{zz}(q, \omega)$ at fixed $q = 4\pi/5$ as a function of ω for various anisotropies. In the XY limit $\alpha = 0$ $M(q, \omega)$ is a constant and $S_{zz}(q, \omega)$ reflects the SWC density of states (5) with a square-root divergence at $\epsilon_2(q)$ and a finite step at $\epsilon_1(q)$. For intermediate anisotropies ($0 < \alpha < 1/2$) $M(q, \omega)$ is no longer a constant, but diverges at $\epsilon_1(q)$ and vanishes at $\epsilon_2(q)$. This leads to a weaker singularity in $S_{zz}(q, \omega)$ at $\epsilon_2(q)$ and gives rise to a second divergence at $\epsilon_2(q)$. We therefore expect that inelastic neutron-scattering experiments will reveal a two-peak structure at low T . In the HB AF limit $\alpha = 1/2$ the matrix elements (5) vanish sufficiently rapidly at $\epsilon_2(q)$ in order to overcome the density-of-states singularity. Most of the spectral weight is now concentrated at $\epsilon_1(q)$. It should be remarked that the spectrum contributing to $S_{zz}(q, \omega)$ has no gap at $q = 0$ or $q = \pi$ in the whole range $0 \leq J_z/J_\perp \leq 1$. This is a clear quantum effect, for it is in marked contrast to classical spin-wave theory. Here, the excitations contributing to $S_{zz}(q, \omega)$ consist of a single branch (in the extended zone) [5]

$$\epsilon_{cl}(q) = 2Js\{(1 - \cos q)[1 + (J_z/J) \cos q]\}^{1/2}, \quad (7)$$

which has a gap at $q = \pi$ increasing with planar anisotropy. Note that in the quantum case the spectral weight of $S_{zz}(\pi, \omega)$ is also redistributed, with increasing anisotropy, towards higher energies, but rather in terms of a varying exponent than in terms of an energy gap.

An important check for the validity of our result (6) is the application of the sum rule [6]

$$K_{zz}(q) \equiv \int_0^\infty \frac{d\omega}{2\pi} \omega S_{zz}(q, \omega) = -2JF_x(1 - \cos q), \quad (8)$$

where the nearest-neighbor correlation function $F_x \equiv \langle S_i^x S_{i+1}^x \rangle$ has been calculated exactly by Yang and Yang [7]. The first frequency moment of (6)

$$K_{zz}(q) = \frac{AJ \sin \theta}{4\theta} (1 - \cos q) \quad (9)$$

Figure 1. Out-of-plane structure function $S_{zz}(q, \omega)$ of Eq. (6) versus ω at fixed $q = 4\pi/5$ for various anisotropies $J_z/J_\perp$. The prefactor A has been determined for this figure via the sum rule.

reproduces the exact q -dependence irrespective of the anisotropy. The integrated intensity is obtained as

$$I_{zz}(q) \equiv \int_0^\infty \frac{d\omega}{2\pi} S_{zz}(q, \omega) = \frac{A}{2\pi} \tan \frac{q}{2} {}_2F_1 \left(\frac{1}{2}, 1 - \alpha, \frac{3}{2}, -\tan^2 \frac{q}{2} \right), \quad (10)$$

$$I_{zz}(q) \simeq \frac{A}{4\pi} q, \quad (q \ll 1); \quad I_{zz}(\pi) = \frac{A}{4\sqrt{\pi}} \frac{\Gamma(1/2 - \alpha)}{\Gamma(1 - \alpha)}. \quad (11)$$

In the XY limit the linear term is already the exact result for all q . In the HB AF limit $I_{zz}(q)$ has a logarithmic divergence at $q = \pi$. The asymptotic behaviour of the correlations in real space is $\langle S_l^z S_{l+R}^z \rangle \sim (-1)^R / R^{2(1-\alpha)}$. Thus, the out-of-plane correlations are strongest in the isotropic limit. With increasing anisotropy the spins become more and more correlated in the easy plane, and hence the fluctuations out of the plane are increasingly suppressed. Another important static quantity is the q -dependent susceptibility

$$\chi_{zz}(q) \equiv \int_0^\infty \frac{d\omega}{2\pi} \omega^{-1} S_{zz}(q, \omega) = \frac{A\theta}{\pi^2 J \sin \theta} {}_2F_1 \left(1, \alpha + \frac{1}{2}, \frac{3}{2}, \sin^2 \frac{q}{2} \right), \quad (12)$$

$$\chi_{zz}(0) = \frac{A\theta}{\pi^2 J \sin \theta}, \quad \chi_{zz}(q) \sim (\pi - q)^{-2\alpha} \quad (q \rightarrow \pi). \quad (13)$$

The exact result for the uniform susceptibility is already available [7]: $\chi_{zz} = \theta / [J\pi(\pi - \theta) \sin \theta]$. Hence, with the sum rule result (8), (9) and with the uniform susceptibility (13), we have two independent ways of determining A , yielding

$$A_K(\theta) = -8\theta F_x(\theta) / \sin \theta, \quad A_\chi(\theta) = \pi / (\pi - \theta). \quad (14)$$

The difference between the two values is not greater than a few percent, except close to the HB AF limit, where it reaches 15%. This discrepancy in the isotropic limit is understood and extensively discussed in Ref. [1].

We conclude by noting that the present approach also works for the in-plane component of the structure function $S_{xx}(q, \omega)$. In this case there are two dominating spin-wave continua, one of which is again confined between the branches $\epsilon_1(q)$ and $\epsilon_2(q)$ of Eq. (3) and the second one between the boundaries $\epsilon_1(\pi - q) = \epsilon_1(q)$ and $\epsilon_2(\pi - q)$. We find that the SWC contributions to $S_{xx}(q, \omega)$ can be written as a sum of two terms of the form (6) with the boundaries in one of them replaced accordingly. The prefactors and the exponents in the two terms are different functions of the planar anisotropy. Hence the corresponding lineshapes are predicted to be even more complex than those of $S_{zz}(q, \omega)$. Furthermore we can show that an ansatz of the form (6) also holds for the isotropic $s > \frac{1}{2}$ HB AF with an exponent α which in this case depends on the spin quantum number s . Here we find that with increasing s the lineshape of $S_{\mu\mu}(q, \omega)$ becomes more symmetric. In the limit $s \rightarrow \infty$ the exact classical result is recovered. More details will be published elsewhere.

Acknowledgments

This work is supported by the Swiss National Science Foundation. We have used a modified cmpj.sty style file.

References

1. G. Müller, H. Thomas, H. Beck and J.C. Bonner, submitted to Phys.Rev. B.
2. I.V. Heilmann, G. Shirane, Y. Endoh, R.J. Birgenau, and S.L. Holt, Phys. Rev. B **18**, 3530 (1978).
3. J. des Cloizeaux and M. Gaudin, J.Math.Phys. **7**, 1384 (1966).
4. A. Luther and I. Peschel, Phys.Rev. B **12**, 3908 (1975).
5. R.E. Dietz, L.R. Walker, F.S.L. Hsu, W.H. Haemmerle, B. Vis, C.K. Chau, and H. Weinstock, Sol.State Comm. **12**, 1185 (1974).
6. P.C. Hohenberg and W.F. Brinkman, Phys.Rev. B **10**, 128 (1974).
7. C.N. Yang and C.P. Yang, Phys.Rev. **151**, 258 (1966).