2020

Weak biofilm formation among carbapenem-resistant *Klebsiella pneumoniae*

Jaclyn A. Cusumano
University of Rhode Island

R. Caffrey
University of Rhode Island, aisling_caffrey@uri.edu

Kathryn E. Daffinee

Megan K. Luther
University of Rhode Island

Vrishali Lopes

Follow this and additional works at: https://digitalcommons.uri.edu/php_facpubs

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Cusumano, J. A., Caffrey, A. R., Daffinee, K. E., Luther, M. K., Lopes, V., & LaPlante, K. L. (2019). Weak biofilm formation among carbapenem-resistant *Klebsiella pneumoniae*. *Diagnostic Microbiology and Infectious Disease*, 95(4), 114877. doi: 10.1016/j.diagmicrobio.2019.114877
Available at: <https://doi.org/10.1016/j.diagmicrobio.2019.114877>

This Article is brought to you for free and open access by the Pharmacy Practice at DigitalCommons@URI. It has been accepted for inclusion in Pharmacy Practice Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Authors

Jaclyn A. Cusumano, R. Caffrey, Kathryn E. Daffinee, Megan K. Luther, Vrishali Lopes, and Kerry L. LaPlante

Title: Weak biofilm formation among carbapenem-resistant *Klebsiella pneumoniae*

Running title: *Klebsiella pneumonia* biofilm formation

Abstract word count: 112

Manuscript word count: 1763

Jaclyn A. Cusumano^{1,2}, Aisling R. Caffrey^{1,2,3}, Kathryn E. Daffinee,¹ Megan K. Luther^{1,2,3}, Vrishali Lopes¹, Kerry L. LaPlante^{1,2,3,4}

1. Infectious Diseases Research Program, Providence Veterans Affairs Medical Center, Providence, RI, United States

2. College of Pharmacy, University of Rhode Island, Kingston, RI, United States

3. Center of Innovation in Long-Term Support Services, Providence Veterans Affairs Medical Center, Providence, RI, United States

4. Warren Alpert Medical School of Brown University, Division of Infectious Diseases, Providence, RI

Corresponding author: Kerry L. LaPlante, Pharm.D., FCCP, FIDSA, Professor, University of Rhode Island, College of Pharmacy, 7 Greenhouse Rd, Suite 295A, Kingston, RI 02881, 401-874-5560 (office); KerryLaPlante@uri.edu

Declarations of Interest: Kerry L. LaPlante receives research funding or is advisor/consultant for Merck, Pfizer Pharmaceuticals, Ocean Spray Cranberries, Inc., Nabriva Therapeutics US, Inc., Melinta Therapeutics, Inc., and Tetrphase Pharmaceuticals. Aisling Caffrey has received research funding from Pfizer, and Merck. Jaclyn A. Cusumano, Kathryn E. Daffinee, Vrishali Lopes, and Megan K. Luther report no financial relationships.

Abstract

Biofilm formation of multidrug and extensively drug resistant *Klebsiella pneumoniae* isolates is poorly understood. We investigated 139 diverse clinical *K. pneumoniae* isolates that possess various resistance patterns to evaluate the relationship between biofilm formation and resistance. Antimicrobial resistance was compared among a diverse collection of weak versus strong biofilm-forming *K. pneumoniae*, and predictors of strong biofilm formation were identified. Multi-drug resistant isolates were more common among weak (97.9%) versus strong biofilm formers (76%; $p=0.002$). Carbapenem-resistant *K. pneumoniae* were 91% less likely to form strong biofilm (odds ratio 0.09; 95% confidence interval 0.02-0.33). The statistically significant inverse relationship between biofilm formation and antibiotic resistance suggests that virulence may be a trade-off for survival.

Keywords: *Klebsiella pneumoniae*; biofilm; carbapenem resistance

INTRODUCTION

Klebsiella pneumoniae, the most common and most concerning carbapenem-resistant *Enterobacteriaceae* (CRE) (1), is associated with mortality rates up to 50% (2). Adding to this challenging infection is *K. pneumoniae*'s high propensity to form biofilms (3, 4). Biofilm-forming *K. pneumoniae* are associated with foreign indwelling device related infections (4), as well as urinary stones (5-7). The most common *K. pneumoniae* infections include urinary tract infections, pneumonia, as well as intra-abdominal infections, which are prone to biofilm formation(4, 8). Biofilm eradication requires high antimicrobial concentrations (9), which often cannot be physiologically achieved in the blood stream or at the site of infection, thus potentially leading to infection recurrence (4). Unfortunately, clinical microbiology labs cannot routinely test for biofilm formation; however, tested phenotypic characteristics may help clinicians predict biofilm potential.

Multidrug-resistant (MDR) organisms have been associated with biofilm formation when *Klebsiella pneumoniae* (10, 11), *Staphylococcus aureus* (11), *Acinetobacter spp.* (10, 11), *Pseudomonas aeruginosa* (10, 11), *Escherichia coli* (10, 11), coagulase-negative staphylococci (10), or *Enterococcus spp.* (10) are assessed together (10, 11). However, the relationship between biofilm-forming *K. pneumoniae* alone and antimicrobial resistance has not been fully elucidated (12-14). The study objective was to determine whether certain antimicrobial class resistance in *K. pneumoniae* was predictive of strong biofilm formation.

MATERIALS AND METHODS:

Our study included 139 unique *K. pneumoniae* clinical isolates obtained from the Centers for Disease Control and Prevention (CDC; n=66), Biodefense and Emerging Infections (BEI; n=36), American Type Culture Collection (ATCC; n=3), and Providence Veterans Affairs (VA) Medical Center and Rhode Island Hospital (n=34). These isolates were selected because they are known to be resistant to a range of antibiotic classes. Isolates from the Providence VA Medical Center

were collected per the VA approved Institutional Review Board (IRB), Research and Development (R&D), and safety committee protocols.

A previously described biofilm assay was performed to assess biofilm formation (15-19). This assay is considered the standard for evaluation of bacterial attachment and biofilm formation in vitro (20). Isolates were obtained from culture stocks, stored at -80°C. After streaking on tryptic soy agar and incubating for 18 to 24 hours, an inoculum of 6 log₁₀ CFU/mL in tryptic soy broth with added 25 mg/L calcium, 12.5 mg/L magnesium, and 1.25% total dextrose (21). Media was selected and validated based on the greatest biofilm formation, which is consistent with previously published data for *Escherichia coli* (22) and *Klebsiella pneumoniae* (17, 18). Each isolate was incubated in a 96-well plate (Costar 3596) for 24 hours at 37°C and 120 rpm in octuplicate (17, 18). Wells were stained with 0.1% crystal violet (CV) and resolubilized with 33% glacial acetic acid (19). *K. pneumoniae* (ATCC 700603) served as the biofilm positive control and media alone was the negative control (12, 23-25).

Biofilm formation was measured as an optical density (OD₅₇₀). We further categorized isolates as either weak, moderate, or strong biofilm formers based on tertiles of OD₅₇₀. In order to assess differences between the highest and lowest OD₅₇₀, we removed isolates in the moderate range (19, 26).

Organism susceptibility was obtained from the site of collection (eg. CDC, BEI). When results were unavailable testing was performed by E-test or Kirby-Bauer disc diffusion on Mueller-Hinton agar, and were interpreted according to 2017 CLSI susceptibility breakpoints (27, 28). The FDA package insert for tigecycline (E-test MIC ≤2) (29) and EUCAST breakpoints for colistin (E-test MIC ≤2) and fosfomycin (E-test MIC ≤32 and disc diffusion ≥24; both contained glucose-6-phosphate) (30) were used as CLSI breakpoints were not available. Isolates were categorized as multi-drug resistant (MDR), extensively drug-resistant (XDR), or resistant to specific

antimicrobial classes/agents, according to CDC and European CDC expert consensus definitions for *Enterobacteriaceae* (31). MDR isolates demonstrated non-susceptibility to at least one agent in three or more antimicrobial categories out of 16 antimicrobial categories and XDR isolates demonstrated susceptibility to at least one agent in less than or equal to two out of 16 antimicrobial categories (31).

To assess the relationship between biofilm formation and resistance to specific antimicrobial classes/agents, we grouped the 16 antimicrobial categories into 12 categories based on mechanism of action (Table 1). Piperacillin/tazobactam and penicillin/ β -lactamase inhibitors were grouped as penicillins plus β -lactamase inhibitors, and non-extended spectrum cephalosporins, extended-spectrum cephalosporins, cephamycins, and ceftaroline were grouped as cephalosporins. This allowed us to avoid collinearity in our statistical models due to overlap in resistance between antimicrobial categories.

Differences in antimicrobial resistance among the weak and strong biofilm formation groups were assessed with chi-square, Fisher's exact, or t-test as appropriate. Predictors of strong biofilm formation were identified from a logistic regression model. A p-value of 0.1 was used for initial inclusion in the model (Table 1) and stepwise backward elimination was used to identify statistically significant predictors of strong biofilm formation (all p-values <0.05). We assessed multicollinearity between potential predictors in the initial model from variance inflation factors, and confirmed the absence of collinearity. A sensitivity analysis was conducted to identify predictors of biofilm formation as a continuous measure (OD₅₇₀) using linear regression.

RESULTS

Optical density (OD₅₇₀) for all 139 isolates were divided using tertiles as follows: weak (n=47; OD₅₇₀ ≤ 0.16), moderate (n=46; 0.16 < OD₅₇₀ < 0.59), and strong biofilm formers (n=46; OD₅₇₀ ≥ 0.59) (26). This method was internally validated as the positive control was consistently

categorized as a strong biofilm former ($OD_{570} \geq 0.59$) (26). Moderate isolates were removed for a total cohort of 93 isolates (Table 1) to best predict biofilm formation extremes (26).

MDR isolates (n=81) were more common among weak biofilm formers (n=46, 97.9%) versus strong biofilm formers (n=35, 76.1%; $p=0.002$), and XDR (n=25) isolates were similar between the groups (n=12, 25.5% vs. n=13, 28.3% $p=0.77$). Number of resistant antimicrobial categories and OD_{570} are shown in Figure 1. Resistance to all classes of beta-lactams (i.e. penicillins plus β -lactamase inhibitors, cephalosporins, monobactams, and carbapenems), aminoglycosides, chloramphenicol, and fluoroquinolones were more common among weak biofilm formers ($p<0.05$). In the multivariate model, the only predictor of biofilm formation was carbapenem resistance, which was inversely associated with strong biofilm formation (odds ratio, OR 0.09; 95% confidence interval, CI 0.02-0.33). Therefore, carbapenem-resistant *K. pneumoniae* were 91% less likely to form strong biofilm.

As the proportion of XDR isolates did not vary between weak and strong biofilm formers, we conducted a post-hoc sensitivity subgroup analysis excluding XDR isolates (n=68) (Table 2). Predictors of strong biofilm formation were again identified from a stepwise backward elimination logistic regression model, with a p-value of 0.1 used for initial inclusion in the model (Table 2). The only predictor of strong biofilm formation was the number of resistant categories, with an odds ratio of 0.70 (95% CI 0.56-0.86), where the odds of strong biofilm formation decreased by 30% with each increase in the number of resistant categories.

In the sensitivity analysis of the continuous measure of biofilm formation, only fluoroquinolone resistance was predictive of the OD_{570} , with a parameter estimate of -0.44 and an intercept of 0.86 ($p<0.001$). According to this model, fluoroquinolone susceptibility had an OD_{570} of 0.86, while fluoroquinolone resistance had an OD_{570} of 0.42. In other words, fluoroquinolone susceptible

isolates were predicted to be strong biofilm formers, and fluoroquinolone resistant isolates were predicted to be moderate biofilm formers.

DISCUSSION

This is the first study to our knowledge, to identify a statistically significant inverse relationship between *K. pneumoniae* antimicrobial resistance and biofilm formation, where carbapenem-resistant *K. pneumoniae* isolates were 91% less likely to be strong biofilm formers. Several published studies have described higher biofilm formation in resistant *K. pneumoniae* isolates, however, these descriptive studies did not assess whether resistance was predictive of biofilm formation in multivariate analyses (10-14). This is also the first study to assess resistance and biofilm formation of a diverse collection of *K. pneumoniae* isolates from multiple sources and centers. Inclusion of isolates only from a single-center introduces potential bias if patients are infected with the same organism, especially if isolates are from an outbreak (14). Additional rationale for the difference in findings requires further research.

Potential limitations of our study include the overall resistance patterns of our isolates. The majority of our isolates (n=81, 87.1%) were MDR, with the most isolates (n=64, 68.8%) resistant to at least 12 out of 16 antimicrobial classes, but only 26.9% (n=25) were XDR. Inclusion of more susceptible or XDR isolates may have resulted in different predictors of biofilm formation. As a post-hoc sensitivity analysis we excluded XDR isolates, which support the findings from the weak versus strong analysis, where strong biofilm formation was 30% less likely with each increase in the number of resistant categories.

Findings from previous studies may also be limited by misclassification of biofilm formation, which may affect conclusions that biofilm formation is more common among resistant isolates. The definition of biofilm formation varies across studies but the most common definition was originally

described for *Staphylococcus spp.* (15, 16). This method utilizes an OD cut-off (OD_c), defined as three standard deviations above the average OD of the negative control, to determine biofilm formation. Isolates are either non-adherent (OD ≤ OD_c), weakly adherent (OD_c < OD ≤ 2xOD_c), moderately adherent (2xOD_c < OD ≤ 4xOD_c), or strongly adherent (4x OD_c < OD). Categorization by this method however, is limited when the negative control has a negligible OD reading, which was the case for our study and previously described literature (12, 13). Applying this method to our cohort, zero isolates were non-adherent, two weak, 12 moderate, and 125 strong biofilm formers. Therefore, we divided biofilm formation into tertiles (26), to overcome potential bias of overestimating strong biofilm formation (12). Previous utilization of tertile biofilm categorization was also utilized for *S. aureus*, however categorization should not be affected by organism as our biofilm quantification method utilized was adapted for *K. pneumoniae*. We also assessed biofilm formation as a continuous variable since there is no standard categorization. However, interpretation of resulting odds ratios is challenging when compared to a dichotomous variable of resistance versus susceptible, and is less clinically meaningful.

Optical density remains an indirect measurement of biofilm formation and standardized methods for both quantification and categorization of *K. pneumoniae* biofilm formation are needed. Varied definitions of biofilm formation are utilized, making direct comparisons across studies difficult. It is imperative to standardize biofilm quantification to allow for accurate assessment of predictors of biofilm across settings and to quantify the relationship between biofilm forming isolates and clinical outcomes.

In our study, strong biofilm formation was 91% less likely with carbapenem-resistant *K. pneumoniae*, which allows clinicians to better predict *K. pneumoniae*'s ability to produce biofilm by phenotypic resistance. This inverse relationship between biofilm formation and antibiotic resistance suggests that virulence may be a trade-off for bacterial survival.

Acknowledgements: The information provided in this manuscript are those of the authors and do not necessarily reflect the position or policy of the United States Department of Veterans Affairs. This work was supported in part, by the Office of Academic Affiliations (OAA) at the Department of Veterans Affairs, and by the COIN: Center of Innovation in Long-Term Services and Supports for Vulnerable Veterans; Providence, RI.

Funding: This work was unfunded and the authors received no outside funding for this work.

References

1. Won SY, Munoz-Price LS, Lolans K, Hota B, Weinstein RA, Hayden MK, Centers for Disease C, Prevention Epicenter P. 2011. Emergence and rapid regional spread of *Klebsiella pneumoniae* carbapenemase-producing Enterobacteriaceae. *Clin Infect Dis* 53:532-40.
2. Patel G, Huprikar S, Factor SH, Jenkins SG, Calfee DP. 2008. Outcomes of carbapenem-resistant *Klebsiella pneumoniae* infection and the impact of antimicrobial and adjunctive therapies. *Infect Control Hosp Epidemiol* 29:1099-106.
3. Ribeiro SM, Cardoso MH, Candido Ede S, Franco OL. 2016. Understanding, preventing and eradicating *Klebsiella pneumoniae* biofilms. *Future Microbiol* 11:527-38.
4. Clegg S, Murphy CN. 2016. Epidemiology and Virulence of *Klebsiella pneumoniae*. *Microbiol Spectr* 4.
5. Schwaderer AL, Wolfe AJ. 2017. The association between bacteria and urinary stones. *Ann Transl Med* 5:32.
6. Nickel JC, Reid G, Bruce AW, Costerton JW. 1986. Ultrastructural microbiology of infected urinary stone. *Urology* 28:512-5.
7. Lebeaux D, Chauhan A, Rendueles O, Beloin C. 2013. From in vitro to in vivo Models of Bacterial Biofilm-Related Infections. *Pathogens* 2:288-356.
8. De Rosa FG, Corcione S, Cavallo R, Di Perri G, Bassetti M. 2015. Critical issues for *Klebsiella pneumoniae* KPC-carbapenemase producing *K. pneumoniae* infections: a critical agenda. *Future Microbiol* 10:283-94.

- 251 9. Hoiby N, Bjarnsholt T, Moser C, Bassi GL, Coenye T, Donelli G, Hall-Stoodley L, Høla
252 V, Imbert C, Kirketerp-Møller K, Lebeaux D, Oliver A, Ullmann AJ, Williams C,
253 Biofilms ESGf, Consulting External Expert Werner Z. 2015. ESCMID guideline for the
254 diagnosis and treatment of biofilm infections 2014. Clin Microbiol Infect 21 Suppl 1:S1-
255 25.
- 256 10. Subramanian P, Shanmugam N, Sivaraman U, Kumar S, Selvaraj S. 2012. Antibiotic
257 resistance pattern of biofilm-forming uropathogens isolated from catheterised patients in
258 Pondicherry, India. Australas Med J 5:344-8.
- 259 11. Sanchez CJ, Jr., Mende K, Beckius ML, Akers KS, Romano DR, Wenke JC, Murray CK.
260 2013. Biofilm formation by clinical isolates and the implications in chronic infections.
261 BMC Infect Dis 13:47.
- 262 12. Vuotto C, Longo F, Pascolini C, Donelli G, Balice MP, Libori MF, Tiracchia V, Salvia
263 A, Varaldo PE. 2017. Biofilm formation and antibiotic resistance in *Klebsiella*
264 *pneumoniae* urinary strains. J Appl Microbiol 123:1003-1018.
- 265 13. Yang D, Zhang Z. 2008. Biofilm-forming *Klebsiella pneumoniae* strains have greater
266 likelihood of producing extended-spectrum beta-lactamases. J Hosp Infect 68:369-71.
- 267 14. Bocanegra-Ibarias P, Garza-Gonzalez E, Morfin-Otero R, Barrios H, Villarreal-Trevino
268 L, Rodriguez-Noriega E, Garza-Ramos U, Petersen-Morfin S, Silva-Sanchez J. 2017.
269 Molecular and microbiological report of a hospital outbreak of NDM-1-carrying
270 Enterobacteriaceae in Mexico. PLoS One 12:e0179651.
- 271 15. Christensen GD, Simpson WA, Younger JJ, Baddour LM, Barrett FF, Melton DM,
272 Beachey EH. 1985. Adherence of coagulase-negative staphylococci to plastic tissue

culture plates: a quantitative model for the adherence of staphylococci to medical devices. *J Clin Microbiol* 22:996-1006.

16. Stepanovic S, Vukovic D, Hola V, Di Bonaventura G, Djukic S, Cirkovic I, Ruzicka F. 2007. Quantification of biofilm in microtiter plates: overview of testing conditions and practical recommendations for assessment of biofilm production by staphylococci. *APMIS* 115:891-9.
17. de Campos PA, Royer S, Batistao DW, Araujo BF, Queiroz LL, de Brito CS, Gontijo-Filho PP, Ribas RM. 2016. Multidrug Resistance Related to Biofilm Formation in *Acinetobacter baumannii* and *Klebsiella pneumoniae* Clinical Strains from Different Pulsotypes. *Curr Microbiol* 72:617-27.
18. Silva A, Sousa AM, Alves D, Lourenco A, Pereira MO. 2016. Heteroresistance to colistin in *Klebsiella pneumoniae* is triggered by small colony variants sub-populations within biofilms. *Pathog Dis* 74.
19. Luther MK, Parente DM, Caffrey AR, Daffinee KE, Lopes VV, Martin ET, LaPlante KL. 2018. Clinical and Genetic Risk Factors for Biofilm-Forming *Staphylococcus aureus*. *Antimicrob Agents Chemother* 62.
20. O'Toole GA. 2011. Microtiter dish biofilm formation assay. *J Vis Exp* doi:10.3791/2437.
21. Luther MK, Bilida S, Mermel LA, LaPlante KL. 2015. Ethanol and Isopropyl Alcohol Exposure Increases Biofilm Formation in *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Infect Dis Ther* 4:219-26.
22. Naves P, del Prado G, Huelves L, Gracia M, Ruiz V, Blanco J, Rodriguez-Cerrato V, Ponte MC, Soriano F. 2008. Measurement of biofilm formation by clinical isolates of *Escherichia coli* is method-dependent. *J Appl Microbiol* 105:585-90.

- 296 23. Naparstek L, Carmeli Y, Navon-Venezia S, Banin E. 2014. Biofilm formation and
297 susceptibility to gentamicin and colistin of extremely drug-resistant KPC-producing
298 *Klebsiella pneumoniae*. *J Antimicrob Chemother* 69:1027-34.
- 299 24. Liu C, Zhou Y, Wang L, Han L, Lei J, Ishaq HM, Nair SP, Xu J. 2016. Photodynamic
300 inactivation of *Klebsiella pneumoniae* biofilms and planktonic cells by 5-aminolevulinic
301 acid and 5-aminolevulinic acid methyl ester. *Lasers Med Sci* 31:557-65.
- 302 25. Baldiris R TV, Vivas-Reyes R, Montes A, Arzuza O. 2016. Anti-biofilm activity of
303 ibuprofen and diclofenac against some biofilm producing *Escherichia coli* and *Klebsiella*
304 *pneumoniae* uropathogens. *Afr J Microbiol Res* 10:1675-1684.
- 305 26. Guembe M, Alonso B, Lucio J, Perez-Granda MJ, Cruces R, Sanchez-Carrillo C,
306 Fernandez-Cruz A, Bouza E. 2017. Biofilm production is not associated with poor
307 clinical outcome in 485 patients with *Staphylococcus aureus* bacteraemia. *Clin Microbiol*
308 *Infect* doi:10.1016/j.cmi.2017.10.018.
- 309 27. Anonymous. Clinical and Laboratory Standards Institute. Performance standards for
310 antimicrobial susceptibility testing: 27th informational supplement, (M100-S27). Wayne,
311 PA: CLSI; 2017.
- 312 28. Anonymous. Clinical and Laboratory Standards Institute. Methods for dilution
313 antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard:
314 tenth edition, (M07-A10):CLSI; 2015.
- 315 29. Tygacil. (tigecycline) [package insert]. Wyeth Pharmaceuticals Inc; Philadelphia
316 (PA):2005.

30. Anonymous. The European Committee on Antimicrobial Susceptibility Testing.
Breakpoint tables for interpretation of MICs and zone diameters. Version 7.1, 2017.
<http://www.eucast.org>.
31. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ, Vatopoulos A, Weber JT, Monnet DL. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. Clin Microbiol Infect 18:268-81.

342 **Table 1.** *Klebsiella pneumoniae*: antimicrobial resistance and biofilm formation

Variable	Total Cohort (n=93)	Weak Biofilm Formation (n=47)	Strong Biofilm Formation (n=46)	p-value
Number of Resistant Categories (n=16), Median, (IQR)	13 (11-14)	13 (12-14)	11.5 (3-14)	0.01
Multidrug-resistant (MDR), n (%) [*]	81 (87.1)	46 (97.9)	35 (76.1)	0.002
Extensively drug-resistant (XDR), n (%) ^{**}	25 (26.9)	12 (25.5)	13 (28.3)	0.77
Penicillins + β -lactamase inhibitors, n (%) [†]	79 (84.9)	46 (97.9)	33 (71.7)	0.0004
Cephalosporins, n (%)	82 (88.2)	46 (97.9)	36 (78.3)	0.003
Monobactam, n (%)	73 (78.5)	45 (95.7)	28 (60.9)	<0.0001
Carbapenems, n (%) ^{††}	70 (75.3)	44 (93.6)	26 (56.5)	<0.0001
Aminoglycosides, n (%)	72 (77.4)	43 (91.5)	29 (63.0)	0.001
Chloramphenicol, n (%)	65 (69.9)	38 (80.9)	27 (58.7)	0.02
Fluoroquinolones, n (%)	73 (78.5)	45 (95.7)	28 (60.9)	<0.0001
Tigecycline, n (%)	13 (14.0)	6 (12.8)	7 (15.2)	0.73
Tetracyclines, n (%)	44 (47.3)	23 (48.9)	21 (45.7)	0.75
Folate pathway inhibitor, n (%)	66 (71.0)	37 (78.7)	29 (63.0)	0.09
Fosfomycin, n (%)	61 (65.6)	29 (61.7)	32 (69.6)	0.42
Colistin, n (%)	11 (11.8)	8 (17.0)	3 (6.5)	0.12

343 **Bolded p-values indicate potential predictors of strong biofilm formation included in the initial**
344 **logistic regression model**

*MDR isolates demonstrated non-susceptibility to at least one agent in three or more antimicrobial categories out of 16 antimicrobial categories

**XDR isolates demonstrated susceptibility to at least one agent in less than or equal to two out of 16 antimicrobial categories

*penicillin + β -lactamase inhibitors category includes piperacillin/tazobactam and penicillin/ β -lactamase inhibitors

*cephalosporins category includes non-extended spectrum cephalosporins, extended-spectrum cephalosporins, cephamycins, and ceftaroline

Table 2. Sensitivity analysis of non-XDR *Klebsiella pneumoniae*: antimicrobial resistance and biofilm formation

Variable	Total Cohort (n=68)	Weak Biofilm Formation (n=35)	Strong Biofilm Formation (n=33)	p-value
Number of Resistant Categories (n=16), Median, (IQR)	12 (8-13)	13 (12-13)	9 (1-12)	<0.0001
Multidrug-resistant (MDR), n (%) [*]	56 (82.4)	34 (97.1)	22 (66.7)	0.001
Extensively drug-resistant (XDR), n (%) ^{**}	0	0	0	---
Penicillins + β -lactamase inhibitors, n (%)	54 (79.4)	34 (97.1)	20 (60.6)	0.0002
Cephalosporins, n (%)	57 (83.8)	34 (97.1)	23 (69.7)	0.002
Monobactam, n (%)	48 (70.6)	33 (94.3)	15 (45.5)	<0.0001
Carbapenems, n (%)	45 (66.2)	32 (91.4)	13 (39.4)	<0.0001
Aminoglycosides, n (%)	48 (70.6)	32 (91.4)	16 (48.5)	0.001
Chloramphenicol, n (%)	40 (58.8)	26 (74.3)	14 (42.4)	0.008
Fluoroquinolones, n (%)	48 (70.6)	33 (94.3)	15 (45.5)	<0.001
Tigecycline, n (%)	6 (8.8)	4 (11.4)	2 (6.1)	0.67
Tetracyclines, n (%)	20 (29.4)	12 (34.3)	8 (24.2)	0.36
Folate pathway inhibitor, n (%)	41 (60.3)	25 (71.4)	16 (48.5)	0.053
Fosfomycin, n (%)	41 (60.3)	19 (54.3)	22 (66.7)	0.30
Colistin, n (%)	6 (8.8)	4 (11.4)	2 (6.1)	0.44

Bolded p-values indicate potential predictors of strong biofilm formation included in the initial logistic regression model

372 *MDR isolates demonstrated non-susceptibility to at least one agent in three or more
373 antimicrobial categories out of 16 antimicrobial categories
374 **XDR isolates demonstrated susceptibility to at least one agent in less than or equal to two out
375 of 16 antimicrobial categories
376 ‡penicillin + β -lactamase inhibitors category includes piperacillin/tazobactam and penicillin/ β -
377 lactamase inhibitors
378 ††cephalosporins category includes non-extended spectrum cephalosporins, extended-
379 spectrum cephalosporins, cephamycins, and ceftaroline
380
381
382
383
384
385
386
387

Figure 1. *Klebsiella pneumoniae* biofilm formation and resistance (n=139)

388

389 (2 column fitting image)