

2017

## Aquatic Global Passive Sampling (AQUA-GAPS) Revisited – First Steps towards a Network of Networks for Organic Contaminants in the Aquatic Environment

Rainer Lohmann

*University of Rhode Island, rlohmann@uri.edu*

Derek Muir

Eddy Y. Zeng

Lian-Jun Bao

Ian J. Allan

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

---

### Citation/Publisher Attribution

Lohmann, Rainer; Muir, Derek; Zeng, Eddy; Bao, Lian-Jun; Allan, Ian; Arinaitwe, Kenneth; Booij, Kees; Helm, Paul; Kaserzon, Sarit; Mueller, Jochen; Shibata, Yasuyuki; Smedes, Foppe; Tsapakis, Manolis; Wong, Charles; You, Jing. "Aquatic Global Passive Sampling (AQUA-GAPS) Revisited – First Steps towards a Network of Networks for Organic Contaminants in the Aquatic Environment". *Environ Sci Technol* 2017, 51, 1060-1067. DOI: 10.1021/acs.est.6b05159

Available at: <http://dx.doi.org/10.1021/acs.est.6b05159>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons-group@uri.edu](mailto:digitalcommons-group@uri.edu). For permission to reuse copyrighted content, contact the author directly.

---

# Aquatic Global Passive Sampling (AQUA-GAPS) Revisited – First Steps towards a Network of Networks for Organic Contaminants in the Aquatic Environment

## Authors

Rainer Lohmann, Derek Muir, Eddy Y. Zeng, Lian-Jun Bao, Ian J. Allan, Kenneth Arinaitwe, Kees Booij, Paul Helm, Sarit Kaserzon, Jochen F. Mueller, Yasukuki Shibata, Foppe Smedes, Manolis Tsapakis, Charles S. Wong, and Jing You

The University of Rhode Island Faculty have made this article openly available.  
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

## Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

1 **Aquatic Global Passive Sampling (AQUA-GAPS) Revisited – First**  
2 **Steps towards a Network of Networks for Organic Contaminants in**  
3 **the Aquatic Environment**

4  
5 Rainer Lohmann,<sup>1,\*</sup> Derek Muir,<sup>2,3</sup> Eddy Y. Zeng,<sup>3</sup> Lian-Jun Bao,<sup>3</sup> Ian J. Allan,<sup>4</sup>  
6 Kenneth Arinaitwe,<sup>5</sup> Kees Booij,<sup>6</sup> Paul Helm,<sup>7</sup> Sarit Kaserzon,<sup>8</sup> Jochen F. Mueller,<sup>8</sup>  
7 Yasuyuki Shibata,<sup>9</sup> Foppe Smedes,<sup>10</sup> Manolis Tsapakis,<sup>11</sup> Charles S. Wong,<sup>3,12</sup> Jing  
8 You<sup>3</sup>

9  
10 <sup>1</sup> Graduate School of Oceanography, University of Rhode Island, Narragansett, Rhode Island  
11 02882-1197, United States.

12 <sup>2</sup> Environment and Climate Change Canada, Aquatic Contaminants Research Division, 867  
13 Lakeshore Road, Burlington, Ontario, Canada L7S 1A1

14 <sup>3</sup> School of Environment, Guangzhou Key Laboratory of Environmental Exposure and Health  
15 and Guangdong Key Laboratory of Environmental Pollution and Health, Jinan University,  
16 Guangzhou 510632, China

17 <sup>4</sup> Norwegian Institute for Water Research (NIVA), Gaustadalleen 21, NO-0349 Oslo, Norway

18 <sup>5</sup> Department of Chemistry, College of Natural Sciences, Makerere University, P. O. Box 7062,  
19 Kampala, Uganda.

20 <sup>6</sup> PaSOC, Greate Pierwei 25, 8821 LV Kimswerd, The Netherlands

21 <sup>7</sup> Environmental Monitoring and Reporting Branch, Ontario Ministry of the Environment and  
22 Climate Change, Toronto, Ontario, Canada M9P 3V6

23 <sup>8</sup> The Queensland Alliance for Environmental Health Sciences (QAEHS), The University of  
24 Queensland, 39 Kessels Road, Coopers Plains, QLD 4108, Australia.

25 <sup>9</sup> Center for Environmental Measurement and Analysis, National Institute for Environmental  
26 Studies, 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan

27 <sup>10</sup> Research Center for Toxic Compounds in the Environment (RECETOX), Masaryk University,  
28 Kamenice 5/573, 62500 Brno, Czech Republic

29 <sup>11</sup> Hellenic Centre for Marine Research, Institute of Oceanography, Gournes, Crete, Greece

30 <sup>12</sup> Richardson College for the Environment, University of Winnipeg, Winnipeg MB R3B 2E9  
31 Canada

32 \*Corresponding author. E-mail address: [rlohmann@uri.edu](mailto:rlohmann@uri.edu);

33 Phone: 401-874-6612; Fax 401-874-6811

34

## 35 **ABSTRACT**

36 Organic contaminants, in particular persistent organic pollutants (POPs), adversely affect water  
37 quality and aquatic food webs across the globe. As of now, there is no globally consistent  
38 information available on concentrations of dissolved POPs in water bodies. The advance of  
39 passive sampling techniques has made it possible to establish a global monitoring program for  
40 these compounds in the waters of the world, which we call the Aquatic Global Passive Sampling  
41 (AQUA-GAPS) network. A recent expert meeting discussed the background, motivations, and  
42 strategic approaches of AQUA-GAPS, and its implementation as a network of networks for  
43 monitoring organic contaminants (e.g., POPs and others contaminants of concern). Initially,  
44 AQUA-GAPS will demonstrate its operating principle via two proof-of-concept studies focused

45 on the detection of legacy and emerging POPs in freshwater and coastal marine sites using both  
46 polyethylene and silicone passive samplers. AQUA-GAPS is set-up as a decentralized network,  
47 which is open to other participants from around the world to participate in deployments and to  
48 initiate new studies. In particular, participants are sought to initiate deployments and studies  
49 investigating the presence of legacy and emerging POPs in Africa, Central and South America.

50

## 51 ■ INTRODUCTION

52 Recognizing the achievements of the Global Atmospheric Passive Sampling program (GAPS),<sup>1,2</sup>  
53 Lohmann and Muir (2010) called for the establishment of Aquatic Global Passive Sampling  
54 (AQUA-GAPS), aiming to understand better the geographical distributions and temporal trends  
55 of organic contaminants, such as persistent organic pollutants (POPs), polycyclic aromatic  
56 hydrocarbons (PAHs), novel flame retardants and other contaminants of emerging concern.<sup>3</sup>  
57 AQUA-GAPS has the potential to facilitate the implementation of the Stockholm Convention  
58 (SC) on POPs, a global treaty under the United Nations Environmental Programme (UNEP) with  
59 the objective to protect human health and the environment from hazardous, long-lasting,  
60 bioaccumulative chemicals with long-range transport potential by restricting and ultimately  
61 eliminating their production, use, trade, and release.<sup>4</sup> Yet the scope of AQUA-GAPS goes  
62 beyond existing POPs by enabling studies into a wide range of organic contaminants.

63 So far, the SC, through its Global Monitoring Plan, measures POPs in air (active and  
64 passive samplers) for capturing their status of emissions and long-range transport, and in human  
65 samples (blood, milk) for assessing exposure status. Water monitoring was added to the Global  
66 Monitoring Plan for PFOS, which is far more water-soluble than legacy POPs; unlike other  
67 POPs, its emission and transport through water and not just air are thought to be significant.<sup>5-7</sup>

68 Reliance upon passive samplers has already been established via the GAPS program, as well as  
69 the Europe/Africa/Asian Monitoring NETWORKS (MONET), Latin American Passive  
70 Atmospheric Sampling (LAPAN), and UNEP/Global Environmental Facility (GEF) projects, all  
71 of which utilize passive air sampling devices at monitoring sites on all continents, mostly in  
72 remote regions, demonstrating the potential for global coverage.<sup>7,8</sup> While data from GAPS does  
73 address the atmospheric compartment and potentially plants and soils exchanging with air, it  
74 does not readily address prevailing concentrations or trends in aquatic environments. The  
75 aquatic environment represents a key compartment for many POPs, most notably for the HCH  
76 isomers and endosulfan<sup>9</sup>, and dissolved concentrations can be used to estimate human and  
77 wildlife exposure using bioaccumulation factors and food chain models.<sup>10-12</sup>

78 Passive samplers offer key benefits for global monitoring of aqueous contaminants,  
79 because of their high enrichment of their target analytes, and the ability to measure time-  
80 weighted average concentrations.<sup>13-15</sup> Most importantly, a key benefit consists of being able to  
81 expose the same sampler in all waters of the world, which cannot be achieved with any  
82 biological or other abiotic matrix. Passive samplers are also more cost-effective and relatively  
83 easier to handle for shipment and deployment than active sampling of large volumes of water.

84 The atmospheric GAPS program is based on monitoring sites at a defined height above  
85 ground that are relatively easy to access. The logistic requirements for AQUA-GAPS sites are  
86 inherently more challenging for on- and off-shore deployments and retrieval, requiring moorings  
87 and boat time, among other practical issues. The biggest hurdle for establishing a realistic  
88 AQUA-GAPS program is perhaps whether enough willing and capable participants from around  
89 the world can be secured to agree on and perform the logistics of field and laboratory work.  
90 GAPS samplers are often deployed at already established and protected atmospheric monitoring

91 sites that are part of the World Meteorological Organization (WMO) network.<sup>16</sup> In parallel,  
92 AQUA-GAPS intends to deploy passive samplers at selected remote/background sites in water  
93 bodies around the globe. Similar to the GAPS program, though, AQUA-GAPS will also monitor  
94 selected urban/industrially impacted sites in an attempt to examine the impacts of anthropogenic  
95 activities on aquatic environments at a global scale.<sup>17</sup>

96 A meeting of 15 passive sampling and monitoring experts from 10 countries covering 5  
97 continents was organized at Jinan University, Guangzhou, China, on 21–22 January, 2016,  
98 aiming to make progress towards the establishment of AQUA-GAPS. In particular, the group  
99 was tasked with addressing whether and how it will be feasible to make the assessment of global  
100 POP distributions through an analysis of global passive sampling devices in waters. The group  
101 then outlined the key steps for implementing them during the meeting. The aim of this feature  
102 article is to detail the framework, approach, and expectation of AQUA-GAPS, and solicit  
103 additional participation to cover extended sampling areas and initiate new global studies.

104

## 105 ■ THE MOTIVATIONS BEHIND AQUA-GAPS

106 To launch AQUA-GAPS successfully, it was the consensus of the workshop participants that  
107 prior experiences of GAPS must be learned and assimilated. The GAPS program has  
108 demonstrated that a global monitoring network using passive samplers is feasible and can be  
109 successfully implemented. GAPS was successful in establishing spatial distributions of targeted  
110 chemicals, while the identification of temporal trends requires longer time and continuing  
111 resource commitment. GAPS has been particularly impressive by making use of their samples  
112 for a wide range of contaminants, such as polychlorinated biphenyls (PCBs), polychlorinated  
113 naphthalenes (PCNs), polybrominated diphenylethers (PBDEs), neutral and ionic perfluoroalkyl

114 substances (PFASs).<sup>2,18,19</sup> GAPS has also demonstrated flexibility with their sampling matrix.  
115 Initially, GAPS relied on the use of polyurethane foam (PUF) disks, which were later modified  
116 to include sampling of compounds with higher volatility in accordance with the increase of list of  
117 POPs under the SC.<sup>20,21</sup> The key lesson here is that flexibility in the type of passive samplers  
118 may be required to respond to changing regulatory and scientific needs and interests (new  
119 compounds of emerging concern, novel samplers, etc.).

120 The success of GAPS and its relevance to the SC are a prime motivating factor for  
121 establishing AQUA-GAPS. Although the current priority sampling matrices under the SC are  
122 limited to air and human samples (and water in the case of PFOS), the workshop participants felt  
123 that AQUA-GAPS would have significant added value in yielding highly comparable  
124 concentration data that allow for better assessing and understanding of the role of water in the  
125 global fate of POPs, and in human and wildlife exposure to these chemicals. Both the data  
126 collection and the enhanced understanding of global distributions and trends in the aquatic  
127 environment would be beneficial for the SC, its Regional Organizational Groups and potentially  
128 to other international conventions such as the International Marine Organization's London  
129 Convention on prevention of marine pollution by dumping of wastes<sup>22</sup>, and Convention on the  
130 Control of Harmful Anti-fouling Systems on Ships.<sup>23</sup> The data would also benefit regional and  
131 national legislative frameworks, such as the EU Water Framework Directive<sup>24</sup> and Marine  
132 Strategy Framework Directive<sup>25</sup>, and the 18 Regional Sea Programs under UNEP<sup>26</sup>, the United  
133 States Toxic Substances Control Act (recently updated and renamed the Frank R. Lautenberg  
134 Chemical Safety for the 21st Century Act)<sup>27</sup>, the EU REACH legislation<sup>28</sup>, and environmental  
135 protection and clean water legislation in many other countries.



136 The workshop proposed to have AQUA-GAPS focus on legacy and emerging POPs, as well as  
137 on other compounds of emerging concern both hydrophilic (e.g., pharmaceuticals and personal  
138 care products) and hydrophobic (novel flame retardants). Currently, the SC's main interest  
139 would be the development and deployment of passive samplers for perfluorinated compounds,  
140 such as PFOS and PFOA (which is currently under review for inclusion in the SC)<sup>29</sup> in waters  
141 around the globe. The SC has developed guidance for sample collection and determination of  
142 baseline levels of PFOS in water, caused by global dispersion/diffusion.<sup>30</sup> Yet the availability of  
143 high quality, consistent global concentration maps and trends on POPs and other contaminants  
144 will certainly support the SC and its regional programs.

145

#### 146 ■ A NETWORK OF NETWORKS

147 The GAPS program has been established around one central laboratory at Environment Canada,  
148 from which samplers are prepared, shipped, returned to, analyzed, and interpreted. This model is  
149 unlikely to be repeatable for AQUA-GAPS<sup>1</sup>. Instead, we propose to establish a '**network of**  
150 **networks**' open to anybody to participate in, but clustered around a central laboratory for sampler  
151 preparation and core analysis, the Research Center for Toxic Compounds in the Environment  
152 (RECETOX, Masaryk University, Czech Republic). Initially, the network consists of a group of  
153 scientists with experience in working with passive samplers. The proposed *modus operandi* of the  
154 AQUA-GAPS network consists of in-kind contributions of participating scientists to deploy  
155 passive samplers to the best of their abilities, and share ancillary data with respect to their sites.  
156 In return, the expectation is one of data sharing by the leading team and inclusion in the data  
157 discussion and interpretation. Possible authorship will depend on contributions to the interpretation

158 and discussion of results. We foresee AQUA-GAPS to be a platform in which scientists offer  
159 mutual help in deploying samplers at specified locations (e.g., wastewater treatment plants, rivers,  
160 freshwater lakes, coastal seas, and oceans) and site characterization (e.g., urban, industrial, and  
161 remote). Examples of potential aquatic networks include:

162 *An oceans network.* This is logistically the most challenging, and would often require  
163 deployment time of several months up to 1 year, which is typical for open ocean mooring  
164 turnaround time.<sup>31,32</sup> In view of low concentrations, samplers would need to be designed to  
165 maximize the uptake of the target compounds to overcome detection limits for as many  
166 compounds as possible. Most likely target compounds are legacy POPs, non-polar current use  
167 pesticides, organophosphorus flame retardants, perfluorinated compounds, and other chemicals  
168 of interest that accumulate in samplers for hydrophobic POPs, such as hydrocarbons or natural  
169 halogenated compounds. The benefits of working with the oceanographic community's set of  
170 moorings is the general availability of ship-time and access to ancillary data.

171 *A coastal/estuarine network.* Coastal and estuarine sites are easier to reach for  
172 deployments, and often coincide with major fishing grounds, which makes them relevant for  
173 human exposure and links to biomonitoring data. Deployment time can be shorter, in view of  
174 greater concentrations and challenges linked to biofouling of samplers during deployments in  
175 productive water bodies.

176 *A lakes network.* Lakes and reservoirs are of high relevancy for human and ecosystem  
177 exposure as they are regularly used for aquaculture and irrigation, and often serve as a source of  
178 drinking water. Remote lakes (e.g., Experimental Lakes Area in Canada) can serve to quantify  
179 background concentrations associated with minimal anthropogenic impacts.<sup>30</sup> For both estuarine

180 and freshwater, sampler deployment at near-shore sites is straightforward, but carries a risk of  
181 sampler loss through theft, vandalism and loss from accidental ship strikes and fishing efforts.

182 *A network of source waters (waste water treatment plant effluents and rivers).* This  
183 network could be used to identify the compounds being introduced into lakes and oceans, before  
184 they become of concern. This network could act as an early warning system to identify  
185 chemicals of concern through their release to the aquatic environment from human activity and  
186 global spread. As concentrations of contaminants in such a network are likely to be much  
187 greater than in the other networks mentioned, this network would lend itself to target less  
188 persistent compounds, including breakdown products. Identification of major sources and  
189 establishment of reliable inventories are indispensable for the efficient and effective management  
190 of chemicals on the national, regional and global scale. The various AQUA-GAPS networks  
191 should each aim to collaborate and communicate with relevant stakeholders in local, national, or  
192 regional levels.

193

#### 194 ■ TECHNICAL FOUNDATION FOR AQUA-GAPS

195 Passive sampling in the water gives a direct measure of a chemicals' activity (or fugacity) in the  
196 water, as only freely dissolved contaminants diffuse into the passive sampler material.<sup>33</sup> Thus the  
197 freely dissolved concentration derived for passive sampler accumulated pollutants can be used to  
198 assess the gradient of chemical activities between different media (air, water, sediment, and  
199 biota), and these freely dissolved contaminant concentrations and chemical activities are more  
200 useful to assess net fluxes among environmental compartments and bioaccumulation in  
201 organisms.<sup>34-38</sup> Passive sampling derived dissolved pollutant concentrations are therefore

202 fundamentally different from total concentrations reported for e.g., sediment or active water  
203 samples.

204         The novelty that AQUA-GAPS introduces is a coordinated effort at pollutant sampling  
205 with a chosen passive sampler for worldwide deployments towards the generation of globally  
206 comparative data sets. Passive sampling with a well-characterized polymer/sampler can help  
207 achieve a level of standardization on a global scale that cannot really be obtained with other  
208 environmental matrices (e.g. biota, sediments, etc.) due to their variable properties. Nowadays,  
209 silicone rubber and polyethylene are the two most widely used polymers for passive sampling of  
210 hydrophobic organic contaminants in water.<sup>39</sup> The increased use of these polymers is partly due  
211 to the availability of calibration data, i.e., polymer diffusion coefficients and polymer-water  
212 partition coefficients for a number of non-ionised hydrophobic chemicals <sup>33,40–42</sup> including  
213 chemicals of emerging concern.<sup>43</sup> These absorption-based passive samplers offer the opportunity  
214 to use performance reference compounds (PRCs) to assess contaminant exchange kinetics  
215 between water and the polymer in situ for every deployment location and exposure period.<sup>44</sup> In  
216 addition, these polymers facilitate the comparison of contaminant levels in different  
217 environmental compartments (i.e. air, biota, sediment or water). The critical review by Booij et  
218 al. demonstrated that absorption-based passive sampling is today the best available tool for  
219 chemical monitoring of non-ionised hydrophobic chemicals in the aquatic environment.<sup>45</sup> While  
220 a lack of robust quality assurance was identified as a weakness of passive sampling in water,  
221 recent results from the QUASIMEME Proficiency Testing schemes conducted using silicone  
222 rubber were very encouraging.<sup>39</sup> These results show that the analysis of passive samplers within  
223 the AQUA-GAPS network may not ultimately require analysis of all samplers by a single  
224 laboratory, so long as proficiency testing schemes are organized regularly to evaluate the

225 performance of participating laboratories. At least initially, though, AQUA-GAPS studies will be  
226 organized around one central laboratory, RECETOX, for the above mentioned legacy and  
227 emerging pollutants from using SR and PE passive samplers prepared and analyzed there.

228

## 229 ■ STRATEGIES FOR FIELD SAMPLING AND DATA ASSIMILATION

230 The unique feature of AQUA-GAPS is that studies can be initiated by anybody with an interest  
231 in answering a global question linked to contaminants in water. The lead team initiating an  
232 AQUA-GAPS sampling campaign needs to have sufficient resources to organize sampler  
233 preparation, distribution/retrieval, analysis, and interpretation. AQUA-GAPS is intended to be  
234 flexible on which sampler to use, how to deploy it, for how long, and where. This will all be  
235 decided by the leading team, who will ask others to participate in-kind by deploying in their  
236 water body (Figure 1). It may be cost efficient to deploy different types of samplers targeting a  
237 wide range of compounds simultaneously (e.g., nonpolar, hydrophilic neutral, positively and  
238 negatively charged, etc.).<sup>46</sup> Passive samplers are relatively inexpensive, and have the potential to  
239 be archived.<sup>47</sup>

240 Similar to the GAPS deployments, AQUA-GAPS' challenge will be on how to identify  
241 sites in aquatic environments suitable for evaluating spatial and temporal changes in contaminant  
242 levels, that would therefore help assess the effectiveness of the control measures implemented  
243 under the SC and/or regional efforts. AQUA-GAPS networks will benefit greatly by selecting  
244 sites overlapping with other continuous sampling efforts and programs, such as existing GAPS  
245 stations. Many GAPS samplers were strategically placed at WMO sites, such that GAPS had  
246 site-specific meteorological data available. It will be important to work with sites and groups that  
247 are capable of conducting repeatable, long-term deployments.

248 Beyond GAPS sites, other examples for AQUA-GAPS include lakes, ocean and coastal sea  
249 monitoring initiatives using regularly serviced buoys and moorings, which can provide ancillary  
250 data (temperature profiles, salinity, and current data), and potentially ships of opportunity. There  
251 are also on-going contaminant sampling initiatives or networks (Canada's National Water  
252 Quality Monitoring Program <sup>48</sup>, The Great Barrier Reef Marine Monitoring Program <sup>49</sup>, etc.)  
253 which can contribute to AQUA-GAPS for mutual benefits. AQUA-GAPS will of course be open  
254 to other programs as it evolves.

255         Across the globe various types of passive samplers have been successfully used for the  
256 detection of a range of organic contaminants in waters, resulting in a global ISO standard  
257 protocol.<sup>50</sup> Semi-Permeable Membrane Devices (SPMDs) were arguably the first passive  
258 samplers that were used on a wide geographical scale in water sampling.<sup>51</sup> Over time, other  
259 sampler types, often single-phase polymers, have become more commonplace. Notable  
260 examples include the use of silicone rubber (SR) in an OSPAR-lead initiative across Europe and  
261 Australia <sup>52</sup>, The Great Barrier Reef Marine Monitoring Sampling Campaign <sup>53</sup>, and several  
262 years of polyethylene (PE) deployments across the Great Lakes <sup>54,55</sup> and in the Canadian  
263 Arctic.<sup>56</sup>

264

## 265 ■ **PROOF OF CONCEPT STUDIES: FRESHWATER AND COASTAL AQUA-GAPS**

266 Initially, it will be necessary to establish a proof of concept for AQUA-GAPS to show that this  
267 network of networks can actually achieve meaningful results and global coverage. At the proof  
268 of concept stage, we aim to demonstrate that it is feasible to ship and deploy passive samplers to  
269 participating volunteers around the globe, have them deployed, returned for analysis, and yield  
270 meaningful results. At this stage, two proof-of-concept studies are being planned and performed

271 (Figure 1). In both cases, both polyethylene (PE) and silicone rubber (SR) samplers will be co-  
272 deployed, such that spare samplers are available for archiving. While actual sampler designs  
273 differ between the proof-of-concept studies (Figure 2), both are designed to be easily deployed,  
274 provide basic shelter and house several PE and SR sheets simultaneously. Initially, the analytical  
275 target compounds include polychlorinated biphenyls (PCBs), organochlorine pesticides (OCPs),  
276 polycyclic aromatic hydrocarbons (PAHs), polybrominated diphenyl ethers (PBDEs) and various  
277 hydrophobic novel flame retardants (NFRs). The first proof-of-concept study will focus on  
278 legacy and emerging POPs in lakes (Figure 3); the second study will focus on global passive  
279 sampling deployments at coastal sites. The two proof-of-concept studies illustrate the flexibility  
280 within the AQUA-GAPS network. The first freshwater study shares the responsibilities of  
281 logistics, analysis and interpretation among three research groups, while the coastal study will be  
282 performed by two academic/research laboratories (Figure 1).

283

#### 284 ■ **BEYOND ROUTINE MONITORING**

285 Beyond its focus on sampling of contaminants in water, AQUA-GAPS could be well positioned  
286 to address additional research questions, such as air–water exchange and/or sediment porewater–  
287 overlying water gradients by including additional samplers in adjacent media. As mentioned  
288 above, expected or measured equilibrium polymer concentrations are directly proportional to the  
289 activity of the chemical in the medium being sampled, and can help compare contaminant levels  
290 and gradients in/between various environmental media.<sup>57</sup> This holds true whether equilibrium  
291 between the contaminant concentration in the medium being sampled and the polymer is reached  
292 (e.g. during sediment or biota exposures) or not (e.g. when sampling air or water).<sup>58,59</sup> Having  
293 passive samplers measure freely dissolved concentrations in close proximity to biomonitoring

294 locations can help understand bioaccumulation potential and chemical concentration gradients.  
295 Passive samplers can also be used for non-target screening to detect the presence of other  
296 chemicals, derived from industrial, natural or transformation products. Linking to regional  
297 efforts, particularly biomonitoring programs (e.g., Mussel Watch<sup>60</sup>), AMAP<sup>61</sup> or GAPS<sup>1</sup> seems  
298 particularly useful to enable a comparison of POP concentrations across media. Results from  
299 passive sampling will represent time-weighted-average concentrations in water and thus support  
300 model development and validation, while biological monitoring has a longer history including  
301 archived samples and is often better suited for assessing human exposure (particularly in the case  
302 of edible fishes, shellfish, and other aquatic biota).

303 GAPS has derived part of its strength by having a centralized laboratory (i.e., Environment  
304 Canada) initiating deployments and analyzing all samples within a particular study, in order to  
305 enhance data comparability. As noted, AQUA-GAPS will operate slightly differently, though  
306 RECETOX will perform the sampler preparation and analysis for the routine suite of  
307 hydrophobic compounds (PAHs, PCBs, OCPs, PBDEs and NFRs). Yet AQUA-GAPS will have  
308 different research groups leading deployments, and potentially extra analyses for a specific  
309 project. The leading group will provide passive samplers that are suitable for the specific  
310 compounds of interest to all participating scientists; the samplers will be returned to the same  
311 lead group and analyzed in a single laboratory. Additional samplers can be shared with the local  
312 deploying groups, as a secondary aim of enhancing QA/QC, to enable cross-validation of results,  
313 and the assessment of inter-laboratory variability. This can also lead to capacity-building (see  
314 below). An AQUA-GAPS deployment can be shared/initiated by 2 or more groups, such that  
315 different samplers can be exposed during the same deployment. The leading group needs not  
316 perform all tasks themselves; it could finance another team to produce samplers and deployment



317 cages, organize the distribution, perform the analysis, and calculate the dissolved concentrations,  
318 etc.

319 For AQUA-GAPS to become successful and global, it should lead to global capacity  
320 building linked to passive sampling. There are several regions where little to no information  
321 exists on organic contaminants in water, in particular from Africa, South Asia, Central and South  
322 America. Additionally, non-traditional deployment opportunities can also be leveraged,  
323 including ferrybox samplers, towing samplers and ships of opportunity (expedition vessels;  
324 regular cargo or ferry routes) to target remote locations.<sup>62,63</sup> The network will become more  
325 useful in addressing scientific questions only if more support of AQUA-GAPS deployments is  
326 secured. It would be ideal if the Stockholm Convention could support the capacity building with  
327 their own efforts, via the Global Environment Facility, or other funding systems. For the GEF  
328 passive sampling of PFOS and PFOA could be of interest, as these are compounds for which  
329 water is a matrix of concern.

330

### 331 ■ GOING FORWARD

332 The organization of sampler deployments, and the deployments themselves are among the  
333 biggest cost for AQUA-GAPS. Hence the more samplers can be deployed at the same time, the  
334 better. Spare samplers should be archived to enable retrospective analysis. Expansion to include  
335 passive samplers designed for other contaminants (e.g., polar and nonpolar chemicals of  
336 emerging concern) would enhance the utility of the program. Samplers, and/or extracts, could be  
337 analyzed for possible temporal trends later. Spare samplers will be stored in a specimen bank  
338 operated by RECETOX. To enable quality control over time, specific samplers for QA/QC  
339 purposes will also be made available. Scientists interested in new studies, retrospective analysis

340 of extracts or samplers can request this by contacting the AQUA-GAPS co-chairs (email: aqua-  
341 gaps@passivesampling.net).

342 A welcome side-effect of AQUA-GAPS is the opportunity to increase awareness of the  
343 benefits and uncertainties of passive sampling of aqueous organic contaminants on the global  
344 scale. This might help regulatory agencies, academics, and industries still unfamiliar or hesitant  
345 to use passive sampling techniques for their own monitoring programs and other purposes. The  
346 roster of AQUA-GAPS thus also becomes a network of experts who can serve as points of  
347 contact within their regions. News and results from AQUA-GAPS will be shared via its own  
348 website (www.aqua-gaps.passivesampling.net), publications, and presentations.

349

#### 350 ■ **ACKNOWLEDGEMENTS**

351 We acknowledge support from Jinan University for workshops in Guangzhou on 21–22 January,  
352 2016 and 28-29 November, 2016 to advance AQUA-GAPS. AQUA-GAPS is supported by the  
353 National Sustainability Programme of the Czech Ministry of Education, Youth and Sports  
354 (LO1214) and the RECETOX research infrastructure (LM2011028).

355

#### 356 ■ **REFERENCES**

- 357 (1) Pozo, K.; Harner, T.; Wania, F.; Muir, D. C. G.; Jones, K. C.; Barrie, L. A. Toward a  
358 global network for persistent organic pollutants in air: Results from the global atmospheric  
359 passive sampling study. *Environ. Sci. Technol.* **2006**, *40* (16), 4867–4873.
- 360 (2) Pozo, K.; Harner, T.; Lee, S. C.; Wania, F.; Muir, D. C. G.; Jones, K. C. Seasonally  
361 resolved concentrations of persistent organic pollutants in the global atmosphere from the  
362 first year of the GAPS study. *Environ. Sci. Technol.* **2009**, *43* (3), 796–803.
- 363 (3) Lohmann, R.; Muir, D. Global aquatic passive sampling (AQUA-GAPS): Using passive  
364 samplers to monitor POPs in the waters of the world. *Environ. Sci. Technol.* **2010**, *44* (3),  
365 860–864.
- 366 (4) UNEP. *Final act of the plenipotentiaries on the Stockholm Convention on persistent*  
367 *organic pollutants.*; Geneva, Switzerland, 2001 (accessed Sep 29, 2016).
- 368 (5) UNEP. *Global monitoring plan for persistent organic pollutants as amended after the*

- 369 *fourth meeting of the Conference of the Parties to the Stockholm Convention.*  
370 *UNEP/POPS/COP.6/INF/31/Add.1.*; Geneva, Switzerland, 2013 (accessed Sep 29, 2016).
- 371 (6) UNEP. *Guidance on the global monitoring plan for persistent organic pollutants.*  
372 *UNEP/POPS/COP.6/INF/31.*; Geneva, Switzerland, 2015 (accessed Sep 29, 2016).
- 373 (7) UNEP. *Global Monitoring Plan For Persistent Organic Pollutants Under The Stockholm*  
374 *Convention Article 16 On Effectiveness Evaluation. Second Global Monitoring Report. In*  
375 *preparation.*; Geneva, Switzerland, 2016 (accessed Sep 29, 2016).
- 376 (8) Conference of Parties to the Stockholm Convention. Global Monitoring Report  
377 <http://www.pops-gmp.org/index.php?pg=gmp-data-warehouse>. (accessed Sep 29, 2016).
- 378 (9) Muir, D.; Lohmann, R. Water as a new matrix for global assessment of hydrophilic POPs.  
379 *TrAC - Trends Anal. Chem.* **2013**, *46*, 162–172.
- 380 (10) Czub, G.; McLachlan, M. S. A food chain model to predict the levels of lipophilic organic  
381 contaminants in humans. *Environ. Toxicol. Chem.* **2004**, *23* (10), 2356–2366.
- 382 (11) Arnot, J. A.; Gobas, F. A. P. C. A review of bioconcentration factor (BCF) and  
383 bioaccumulation factor (BAF) assessments for organic chemicals in aquatic organisms.  
384 *Environ. Rev.* **2006**, *14* (4), 257–297.
- 385 (12) Borgå, K.; Fisk, A. T.; Hargrave, B.; Hoekstra, P. F.; Swackhamer, D.; Muir, D. C. G.  
386 Bioaccumulation Factors for PCBs Revisited. *Environ. Sci. Technol.* **2005**, *39* (12), 4523–  
387 4532.
- 388 (13) Vrana, B.; Mills, G. A.; Allan, I. J.; Dominiak, E.; Svensson, K. .; Knutsson, J.; Morrison,  
389 G.; Greenwood, R. Passive sampling techniques for monitoring pollutants in water. *TrAC*  
390 *- Trends Anal. Chem.* **2005**, *24*, 845–868.
- 391 (14) Mayer, P.; Tolls, J.; Hermens, J. L. M.; Mackay, D. Equilibrium sampling devices: An  
392 emerging strategy for monitoring exposure to hydrophobic organic chemicals. *Environ.*  
393 *Sci. Technol.* **2003**, *37* (9), 184A– 191A.
- 394 (15) Lohmann, R.; Booij, K.; Smedes, F.; Vrana, B. Use of passive sampling devices for  
395 monitoring and compliance checking of POP concentrations in water. *Environ. Sci. Pollut.*  
396 *Res.* **2012**, *19*, 1885–1895.
- 397 (16) Hung, H.; MacLeod, M.; Guardans, R.; Scheringer, M.; Barra, R.; Harner, T.; Zhang, G.  
398 Toward the next generation of air quality monitoring: Persistent organic pollutants. *Atmos.*  
399 *Environ.* **2013**, *80*, 591–598.
- 400 (17) Bao, L.; Zeng, E. Y. Comment on “Global Aquatic Passive Sampling (AQUA-GAPS):  
401 Using Passive Samplers to Monitor POPs in the Waters of the World.” *Environ. Sci.*  
402 *Technol.* **2010**, *44* (12), 4385–4385.
- 403 (18) Lee, S. C.; Harner, T.; Pozo, K.; Shoeib, M.; Wania, F.; Muir, D. C. G.; Barrie, L. A.;  
404 Jones, K. C. Polychlorinated naphthalenes in the Global Atmospheric Passive Sampling  
405 (GAPS) study. *Environ. Sci. Technol.* **2007**, *41* (8), 2680–2687.
- 406 (19) Genualdi, S.; Lee, S. C.; Shoeib, M.; Gawor, A.; Ahrens, L.; Harner, T. Global Pilot Study  
407 of Legacy and Emerging Persistent Organic Pollutants using Sorbent-Impregnated  
408 Polyurethane Foam Disk Passive Air Samplers. *Environ. Sci. Technol.* **2010**, *44* (14),

- 409 5534–5539.
- 410 (20) Schuster, J. K.; Gioia, R.; Harner, T.; Lee, S. C.; Breivik, K.; Jones, K. C. Assessment of  
411 sorbent impregnated PUF disks (SIPs) for long-term sampling of legacy POPs. *J. Environ.*  
412 *Monit.* **2012**, *14* (1), 71–78.
- 413 (21) Koblizkova, M.; Genualdi, S.; Lee, S. C.; Harner, T. Application of Sorbent Impregnated  
414 Polyurethane Foam (SIP) Disk Passive Air Samplers for Investigating Organochlorine  
415 Pesticides and Polybrominated Diphenyl Ethers at the Global Scale. *Environ. Sci.*  
416 *Technol.* **2012**, *46* (1), 391–396.
- 417 (22) IMO. Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other  
418 Matter. [http://www.imo.org/en/About/conventions/](http://www.imo.org/en/About/conventions/listofconventions/pages/convention-on-the-prevention-of-marine-pollution-by-dumping-of-wastes-and-other-matter.aspx)  
419 [listofconventions/pages/convention-](http://www.imo.org/en/About/conventions/listofconventions/pages/convention-on-the-prevention-of-marine-pollution-by-dumping-of-wastes-and-other-matter.aspx)  
420 [on-the-prevention-of-marine-pollution-by-dumping-of-wastes-and-other-matter.aspx](http://www.imo.org/en/About/conventions/listofconventions/pages/convention-on-the-prevention-of-marine-pollution-by-dumping-of-wastes-and-other-matter.aspx).  
(accessed Sep 28, 2016).
- 421 (23) IMO. Convention on the Control of Harmful Anti-fouling Systems on Ships.  
422 [http://www.imo.org/en/About/conventions/listofconventions/pages/international-](http://www.imo.org/en/About/conventions/listofconventions/pages/international-convention-on-the-control-of-harmful-anti-fouling-systems-on-ships-(afs).aspx)  
423 [convention-on-the-control-of-harmful-anti-fouling-systems-on-ships-\(afs\).aspx](http://www.imo.org/en/About/conventions/listofconventions/pages/international-convention-on-the-control-of-harmful-anti-fouling-systems-on-ships-(afs).aspx) (accessed  
424 Sep 28, 2016)
- 425 (24) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000  
426 establishing a framework for Community action in the field of water policy.  
427 <http://data.europa.eu/eli/dir/2000/60/oj> (accessed Sep 28, 2016)
- 428 (25) Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008  
429 establishing a framework for community action in the field of marine environmental  
430 policy (Marine Strategy Framework Directive). <http://data.europa.eu/eli/dir/2008/56/oj>  
431 (accessed Sep 28, 2016)
- 432 (26) UNEP. Regional Seas Programme (<http://www.unep.org/regionalseas/about/default.asp>)  
433 (accessed Sep 28, 2016)
- 434 (27) US EPA. The Frank R. Lautenberg Chemical Safety for the 21st Century Act: First Year  
435 Implementation Plan. [https://www.epa.gov/assessing-and-managing-chemicals-under-](https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/frank-r-lautenberg-chemical-safety-21st-century-act-2)  
436 [tsca/frank-r-lautenberg-chemical-safety-21st-century-act-2](https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/frank-r-lautenberg-chemical-safety-21st-century-act-2) (accessed Sep 28, 2016)
- 437 (28) *Corrigendum to Regulation (EC) No 1907/2006 of the European Parliament and of the*  
438 *Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and*  
439 *Restriction of Chemicals (REACH), establishing a European Chemicals Agency; 2014*  
440 (accessed Sep 28, 2016)
- 441 (29) UNEP. *Proposal to list pentadecafluorooctanoic acid (CAS No: 335-67-1, PFOA,*  
442 *perfluorooctanoic acid), its salts and PFOA-related compounds in Annexes A, B and/or C*  
443 *to the Stockholm Convention on Persistent Organic Pollutants; Geneva, Switzerland, 2015*  
444 (accessed Sep 28, 2016)
- 445 (30) Weiss, J.; de Boer, J.; Berger, U.; Muir, D. C. G.; Ruan, T.; Torre, A.; Smedes, F.; Vrana,  
446 B.; Clavien, F.; Fiedler, H. *PFAS analysis in water for the Global Monitoring Plan of the*  
447 *Stockholm Convention. Set-up and guidelines for monitoring; Geneva, Switzerland, 2015.*
- 448 (31) Booij, K.; van Bommel, R.; van Aken, H. M.; van Haren, H.; Brummer, G.-J. A.;  
449 Ridderinkhof, H. Passive sampling of nonpolar contaminants at three deep-ocean sites.

- 450 *Environ. Pollut.* **2014**, *195*, 101–108.
- 451 (32) Sun, C.; Soltwedel, T.; Bauerfeind, E.; Adelman, D. A.; Lohmann, R. Depth Profiles of  
452 Persistent Organic Pollutants in the North and Tropical Atlantic Ocean. *Environ. Sci.*  
453 *Technol.* **2016**, *50* (12), 6172–6179.
- 454 (33) Gilbert, D.; Witt, G.; Smedes, F.; Mayer, P. Polymers as Reference Partitioning Phase:  
455 Polymer Calibration for an Analytically Operational Approach To Quantify Multimedia  
456 Phase Partitioning. *Anal. Chem.* **2016**, *88* (11), 5818–5826.
- 457 (34) Reichenberg, F.; Mayer, P. Two complementary sides of bioavailability: Accessibility  
458 and chemical activity of organic contaminants in sediments and soil. *Environ. Toxicol.*  
459 *Chem.* **2006**, *25* (5), 1239–1245.
- 460 (35) Joyce, A. S.; Portis, L. M.; Burgess, R. M. Evaluating the Relationship between  
461 Equilibrium Passive Sampler Uptake and Aquatic Organism Bioaccumulation. *Env. Sci*  
462 *Technol* **2016**, *50* (21), 11437–11451.
- 463 (36) Smedes, F. Monitoring of chlorinated biphenyls and polycyclic aromatic hydrocarbons by  
464 passive sampling in concert with deployed mussels. In *Passive sampling techniques in*  
465 *environmental monitoring*; Greenwood, R.; Mills, G.A.; Vrana, B., E., Ed.; Elsevier:  
466 Amsterdam, 2007; pp 407–448.
- 467 (37) Lohmann, R.; Burgess, R. M.; Cantwell, M. G.; Ryba, S. A.; MacFarlane, J. K.;  
468 Gschwend, P. M. Dependency of polychlorinated biphenyl and polycyclic aromatic  
469 hydrocarbon bioaccumulation in *Mya arenaria* on both water column and sediment bed  
470 chemical activities. *Environ. Toxicol. Chem.* **2004**, *23* (11), 2551–2562.
- 471 (38) Jahnke, A.; MacLeod, M.; Wickstrom, H.; Mayer, P. Equilibrium sampling to determine  
472 the thermodynamic potential for bioaccumulation of persistent organic pollutants from  
473 sediment. *Environ. Sci. Technol.* **2014**, *48* (19), 11352–11359.
- 474 (39) Booij, K.; Smedes, F.; Crum, S. Laboratory performance study for passive sampling of  
475 nonpolar chemicals in water. *Environ. Toxicol. Chem.* **2016**, DOI: 10.1002/etc.3657.
- 476 (40) Lohmann, R. Critical Review of Low-Density Polyethylene's Partitioning and Diffusion  
477 Coefficients for Trace Organic Contaminants and Implications for Its Use As a Passive  
478 Sampler. *Environ. Sci. Technol.* **2012**, *46* (2), 606–618.
- 479 (41) Rusina, T. P.; Smedes, F.; Klanova, J. Diffusion Coefficients of Polychlorinated  
480 Biphenyls and Polycyclic Aromatic Hydrocarbons in Polydimethylsiloxane and Low-  
481 Density Polyethylene Polymers. *J. Appl. Polym. Sci.* **2010**, *116* (3), 1803–1810.
- 482 (42) Smedes, F.; Geertsma, R. W.; Von der Zande, T.; Booij, K. Polymer - Water Partition  
483 Coefficients of Hydrophobic Compounds for Passive Sampling : Application of Cosolvent  
484 Models for Validation. *Environ. Sci. Technol.* **2009**, *43* (18), 7047–7054.
- 485 (43) Pintado-Herrera, M. G.; Lara-Martín, P. A.; González-Mazo, E.; Allan, I. J. Determination  
486 of silicone rubber and low-density polyethylene diffusion and polymer/water partition  
487 coefficients for emerging contaminants. *Environ. Toxicol. Chem.* **2016**, *35* (9), 2162–  
488 2172.
- 489 (44) Huckins, J. N.; Petty, J. D.; Lebo, J. A.; Almeida, F. V.; Booij, K.; Alvarez, D. A.; Clark,  
490 R. C.; Mogensen, B. B. Development of the Permeability/Performance Reference

- 491 Compound Approach for in situ Calibration of Semipermeable Membrane Devices.  
492 *Environ. Sci. Technol.* **2002**, *36* (1), 85–91.
- 493 (45) Booij, K.; Robinson, C. D.; Burgess, R. M.; Mayer, P.; Roberts, C. A.; Ahrens, L.; Allan,  
494 I. J.; Brant, J.; Jones, L.; Kraus, U. R.; et al. Passive Sampling in Regulatory Chemical  
495 Monitoring of Nonpolar Organic Compounds in the Aquatic Environment. *Env. Sci*  
496 *Technol* **2016**, *50* (1), 3–17.
- 497 (46) Rusina; T.P.; Smedes, F.; Klanova, J.; Booi, K.; Holoubek, I. Polymer selection for  
498 passive sampling : A comparison of critical properties. *Chemosphere* **2007**, *68* (7), 1344–  
499 1351.
- 500 (47) Carlson; J. C. ; Challis, J. K.; Hanson, M. L.; Wong, C. S. Stability of pharmaceuticals  
501 and other polar organic compounds stored on polar organic chemical integrative samplers  
502 and solid-phase extraction cartridges. *Environ. Toxicol. Chem.* **2013**, *32* (2), (334–337).
- 503 (48) Gewurtz, S. B.; Backus, S. M.; De Silva, A. O.; Ahrens, L.; Armellin, A.; Evans, M.;  
504 Fraser, S.; Gledhill, M.; Guerra, P.; Harner, T.; et al. Perfluoroalkyl acids in the Canadian  
505 environment: Multi-media assessment of current status and trends. *Environ. Int.* **2013**, *59*,  
506 183–200.
- 507 (49) Kennedy, K.; Schroeder, T.; Shaw, M.; Haynes, D.; Lewis, S.; Bentley, C.; Paxman, C.;  
508 Carter, S.; Brando, V.; Bartkow, M.; et al. Long-term monitoring of photosystem-II  
509 herbicides on the Great Barrier Reef – trends and correlation to remotely sensed water  
510 quality. *Mar. Pollut. Bull.* **2011**, *65* (4-9), 295–305.
- 511 (50) International Organization for Standardisation. *ISO 5667-23:2011(en) Water quality —*  
512 *Sampling — Part 23: Guidance on passive sampling in surface waters*; 2011 (accessed  
513 Sep 28, 2016)
- 514 (51) Huckins, J. N. ; Booij, K.; Petty. *Monitors of Organic Chemicals in the Environment*;  
515 Springer-Verlag, 2006.
- 516 (52) Smedes, F.; van der Zande, A.; Roose, P.; Davies, I. ICES passive sampling trial survey  
517 for water and sediment (PSTS) 2006-2007. Part 3: preliminary interpretation of field data.  
518 Available from <http://www.ices.dk/sites/pub/CM Documents/CM-2007/J/J0407.pdf>  
519 (accessed Oct 7, 2016).
- 520 (53) Kuhnert, P.; Liu, Y.; Henderson, B.; Dambacher, J.; Lawrence, E.; Kroon, F. *Review of the*  
521 *Marine Monitoring Program (MMP)*;2015, CSIRO Digital Productivity Flagship, CSIRO  
522 Land and Water Flagship. Report No. EP149350.
- 523 (54) Ruge, Z.; Muir, D.; Helm, P.; Lohmann, R. Concentrations, Trends, and Air – Water  
524 Exchange of PAHs and PBDEs Derived from Passive Samplers in Lake Superior in 2011.  
525 *Environ. Sci. Technol.* **2015**, *49* (23), 13777–13786.
- 526 (55) Liu, Y.; Wang, S.; Mcdonough, C. A.; Khairy, M.; Muir, D. C. G.; Helm, P. A.; Lohmann,  
527 R. Gaseous and Freely-Dissolved PCBs in the Lower Great Lakes Based on Passive  
528 Sampling: Spatial Trends and Air – Water Exchange. *Environ. Sci. Technol.* **2016**, *50*  
529 (10), 4932–4939.
- 530 (56) Muir, D. C. G.; Fisk, A.; Lehnher, I.; Lohmann, R.; Amarualik, P. Community based  
531 seawater monitoring for organic contaminants and mercury in the Canadian Arctic. In  
532 *Synopsis of research conducted under the 2014-2015, Northern Contaminants Program*,



- 533 *Aboriginal Affairs and Northern Development Canada, Ottawa.*; Aboriginal Affairs and  
534 Northern Development Canada: Ottawa, ON (Canada), 2015; pp 289–296.
- 535 (57) McDonough, C. A.; Puggioni, G.; Helm, P. A.; Muir, D.; Lohmann, R. Spatial  
536 Distribution and Air – Water Exchange of Organic Flame Retardants in the Lower Great  
537 Lakes. *Environ. Sci. Technol.* **2016**, *50* (17), 9133–9141.
- 538 (58) Morgan, E. J.; Lohmann, R. Detecting air– water and surface– deep water gradients of  
539 PCBs using polyethylene passive samplers. *Environ. Sci. Technol.* **2008**, *42* (19), 7248–  
540 7253.
- 541 (59) Morgan, E. J.; Lohmann, R. Dietary uptake from historically contaminated sediments as a  
542 source of pcbs to migratory fish and invertebrates in an urban estuary. *Environ. Sci.*  
543 *Technol.* **2010**, *44* (14), 5444–5449.
- 544 (60) Melwani, A. R.; Gregorio, D.; Jin, Y.; Stephenson, M.; Ichikawa, G.; Siegel, E.; Crane,  
545 D.; Lauenstein, G.; Davis, J. A. Mussel watch update: Long-term trends in selected  
546 contaminants from coastal California, 1977–2010. *Mar. Pollut. Bull.* **2014**, *81* (2), 291–  
547 302.
- 548 (61) Hung, H.; Kallenborn, R.; Breivik, K.; Su, Y.; Brorstrom-Lunden, E.; Olafsdottir, K.;  
549 Thorlacius, J. M.; Leppanen, S.; Bossi, R.; Skov, H.; et al. Atmospheric monitoring of  
550 organic pollutants in the Arctic under the Arctic Monitoring and Assessment Programme  
551 (AMAP): 1993-2006. *Sci. Total Environ.* **2010**, *408* (15), 2854–2873.
- 552 (62) Lohmann, R.; Klanova, J.; Kukucka, P.; Yonis, S.; Bollinger, K. PCBs and OCPs on a  
553 east-to-west transect: The importance of major currents and net volatilization for PCBs in  
554 the atlantic ocean. *Environ. Sci. Technol.* **2012**, *46* (19), 10471–10479.
- 555 (63) Allan, I. J.; Harman, C. Global Aquatic Passive Sampling : Maximizing Available  
556 Resources Using a Novel Exposure Procedure. *Env. Sci Technol* **2011**, *45* (15), 6233–  
557 6234
- 558

559 **FIGURE CAPTIONS**

560 **Figure 1.** **General approach of an AQUA-GAPS campaign for dissolved organic**  
561 **pollutants, with the proof-of-concept campaigns for a freshwater and a**  
562 **coastal water deployments.** (OMoE-CC – Ontario Ministry of the Environment  
563 and Climate Change; RECETOX – Research Center for Toxic Compounds in the  
564 Environment, Masaryk University; Jinan U – Jinan University; PCBs –  
565 polychlorinated biphenyls; OCPs – organochlorine pesticides; PAHs – polycyclic  
566 aromatic hydrocarbons; PBDEs – polybrominated diphenylethers; NFRs – novel  
567 flame retardants; PE – polyethylene; SR – silicone rubber).


568 **Figure 2.** Passive sampling holders to be deployed during AQUA-GAPS proof-of-concept  
569 studies in freshwater (left) and coastal water (right) equipped with both  
570 polyethylene and silicone rubber samplers.

571 **Figure 3.** Projected sites for freshwater and coastal water AQUA-GAPS proof-of-concept  
572 deployments.

573



## Global Aquatic Passive Sampling (AQUA-GAPS) Flow Diagram



574

575

**Figure 1. General approach of an AQUA-GAPS campaign for dissolved**

576

**organic pollutants, followed by details about the first two proof-of-concept**

577

**campaigns for a freshwater and a coastal water deployments. (OMoE-CC –**

578

Ontario Ministry of the Environment and Climate Change; RECETOX – Research

579

Center for Toxic Compounds in the Environment, Masaryk University; Jinan U –

580

Jinan University; PCBs – polychlorinated biphenyls; OCPs – organochlorine

581

pesticides; PAHs – polycyclic aromatic hydrocarbons; PBDEs – polybrominated


582

diphenylethers; NFRs – novel flame retardants; PE – polyethylene; SR – silicone

583

rubber).

584



**Freshwater Passive Sampler**



**Coastal Water Passive Sampler**

585


586

**Figure 2. Passive sampling holders to be deployed during AQUA-GAPS proof-of-concept studies in freshwater (left) and coastal water (right) equipped with both polyethylene and silicone rubber samplers.**

587

588

589



590

591

**Figure 3. Projected sites for a freshwater and coastal water AQUA-GAPS proof-of-concept deployments.** The map was created using ArcGIS 10.2.

592

593

## Global Aquatic Passive Sampling (AQUA-GAPS)

*The Network for Monitoring Organic Contaminants in the Aquatic Environment*

