

2020

Chronic Opioid Use in Women Following Hysterectomy: Patterns and Predictors

Xuerong Wen

Stephen Jon Kogut

Hilary Aroke

Lynn E. Taylor

Kristen A. Matteson

Follow this and additional works at: https://digitalcommons.uri.edu/php_facpubs

**The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.**

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Chronic Opioid Use in Women Following Hysterectomy: Patterns and Predictors

Journal:	<i>Pharmacoepidemiology and Drug Safety</i>
Manuscript ID	PDS-18-0298.R2
Wiley - Manuscript type:	Original Report
Date Submitted by the Author:	n/a
Complete List of Authors:	Wen, Xuerong; University of Rhode Island - College of Pharmacy, Health Outcomes, Department of Pharmacy Practice Kogut, Stephen ; University of Rhode Island - College of Pharmacy, Health Outcomes, Department of Pharmacy Practice Aroke, Hilary; University of Rhode Island - College of Pharmacy, Health Outcomes, Department of Pharmacy Practice Tayler, Lynn; University of Rhode Island - College of Pharmacy, Health Outcomes, Department of Pharmacy Practice Matteson, Kristen; The Warren Alpert Medical School, Brown University., Obstetrics and Gynecology
Keywords:	hysterectomy, predictors, chronic opioid use
Abstract:	<p>Background: Most women are prescribed an opioid after hysterectomy. The objective of this study was to determine the association between initial opioid prescribing characteristics and chronic opioid use after hysterectomy.</p> <p>Methods: This study included women enrolled in a commercial health plan who had a hysterectomy between July 01, 2010 and March 31, 2015. We used trajectory models to define chronic opioid use as patients with the highest probability of having an opioid prescription filled during the 6 months post-surgery. We used multivariable logistic regression to examine the association between initial opioid dispensing (amount prescribed and duration of treatment) and chronic opioid use after adjusting for potential confounders.</p> <p>Results: A total of 693 of 50,127 (1.38%) opioid naïve women met the criteria for chronic opioid use following hysterectomy. The baseline variables and initial opioid prescription characteristics predicted the pattern of long-term opioid use with moderate discrimination (c statistic = 0.70). Significant predictors of chronic opioid use included initial opioid daily dose (≥ 60 MME vs < 40 MME, aOR: 1.43, 95%CI: 1.14-1.79), and days' supply 4-7 days vs 1-3 days, aOR: 1.28, 95%CI: 1.06-1.54; ≥ 8 days vs 1-3 days, aOR: 1.41, 95%CI: 1.05-1.89). Other significant baseline predictors included older age, abdominal or laparoscopic/robotic hysterectomy, tobacco use, psychiatric medication use, back pain, and headache.</p> <p>Conclusion: Initial opioid prescribing characteristics are associated with risk of chronic opioid use after hysterectomy. Prescribing lower daily doses and shorter days' supply of opioids to women after hysterectomy</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	may result in lower risk of chronic opioid use.

SCHOLARONE™
Manuscripts

Chronic Opioid Use in Women following Hysterectomy: Patterns and Predictors

Xuerong Wen,¹ PhD, MPH, Stephen Kogut,¹ PhD, Hilary Aroke,¹ MD, PhD, MPH, Lynn Taylor¹,
MD, Kristen A. Matteson,² MD, MPH,

1. Health Outcomes Research, Department of Pharmacy Practice, College of Pharmacy, University of Rhode Island.
2. Obstetrics and Gynecology, Women & Infants Hospital and the Warren Alpert Medical School, Brown University.

For Review Only

Abstract

Background: Most women are prescribed an opioid after hysterectomy. The objective of this study was to determine the association between initial opioid prescribing characteristics and chronic opioid use after hysterectomy.

Methods: This study included women enrolled in a commercial health plan who had a hysterectomy between July 01, 2010 and March 31, 2015. We used trajectory models to define chronic opioid use as patients with the highest probability of having an opioid prescription filled during the 6 months post-surgery. We used multivariable logistic regression to examine the association between initial opioid dispensing (amount prescribed and duration of treatment) and chronic opioid use after adjusting for potential confounders.

Results: A total of 693 of 50,127 (1.38%) opioid naïve women met the criteria for chronic opioid use following hysterectomy. The baseline variables and initial opioid prescription characteristics predicted the pattern of long-term opioid use with moderate discrimination (c statistic = 0.70). Significant predictors of chronic opioid use included initial opioid daily dose (≥ 60 MME vs < 40 MME, aOR: 1.43, 95%CI: 1.14-1.79), and days' supply 4-7 days vs 1-3 days, aOR: 1.28, 95%CI: 1.06-1.54; ≥ 8 days vs 1-3 days, aOR: 1.41, 95%CI: 1.05-1.89). Other significant baseline predictors included older age, abdominal or laparoscopic/robotic hysterectomy, tobacco use, psychiatric medication use, back pain, and headache.

Conclusion: Initial opioid prescribing characteristics are associated with risk of chronic opioid use after hysterectomy. Prescribing lower daily doses and shorter days' supply of opioids to women after hysterectomy may result in lower risk of chronic opioid use.

Introduction

The rapid increase in the incidence of opioid-related overdoses and deaths has become a major public health crisis in the United States. Overdose and death from prescription opioid overdose increased 400% in women between 1999 and 2013.¹⁻³ For many women, their first exposure to prescription opioids often occurs during the post-operative period, which makes this a potential target for strategies to reduce the risk of chronic opioid use.⁴⁻⁶ Several observational studies suggest that surgery is a risk factor for chronic opioid use.⁵⁻¹⁸ Two studies have examined the relationship between the initial opioid prescribing characteristics and chronic opioid use in the postoperative setting and arrived at contradictory conclusions.^{6,12} One study reported that initial exposure to prescription opioids after minor surgery increases the risk of chronic opioid use by 44%,¹² while the other suggested that the initial opioid prescribing characteristics, including type of opioid dispensed, days' supply, or daily dose (in morphine milligram equivalents), were not associated with chronic opioid use after cesarean delivery.⁶

Hysterectomy, the most commonly performed non-obstetric surgery among women in the United States,¹⁹⁻²¹ poses a potential risk for chronic opioid use because an estimated 82% of patients receive an opioid prescription after the hysterectomy surgery.^{20,22} However, the relationship between the initial opioid prescribing characteristics and chronic opioid use after hysterectomy remains largely unknown. The objective of this study was to identify baseline risk factors for chronic opioid use after hysterectomy and examine the association between initial opioid prescribing characteristics after hysterectomy and subsequent chronic use of opioids among opioid naïve women.

Methods

Data Sources: Study data were derived from the national OptumInsight Clinformatics Data Mart™ (OptumInsight, Eden Prairie, MN). The Optum Clinformatics Data Mart is an administrative health claims database from a large national insurer which includes

1
2
3 approximately 35 million beneficiaries. The dataset contains transactional reimbursement data
4 for health care utilization including outpatient pharmacy dispensing, and inpatient and outpatient
5 medical claims.²³ The administrative enrollment file has eligibility information, the outpatient
6 pharmacy file has the national drug code (NDC) for each drug dispensed, and the medical files
7 have the Current Procedural Terminology (CPT) code for medical procedure, and International
8 Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code for medical
9 procedure and diagnosis. Patients included in the database had both medical and prescription
10 coverage. The data was used under license agreement between the University of Rhode Island
11 and OptumInsight Inc.

12
13
14
15
16
17
18
19
20
21
22 **Study Population:** This study included women (≥ 18 years) who had a hysterectomy between
23 July 01, 2010 and March 30, 2015. Women were required to have at least 6 months of
24 continuous enrollment prior to hysterectomy. To identify an opioid-naïve study cohort, we
25 excluded women with a diagnosis of opioid use disorders or any opioids filled 6 months before
26 the date of the hysterectomy (index date) (eTable 1). The hysterectomy procedures were
27 identified using ICD-9-CM procedure and CPT codes from both the inpatient and outpatient
28 claims (eTable 1).^{21,24} We excluded patients who had a radical hysterectomy and patients for
29 whom we could not rule-out a cancer diagnosis which was defined as ≥ 2 separate medical
30 claims with a cancer diagnosis occurred ≥ 42 days apart or receipt of chemotherapy, radiation,
31 or other cancer-related surgery (eTable 1).²⁵ Women who filled at least one opioid prescription
32 at a retail pharmacy within 7 days of discharge were considered exposed to prescription opioids
33 and included in the final study cohort. The 7-day window was based on the assumption that an
34 opioid prescription filled by an opioid naïve individual during this period was likely used to treat
35 acute pain after hysterectomy. This study was approved by the University of Rhode Island
36 Institutional Review Board (IRB#957873-2).

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 **Exposure to opioids:** Opioid prescriptions were identified using national drug codes (NDC)
55 from the outpatient pharmacy claims. Opioid prescriptions were classified as hydrocodone,
56
57
58
59
60

1
2
3 oxycodone and others (including codeine, fentanyl, hydromorphone, meperidine, morphine,
4 pentazocine, tapentadol, and tramadol). Duration of the initial prescription was categorized as 1-
5 3, 4-7, ≥ 8 days. We calculated the morphine milligram equivalent (MME) for the initial opioid
6 prescription using the Center for Disease Control (CDC) conversion Tables (2016 version). The
7 average daily MME dose was categorized into tertiles (≤ 40 mg/day, 40.50-58.9mg/day, and
8 ≥ 60 mg/day).
9

10
11 **Outcome Assessment:** We used trajectory models to generate our dichotomous primary study
12 outcome as chronic opioid use after hysterectomy present or absent.¹¹ The trajectory models
13 allow the use of observed longitudinal data to determine distinct opioid prescription filling
14 patterns in the study population during the six months period after hysterectomy. This approach
15 classifies patients into groups with similar opioid prescription filling patterns during follow-up
16 without relying on a prespecified cutoff value for the definition of chronic opioid use.^{26,27} In order
17 to classify the trajectory groups for opioid use during the defined follow-up window (6 months
18 after hysterectomy), we first generated 6 dichotomous variables to indicate if a study participant
19 filled a prescription of an opioid analgesic during each of 6 consecutive 30-day follow-up
20 periods, e.g., month 1 is from 7 to 37 days, month 2 is from 37 to 67 days, and month 6 is from
21 157 to 187 days.¹¹ We then modeled these 6 binary indicators of opioid use in each 30-day
22 follow-up period as a longitudinal response in a logistic group-based trajectory.^{28,29} Using the
23 trajectory model, we estimated the probability of membership of patients in each group, and the
24 probability of filling an opioid prescription over time as a smooth function of time. The model was
25 fitted using 2 to 5 opioid exposure groups and the number of groups was chosen based on the
26 value of the Bayesian Information Criterion.³⁰ In each group, a third-order polynomial (including
27 linear, squared, and cubic terms) of time were used to model the probability of filling a
28 prescription for opioids. Patients were assigned to the trajectory group in which they had the
29 highest probability of membership. Based on the model results, the trajectory group with the
30 highest probability of filling any opioid prescription beyond the initial prescription was defined as
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 “chronic opioid users”. All other trajectory groupings were classified as non-chronic opioid users.

4
5 The percentage of patients who filled an opioid prescription was reported in each trajectory for
6
7 every consecutive 30-day follow-up period.
8

9 **Potential Predictors of Chronic Opioid Use:** We collected data on potential confounding
10
11 variables that may be related to both initial opioid prescribing characteristics and chronic opioid
12
13 use after hysterectomy.^{6-11,31} Previous studies demonstrated that age, certain pain conditions,
14
15 and psychiatric disorders are associated with both postsurgical chronic pain and chronic
16
17 prescription opioid use.^{6,21,31-36} Covariates assessed in our analyses included age at
18
19 hysterectomy, Charlson comorbidity index, hospital length of stay, smoking status, use of
20
21 alcohol and illicit substances, psychiatric medication use, pre-operative pain conditions, types of
22
23 hysterectomy procedure performed, US census region, type of insurance (Medicare or private
24
25 insurance) and health plan (exclusive provider organization, health maintenance organization,
26
27 point of service, and others including indemnity and preferred provider organization), year of
28
29 surgery, and initial opioid prescribing characteristics (medication type, average daily dose, and
30
31 number of days' supplied).
32
33

34 **Statistical Analysis:** Categorical variables were examined and compared using Chi-squared or
35
36 Fisher exact tests. Continuous variables were compared using the student *t* test. Statistical tests
37
38 and modeling were conducted to identify risk factors for chronic opioid use among women after
39
40 hysterectomy. A multivariable logistic regression model was used to estimate the association
41
42 between baseline demographic and clinical variables, initial opioid prescribing characteristics,
43
44 and chronic opioid use after hysterectomy among opioid-naive patients who filled at least one
45
46 opioid prescription within 7 days after hysterectomy.
47
48

49 Sensitivity studies were conducted using multivariate logistic regression models to
50
51 examine if the significant association between initial prescribing pattern and subsequent chronic
52
53 opioid use were upheld with a different definition of chronic opioid use post hysterectomy or a
54
55 more restricted definition for opioid naivety after adjusting for other covariates.
56
57
58
59
60

1
2
3 A probability of type 1 error (α) = 0.05 was set as the threshold of statistical
4 significance. The trajectory models were conducted using “Proc Traj”. All statistical analyses
5 were performed using Statistical Analysis Software (SAS), version 9.4 (SAS Institute Inc,
6 Cary, NC).
7
8
9
10

11 12 13 14 **Results**

15
16 **Study cohort:** A total of 157,934 women underwent hysterectomy between 07/01/2010 and
17 03/31/2015. Of these patients, 22,186 (14%) patients did not meet the continuous enrollment
18 criteria, 14,696 (9%) had suspected cancer diagnoses or radical hysterectomy, 18,202 (12%)
19 didn't fill an opioid prescription within 7 days of surgery, 3,918 (2%) were missing important
20 demographic data, 48,778 (31%) used opioids in the 6 months prior to surgery and 232 (0.1%)
21 were diagnosed with OUD within 6 months prior to hysterectomy, leaving 50,127 (32%) patients
22 for the analysis. A total of 12,985 (26%) patients had an abdominal hysterectomy; 29,640
23 (59%) patients had a laparoscopic or robotic assisted laparoscopic hysterectomy; and 7,502
24 (15%) patients had a vaginal hysterectomy (Figure 1).
25
26
27
28
29
30
31
32
33
34

35 The trajectory model identified 4 distinct trajectories of prescription opioid use after
36 hysterectomy, with total 39,677 (79.15%) patients in trajectory 1, 3,304 (6.59%) in trajectory 2,
37 6,453 (12.87%) in trajectory 3, and 693 (1.38%) in trajectory 4. (Figure 2). Among the patients in
38 trajectory 4 (n=693, 1.38%), 57%, 55%, 57%, 55%, 48%, and 41% filled an opioid prescription
39 at 1 month, 2 months, 3 months, 4 months, 5 months, and 6 months following an initial
40 prescription in the 7 days post-hysterectomy, respectively. Patients in trajectory 4 were
41 classified as chronic opioid users. No patient in trajectory 1 (n=39,677, 79.15%) filled any opioid
42 prescription (0%) from month 1 to 6. Among patients in trajectory 2 (n=3,304, 6.59%), 0.3%
43 filled a prescription at month 1, which increased to 2.6% at month 2, 8% at month 3, and then
44 followed by 10% with opioid fills for the subsequent 3 months. Thirty-five percent of patients in
45 trajectory 3 (n=6,543, 12.87%) filled an opioid prescription at month 1, which decreased to 8%
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 at month 2, and then gradually decreased to 4% at month 3, and 3% at months 4, 5, and 6. A
4 description of the demographic and clinical characteristics of the patients assigned to each of
5 these 4 trajectories is included in eTable2.
6
7
8

9
10 **Characteristics of patients and initial prescriptions:** Among the total 50,127 opioid naïve
11 patients who filled at least one opioid prescription within 7 days after hysterectomy, 693 (1.38%)
12 patients were categorized as chronic opioid users and 48,434(98.62%) were categorized as
13 non-chronic users. Baseline characteristics among chronic opioid users after hysterectomy were
14 significantly different from non-chronic opioid users (Table 1). Compared to the women without
15 chronic opioid use, women with chronic opioid use were older and more likely to have a history
16 of tobacco use, and a history of alcohol use. Chronic opioid users also were also more likely to
17 be taking psychiatric medications, including benzodiazepines, antidepressants, and stimulants.
18 Back pain and headache syndromes were more prevalent in the chronic opioid use group.
19 Compared to women with commercial insurance, women with Medicare insurance had higher
20 rates of chronic opioid use. About 1.5 percent of Medicare patients who were chronic opioid
21 users after hysterectomy were younger than 65.
22
23
24
25
26
27
28
29
30
31
32
33
34

35 In unadjusted analyses, the average daily MME/day and the type of the initial opioid
36 prescription were similar between women categorized as having chronic use and not having
37 chronic opioid use. The days' supplied of the initial opioid was marginally greater among chronic
38 users than for non-chronic users (5.12 versus 4.85 days, respectively, $p = 0.02$) (Table 1).
39 Overall, oxycodone was more frequently prescribed hydrocodone or other opioids.
40
41
42
43
44

45 **Predictors of chronic opioid use among all patients in the study:** The results of
46 multivariable logistic regression analyses are shown in Table 2. A number of factors were
47 significantly associated with chronic opioid use after hysterectomy including older age,
48 abdominal or laparoscopic/robotic hysterectomy, Charlson comorbidity index, tobacco use,
49 alcohol use, and prescribed psychotropic medications. Pain conditions, including headache
50 syndromes and back pain at baseline, were also identified as significant risk factors for chronic
51
52
53
54
55
56
57
58
59
60

1
2
3 opioid use after hysterectomy. Health plan type, census region, surgical setting, fibromyalgia,
4 and other substance use disorders were not included in the final multivariable analyses due to
5 lack of significance.
6
7

8
9 The characteristics of the initial opioid prescription, including type of opioid dispensed,
10 days' supply, and daily dose in MME, were fitted in the model. Among initial opioid prescribing
11 characteristics, significant predictors of chronic opioid use were initial prescription of
12 hydrocodone (compared to oxycodone, aOR: 1.31, 95%CI: 1.10-1.57), days' supply (4-7 days
13 vs ≤ 3 days, aOR: 1.28, 95%CI: 1.06-1.54; ≥ 8 days vs ≤ 3 days, aOR: 1.41, 95%CI: 1.05-1.89),
14 and daily dose in MME (≥ 60 mg/day vs ≤ 40 mg/day, aOR: 1.43, 95%CI: 1.14-1.79). The C-
15 statistic for the fitted full model was 0.70, indicating moderate predictability (Table 2)
16
17
18
19
20
21
22
23

24 The frequencies of post-surgery complications and other conditions associated with pain
25 during the 6-month follow-up were examined in the study cohort (Table 3). Compared with non-
26 chronic opioid users, women who developed chronic opioid use after hysterectomy were more
27 frequently diagnosed with chronic pancreatitis, headache syndrome, back pain, fibromyalgia,
28 and underwent additional anesthetic procedures.
29
30
31
32
33
34
35
36

37 **Sensitivity Studies**

38
39 We redefined chronic opioid use as a total duration of opioid use (total days' supply) that
40 exceeded 90 days during 6-month follow-up period, finding that the days supplied and MME of
41 the initial opioid prescription were associated with chronic opioid use according to this definition.
42
43 An initial days supply was associated with an adjusted 1.81 fold increase in chronic use as
44 compared with a supply of less than 3 days (95%CI: 1.21-2.70); while an initial MME/day of 60
45 or greater was associated with an adjusted 1.37 fold increase in chronic use (95%CI: 1.00-1.87)
46
47 as compared with an initial MME of ≤ 39.64 mg/day). These results are similar as the results
48
49
50
51
52
53
54 which defined chronic use using trajectory models.
55
56
57
58
59
60

1
2
3 In another sensitivity analysis, we applied a more restricted definition of opioid naivety by only
4 including patients having 12-month continuous eligibility and without any opioid dispensing or
5 diagnosis of OUD during the 12 months prior to hysterectomy. In this analysis, the highest tertile
6 of MME/day (exceeding 58.92 mg) for the initial opioid prescription remained predictive of
7 subsequent chronic opioid use as compared with the lowest tertile of ≤ 39.64 mg/day (aOR:
8 1.38; 95%CI: 1.07-1.80); while the trend for days supplied also remained consistent, albeit
9 failing to reach statistical significance. (aOR: 1.38, 95%CI: 1.0-1.90 for 8+ days' supply vs ≤ 3
10 days' supply)
11
12
13
14
15
16
17
18
19
20
21

22 Discussion

23
24 Hysterectomy is the most common surgical procedure among non-pregnant women. In
25 our study, among the 50,127 opioid naïve women who had a hysterectomy and were dispensed
26 at least one opioid prescription within 7 days after hysterectomy, 1.38% became chronic opioid
27 users during the 6 months after surgery. Several demographic factors, such as age, mode of
28 hysterectomy, tobacco use, alcohol use, and psychiatric use were associated with chronic
29 opioid use after hysterectomy. More importantly, however, we found that characteristics of the
30 initial opioid prescription, including type, dose, and duration affected the probability of chronic
31 opioid use. An initial supply of 8 or more days, and an initial MME of at least 60 mg per day
32 were each associated with a greater than 40% increase in the risk of subsequent chronic use of
33 opioids (Table 2). Each year in the United States, there are approximately 430,000 inpatient
34 hysterectomy cases.³⁷ The findings of this study suggest that lessening the duration and amount
35 of opioids in the initial opioid prescription may decrease in the risk of chronic opioid use after
36 hysterectomy.
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Multiple demographic and clinical factors were associated with increased risk of chronic
52 opioid use after hysterectomy, including older age, abdominal or laparoscopic/robotic
53 hysterectomy, comorbidities, tobacco use, substance use disorders, certain pain conditions, and
54
55
56
57
58
59
60

1
2
3 use of prescription psychiatric medications use (Table 2). Our findings were similar to other
4
5 studies that showed that these factors increase the risk of chronic opioid use after major
6
7 cardiac, thoracic, abdominal and pelvic procedures, or specific surgeries, including cesarean
8
9 delivery, hysterectomy, hip or knee arthroplasty, spine, or bariatric surgeries.^{5-17, 38, 39} These
10
11 findings underscore the importance of considering the individual patient's pain management
12
13 needs, risk factors for opioid misuse at the time of opioid prescribing after hysterectomy and
14
15 other surgical procedures, and importance of medical care, monitoring and follow-up post-
16
17 operatively .
18
19

20 The literature describing the relationship between the initial opioid prescribing
21
22 characteristics and the risk of chronic opioid use after hysterectomy among opioid naïve
23
24 patients is limited. Prior observational studies have produced inconclusive results, which may be
25
26 attributed to the study sample (inclusion of patients with minor surgical procedures) or small
27
28 study sample size.^{11,38} Our results showed that initial opioid prescribing characteristics, including
29
30 opioid type, daily dose, and duration of days' supply were associated with chronic opioid use in
31
32 the 6 months following hysterectomy among opioid naïve patients. The odds of chronic opioid
33
34 use following initiation after hysterectomy were 31% higher among patients starting
35
36 hydrocodone compared to patients who began therapy with oxycodone even though prior
37
38 research has demonstrated that oxycodone is associated with risk of addiction, morbidity and
39
40 mortality.⁴⁰⁻⁴² One explanation for this finding is that for many years hydrocodone was the most
41
42 commonly prescribed opioid medication in the United States, in part because it was a schedule
43
44 III drug during the study period and was considered to be lesser risk than schedule II opioids
45
46 such as oxycodone, therefore, hydrocodone may have been prescribed more liberally,
47
48 especially to patients with a higher tendency of opioid overuse, which is a possible confounding
49
50 by contraindication. In August 21, 2014, the U.S. Drug Enforcement Administration (DEA)
51
52 issued stricter prescribing requirements and moved hydrocodone-containing medications from a
53
54 Schedule III to a Schedule II controlled substance.⁴³ Notably, the initial opioid prescribing
55
56
57
58
59
60

1
2
3 characteristics, such as longer days' supply and higher daily morphine milligram equivalent
4 dose, that were associated with post-operative chronic opioid use after hysterectomy, are also
5 risk factors for opioid misuse and opioid-related mortality.⁴⁴⁻⁴⁷ By identifying modifiable risk
6 factors for chronic opioid use after hysterectomy, such as pre-operative opioid prescribing,
7 abdominal or laparoscopic/robotic hysterectomy, and characteristics of opioid prescribing after
8 surgery, we may be able to develop strategies and interventions to decrease the likelihood of
9 chronic opioid use.
10
11
12
13
14
15
16
17

18 There is no standard definition for chronic opioid use after surgery. We selected the
19 trajectory modeling approach because of its advantages over other methods for describing
20 longitudinal trajectories and identifying patients who used opioids consistently over a longer
21 duration.^{29,30} Traditional medication adherence measures, such as the proportion of days
22 covered or the number of months of continuous medication use, may not distinguish between
23 consistent users in follow-up periods or between patients who discontinue medication
24 completely versus those who simply have a gap in use. The trajectory models repeatedly
25 assess medication use throughout the entire follow-up period and summarize long-term
26 medication adherence accounting for the time-varying nature of adherence.²⁹ We found that
27 1.38% of hysterectomy patients became chronic opioid users in the 6 months after surgery.
28 Using Optum data from January 01, 2011 to December 31, 2014, Swenson et al. reported 0.5%
29 (122 out of 24,331) of women who had a hysterectomy had new persistent opioid use following
30 hysterectomy.⁵ In Swenson's study, persistent opioid use in the 6 months post-hysterectomy
31 was defined as a minimum of 2 opioid prescriptions in the 6 months post-surgery (one between
32 15-90 days and one at least 91-180 days) and top quartile use (amount and duration).⁵ The
33 difference in incidence of chronic use/new persistent use between these studies is attributable
34 to the use of different definitions for chronic opioid use.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 Our findings suggest that prescribing lower doses for shorter days' supply may reduce the
54 risk of subsequent chronic opioid use, and is consistent with the strategies published in
55
56
57
58
59
60

1
2
3 guidelines advocated by multiple federal and state agencies, professional societies, and
4
5 advocacy groups. Although specific guidelines differ in exact wording and specific
6
7 recommendations, prescribing the lowest dose for the shortest amount of time and screening for
8
9 substance use and opioid dependence are clearly defined themes across multiple guidelines.⁴⁸⁻
10
11 ⁵⁰ However, practice change solely on the basis of guidelines can be slow, and current evidence
12
13 suggests that physicians are prescribing excess opioids to patients to control acute pain post-
14
15 surgery at hospital discharge. Studies from the surgical fields of urology, orthopedics, and
16
17 obstetrics and gynecology have shown that physicians prescribe more than twice the amount of
18
19 opioid medication patients actually consume post-discharge which adds up to millions of excess
20
21 unused opioid tablets available for diversion and abuse.⁵¹⁻⁵⁴ A recent study showed that among
22
23 obstetricians and gynecologists, only 62% reported tailoring prescriptions to the individual
24
25 patient and only 22% reported they performed an opioid dependence screen prior to
26
27 prescribing.⁵⁵ Improving adherence to best practices for opioid prescribing, including tailoring
28
29 prescriptions to the individual patient's pain management needs and risks of future opioid
30
31 misuse and abuse, could play a major role in reducing the magnitude of the opioid epidemic.
32
33
34
35
36
37
38
39

40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Limitations

First, the study subjects included in this study are commercially insured in the United States with 9% covered by Medicare Advantage (Table1). The demographic characteristics, clinical conditions, and opioid preoperative use patterns may differ from uninsured, Medicaid or Medicare covered populations. Thus, the study results might not be generalizable to patients enrolled in Medicaid or Medicare.

Our study is subject to several limitations due to the nature of observational studies using claims data. Some important medical indications, such as severity of pain, which may be related to the hysterectomy procedure or underlying conditions leading to hysterectomy, were

1
2
3 not captured in the data. Other unmeasured confounding factors included social and economic
4 factors during follow up. The low prevalence of OUDs or methadone use excluded in figure 1,
5 and other substance use disorders, tobacco use, and alcohol use presented in table 1 might be
6 due to poor sensitivity of ICD-9 diagnosis codes or CPT codes. However, the impact may be
7 non-differential if the unmeasured variables are similarly distributed in two comparison groups.
8
9
10
11
12

13
14 Opioid analgesics were assessed using pharmacy claims, which only captured opioid
15 medications legally obtained and filled at outpatient pharmacies. Patients not filling the initial
16 opioid prescription post-hysterectomy may be attributed to either not using opioids for post-
17 surgical pain or to receiving their opioid prescription from an inpatient pharmacy. The actual
18 patterns of opioid use (actual consumption) was not measured and, based on prior studies,
19 differs substantially from the amount of opioid medication prescribed. Furthermore, the result
20 that hydrocodone has a higher risk of chronic use after hysterectomy compared to oxycodone
21 could be due to a possible confounding by contraindication.
22
23
24
25
26
27
28
29

30
31 The outcome assessed in this study was chronic opioid use during the first 6 months
32 after hysterectomy. Although chronic opioid use within 6 months has been associated with
33 opioid misuse and opioid-related death, it was not possible to differentiate whether patients
34 persistently used opioids for 6 months as a treatment for pain control or because they became
35 dependent upon opioids after their post-surgical pain had abated.⁵⁶ Additionally, it was not
36 possible to determine if the women were continuing to receive their opioid for the pain related
37 to their hysterectomy or for other pain; It is possible that the reason for filling opioid
38 prescriptions in the 6 months following hysterectomy was not for personal use but for
39 diversion.⁵⁷ Our findings that chronic opioid use post hysterectomy was significantly associated
40 with initial opioid prescribing could be due to complex relationships between chronic pain post-
41 surgery, acute pain post-surgery, and chronic pain pre-surgery.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusions

In this study, approximately 3 in 200 opioid naïve women appear to become chronic opioid users after hysterectomy. Besides commonly recognized demographic and clinical risk factors, chronic opioid use after hysterectomy was associated with initial opioid prescribing characteristics, such as longer days' supply and higher daily opioid dose. Although prescription opioid medications provide effective analgesia after surgery, they must be used with caution given the potential risk for subsequent chronic opioid use that is associated with opioid misuse and overdose-related mortality. Our findings support the need to manage post-surgical pain with the least amount of opioid medication possible to effectively control a patient's symptoms, and close clinical post-op follow-up for those patients who prescribed opioids.

References

1. The U.S. Department of Health and Human Services Office on Women's Health. White Paper: Opioid Use, Misuse, and Overdose in Women.
<https://www.womenshealth.gov/files/documents/white-paper-opioid-508.pdf>. Accessed 01/18/17.
2. McCarthy M. Opioid overdose deaths rose fivefold among US women in 10 years. *BMJ* 2013;347:f4415.
3. Chou R, Ballantyne JC, Fanciullo GJ, Fine PG, Miaskowski C. Research Gaps on Use of Opioids for Chronic Noncancer Pain: Findings From a Review of the Evidence for an American Pain Society and American Academy of Pain Medicine Clinical Practice Guideline. *The Journal of Pain*, Vol 10, No 2 (February), 2009: pp 147-159
4. An analysis of the impact of opioid overprescribing in America. United States for non-dependence. The role of opioids in treating postsurgical pain. Pg 9-13.
https://www.planagainstpain.com/wp-content/uploads/2017/09/PlanAgainstPain_USND.pdf. The Plan Against Pain. September 26, 2017. Accessed December 20, 2018.
5. Swenson CW, Kamdar NS, Seiler K, Morgan DM, Lin P, As-Sanie S. Definition development and prevalence of new persistent opioid use following hysterectomy. *Am J Obstet Gynecol*. 2018 Nov;219(5):486.e1-486.e7.
6. Bateman BT, Franklin JM, Bykov K, et al. Persistent opioid use following cesarean delivery: patterns and predictors among opioid-naïve women. *Am J Obstet Gynecol*. 2016; S0002-9378(16)00478-6.
7. Hansen CA, Inacio MC, Pratt NL, Roughead EE, Graves SE. Chronic Use of Opioids Before and After Total Knee Arthroplasty: A Retrospective Cohort Study. *J Arthroplasty*. 2016 Oct 4.

- 1
2
3 8. Clarke H, Soneji N, Ko DT, Yun L, Wijeyesundera DN. Rates and risk factors for
4 prolonged opioid use after major surgery: population based cohort study. *BMJ*.
5
6 2014;348:g1251.
7
8
- 9 9. Hetmann F, Kongsgaard UE, Sandvik L, Schou-Bredal I. Prevalence and predictors
10 of persistent post-surgical pain 12 months after thoracotomy. *Acta Anaesthesiol Scand*.
11
12 2015;59(6):740-748.
13
14
- 15 10. Sun EC, Darnall BD, Baker LC, Mackey S. Incidence of and Risk Factors for chronic
16 opioid use among opioid-naive patients in the postoperative period. *JAMA Intern Med*.
17
18 2016 Sep 1;176(9):1286-93.
19
20
- 21 11. Raebel MA, Newcomer SR, Reifler LM, Boudreau D, Elliott TE, DeBar L, Ahmed A,
22 Pawloski PA, Fisher D, Donahoo WT, Bayliss EA. Chronic use of opioid medications
23 before and after bariatric surgery. *JAMA*. 2013 Oct 2;310(13):1369-76.
24
25
26
27
- 28 12. Alam A, Gomes T, Zheng H, Mamdani MM, Juurlink DN, Bell CM. Long-term
29 analgesic use after low-risk surgery: a retrospective cohort study. *Arch Intern Med*.
30
31 2012;172(5):425-430.
32
33
- 34 13. Calcaterra SL, Scarbro S, Hull ML, Forber AD, Binswanger IA, Colborn KL.
35
36 Prediction of Future Chronic Opioid Use Among Hospitalized Patients. *J Gen Intern*
37
38 *Med*. 2018 Jun;33(6):898-905. doi: 10.1007/s11606-018-4335-8. Epub 2018 Feb 5.
39
40
- 41 14. Rao AG, Chan PH, Prentice HA, Paxton EW, Navarro RA, Dillon MT, Singh A. Risk
42 factors for postoperative opioid use after elective shoulder arthroplasty. *J Shoulder*
43
44 *Elbow Surg*. 2018 Jun 8.
45
46
- 47 15. Jafari A, Shen SA, Bracken DJ, Pang J, DeConde AS. Incidence and predictive factors
48 for additional opioid prescription after endoscopic sinus surgery. *Int Forum Allergy*
49
50 *Rhinol*. 2018 May 31.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 16. Namba RS, Singh A, Paxton EW, Inacio MCS. Patient Factors Associated With
4
5 Prolonged Postoperative Opioid Use After Total Knee Arthroplasty. *J Arthroplasty*. 2018
6
7 Apr 9.
- 8
9
10 17. Pugely AJ, Bedard NA, Kalakoti P, Hendrickson NR, Shillingford JN, Laratta JL, Saifi C,
11
12 Lehman RA, Riew KD. Opioid use following cervical spine surgery: trends and factors
13
14 associated with long-term use. *Spine J*. 2018 Apr 10. pii: S1529-9430(18)30129-3.
- 15
16 18. Dunn LK, Yerra S, Fang S, Hanak MF, Leibowitz MK, Tsang S, Durieux ME, Nemergut
17
18 EC, Naik BI. Incidence and Risk Factors for Chronic Postoperative Opioid Use After
19
20 Major Spine Surgery: A Cross-Sectional Study With Longitudinal Outcome. *Anesth*
21
22 *Analg*. 2018 Jul;127(1):247-254.
- 23
24 19. Merrill RM. Hysterectomy surveillance in the United States, 1997 through 2005. *Med*
25
26 *Sci Monit*. Jan; 2008 14(1):CR24–31.
- 27
28 20. Darnall B, Li H. Hysterectomy and predictors for opioid prescription in a chronic pain
29
30 clinic sample. *Pain Med*. 2011 Feb;12(2):196-203.
- 31
32 21. Merrill RM, Layman AB, Oderda G, Asche C. Risk estimates of hysterectomy and
33
34 selected conditions commonly treated with hysterectomy. *Ann Epidemiol*. Mar; 2008
35
36 18(3):253–260.
- 37
38 22. Brandsborg B, Nikolajsen L, Hansen CT, Kehlet H, Jensen TS. Risk factors for chronic
39
40 pain after hysterectomy: a nationwide questionnaire and database study.
41
42 *Anesthesiology*. May; 2007 106(5): 1003–1012.
- 43
44 23. Optum. Retrospective database analysis. Optum Inc. 2013. Available at:
45
46 [https://www.optum.com/content/dam/optum/resources/productSheets/Retrospective-](https://www.optum.com/content/dam/optum/resources/productSheets/Retrospective-Database-Analysis.pdf)
47
48 [Database-Analysis.pdf](https://www.optum.com/content/dam/optum/resources/productSheets/Retrospective-Database-Analysis.pdf). Accessed 01/28/2017.
- 49
50 24. Keshavarz H, Hillis SD, Kieke BA. Hysterectomy surveillance—United States, 1994–
51
52 1999. *MMWR*. 2002;1(SS-5):1–8.
- 53
54
55
56
57
58
59
60

- 1
2
3 25. Chastek B, Harley C, Kallich J, Newcomer L, Paoli CJ, Teitelbaum AH. Health care
4 costs for patients with cancer at the end of life. *J Oncol Pract*. 2012 Nov;8(6):75s-80s.
5
6
7 26. Jones BL, Nagin DS. Advances in group-based trajectory modeling and an SAS
8 procedure for estimating them. *Socio Meth Res* 2007; 35(4): 542–571.
9
10
11 27. Schwarz G. Estimating the dimension of a model. *Ann Stat* 1978; 6(2): 461–464.
12
13 28. Nagin DS. Analyzing developmental trajectories: a semiparametric, group-based
14 approach. *Psychol Methods* 1999;4:139.
15
16
17 29. Franklin JM, Shrank WH, Pakes J, et al. Group-based trajectory models: a new
18 approach to classifying and predicting long-term medication adherence. *Med Care*
19 2013;51:789-96.
20
21
22 30. Jones BL, Nagin DS. Advances in group-based trajectory modeling and an SAS
23 procedure for estimating them. *Socio Meth Res* 2007; 35(4): 542–571.
24
25
26 31. Turk DC, Okifuji A. What factors affect physicians' decisions to prescribe opioids for
27 chronic noncancer pain patients? *Clin J Pain*. Dec; 1997 13(4):330–336.
28
29
30 32. Sullivan MD, Edlund MJ, Zhang L, Unutzer J, Wells KB. Association between mental
31 health disorders, problem drug use, and regular prescription opioid use. *Arch Intern*
32 *Med*. Oct 23; 2006 166(19):2087–2093.
33
34
35 33. Hooten WM, Townsend CO, Bruce BK, et al. Effects of smoking status on immediate
36 treatment outcomes of multidisciplinary pain rehabilitation. *Pain Med*. Mar; 2009
37 10(2):347–355.
38
39
40 34. Rudd RA, Aleshire N, Zibbell JE, Gladden MR, Increases in Drug and Opioid Overdose
41 Deaths – United States, 2000-2014.
42
43 <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6450a3.htm>. Accessed 01/26/18.
44
45
46 35. Brandborg B, Dueholm M, Nikolajsen L, Kehlet H, Jensen TS. A prospective study of
47 risk factors for pain persisting 4 months after hysterectomy. *Clin J Pain*. 2009
48 May;25(4):263-8.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 36. Calderon M, Castorena G, Pasic E. Postoperative pain management after hysterectomy
4 – a simple approach. [http://cdn.intechopen.com/pdfs/35360/InTech-](http://cdn.intechopen.com/pdfs/35360/InTech-Postoperative_pain_management_after_hysterectomy_a_simple_approach.pdf)
5
6 [Postoperative pain management after hysterectomy a simple approach.pdf](http://cdn.intechopen.com/pdfs/35360/InTech-Postoperative_pain_management_after_hysterectomy_a_simple_approach.pdf).
7
8 [Accessed 01/24/17](#).
9
10
11 37. Wright JD, Herzog TJ, Tsui J, et al. Nationwide trends in the performance of inpatient
12 hysterectomy in the United States. *Obstet Gynecol*. 2013;122(2 Pt 1):233-41.
13
14 38. Kim SC, Choudhry N, Franklin JM, Bykov K, Eikermann M, Lii J, Fischer MA, Bateman
15 BT. Patterns and predictors of persistent opioid use following hip or knee arthroplasty.
16 *Osteoarthritis Cartilage*. 2017 Sep;25(9):1399-1406. doi: 10.1016/j.joca.2017.04.002.
17
18 Epub 2017 Apr 19.
19
20 39. Soneji N, Clarke HA, Ko DT, Wijeyesundera DN. Risks of developing persistent opioid
21 use after major surgery. *JAMA Surg*. 2016 Nov 1;151(11):1083-1084. doi:
22
23 10.1001/jamasurg.2016.1681
24
25 40. Solomon DH, Rassen JA, Glynn RJ, et al. The comparative safety of opioids for
26 nonmalignant pain in older adults. *Arch Intern Med*. 2010;170(22):1979-1986.
27
28 41. Dhalla IA, Mamdani MM, Sivilotti ML, Kopp A, Qureshi O, Juurlink DN. Prescribing of
29 opioid analgesics and related mortality before and after the introduction of long-acting
30
31 oxycodone. *CMAJ*. 2009;181(12):891-896.
32
33 42. Madsen AM, Stark LM, Has P, Emerson JB, Schulkin J, Matteson KA. Opioid
34 Knowledge and Prescribing Practices Among Obstetrician-Gynecologists. *Obstet*
35
36 *Gynecol*. 2018 Jan;131(1):150-157.
37
38 43. Coleman JJ. Rescheduling Hydrocodone Combination Products: Addressing the Abuse
39
40 of America's Favorite Opioid. ASAM.
41
42 <https://www.asam.org/resources/publications/magazine/read/article/2015/04/10/resched>
43
44 [uling-hydrocodone-combination-products-addressing-the-abuse-of-america-s-favorite-](https://www.asam.org/resources/publications/magazine/read/article/2015/04/10/resched)
45
46 [opioid](https://www.asam.org/resources/publications/magazine/read/article/2015/04/10/resched). Accessed July 12, 2018.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 44. Bohnert AS, Valenstein M, Bair MJ, et al. Association between opioid prescribing
4 patterns and opioid overdose-related deaths. *Jama*. Apr 6 2011;305(13):1315-1321.
5
6
7 45. Jones JD, Mogali S, Comer SD. Polydrug abuse: A review of opioid and
8 benzodiazepine combination use. *Drug and alcohol dependence*. 9/1/ 2012;125(1–2):8-
9 18.
10
11
12
13 46. Paulozzi LJ, Zhang K, Jones CM, Mack KA. Risk of adverse health outcomes with
14 increasing duration and regularity of opioid therapy. *Journal of the American Board of*
15 *Family Medicine : JABFM*. May-Jun 2014;27(3):329-338.
16
17
18 47. Gwira Baumblatt JA, Wiedeman C, Dunn JR, Schaffner W, Paulozzi LJ, Jones TF.
19 High-risk use by patients prescribed opioids for pain and its role in overdose deaths.
20 *JAMA internal medicine*. May 2014;174(5):796-801.
21
22
23
24
25 48. Washington State Agency Medical Directors' Group. Interagency Guideline on
26 Prescribing Opioids for Pain 2015. Available from:
27 <http://www.agencymeddirectors.wa.gov/Files/2015AMDGOpioidGuideline.pdf>.
28
29
30
31
32 49. Alexander GC, Frattaroli S, Gielen AC, eds. The Prescription Opioid Epidemic: An
33 Evidence-Based Approach. Johns Hopkins Bloomberg School of Public Health,
34 Baltimore, Maryland: 2015. Available from: [https://www.drugabuse.gov/nidamed-](https://www.drugabuse.gov/nidamed-medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-safe-prescribing-pain)
35 [medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-](https://www.drugabuse.gov/nidamed-medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-safe-prescribing-pain)
36 [safe-prescribing-pain](https://www.drugabuse.gov/nidamed-medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-safe-prescribing-pain). Accessed: July 24, 2018.
37
38
39
40
41
42
43 50. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic
44 Pain - United States, 2016. *MMWR Recommendations and Reports: Morbidity and*
45 *Mortality Weekly Report*. 2016;65(1):1-49. PMID: 26987082.
46
47
48
49 51. Bartels K, Mayes LM, Dingmann C, Bullard KJ, Hopfer CJ, Binswanger IA. Opioid use
50 and storage patterns by patients after hospital discharge following surgery. *PloS one*.
51 2016;11(1):e0147972. PMID: 26824844; PMCID: PMC4732746.
52
53
54
55
56
57
58
59
60

- 1
2
3 52. Bates C, Laciak R, Southwick A, Bishoff J. Overprescription of postoperative narcotics:
4 a look at postoperative pain medication delivery, consumption and disposal in urological
5 practice. *J Urol*. 2011;185(2):551-5. PMID: 21168869.
6
7
8
9 53. Chapman T, Kim N, Maltenfort M, Ilyas AM. Prospective evaluation of opioid
10 consumption following carpal tunnel release surgery. *Hand*. 2017;12(1):39-42. PMID:
11 28082841; PMCID: PMC5207284.
12
13
14
15 54. Kim N, Matzon JL, Abboudi J, Jones C, Kirkpatrick W, Leinberry CF, et al. A
16 prospective evaluation of opioid utilization after upper-extremity surgical procedures:
17 identifying consumption patterns and determining prescribing guidelines. *JBJS*
18 *American*. 2016;98(20):e89. PMID: 27869630.
19
20
21
22 55. Madsen AM, Stark LM, Has P, Emerson JB, Schulkin J, Matteson KA. Opioid
23 Knowledge and Prescribing Practices Among Obstetrician-Gynecologists. *Obstet*
24 *Gynecol*. 2018 Jan;131(1):150-157. PubMed PMID:29215508.
25
26
27
28 56. Paulozzi LJ, Strickler GK, Kreiner PW, Koris CM, Centers for Disease C, Prevention.
29 Controlled Substance Prescribing Patterns--Prescription Behavior Surveillance System,
30 Eight States, 2013. *Morbidity and mortality weekly report. Surveillance summaries*. Oct
31 16 2015;64(9):1-14.
32
33
34
35
36
37
38 57. Neuman MD, Bateman BT, Wunsch H, Postoperative pain management and opioids 2
39 Inappropriate opioid prescription after surgery. *The Lancet*. Vol 393, Issue 10180, 13-19
40 April 2019, Pg 1547-1557.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Flow Chart of the Study Population.





Figure 2. Trajectories of filling opioid prescriptions in 6 months post hysterectomy. The solid or dashed lines represent the distinct estimated opioid prescription filling trajectories. The dot symbols represent the mean estimated proportion in each trajectory group dispensed an opioid prescription at each month. These indicate the mean estimated probability of filling an opioid prescription for patients in each group at each time point. Overall counts and percentages of patients in each group are presented in the underneath table. Patients in trajectory 4 were classified as opioid chronic users. Patients in trajectories 1, 2, 3 were classified as opioid non-chronic users. The x-axis indicates each 30-day interval during the 6-month follow-up period, while the y-axis indicates the estimated proportion of patients filled a prescription opioid within each 30-day interval.



Trajectories	Trajectory 1	Trajectory 2	Trajectory 3	Trajectory 4
N (%)	39,677 (79.15%)	3,304 (6.59%)	6,453 (12.87%)	693 (1.38%)

Table 1. Comparison of baseline patient characteristics among opioid-naïve patients who received prescription opioids within 7 days after hysterectomy, N=50,127

Patient characteristics	Non-chronic Users [†] (N=48,434)	Chronic Opioid Users [‡] (N= 693)	P-value
Charlson Comorbidity Index, mean (95% CI)	1.60 (1.57, 1.64)	1.74 (1.45, 2.02)	0.3479
Length of stay in days, mean (95% CI)	2.32 (2.30, 2.35)	3.79 (3.39, 4.19)	<0.0001
Age, years, mean (95% CI)	48.1 (48.0, 48.2)	49.8 (48.9,50.7)	0.0004
Age at hysterectomy, years			
<40	9,100 (18.41)	126 (18.18)	<.0001
40-49	22,538 (45.59)	255 (36.80)	
50+	17,796 (36.00)	312 (45.02)	
Substance use and abuse, N (%)			
Tobacco	2,637 (5.33)	60 (8.66)	0.0001
Alcohol	177 (0.36)	9 (1.30)	<.0001
Other substance use disorders [§]	55 (0.11)	4 (0.58)	0.0004
Psychiatric medications, N (%)			
Benzodiazepines	4,201 (8.50)	125 (18.04)	<.0001
Antidepressants	9,984 (20.20)	233 (33.62)	< .0001
Stimulants	767 (1.55)	25 (3.61)	< .0001
Pain conditions, N (%)			
Fibromyalgia	1,530 (3.10)	30 (4.33)	0.0632
Headache syndromes	4,019 (8.13)	82 (11.83)	0 .0004
Back pain	4,659 (9.42)	110 (15.87)	< .0001
Type of hysterectomy, N (%)			
Abdominal hysterectomy	12,757 (25.81)	228 (32.90)	<.0001
Vaginal hysterectomy	7,428 (15.03)	74 (10.68)	
Laparoscopic/Robotic	29,249 (59.17)	391 (56.42)	
Surgery setting, N (%)			
Inpatient	19,805 (40.06)	352 (50.79)	<.0001
Outpatient	29,629 (59.94)	341 (49.21)	
US census region, N (%)			
Midwest	13,705 (27.72)	186 (26.84)	0.6351
Northeast	5,075 (10.27)	66 (9.52)	
South	23,642 (47.83)	332 (47.91)	
West	7,012 (14.18)	109 (15.73)	
Insurance type, N (%)			
Commercial	45,107 (91.25)	586 (84.56)	< .0001
Medicare	4,327 (8.75)	107 (15.44)	
Type of health plan, N (%)			
Exclusive provider organization	5,990 (12.12)	70 (10.10)	0.0008
Health maintenance organization	4,913 (9.94)	89 (12.84)	
Point of service	34,004 (68.79)	449 (64.79)	
Others [¶]	4,527 (9.16)	85 (12.27)	
Year of hysterectomy			
2010-2013	21,949 (44.40)	287 (41.41)	0.1161
2014-2015	27,485 (55.60)	406 (58.59)	

Initial Opioid Prescription Characteristics			
Days' supply, days	4.85 (4.82, 4.87)	5.12 (4.89, 5.34)	0.0205
Average daily MME, mg/day	54.7 (54.3, 55.0)	58.4 (54.6, 62.2)	0.0571
Type of Opioid, N(%)			
Hydrocodone	17,555 (35.51)	266 (38.38)	0.0553
Oxycodone	27,434 (55.50)	354 (51.08)	
Other	4,445 (8.99)	73 (10.53)	
Tertile MME, mg/day			
≤40	16,154 (32.68)	219 (31.60)	0.2425
40.5-58.9	16,718 (33.82)	221 (31.89)	
60.0+	16,562 (33.50)	253 (36.51)	
Days' supply, days			
1-3	15,667 (31.69)	193 (27.85)	0.0566
4-7	29,098 (58.86)	423 (61.04)	
8+	4,669 (9.44)	77 (11.11)	

Abbreviation: MME=Morphine Milligram Equivalent

[†]Non-chronic users refer to patients in trajectory 1, 2, and 3 in Figure 2.

[‡]Chronic Opioid users refer to patients in trajectory 4 in Figure 2.

[§]Other substance use includes cocaine, marijuana, anxiolytic, stimulant, hallucinogenic drugs, or abuse of unspecified drugs/medications.

[¶]Others include Indemnity and Preferred provider organization.

Table 2. Baseline and initial opioid prescription characteristics that predict chronic opioid use following hysterectomy among opioid naive patients, 2010-2015; N=50, 127

Characteristics	Unadjusted OR (95% CI)	Adjusted OR [†] (95% CI)
Age group, years		
40-49	Ref.	Ref.
< 40	1.22 (0.99, 1.52)	1.20 (0.97, 1.49)
≥ 50	1.55 (1.31, 1.83)	1.37 (1.14, 1.65)
Insurance type		
Commercial	Ref.	Ref.
Medicare	1.90 (1.55, 2.35)	1.98 (1.37, 2.87)
Type of health plan, N (%)		
Exclusive provider organization	Ref.	Ref.
Health maintenance organization	1.55 (1.13, 2.13)	1.19 (0.84, 1.68)
Point of service	1.13 (0.88, 1.46)	1.10 (0.85, 1.42)
Others [‡]	1.61 (1.17, 2.21)	0.75 (0.48, 1.18)
Type of hysterectomy		
Vaginal hysterectomy	Ref.	Ref.
Laparoscopic/Robotic	1.79 (1.38, 2.34)	1.52 (1.16, 2.00)
Abdominal hysterectomy	1.34 (1.05, 1.72)	1.40 (1.09, 1.80)
Year (2010-2013 vs 2014-2015)	1.13 (0.97, 1.32)	1.24 (1.06, 1.46)
Length of stay (days)	1.14 (1.11, 1.16)	1.12 (1.10, 1.15)
Charlson comorbidity index	1.18 (1.06, 1.31)	1.16 (1.04, 1.30)
Pain conditions		
Headache (Yes vs. No)	1.52 (1.20, 1.91)	1.33 (1.04, 1.69)
Back pain (Yes vs. No)	1.81 (1.48, 2.23)	1.57 (1.27, 1.94)
Substance use		
Alcohol	3.66 (1.87, 7.18)	1.84 (0.91, 3.74)
Tobacco	1.68 (1.29, 2.20)	1.47 (1.12, 1.93)
Psychiatric medications		
Antidepressants	2.00 (1.71, 2.35)	1.71 (1.45, 2.03)
Benzodiazepines	2.37 (1.95, 2.88)	1.89 (1.54, 2.33)
Psychostimulants	2.38 (1.58, 3.56)	1.84 (1.21, 2.80)
Initial Opioid Prescription Characteristics		
Type of opioid		
Oxycodone	Ref.	Ref.
Hydrocodone	1.17 (1.00, 1.38)	1.31 (1.10, 1.57)
Other	1.27 (0.99, 1.64)	1.26 (0.96, 1.64)
Days' supply, days		
≤ 3	Ref.	Ref.
4-7	1.18 (0.99, 1.40)	1.28 (1.06, 1.54)
≥ 8	1.34 (1.03, 1.75)	1.41 (1.05, 1.89)
Tertile MME, mg/day		
< 40	Ref.	Ref.
40 -58.9	0.98 (0.81, 1.18)	1.14 (0.93, 1.40)
≥ 60.0	1.13 (0.94, 1.35)	1.43 (1.14, 1.79)

Abbreviation: MME=Morphine Milligram Equivalent

Adjusted model: c-statistic = 0.70; Hosmer-Lemeshow Goodness-of-fit Test: $\chi^2 = 19.30$, $df=8$, $p=0.0133$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

†Adjusted for baseline patient demographic and clinical characteristics listed in this table.

‡Others include Indemnity and Preferred provider organization.

For Review Only

Table 3. Distribution of possible indications for chronic opioid use during the 6-month follow-up period after hysterectomy: shown as N (%).

Possible indications	All patients in cohort (N = 50,127)	Non-chronic Users (N = 49,434)	Chronic Opioid Users (N = 693)	P-value
Chronic pancreatitis	70 (0.14)	63 (0.13)	7 (1.01)	<.0001
Headache syndromes	3,284 (6.55)	3,196 (6.47)	88 (12.70)	< .0001
Back pain	4,477 (8.93)	4,287 (8.67)	190 (27.42)	< .0001
Fibromyalgia	1,470 (2.93)	1,407 (2.85)	63 (9.09)	< .0001
Additional anesthetic procedures [†]				< .0001
0	46,132 (92.03)	45,768 (92.58)	364 (52.53)	
1	3,310 (6.60)	3,099 (6.27)	211 (30.45)	
2	525 (1.05)	459 (0.93)	66 (9.52)	
≥ 3	160 (0.32)	108 (0.22)	52 (7.50)	

[†]Number of additional procedures requiring anesthesia.

eTable 1. Operational definitions of essential baseline covariates

Patient characteristics	Operational definitions
Age	Age in years as of the date of the hysterectomy (index date). Age was categorized as <40, 40–49, and ≥50 years.
Charlson comorbidity index	The Charlson Comorbidity Index was to categorize comorbidities based on the presence of ICD-9 and ICD-10 diagnosis codes. A weighted sum provided in a single comorbidity score for a patient. We used a Quan's enhanced Charlson and Elixhauser comorbidity index.
Length of stay	Length of stay refers to the duration of hospitalization and clinical stay for the index hysterectomy procedure.
Year	The calendar year of the index date was divided into early (2010 to 2012) and late (2013 to 2015) and included in the model as a dichotomous variable.
Type of hysterectomy	Identified using ICD9 procedure codes and CPT codes
Opioid use disorder	ICD9 diagnosis codes for opioid use disorder and buprenorphine and methadone used for medication-assisted treatment.
Non-rule-out cancer diagnosis	Defined as ≥ 2 separate cancer diagnoses occurred ≥ 42 days apart or 1 cancer diagnosis with ≥ 1 procedures, including chemotherapy, radiation, or cancer-related surgery. ²⁵
Chronic opioid use	To identify chronic opioid use after hysterectomy group-based trajectory models were used to group together patients with similar patterns of medication filling during follow-up. The group with the highest filling probability was classified as chronic users.
Psychiatric medications	Psychotropic medication use during baseline identified using the American Hospital Formulary System (AHFS) Therapeutic codes and national drug codes
Pain conditions (fibromyalgia, headache syndromes, back pain)	Identified using ICD-9 diagnosis codes
Substance use and abuse (tobacco, alcohol, marijuana, cocaine, etc.)	Identified using ICD-9 diagnosis codes
US census region	US census region refers to the region where patient had the hysterectomy procedure as defined by the US Census Bureau
Insurance type	All patients in the database are insured. There were two types of insurance: commercial or Medicare.
Characteristics of initial opioid prescription	
Type of opioid	The first opioid prescriptions filled were classified as Oxycodone, Hydrocodone, or other (Codeine, Fentanyl, Hydromorphone, Meperidine, Morphine, Pentazocine, Tapentadol, and Tramadol).
Number of days' supply	Number of days' supply for the first opioid prescription after hysterectomy was divided into 3 categories (1-3 days, 4-7 days, and 8 or more days).
MME tertiles	The morphine milligram equivalent daily dose for the initial opioid prescription was determined, and the daily dose was

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	divided into tertiles for the analysis (corresponding to thresholds of ≤ 40 mg, 40.50 mg to 57.86 mg and ≥ 58.33 mg of oral morphine equivalent).
--	---

eTable 2. Comparison of baseline patient characteristics among four trajectory-based model groups, N=50,127

Patient characteristics	TBM Group 1 (N=39,677)	TBM Group 2 (N= 3304)	TBM Group 3 (N= 6453)	TBM Group 4 (N=693)
LOS in days, mean (95% CI)	2.23 (2.20, 2.25)	2.46 (2.34, 2.58)	2.77 (2.68, 2.86)	3.79 (3.39, 4.19)
CCI, mean (95% CI)	1.59 (1.54, 1.63)	1.71 (1.57, 1.85)	1.63 (1.52, 1.75)	1.74 (1.45, 2.02)
Age (years), mean (95% CI)	48.27 (48.17, 48.38)	48.30 (47.92, 48.67)	47.17 (46.92, 47.42)	49.77 (48.87, 50.67)
Age at hysterectomy (years), N (%)				
< 40	7086 (17.86)	666 (20.16)	1348 (20.89)	126 (18.18)
40-49	18119 (45.67)	1411 (42.71)	3008 (46.61)	255 (36.80)
≥ 50	14472 (36.47)	1227 (37.14)	2097 (32.50)	312 (45.02)
Substance use and abuse, N (%)				
Tobacco use	1907 (4.81)	230 (6.96)	500 (7.75)	60 (8.66)
Alcohol abuse	117 (0.29)	12 (0.36)	48 (0.74)	9 (1.30)
Other substance abuse [†]	41 (0.10)	3 (0.09)	11 (0.17)	4 (0.58)
Psychiatric medications				
Benzodiazepines	3071 (7.74)	355 (10.74)	775 (12.01)	125 (18.04)
Antidepressants	7485 (18.86)	878 (26.57)	1621 (25.12)	233 (33.62)
Psychostimulants	556 (1.40)	86 (2.60)	125 (1.94)	25 (3.61)
Pain conditions, N (%)				
Fibromyalgia	1144 (2.88)	155 (4.69)	231 (3.58)	30 (4.33)
Headache syndromes	3012 (7.59)	356 (10.77)	651 (10.09)	82 (11.83)
Back pain	3624 (9.13)	363 (10.99)	672 (10.41)	110 (15.87)
Type of hysterectomy, N (%)				
Vaginal hysterectomy	9779 (24.65)	773 (23.40)	2205 (34.17)	228 (32.90)
Laparoscopic/Robotic	6113 (15.41)	499 (15.10)	816 (12.65)	74 (10.68)
Abdominal hysterectomy	23785 (59.95)	2032 (61.50)	3432 (53.18)	391 (56.42)
US census region, N (%)				
Midwest	11101 (27.98)	900 (27.24)	1704 (26.41)	186 (26.84)
Northeast	4232 (10.67)	313 (9.47)	530 (8.21)	66 (9.52)
South	18700 (47.13)	1654 (50.06)	3288 (50.95)	332 (47.91)
West	5644 (14.22)	437 (13.23)	931 (14.43)	109 (15.73)
Insurance type, N (%)				
Commercial	36225 (91.30)	2964 (89.71)	5918 (91.71)	586 (84.56)
Medicare	3452 (8.70)	340 (10.29)	535 (8.29)	107 (15.44)
Type of health plan, N (%)				
Exclusive provider organization	4779 (12.04)	408 (12.35)	1,061 (12.54)	70 (10.10)
Health maintenance organization	3981 (10.03)	320 (9.69)	867 (10.24)	89 (12.84)
Others [‡]	3646 (9.19)	333 (10.08)	647 (7.65)	85 (12.27)
Point of service	27271 (68.73)	2243 (67.89)	5,888 (69.57)	449 (64.79)

Initial Opioid Prescription Characteristics				
Days' supply (days), mean (95% CI)	4.83 (4.81, 4.86)	4.86 (4.76, 4.95)	4.95 (4.88, 5.02)	5.12 (4.89, 5.34)
Avg. MME, mg/day (95% CI)	54.37 (53.99, 54.75)	55.93 (54.33, 57.54)	56.04 (54.96, 57.11)	58.39 (54.60, 62.18)
Type of Opioid, N (%)				
Hydrocodone	13697 (34.52)	1241 (37.56)	2617 (40.55)	266 (38.38)
Oxycodone	22431 (56.53)	1766 (53.45)	3237 (50.16)	354 (51.08)
Other	3549 (8.94)	297 (8.99)	599 (9.28)	73 (10.53)
Days' supply (days), N (%)				
1-3	12703 (32.02)	1048 (31.72)	1916 (29.69)	193 (27.85)
4-7	23273 (58.66)	1951 (59.05)	3874 (60.03)	423 (61.04)
≥ 8	3701 (9.33)	305 (9.23)	663 (10.27)	77 (11.11)
Tertile MME (mg/day), N (%)				
< 40	13034 (32.85)	1079 (32.66)	2041 (31.63)	219 (31.60)
40 – 58.93	13412 (33.80)	1102 (33.35)	2204 (34.15)	221 (31.89)
≥ 60	13231 (33.35)	1123 (33.99)	2208 (34.22)	253 (36.51)

Abbreviations: TBM=Trajectory based model; CCI= Charlson Comorbidity Index; LOS=Length of stay; MME=Morphine milligram equivalent;

†Other substance use includes marijuana, cocaine, anxiolytic, stimulant, hallucinogenic drugs, or abuse of unspecified drugs/medications.

‡Others include Preferred Provider Organization and Indemnity.

Chronic Opioid Use in Women following Hysterectomy: Patterns and Predictors

Xuerong Wen,¹ PhD, MPH, Stephen Kogut,¹ PhD, Hilary Aroke,¹ MD, PhD, MPH, Lynn Taylor¹,
MD, Kristen A. Matteson,² MD, MPH,

1. Health Outcomes Research, Department of Pharmacy Practice, College of Pharmacy, University of Rhode Island.
2. Obstetrics and Gynecology, Women & Infants Hospital and the Warren Alpert Medical School, Brown University.

For Review Only

Abstract

Background Most women are prescribed an opioid ~~at hospital discharge~~ after hysterectomy.

The objective of this study was to determine the association between initial opioid prescribing characteristics and chronic opioid use after hysterectomy.

Methods: This study included women enrolled in a commercial health plan who had a hysterectomy between July 01, 2010 and March 31, 2015. We used trajectory models to define chronic opioid use as patients with the highest probability of having an opioid prescription filled during the 6 months post-surgery. We used multivariable logistic regression to examine the association between initial opioid dispensing (amount prescribed and duration of treatment) and chronic opioid use after adjusting for potential confounders.

Results: A total of [434,693](#) of [49,844,501,127](#) ([0.9138%](#)) opioid naïve women met the criteria for chronic opioid use following hysterectomy. The baseline variables and initial opioid prescription characteristics predicted the pattern of long-term opioid use with moderate discrimination (c statistic = [0.73070](#)). Significant predictors of chronic opioid use included initial opioid daily dose (≥ 60 MME vs < 40 MME, aOR: [1.70143](#), 95%CI: [1.28114-2.26179](#)), and days' supply 4-7 days vs 1-3 days, aOR: [1.44128](#), 95%CI: [1.11106-1.79154](#); ≥ 8 days vs 1-3 days, aOR: [2.04141](#), 95%CI: [1.43105-2.90189](#)). Other significant baseline predictors included older age, [abdominal or laparoscopic/robotic hysterectomy](#)~~hysterectomy type~~, tobacco use, [co-morbidities](#)~~psychiatric medication use~~, back pain, and [fibromyalgia](#)~~headache~~.

Conclusion: Initial opioid prescribing characteristics are associated with risk of chronic opioid use after hysterectomy. Prescribing lower daily doses and shorter days' supply of opioids to women after hysterectomy may result in lower risk of chronic opioid use.

Introduction

The rapid increase in the incidence of opioid-related overdoses and deaths has become a major public health issue-crisis in the United States. Overdose and death from prescription opioid overdose increased 400% in women between 1999 and 2013.¹⁻³ For many women, their first exposure to prescription opioids often occurs during the post-operative period, which makes this a potential target for strategies to reduce the risk of chronic opioid use.⁴⁻⁶ Several observational studies suggest that surgery is a risk factor for chronic opioid use.⁵⁻¹⁸ Two studies have examined the relationship between the initial opioid prescribing characteristics and chronic opioid use in the postoperative setting and arrived at contradictory conclusions.^{6,12} One study reported that initial exposure to prescription opioids after minor surgery increases the risk of chronic opioid use by 44%,¹² while the other suggested that the initial opioid prescribing characteristics, including type of opioid dispensed, days' supply, or daily dose (in morphine milligram equivalents), were not associated with chronic opioid use after cesarean delivery.⁶

Hysterectomy, the most commonly performed non-obstetric surgery among women in the United States,¹⁹⁻²¹ poses a potential risk for chronic opioid use because an estimated 82% of hysterectomy patients receive an opioid prescription at hospital discharge after the hysterectomy surgery.^{20,22} However, the relationship between the initial opioid prescribing characteristics and chronic opioid use after hysterectomy remains largely unknown. The objective of this study was to identify baseline risk factors for chronic opioid use after hysterectomy and examine the association between initial opioid prescribing characteristics after hysterectomy and subsequent chronic use of opioids among opioid naïve women.

Methods

Data Sources: Study data were derived from the national OptumInsight Clinformatics Data Mart™ (OptumInsight, Eden Prairie, MN). The Optum Clinformatics Data Mart is an

1
2
3 administrative health claims database from a large national insurer which includes
4
5 approximately 35 million beneficiaries. The dataset contains transactional reimbursement data
6
7 for health care utilization including outpatient pharmacy dispensing, and inpatient and outpatient
8
9 medical claims.²³ The administrative enrollment file has eligibility information, the outpatient
10
11 pharmacy file has the national drug code (NDC) for each drug dispensed, and the medical files
12
13 have the Current Procedural Terminology (CPT) code for medical procedure, and International
14
15 Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code for medical
16
17 procedure and diagnosis. Patients included in the database had both medical and prescription
18
19 coverage. The data was used under license agreement between the University of Rhode Island
20
21 and OptumInsight Inc.
22
23

24 **Study Population:** This study included women (≥ 18 years) who had a hysterectomy between
25
26 July 01, 2010 and March 30, 2015. Women were required to have at least 6 months of
27
28 continuous enrollment prior to hysterectomy. To identify an opioid-naïve study cohort, we
29
30 excluded women with a diagnosis of opioid use disorders or any opioids filled 6 months before
31
32 the date of the hysterectomy (index date) (eTable 1). The hysterectomy procedures were
33
34 identified using ICD-9-CM procedure and CPT codes from both the inpatient and outpatient
35
36 claims (eTable 1).^{21,24} We excluded patients who had a radical hysterectomy and patients for
37
38 whom we could not rule-out a cancer diagnosis which was defined as ≥ 2 separate medical
39
40 claims with a cancer diagnosis occurred ≥ 42 days apart or receipt of chemotherapy, radiation,
41
42 or other cancer-related surgery (eTable 1).²⁵ Women who filled at least one opioid prescription
43
44 at a retail pharmacy within 7 days of discharge were considered exposed to prescription opioids
45
46 and included in the final study cohort. The 7-day window was based on the assumption that an
47
48 opioid prescription filled by an opioid naïve individual during this period was likely used to treat
49
50 acute pain after hysterectomy. This study was approved by the University of Rhode Island
51
52 Institutional Review Board (IRB#957873-2).
53
54
55
56
57
58
59
60

1
2
3 **Exposure to opioids:** Opioid prescriptions were identified using national drug codes (NDC)
4 from the outpatient pharmacy claims. Opioid prescriptions were classified as hydrocodone,
5 oxycodone and others (including codeine, fentanyl, hydromorphone, meperidine, morphine,
6 pentazocine, tapentadol, and tramadol). Duration of the initial prescription was categorized as 1-
7 3, 4-7, ≥ 8 days. We calculated the morphine milligram equivalent (MME) for the initial opioid
8 prescription using the Center for Disease Control (CDC) conversion Tables (2016 version). The
9 average daily MME dose was categorized into tertiles (≤ 40 mg/day, 40.50-58.9mg/day, and
10 ≥ 60 mg/day).

11
12 **Outcome Assessment:** We used trajectory models to generate our dichotomous primary study
13 outcome **as** (chronic opioid use after hysterectomy present or absent).¹¹ The trajectory models
14 allow the use of observed **longitudinal** data to determine distinct opioid prescription filling
15 patterns in the study population during the six months period after hysterectomy. This approach
16 classifies patients into groups with similar opioid prescription filling patterns during follow-up
17 without relying on a ~~priori and often subjective prespecified~~ cutoff value for the definition of
18 chronic opioid use.^{26,27} In order to classify the trajectory groups for opioid use during the defined
19 follow-up window (**6 months after hysterectomy**), we first generated 6 dichotomous variables to
20 indicate if a study participant filled a prescription of an opioid analgesic during each of 6
21 consecutive 30-day follow-up periods, **e.g., month 1 is from 7 to 37 days, month 2 is from 37 to**
22 **67 days, and month 6 is from 157 to 187 days.**¹¹ We then modeled these 6 binary indicators of
23 opioid use in each 30-day follow-up period as a longitudinal response in a logistic group-based
24 trajectory.^{28,29} Using the trajectory model, we estimated the probability of membership of patients
25 in each group, and the probability of filling **an** opioid prescription over time as a smooth function
26 of time. The model was fitted using 2 to **6-5** opioid exposure groups and the number of groups
27 was chosen based on the value of the Bayesian Information Criterion.³⁰ In each group, a third-
28 order polynomial (including linear, squared, and cubic terms) of time were used to model the
29 probability of filling a prescription for opioids. Patients were assigned to the trajectory group in
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 which they had the highest probability of membership. Based on the model results, the
4 trajectory group with the highest probability of filling any opioid prescription beyond the initial
5 prescription was defined as “chronic opioid users”. All other trajectory groupings were classified
6 as non-chronic opioid users. The percentage of patients who filled an opioid prescription was
7 reported in each trajectory for every consecutive 30-day follow-up period.

8
9
10
11
12
13 **Potential Predictors of Chronic Opioid Use:** We collected data on potential confounding
14 variables that may be related to both initial opioid prescribing characteristics and chronic opioid
15 use after hysterectomy.^{6-11,31} Previous studies demonstrated that age, certain pain conditions,
16 and psychiatric disorders are associated with both postsurgical chronic pain and chronic
17 prescription opioid use.^{6,21,31-36} Covariates assessed in our analyses included age at
18 hysterectomy, Charlson comorbidity index, hospital length of stay, smoking status, use of
19 alcohol and illicit substances, psychiatric medication use, pre-operative pain conditions, types of
20 hysterectomy procedure performed, US census region, type of insurance (Medicare or private
21 insurance) and health plan (exclusive provider organization, health maintenance organization,
22 point of service, and others including indemnity and preferred provider organization), year of
23 surgery, and initial opioid prescribing characteristics (medication type, average daily dose, and
24 number of days’ supplied).

25
26
27
28
29
30
31
32
33
34
35
36
37
38 **Statistical Analysis:** Categorical variables were examined and compared using Chi-squared or
39 Fisher exact tests. Continuous variables were compared using the student *t* test. Statistical tests
40 and modeling were conducted to identify risk factors for chronic opioid use among women after
41 hysterectomy. A multivariable logistic regression model was used to estimate the association
42 between baseline demographic and clinical variables, initial opioid prescribing characteristics,
43 and chronic opioid use after hysterectomy among opioid-naive patients who filled at least one
44 opioid prescription within 7 days after hysterectomy.

45
46
47
48
49
50
51
52
53
54 Sensitivity studies were conducted using multivariate logistic regression models to
55 examine if the significant association between initial prescribing pattern and subsequent chronic
56

opioid use were upheld with a different definition of chronic opioid use post hysterectomy or a more restricted definition for opioid naivety after adjusting for other covariates.

A probability of type 1 error (α) = 0.05 was set as the threshold of statistical significance. The trajectory models were conducted using “Proc Traj”. All statistical analyses were performed using Statistical Analysis Software (SAS), version 9.4 (SAS Institute Inc, Cary, NC).

Results

Study cohort: A total of 157,934 women underwent hysterectomy between 07/01/2010 and 03/31/2015. Of these patients, 22,186 (14%) patients did not meet the continuous enrollment criteria, 14,696 (9%) had suspected cancer diagnoses or radical hysterectomy, 23,707 (15%) didn't fill an opioid prescription within 7 days of surgery, 3,810 (2%) were missing important demographic data, 43,303 (27%) used opioids in the 6 months prior to surgery and 232 (0.1%) were diagnosed with OUDs within 6 months prior to hysterectomy, leaving 49,844 (32%) patients for the analysis. ~~In the final study cohort, only 3 patients lost eligibility and were censored during the 6-month follow-up period.~~ A total of 12,966 (26%) patients had an abdominal hysterectomy; 29,409 (59%) patients had a laparoscopic or robotic assisted laparoscopic hysterectomy; and 7,469 (15%) patients had a vaginal hysterectomy (Figure 1).

The trajectory model identified 4 distinct trajectories of prescription opioid use after hysterectomy, with total 39,677 (79.15%) patients in trajectory 1, 3,304 (6.59%) in trajectory 2, 6,453 (12.87%) in trajectory 3, and 693 (1.38%) in trajectory 4. (Figure 2). Among the patients in trajectory 4 (n=434 (1.38%), 7057%, 6055%, 6857%, 7655%, 7248%, and 5341% filled an opioid prescription at 1 month, 2 months, 3 months, 4 months, 5 months, and 6 months following an initial prescription in the 7 days post-hysterectomy, respectively. Patients in trajectory 4 were classified as chronic opioid users. No patient in trajectory 1 (n=39,677,

1
2
3 [79.15%](#) filled any opioid prescription (0%) from month 1 to 6. Among patients in trajectory 1-2
4 (n=~~2,8083,304~~, ~~5.66.59%~~), ~~500.3%~~ filled a prescription at month 1, which [increased to 2.6% at](#)
5 [month 2](#), 8% at month 3, and then followed by 10% with opioid fills for the subsequent 3
6 [months](#) gradually decreased in each subsequent month to 0% with an opioid prescription fill at
7 [month 5](#). Thirty-two five percent of patients in trajectory 2-3 (n=~~2,5436,543~~, ~~5.112.87%~~) filled an
8 opioid prescription at month 1, which decreased to ~~58%~~ at month 2, and then gradually
9 [increased-decreased to from](#) 4% at month 3, [and to 293%](#) at months [4, 5, and 6](#). Among
10 [patients seen in trajectory 3 \(n=44,059, 88.4%\), 24% filled an opioid prescription at month 1](#)
11 [followed by 0% with opioid fills for the subsequent 5 months](#). A description of the demographic
12 and clinical characteristics of the patients assigned to each of these 4 trajectories is included in
13 eTable2.

14
15
16
17
18
19
20
21
22
23
24
25
26 **Characteristics of patients and initial prescriptions:** Among the total ~~49,84450,127~~ opioid
27 naïve patients who filled at least one opioid prescription within 7 days after hysterectomy, ~~434~~
28 [693 \(0.871.38%\)](#) patients were categorized as chronic opioid users and
29 [49,41048,434\(99.1398.62%\)](#) were categorized as non-chronic users. Baseline characteristics
30 among chronic opioid users after hysterectomy were significantly different from non-chronic
31 opioid users (Table 1). Compared to the women without chronic opioid use, women with chronic
32 opioid use were older and more likely to have ~~a higher Charlson comorbidity index~~, a history of
33 tobacco use, and a history of alcohol use. Chronic opioid users also were also more likely to be
34 taking psychiatric medications, including benzodiazepines, antidepressants, and stimulants.
35 Back pain and ~~headache syndromes fibromyalgia~~ were more prevalent in the chronic opioid use
36 group. Compared to women with commercial insurance, women with Medicare insurance had
37 higher rates of chronic opioid use. About 1.5 percent of Medicare patients who were chronic
38 opioid users after hysterectomy were younger than 65.

39
40
41
42
43
44
45 [In unadjusted analyses](#), the average daily MME/day and the type of the initial opioid
46 prescription were similar between women categorized as having chronic use and not having
47
48
49
50
51
52

1
2
3 chronic opioid use. The days' supplied of the initial opioid was marginally greater among chronic
4 users than for non-chronic users (5.12 versus 4.85 days, respectively, $p = 0.0295$) (Table 1).

5
6
7 Overall, oxycodone was more frequently prescribed hydrocodone or other opioids.

8
9 **Predictors of chronic opioid use among all patients in the study:** The results of
10 multivariable logistic regression analyses are shown in Table 2. A number of factors were
11 significantly associated with chronic opioid use after hysterectomy including older age,
12 geographic region, abdominal or laparoscopic/robotic hysterectomy mode of hysterectomy,
13 Charlson comorbidity index, tobacco use, alcohol use, and prescribed psychotropic
14 medications other substance use, benzodiazepines use, antidepressants use, antipsychotics
15 use, stimulants use. Pain conditions, including headache syndromes fibromyalgia and back pain
16 at baseline, were also identified as significant risk factors for chronic opioid use after
17 hysterectomy. Health plan type, census region, surgical setting, fibromyalgia, and other
18 substance use disorders were not included in the final multivariable analyses due to lack of
19 significance.

20
21
22 The characteristics of the initial opioid prescription, including type of opioid dispensed,
23 days' supply, and daily dose in MME, were fitted in the model. Days' supply and daily dose in
24 MME were divided into their tertiles. Among initial opioid prescribing characteristics, significant
25 predictors of chronic opioid use were initial prescription of hydrocodone (compared to
26 oxycodone, aOR: 1.3331, 95%CI: 1.4410-1.5957), days' supply (4-7 days vs ≤ 3 days, aOR:
27 1.3528, 95%CI: 1.4406-1.7954; ≥ 8 days vs ≤ 3 days, aOR: 2.041.41, 95%CI: 1.4305-2.901.89),
28 and daily dose in MME (≥ 60 mg/day vs ≤ 40 mg/day, aOR: 1.7043, 95%CI: 1.2814-2.261.79),
29 The C-statistic for the fitted full model was 0.7370, indicating moderate predictability (Table 2)

30
31
32 The frequencies of post-surgery complications and other conditions associated with pain
33 during the 6-month follow-up were examined in the study cohort (Table 3). Compared with non-
34 chronic opioid users, women who developed chronic opioid use after hysterectomy were more
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

frequently diagnosed with chronic pancreatitis, headache syndrome, back pain, fibromyalgia, and underwent additional anesthetic procedures.

Sensitivity Studies

We redefined chronic opioid use as a total duration of opioid use (total days' supply) that exceeded 90 days during 6-month follow-up period, finding that the days supplied and MME of the initial opioid prescription were associated with chronic opioid use according to this definition. An initial days supply was associated with an adjusted 1.81 fold increase in chronic use as compared with a supply of less than 3 days (95%CI: 1.21-2.70); while an initial MME/day of 60 or greater was associated with an adjusted 1.37 fold increase in chronic use (95%CI: 1.00-1.87) as compared with an initial MME of ≤ 39.64 mg/day). These results are similar as the results which defined chronic use using trajectory models.

In another sensitivity analysis, we applied a more restricted definition of opioid naivety by only including patients having 12-month continuous eligibility and without any opioid dispensing or diagnosis of OUD during the 12 months prior to hysterectomy. In this analysis, the highest tertile of MME/day (exceeding 58.92 mg) for the initial opioid prescription remained predictive of subsequent chronic opioid use as compared with the lowest tertile of ≤ 39.64 mg/day (aOR: 1.38; 95%CI: 1.07-1.80); while the trend for days supplied also remained consistent, albeit failing to reach statistical significance. (aOR: 1.38, 95%CI: 1.0-1.90 for 8+ days' supply vs ≤ 3 days' supply)

Discussion

Hysterectomy is the most common surgical procedure among non-pregnant women. In our study, among the ~~49,844~~50,127 opioid naïve women who had a hysterectomy and were dispensed at least one opioid prescription within 7 days after hysterectomy, ~~0.87~~1.38% became chronic opioid users ~~during the within~~ 6 months after surgery. Several demographic factors,

1
2
3 such as age, mode of hysterectomy, tobacco use, alcohol use, and psychiatric use were
4 associated with chronic opioid use after hysterectomy. More importantly, however, we found
5 that characteristics of the initial opioid prescription, including type, dose, and duration affected
6 the probability of chronic opioid use. An initial supply of 8 or more days, and an initial MME of at
7 least 60 mg per day were each associated with a greater than 40% increase in the risk of
8 subsequent chronic use of opioids (Table 2). Each year in the United States, there are
9 approximately 430,000 inpatient hysterectomy cases.³⁷ The findings of this study suggest that
10 lessening the intensity, duration and amount of opioids in the initial focusing on these easily
11 modifiable opioid prescription characteristics may lead to decrease in the risk incidence of
12 chronic opioid use after hysterectomy, this prevalent surgical procedure. The findings of this
13 study suggest that we could focus on these easily modifiable opioid prescription characteristics
14 to decrease the incidence of chronic opioid use after this prevalent surgical procedure.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29 Multiple demographic and clinical factors were associated with increased risk of chronic
30 opioid use after hysterectomy, including older age, abdominal or laparoscopic/robotic
31 hysterectomy mode of hysterectomy, comorbidities, tobacco use, substance abuse use
32 disorders, certain pain conditions, and use of prescription any psychiatric medications use
33 (Table 2). Our findings were similar to other studies that showed that these factors increase the
34 risk of chronic opioid use after major cardiac, thoracic, abdominal and pelvic procedures, or
35 specific surgeries, including cesarean delivery, hysterectomy, hip or knee arthroplasty, spine, or
36 bariatric surgeries.^{5-17, 38, 39} Previous studies have reported that residing in the South or Midwest
37 regions of the U.S. was associated with prescription opioid abuse specifically among women. In
38 our study, the risk of chronic opioid use after hysterectomy was also higher in the South,
39 Midwest, and West compared with the Northeast. (Table 2) These findings underscore the
40 importance of considering the individual patient's pain management needs, and risk factors for
41 opioid misuse at the time of opioid prescribing after hysterectomy and other surgical
42 procedures, and importance of medical care, monitoring and follow-up post-operatively.

1
2
3 The literature describing the relationship between the initial opioid prescribing
4 characteristics and the risk of chronic opioid use after hysterectomy among opioid naïve
5 patients is limited. Prior observational studies have produced inconclusive results, which may be
6 attributed to the study sample (inclusion of patients with minor surgical procedures) or small
7 study sample size.^{11,38} Our results showed that initial opioid prescribing characteristics, including
8 opioid type, daily dose, and duration of days' supply were associated with chronic opioid use in
9 the 6 months following hysterectomy among opioid naïve patients. The odds of chronic opioid
10 use following initiation after hysterectomy were [3531%](#) higher among patients starting
11 hydrocodone compared to patients who began therapy with oxycodone even though prior
12 research has demonstrated that oxycodone is associated with risk of addiction, morbidity and
13 mortality.⁴⁰⁻⁴² One explanation [for this finding](#) is that for many years hydrocodone was the most
14 commonly prescribed opioid medication in the United States, in part because it was a schedule
15 III drug [during the study period](#) and [was considered to be lesser risk than schedule II opioids](#)
16 [such as oxycodone, therefore, hydrocodone may have been prescribed more liberally,](#)
17 [especially to patients with a higher tendency of opioid overuse,](#) which is a possible confounding
18 by contraindication. In August 21, 2014, the U.S. Drug Enforcement Administration (DEA)
19 issued stricter prescribing requirements and moved hydrocodone-containing medications from a
20 Schedule III to a Schedule II controlled substance.⁴³ Notably, the initial opioid prescribing
21 characteristics, such as longer days' supply and higher daily morphine milligram equivalent
22 dose, that were associated with post-operative chronic opioid use after hysterectomy, are also
23 risk factors for opioid misuse and opioid-related mortality.⁴⁴⁻⁴⁷ By identifying modifiable risk
24 factors for chronic opioid use after hysterectomy, such as pre-operative opioid prescribing,
25 [abdominal or laparoscopic/robotic hysterectomy mode of hysterectomy,](#) and characteristics of
26 opioid prescribing after surgery, we may be able to develop strategies and interventions to
27 decrease the likelihood of chronic opioid use.

1
2
3 There is no standard definition for chronic opioid use after surgery. We selected the
4 trajectory modeling approach because of its advantages over other methods for describing
5 longitudinal trajectories and identifying patients who used opioids consistently over a longer
6 duration.^{29,30} Traditional medication adherence measures, such as the proportion of days
7 covered or the number of months of continuous medication use, may not distinguish between
8 consistent users in follow-up periods or between patients who discontinue medication
9 completely versus those who simply have a gap in use. The trajectory models repeatedly
10 assess medication use throughout the entire follow-up period and summarize long-term
11 medication adherence accounting for the time-varying nature of adherence.²⁹ We found that
12 [0.91.38%](#) of hysterectomy patients became chronic opioid users in the 6 months after surgery.
13 Using Optum data from January 01, 2011 to December 31, 2014, Swenson et al. reported 0.5%
14 (122 out of 24,331) of women who had a hysterectomy had new persistent opioid use following
15 hysterectomy.⁵ In Swenson's study, persistent opioid use in the 6 months post-hysterectomy
16 was defined as a minimum of 2 opioid prescriptions in the 6 months post-surgery (one between
17 15-90 days and one at least 91-180 days) and top quartile use (amount and duration).⁵ The
18 difference in incidence of chronic use/new persistent use between these studies is attributable
19 to the use of different definitions for chronic opioid use.

20
21
22 Our findings suggest that prescribing lower doses for shorter days' supply may [reduce the](#)
23 [risk of subsequent chronic opioid use, be a better strategy for post-surgical pain management,](#)
24 [and which](#) is consistent with the strategies published in guidelines advocated by multiple federal
25 and state agencies, professional societies, and advocacy groups. Although specific guidelines
26 differ in exact wording and specific recommendations, prescribing the lowest dose for the
27 shortest amount of time and screening for substance use and opioid dependence are clearly
28 defined themes across multiple guidelines.⁴⁸⁻⁵⁰ However, practice change solely on the basis of
29 guidelines can be slow, and current evidence suggests that physicians are prescribing excess
30 opioids to patients to control acute pain post-surgery at hospital discharge. Studies from the

1
2
3 surgical fields of urology, orthopedics, and obstetrics and gynecology have shown that
4 physicians prescribe more than twice the amount of opioid medication patients actually
5 consume post-discharge which adds up to millions of excess unused opioid tablets available for
6 diversion and abuse.⁵¹⁻⁵⁴ A recent study showed that among obstetricians and gynecologists,
7 only 62% reported tailoring prescriptions to the individual patient and only 22% reported they
8 performed an opioid dependence screen prior to prescribing.⁵⁵ Improving adherence to best
9 practices for opioid prescribing, including tailoring prescriptions to the individual patient's pain
10 management needs and risks of future opioid misuse and abuse, could play a major role in
11 reducing the magnitude of the opioid epidemic.
12
13
14
15
16
17
18
19
20
21
22
23
24

25 **Limitations**

26
27 First, the study subjects included in this study are commercially insured in the United
28 States with 409% covered by Medicare Advantage (Table1). The demographic characteristics,
29 clinical conditions, and opioid preoperative use patterns may differ from uninsured, Medicaid or
30 Medicare covered populations. Thus, the study results might not be generalizable to patients
31 enrolled in Medicaid or Medicare. with government covered insurances.
32
33
34
35
36
37

38 Our study is subject to several limitations due to the nature of observational studies
39 using claims data. Some important medical indications, such as severity of pain, which may be
40 related to the hysterectomy procedure or underlying conditions leading to hysterectomy, were
41 not captured in the data. Other unmeasured confounding factors included social and economic
42 factors during follow up. The low prevalence of OUDs or methadone use excluded in figure 1,
43 and other substance abuse disorders, tobacco use, and alcohol abuse presented in table 1
44 might be due to poor sensitivity of ICD-9 diagnosis codes or CPT codes. However, the impact
45 may be non-differential if the unmeasured variables are similarly distributed in two comparison
46 groups.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Opioid analgesics were assessed using pharmacy claims, which only captured opioid
4 medications legally obtained and filled at outpatient pharmacies. ~~The 69% prevalence of~~
5 ~~Patients not~~ filling the initial opioid prescription post-hysterectomy may be attributed to either
6 ~~some patients~~ not using opioids for post-surgical pain or to ~~patients~~ receiving their opioid
7 prescription from an inpatient pharmacy. The actual patterns of opioid use (actual consumption)
8 was not measured and, based on prior studies, differs substantially from the amount of opioid
9 medication prescribed. Furthermore, the result that hydrocodone has a higher risk of chronic
10 use after hysterectomy compared to oxycodone could be due to a possible confounding by
11 contraindication.
12
13
14
15
16
17
18
19
20
21

22 The outcome assessed in this study was chronic opioid use during the first 6 months
23 after hysterectomy. Although chronic opioid use within 6 months has been associated with
24 opioid misuse and opioid-related death, it was not possible to differentiate whether patients
25 persistently used opioids for 6 months as a ~~legitimate~~ treatment for pain control or because they
26 became dependent upon opioids after their ~~post-surgical pain had abated.~~ ~~use of the~~
27 ~~medications to treat acute pain.~~⁵⁶ Additionally, it ~~was not possible is impossible to determine~~
28 ~~know if the women were continuing to receive their opioid for the pain related to their~~
29 ~~hysterectomy or for other pain; the reason for the opioid prescription fills during this time~~
30 ~~periods;~~ It is possible that the reason for filling opioid prescriptions in the 6 months following
31 hysterectomy was not for personal use but for diversion.⁵⁷ Our findings that chronic opioid use
32 post hysterectomy was significantly associated with ~~preoperative opioid use and~~ initial opioid
33 prescribing could be due to complex relationships between chronic pain post-surgery, acute
34 pain post-surgery, and chronic pain pre-surgery.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 Conclusions

53
54
55
56
57
58
59
60

1
2
3 [In this study, About 0.87% of approximately 3 in 200](#) opioid naïve women appear to
4
5 become chronic opioid users after hysterectomy. Besides commonly recognized demographic
6
7 and clinical risk factors, chronic opioid use after hysterectomy was associated with initial opioid
8
9 prescribing characteristics, such as longer days' supply and higher daily opioid dose. Although
10
11 prescription opioid medications provide effective analgesia after surgery, they must be used with
12
13 caution given the potential risk for subsequent chronic opioid use that is associated with opioid
14
15 misuse and overdose-related mortality. Our findings support the need to manage post-surgical
16
17 pain with the least amount of opioid medication possible to effectively control a patient's
18
19 symptoms, [and close clinical post-op follow-up for those patients who prescribed opioids.](#)
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. The U.S. Department of Health and Human Services Office on Women's Health. White Paper: Opioid Use, Misuse, and Overdose in Women.
<https://www.womenshealth.gov/files/documents/white-paper-opioid-508.pdf>. Accessed 01/18/17.
2. McCarthy M. Opioid overdose deaths rose fivefold among US women in 10 years. *BMJ* 2013;347:f4415.
3. Chou R, Ballantyne JC, Fanciullo GJ, Fine PG, Miaskowski C. Research Gaps on Use of Opioids for Chronic Noncancer Pain: Findings From a Review of the Evidence for an American Pain Society and American Academy of Pain Medicine Clinical Practice Guideline. *The Journal of Pain*, Vol 10, No 2 (February), 2009: pp 147-159
4. An analysis of the impact of opioid overprescribing in America. United States for non-dependence. The role of opioids in treating postsurgical pain. Pg 9-13.
https://www.planagainstpain.com/wp-content/uploads/2017/09/PlanAgainstPain_USND.pdf. The Plan Against Pain. September 26, 2017. Accessed December 20, 2018.
5. Swenson CW, Kamdar NS, Seiler K, Morgan DM, Lin P, As-Sanie S. Definition development and prevalence of new persistent opioid use following hysterectomy. *Am J Obstet Gynecol*. 2018 Nov;219(5):486.e1-486.e7.
6. Bateman BT, Franklin JM, Bykov K, et al. Persistent opioid use following cesarean delivery: patterns and predictors among opioid-naïve women. *Am J Obstet Gynecol*. 2016; S0002-9378(16)00478-6.
7. Hansen CA, Inacio MC, Pratt NL, Roughead EE, Graves SE. Chronic Use of Opioids Before and After Total Knee Arthroplasty: A Retrospective Cohort Study. *J Arthroplasty*. 2016 Oct 4.

- 1
2
3 8. Clarke H, Soneji N, Ko DT, Yun L, Wijeyesundera DN. Rates and risk factors for
4 prolonged opioid use after major surgery: population based cohort study. *BMJ*.
5 2014;348:g1251.
6
7
- 8
9 9. Hetmann F, Kongsgaard UE, Sandvik L, Schou-Bredal I. Prevalence and predictors
10 of persistent post-surgical pain 12 months after thoracotomy. *Acta Anaesthesiol Scand*.
11 2015;59(6):740-748.
12
13
- 14
15 10. Sun EC, Darnall BD, Baker LC, Mackey S. Incidence of and Risk Factors for chronic
16 opioid use among opioid-naive patients in the postoperative period. *JAMA Intern Med*.
17 2016 Sep 1;176(9):1286-93.
18
19
- 20
21 11. Raebel MA, Newcomer SR, Reifler LM, Boudreau D, Elliott TE, DeBar L, Ahmed A,
22 Pawloski PA, Fisher D, Donahoo WT, Bayliss EA. Chronic use of opioid medications
23 before and after bariatric surgery. *JAMA*. 2013 Oct 2;310(13):1369-76.
24
25
- 26
27 12. Alam A, Gomes T, Zheng H, Mamdani MM, Juurlink DN, Bell CM. Long-term
28 analgesic use after low-risk surgery: a retrospective cohort study. *Arch Intern Med*.
29 2012;172(5):425-430.
30
31
- 32
33 13. Calcaterra SL, Scarbro S, Hull ML, Forber AD, Binswanger IA, Colborn KL.
34
35 Prediction of Future Chronic Opioid Use Among Hospitalized Patients. *J Gen Intern*
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
14. Rao AG, Chan PH, Prentice HA, Paxton EW, Navarro RA, Dillon MT, Singh A. Risk
factors for postoperative opioid use after elective shoulder arthroplasty. *J Shoulder
Elbow Surg*. 2018 Jun 8.
15. Jafari A, Shen SA, Bracken DJ, Pang J, DeConde AS. Incidence and predictive factors
for additional opioid prescription after endoscopic sinus surgery. *Int Forum Allergy
Rhinol*. 2018 May 31.

- 1
2
3 16. Namba RS, Singh A, Paxton EW, Inacio MCS. Patient Factors Associated With
4
5 Prolonged Postoperative Opioid Use After Total Knee Arthroplasty. *J Arthroplasty*. 2018
6
7 Apr 9.
- 8
9
10 17. Pugely AJ, Bedard NA, Kalakoti P, Hendrickson NR, Shillingford JN, Laratta JL, Saifi C,
11
12 Lehman RA, Riew KD. Opioid use following cervical spine surgery: trends and factors
13
14 associated with long-term use. *Spine J*. 2018 Apr 10. pii: S1529-9430(18)30129-3.
- 15
16 18. Dunn LK, Yerra S, Fang S, Hanak MF, Leibowitz MK, Tsang S, Durieux ME, Nemergut
17
18 EC, Naik BI. Incidence and Risk Factors for Chronic Postoperative Opioid Use After
19
20 Major Spine Surgery: A Cross-Sectional Study With Longitudinal Outcome. *Anesth*
21
22 *Analg*. 2018 Jul;127(1):247-254.
- 23
24 19. Merrill RM. Hysterectomy surveillance in the United States, 1997 through 2005. *Med*
25
26 *Sci Monit*. Jan; 2008 14(1):CR24–31.
- 27
28 20. Darnall B, Li H. Hysterectomy and predictors for opioid prescription in a chronic pain
29
30 clinic sample. *Pain Med*. 2011 Feb;12(2):196-203.
- 31
32 21. Merrill RM, Layman AB, Oderda G, Asche C. Risk estimates of hysterectomy and
33
34 selected conditions commonly treated with hysterectomy. *Ann Epidemiol*. Mar; 2008
35
36 18(3):253–260.
- 37
38 22. Brandsborg B, Nikolajsen L, Hansen CT, Kehlet H, Jensen TS. Risk factors for chronic
39
40 pain after hysterectomy: a nationwide questionnaire and database study.
41
42 *Anesthesiology*. May; 2007 106(5): 1003–1012.
- 43
44 23. Optum. Retrospective database analysis. Optum Inc. 2013. Available at:
45
46 [https://www.optum.com/content/dam/optum/resources/productSheets/Retrospective-](https://www.optum.com/content/dam/optum/resources/productSheets/Retrospective-Database-Analysis.pdf)
47
48 [Database-Analysis.pdf](https://www.optum.com/content/dam/optum/resources/productSheets/Retrospective-Database-Analysis.pdf). Accessed 01/28/2017.
- 49
50
51 24. Keshavarz H, Hillis SD, Kieke BA. Hysterectomy surveillance—United States, 1994–
52
53 1999. *MMWR*. 2002;1(SS-5):1–8.
- 54
55
56
57
58
59
60

- 1
2
3 25. Chastek B, Harley C, Kallich J, Newcomer L, Paoli CJ, Teitelbaum AH. Health care
4 costs for patients with cancer at the end of life. *J Oncol Pract*. 2012 Nov;8(6):75s-80s.
5
- 6
7 26. Jones BL, Nagin DS. Advances in group-based trajectory modeling and an SAS
8 procedure for estimating them. *Socio Meth Res* 2007; 35(4): 542–571.
9
- 10
11 27. Schwarz G. Estimating the dimension of a model. *Ann Stat* 1978; 6(2): 461–464.
12
- 13
14 28. Nagin DS. Analyzing developmental trajectories: a semiparametric, group-based
15 approach. *Psychol Methods* 1999;4:139.
16
- 17
18 29. Franklin JM, Shrank WH, Pakes J, et al. Group-based trajectory models: a new
19 approach to classifying and predicting long-term medication adherence. *Med Care*
20 2013;51:789-96.
21
- 22
23 30. Jones BL, Nagin DS. Advances in group-based trajectory modeling and an SAS
24 procedure for estimating them. *Socio Meth Res* 2007; 35(4): 542–571.
25
- 26
27 31. Turk DC, Okifuji A. What factors affect physicians' decisions to prescribe opioids for
28 chronic noncancer pain patients? *Clin J Pain*. Dec; 1997 13(4):330–336.
29
- 30
31 32. Sullivan MD, Edlund MJ, Zhang L, Unutzer J, Wells KB. Association between mental
32 health disorders, problem drug use, and regular prescription opioid use. *Arch Intern*
33 *Med*. Oct 23; 2006 166(19):2087–2093.
34
- 35
36 33. Hooten WM, Townsend CO, Bruce BK, et al. Effects of smoking status on immediate
37 treatment outcomes of multidisciplinary pain rehabilitation. *Pain Med*. Mar; 2009
38 10(2):347–355.
39
- 40
41 34. Rudd RA, Aleshire N, Zibbell JE, Gladden MR, Increases in Drug and Opioid Overdose
42 Deaths – United States, 2000-2014.
43
44 <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6450a3.htm>. Accessed 01/26/18.
45
- 46
47 35. Brandborg B, Dueholm M, Nikolajsen L, Kehlet H, Jensen TS. A prospective study of
48 risk factors for pain persisting 4 months after hysterectomy. *Clin J Pain*. 2009
49 May;25(4):263-8.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 36. Calderon M, Castorena G, Pasic E. Postoperative pain management after hysterectomy
4 – a simple approach. [http://cdn.intechopen.com/pdfs/35360/InTech-](http://cdn.intechopen.com/pdfs/35360/InTech-Postoperative_pain_management_after_hysterectomy_a_simple_approach.pdf)
5 [Postoperative pain management after hysterectomy a simple approach.pdf](http://cdn.intechopen.com/pdfs/35360/InTech-Postoperative_pain_management_after_hysterectomy_a_simple_approach.pdf).
6 [Accessed 01/24/17](http://cdn.intechopen.com/pdfs/35360/InTech-Postoperative_pain_management_after_hysterectomy_a_simple_approach.pdf).
7
8
9
10
11 37. Wright JD, Herzog TJ, Tsui J, et al. Nationwide trends in the performance of inpatient
12 hysterectomy in the United States. *Obstet Gynecol*. 2013;122(2 Pt 1):233-41.
13
14 38. Kim SC, Choudhry N, Franklin JM, Bykov K, Eikermann M, Lii J, Fischer MA, Bateman
15 BT. Patterns and predictors of persistent opioid use following hip or knee arthroplasty.
16 *Osteoarthritis Cartilage*. 2017 Sep;25(9):1399-1406. doi: 10.1016/j.joca.2017.04.002.
17
18 Epub 2017 Apr 19.
19
20 39. Soneji N, Clarke HA, Ko DT, Wijeyesundera DN. Risks of developing persistent opioid
21 use after major surgery. *JAMA Surg*. 2016 Nov 1;151(11):1083-1084. doi:
22
23 10.1001/jamasurg.2016.1681
24
25 40. Solomon DH, Rassen JA, Glynn RJ, et al. The comparative safety of opioids for
26 nonmalignant pain in older adults. *Arch Intern Med*. 2010;170(22):1979-1986.
27
28 41. Dhalla IA, Mamdani MM, Sivilotti ML, Kopp A, Qureshi O, Juurlink DN. Prescribing of
29 opioid analgesics and related mortality before and after the introduction of long-acting
30
31 oxycodone. *CMAJ*. 2009;181(12):891-896.
32
33 42. Madsen AM, Stark LM, Has P, Emerson JB, Schulkin J, Matteson KA. Opioid
34
35 Knowledge and Prescribing Practices Among Obstetrician-Gynecologists. *Obstet*
36
37 *Gynecol*. 2018 Jan;131(1):150-157.
38
39 43. Coleman JJ. Rescheduling Hydrocodone Combination Products: Addressing the Abuse
40
41 of America's Favorite Opioid. ASAM.
42
43 <https://www.asam.org/resources/publications/magazine/read/article/2015/04/10/resched>
44
45 [uling-hydrocodone-combination-products-addressing-the-abuse-of-america-s-favorite-](https://www.asam.org/resources/publications/magazine/read/article/2015/04/10/resched)
46
47 [opioid](https://www.asam.org/resources/publications/magazine/read/article/2015/04/10/resched). Accessed July 12, 2018.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 44. Bohnert AS, Valenstein M, Bair MJ, et al. Association between opioid prescribing
4 patterns and opioid overdose-related deaths. *Jama*. Apr 6 2011;305(13):1315-1321.
5
6
7 45. Jones JD, Mogali S, Comer SD. Polydrug abuse: A review of opioid and
8 benzodiazepine combination use. *Drug and alcohol dependence*. 9/1/ 2012;125(1–2):8-
9 18.
10
11
12 46. Paulozzi LJ, Zhang K, Jones CM, Mack KA. Risk of adverse health outcomes with
13 increasing duration and regularity of opioid therapy. *Journal of the American Board of*
14 *Family Medicine : JABFM*. May-Jun 2014;27(3):329-338.
15
16
17 47. Gwira Baumblatt JA, Wiedeman C, Dunn JR, Schaffner W, Paulozzi LJ, Jones TF.
18 High-risk use by patients prescribed opioids for pain and its role in overdose deaths.
19 *JAMA internal medicine*. May 2014;174(5):796-801.
20
21
22 48. Washington State Agency Medical Directors' Group. Interagency Guideline on
23 Prescribing Opioids for Pain 2015. Available from:
24 <http://www.agencymeddirectors.wa.gov/Files/2015AMDGOpioidGuideline.pdf>.
25
26
27 49. Alexander GC, Frattaroli S, Gielen AC, eds. The Prescription Opioid Epidemic: An
28 Evidence-Based Approach. Johns Hopkins Bloomberg School of Public Health,
29 Baltimore, Maryland: 2015. Available from: [https://www.drugabuse.gov/nidamed-](https://www.drugabuse.gov/nidamed-medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-safe-prescribing-pain)
30 [medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-](https://www.drugabuse.gov/nidamed-medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-safe-prescribing-pain)
31 [safe-prescribing-pain](https://www.drugabuse.gov/nidamed-medical-health-professionals/opioid-pain-management-cmes/unaccredited-module-1-safe-prescribing-pain). Accessed: July 24, 2018.
32
33
34 50. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic
35 Pain - United States, 2016. *MMWR Recommendations and Reports: Morbidity and*
36 *Mortality Weekly Report*. 2016;65(1):1-49. PMID: 26987082.
37
38
39 51. Bartels K, Mayes LM, Dingmann C, Bullard KJ, Hopfer CJ, Binswanger IA. Opioid use
40 and storage patterns by patients after hospital discharge following surgery. *PloS one*.
41 2016;11(1):e0147972. PMID: 26824844; PMCID: PMC4732746.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 52. Bates C, Laciak R, Southwick A, Bishoff J. Overprescription of postoperative narcotics:
4 a look at postoperative pain medication delivery, consumption and disposal in urological
5 practice. J Urol. 2011;185(2):551-5. PMID: 21168869.
6
7
8
9 53. Chapman T, Kim N, Maltenfort M, Ilyas AM. Prospective evaluation of opioid
10 consumption following carpal tunnel release surgery. Hand. 2017;12(1):39-42. PMID:
11 28082841; PMCID: PMC5207284.
12
13
14 54. Kim N, Matzon JL, Abboudi J, Jones C, Kirkpatrick W, Leinberry CF, et al. A
15 prospective evaluation of opioid utilization after upper-extremity surgical procedures:
16 identifying consumption patterns and determining prescribing guidelines. JBJS
17 American. 2016;98(20):e89. PMID: 27869630.
18
19
20 55. Madsen AM, Stark LM, Has P, Emerson JB, Schulkin J, Matteson KA. Opioid
21 Knowledge and Prescribing Practices Among Obstetrician-Gynecologists. Obstet
22 Gynecol. 2018 Jan;131(1):150-157. PubMed PMID:29215508.
23
24
25 56. Paulozzi LJ, Strickler GK, Kreiner PW, Koris CM, Centers for Disease C, Prevention.
26 Controlled Substance Prescribing Patterns--Prescription Behavior Surveillance System,
27 Eight States, 2013. Morbidity and mortality weekly report. Surveillance summaries. Oct
28 16 2015;64(9):1-14.
29
30
31
32
33
34
35
36
37
38
39 57. Neuman MD, Bateman BT, Wunsch H, Postoperative pain management and opioids 2
40 Inappropriate opioid prescription after surgery. The Lancet. Vol 393, Issue 10180, 13-19
41 April 2019, Pg 1547-1557.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Flow Chart of the Study Population.






Figure 2. Trajectories of filling opioid prescriptions in 6 months post hysterectomy. The solid or dashed lines represent the distinct estimated opioid prescription filling trajectories. The dot symbols represent the mean estimated proportion in each trajectory group dispensed an opioid prescription at each month. These indicate the mean estimated probability of filling an opioid prescription for patients in each group at each time point. Overall counts and percentages of patients in each group are presented in the underneath table. Patients in trajectory 4 were classified as opioid chronic users. Patients in trajectories 1, 2, 3 were classified as opioid non-chronic users. The x-axis indicates each 30-day interval during the 6-month follow-up period, while the y-axis indicates the estimated proportion of patients filled a prescription opioid within each 30-day interval.



Trajectories	Trajectory 1	Trajectory 2	Trajectory 3	Trajectory 4
N (%)	39,677 (79.15%)	3,304 (6.59%)	6,453 (12.87%)	693 (1.38%)

Figure 2. Proportion in each trajectory group dispensed an opioid prescription in each month.



Trajectories	1	2	3	4
N (%)	2,808 (5.63%)	2,543 (5.10%)	44,059 (88.39%)	434 (0.87%)

Table 1. Comparison of baseline patient characteristics among opioid-naïve patients who received prescription opioids within 7 days after hysterectomy, N=50,127

<u>Patient characteristics</u>	<u>Non-chronic Users[†]</u> (N=48,434)	<u>Chronic Opioid Users[‡]</u> (N= 693)	<u>P-value</u>
<u>Charlson Comorbidity Index, mean (95% CI)</u>	<u>1.60 (1.57, 1.64)</u>	<u>1.74 (1.45, 2.02)</u>	<u>0.3479</u>
<u>Length of stay in days, mean (95% CI)</u>	<u>2.32 (2.30, 2.35)</u>	<u>3.79 (3.39, 4.19)</u>	<u><0.0001</u>
<u>Age, years, mean (95% CI)</u>	<u>48.1 (48.0, 48.2)</u>	<u>49.8 (48.9,50.7)</u>	<u>0.0004</u>
<u>Age at hysterectomy, years</u>			
<u><40</u>	<u>9,100 (18.41)</u>	<u>126 (18.18)</u>	<u><.0001</u>
<u>40-49</u>	<u>22,538 (45.59)</u>	<u>255 (36.80)</u>	
<u>50+</u>	<u>17,796 (36.00)</u>	<u>312 (45.02)</u>	
<u>Substance use and abuse, N (%)</u>			
<u>Tobacco</u>	<u>2,637 (5.33)</u>	<u>60 (8.66)</u>	<u>0.0001</u>
<u>Alcohol</u>	<u>177 (0.36)</u>	<u>9 (1.30)</u>	<u><.0001</u>
<u>Other substance use disorders[§]</u>	<u>55 (0.11)</u>	<u>4 (0.58)</u>	<u>0.0004</u>
<u>Psychiatric medications, N (%)</u>			
<u>Benzodiazepines</u>	<u>4,201 (8.50)</u>	<u>125 (18.04)</u>	<u><.0001</u>
<u>Antidepressants</u>	<u>9,984 (20.20)</u>	<u>233 (33.62)</u>	<u><.0001</u>
<u>Stimulants</u>	<u>767 (1.55)</u>	<u>25 (3.61)</u>	<u><.0001</u>
<u>Pain conditions, N (%)</u>			
<u>Fibromyalgia</u>	<u>1,530 (3.10)</u>	<u>30 (4.33)</u>	<u>0.0632</u>
<u>Headache syndromes</u>	<u>4,019 (8.13)</u>	<u>82 (11.83)</u>	<u>0.0004</u>
<u>Back pain</u>	<u>4,659 (9.42)</u>	<u>110 (15.87)</u>	<u><.0001</u>
<u>Type of hysterectomy, N (%)</u>			
<u>Abdominal hysterectomy</u>	<u>12,757 (25.81)</u>	<u>228 (32.90)</u>	<u><.0001</u>
<u>Vaginal hysterectomy</u>	<u>7,428 (15.03)</u>	<u>74 (10.68)</u>	
<u>Laparoscopic/Robotic</u>	<u>29,249 (59.17)</u>	<u>391 (56.42)</u>	
<u>Surgery setting, N (%)</u>			
<u>Inpatient</u>	<u>19,805 (40.06)</u>	<u>352 (50.79)</u>	<u><.0001</u>
<u>Outpatient</u>	<u>29,629 (59.94)</u>	<u>341 (49.21)</u>	
<u>US census region, N (%)</u>			
<u>Midwest</u>	<u>13,705 (27.72)</u>	<u>186 (26.84)</u>	<u>0.6351</u>
<u>Northeast</u>	<u>5,075 (10.27)</u>	<u>66 (9.52)</u>	
<u>South</u>	<u>23,642 (47.83)</u>	<u>332 (47.91)</u>	
<u>West</u>	<u>7,012 (14.18)</u>	<u>109 (15.73)</u>	
<u>Insurance type, N (%)</u>			
<u>Commercial</u>	<u>45,107 (91.25)</u>	<u>586 (84.56)</u>	<u><.0001</u>
<u>Medicare</u>	<u>4,327 (8.75)</u>	<u>107 (15.44)</u>	
<u>Type of health plan, N (%)</u>			
<u>Exclusive provider organization</u>	<u>5,990 (12.12)</u>	<u>70 (10.10)</u>	<u>0.0008</u>
<u>Health maintenance organization</u>	<u>4,913 (9.94)</u>	<u>89 (12.84)</u>	
<u>Point of service</u>	<u>34,004 (68.79)</u>	<u>449 (64.79)</u>	
<u>Others[¶]</u>	<u>4,527 (9.16)</u>	<u>85 (12.27)</u>	
<u>Year of hysterectomy</u>			
<u>2010-2013</u>	<u>21,949 (44.40)</u>	<u>287 (41.41)</u>	<u>0.1161</u>
<u>2014-2015</u>	<u>27,485 (55.60)</u>	<u>406 (58.59)</u>	
<u>Initial Opioid Prescription Characteristics</u>			

Days' supply, days	4.85 (4.82, 4.87)	5.12 (4.89, 5.34)	0.0205
Average daily MME, mg/day	54.7 (54.3, 55.0)	58.4 (54.6, 62.2)	0.0571
Type of Opioid, N(%)			
Hydrocodone	17,555 (35.51)	266 (38.38)	0.0553
Oxycodone	27,434 (55.50)	354 (51.08)	
Other	4,445 (8.99)	73 (10.53)	
Tertile MME, mg/day			
<=40	16,154 (32.68)	219 (31.60)	0.2425
40.5-58.9	16,718 (33.82)	221 (31.89)	
60.0+	16,562 (33.50)	253 (36.51)	
Days' supply, days			
1-3	15,667 (31.69)	193 (27.85)	0.0566
4-7	29,098 (58.86)	423 (61.04)	
8+	4,669 (9.44)	77 (11.11)	

Abbreviation: MME=Morphine Milligram Equivalent

[†]Non-chronic users refer to patients in trajectory 1, 2, and 3 in Figure 2.

[‡]Chronic Opioid users refer to patients in trajectory 4 in Figure 2.

[§]Other substance use includes cocaine, marijuana, anxiolytic, stimulant, hallucinogenic drugs, or abuse of unspecified drugs/medications.

[¶]Others include Indemnity and Preferred provider organization.

Table 2. Baseline and initial opioid prescription characteristics that predict chronic opioid use following hysterectomy among opioid naive patients, 2010-2015; N=50, 127

Characteristics	Unadjusted OR (95% CI)	Adjusted OR [†] (95% CI)
<u>Age group, years</u>		
<u>40-49</u>	Ref.	Ref.
<u>< 40</u>	<u>1.22 (0.99, 1.52)</u>	<u>1.20 (0.97, 1.49)</u>
<u>≥ 50</u>	<u>1.55 (1.31, 1.83)</u>	<u>1.37 (1.14, 1.65)</u>
<u>Insurance type</u>		
<u>Commercial</u>	Ref.	Ref.
<u>Medicare</u>	<u>1.90 (1.55, 2.35)</u>	<u>1.98 (1.37, 2.87)</u>
<u>Type of health plan, N (%)</u>		
<u>Exclusive provider organization</u>	Ref.	Ref.
<u>Health maintenance organization</u>	<u>1.55 (1.13, 2.13)</u>	<u>1.19 (0.84, 1.68)</u>
<u>Point of service</u>	<u>1.13 (0.88, 1.46)</u>	<u>1.10 (0.85, 1.42)</u>
<u>Others[‡]</u>	<u>1.61 (1.17, 2.21)</u>	<u>0.75 (0.48, 1.18)</u>
<u>Type of hysterectomy</u>		
<u>Vaginal hysterectomy</u>	Ref.	Ref.
<u>Laparoscopic/Robotic</u>	<u>1.79 (1.38, 2.34)</u>	<u>1.52 (1.16, 2.00)</u>
<u>Abdominal hysterectomy</u>	<u>1.34 (1.05, 1.72)</u>	<u>1.40 (1.09, 1.80)</u>
<u>Year (2010-2013 vs 2014-2015)</u>	<u>1.13 (0.97, 1.32)</u>	<u>1.24 (1.06, 1.46)</u>
<u>Length of stay (days)</u>	<u>1.14 (1.11, 1.16)</u>	<u>1.12 (1.10, 1.15)</u>
<u>Charlson comorbidity index</u>	<u>1.18 (1.06, 1.31)</u>	<u>1.16 (1.04, 1.30)</u>
<u>Pain conditions</u>		
<u>Headache (Yes vs. No)</u>	<u>1.52 (1.20, 1.91)</u>	<u>1.33 (1.04, 1.69)</u>
<u>Back pain (Yes vs. No)</u>	<u>1.81 (1.48, 2.23)</u>	<u>1.57 (1.27, 1.94)</u>
<u>Substance use</u>		
<u>Alcohol</u>	<u>3.66 (1.87, 7.18)</u>	<u>1.84 (0.91, 3.74)</u>
<u>Tobacco</u>	<u>1.68 (1.29, 2.20)</u>	<u>1.47 (1.12, 1.93)</u>
<u>Psychiatric medications</u>		
<u>Antidepressants</u>	<u>2.00 (1.71, 2.35)</u>	<u>1.71 (1.45, 2.03)</u>
<u>Benzodiazepines</u>	<u>2.37 (1.95, 2.88)</u>	<u>1.89 (1.54, 2.33)</u>
<u>Psychostimulants</u>	<u>2.38 (1.58, 3.56)</u>	<u>1.84 (1.21, 2.80)</u>
<u>Initial Opioid Prescription Characteristics</u>		
<u>Type of opioid</u>		
<u>Oxycodone</u>	Ref.	Ref.
<u>Hydrocodone</u>	<u>1.17 (1.00, 1.38)</u>	<u>1.31 (1.10, 1.57)</u>
<u>Other</u>	<u>1.27 (0.99, 1.64)</u>	<u>1.26 (0.96, 1.64)</u>
<u>Days' supply, days</u>		
<u>≤ 3</u>	Ref.	Ref.
<u>4-7</u>	<u>1.18 (0.99, 1.40)</u>	<u>1.28 (1.06, 1.54)</u>
<u>≥ 8</u>	<u>1.34 (1.03, 1.75)</u>	<u>1.41 (1.05, 1.89)</u>
<u>Tertile MME, mg/day</u>		
<u>< 40</u>	Ref.	Ref.
<u>40 -58.9</u>	<u>0.98 (0.81, 1.18)</u>	<u>1.14 (0.93, 1.40)</u>
<u>≥ 60.0</u>	<u>1.13 (0.94, 1.35)</u>	<u>1.43 (1.14, 1.79)</u>

Abbreviation: MME=Morphine Milligram Equivalent

Adjusted model: c-statistic = 0.70; Hosmer-Lemeshow Goodness-of-fit Test: $X^2 = 19.30$, $df=8$, $p=0.0133$

1
2
3 [†Adjusted for baseline patient demographic and clinical characteristics listed in this table.](#)

4 [‡Others include Indemnity and Preferred provider organization.](#)
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Table 3. Distribution of possible indications for chronic opioid use during the 6-month follow-up period after hysterectomy: shown as N (%).

<u>Possible indications</u>	<u>All patients in cohort</u> <u>(N = 50,127)</u>	<u>Non-chronic Users</u> <u>(N = 49,434)</u>	<u>Chronic Opioid Users</u> <u>(N = 693)</u>	<u>P-value</u>
<u>Chronic pancreatitis</u>	<u>70 (0.14)</u>	<u>63 (0.13)</u>	<u>7 (1.01)</u>	<u><.0001</u>
<u>Headache syndromes</u>	<u>3,284 (6.55)</u>	<u>3,196 (6.47)</u>	<u>88 (12.70)</u>	<u>< .0001</u>
<u>Back pain</u>	<u>4,477 (8.93)</u>	<u>4,287 (8.67)</u>	<u>190 (27.42)</u>	<u>< .0001</u>
<u>Fibromyalgia</u>	<u>1,470 (2.93)</u>	<u>1,407 (2.85)</u>	<u>63 (9.09)</u>	<u>< .0001</u>
<u>Additional anesthetic procedures[†]</u>				<u>< .0001</u>
<u>0</u>	<u>46,132 (92.03)</u>	<u>45,768 (92.58)</u>	<u>364 (52.53)</u>	
<u>1</u>	<u>3,310 (6.60)</u>	<u>3,099 (6.27)</u>	<u>211 (30.45)</u>	
<u>2</u>	<u>525 (1.05)</u>	<u>459 (0.93)</u>	<u>66 (9.52)</u>	
<u>≥ 3</u>	<u>160 (0.32)</u>	<u>108 (0.22)</u>	<u>52 (7.50)</u>	

[†]Number of additional procedures requiring anesthesia.

Table 1. Comparison of baseline patient characteristics among opioid-naïve patients who received prescription opioids within 7 days after hysterectomy, N=49,844

Patient characteristics	Non-chronic Users** (N=49,410)	Chronic Opioid Users* (N= 434)	P-value
Charlson Comorbidity Index, mean (95% CI)	0.4 (0.4, 0.4)	0.8 (0.7, 0.9)	<.0001
Length of stay in days, mean (95% CI)	1.00 (1.00, 1.02)	2.19 (1.86, 2.51)	<.0001
Age, years, mean (95% CI)	48.1 (48.0, 48.2)	50.5 (49.3, 51.7)	<.0001
Age at hysterectomy, years			
— <40	9,245 (18.71)	84 (19.35)	<.0001
— 40-49	22,461 (45.46)	141 (32.49)	
— 50+	17,704 (35.83)	209 (48.16)	
Substance use and abuse, N (%)			
— Tobacco	2,714 (5.49)	49 (11.29)	<.0001
— Alcohol	181 (0.37)	8 (1.84)	.0003
— Marijuana	5 (0.01)	0 (0.00)	NA
— Cocaine	8 (0.02)	1 (0.23)	0.08
— Other substance abuse***	35 (0.07)	0 (0.0)	NA
Psychiatric medications, N(%)			
— Benzodiazepines	4,403 (8.91)	87 (20.05)	<.0001
— Antidepressants	10,331 (20.91)	159 (36.64)	<.0001
— Stimulants	766 (1.55)	18 (4.15)	<.0001
Pain conditions, N (%)			
— Fibromyalgia	1,692 (3.42)	32 (7.37)	<.0001
— Headache syndromes	4,212 (8.52)	41 (9.45)	0.49
— Back pain	5,053 (10.23)	85 (19.59)	<.0001
Type of hysterectomy, N (%)			
— Abdominal hysterectomy	12,814 (25.93)	152 (35.02)	<.0001
— Vaginal hysterectomy	7,412 (15.00)	57 (13.13)	
— Laparoscopic/Robotic	29,184 (59.06)	225 (51.84)	
US census region, N (%)			
— Midwest	13,907 (28.15)	118 (27.19)	0.07
— Northeast	5,041 (10.20)	36 (8.29)	
— South	23,609 (47.78)	202 (46.54)	
— West	6,853 (13.87)	78 (17.97)	
Insurance type, N (%)			
— Commercial	44,966 (91.01)	354 (81.57)	<.0001
— Medicare	4,444 (8.99)	80 (18.43)	
Type of health plan, N (%)			
— Exclusive provider organization	5,905 (11.95)	43 (9.91)	0.0004
— Health maintenance organization	4,899 (9.91)	55 (12.67)	
— Point of service	33,992 (68.80)	274 (63.13)	
— Others****	4,614 (9.31)	62 (14.29)	
Year of hysterectomy			
— 2010-2013	21,952 (44.43)	180 (41.47)	0.22
— 2014-2015	27,458 (55.57)	254 (58.53)	
Initial Opioid Prescription Characteristics			
Days' supply, days	4.81 (4.78, 4.83)	5.44 (5.07, 5.80)	<0.0001
Average daily MME*****, mg/day	55.0 (54.7, 55.4)	61.1 (55.5, 66.8)	0.03
Type of Opioid, N(%)			
— Hydrocodone	18,738 (37.92)	176 (40.55)	0.16

— Oxycodone	27,332 (55.32)	222 (51.15)	
— Other	3,340 (6.76)	36 (8.29)	
Tertile MME****, mg/day			
— ≤40	16,964 (34.33)	140 (32.26)	0.05
— 40.5-58.9	15,794 (31.97)	124 (28.57)	
— 60.0+	16,652 (33.70)	170 (39.17)	
Days' supply, days			
— 1-3	15,838 (32.05)	116 (26.73)	0.001
— 4-7	29,097 (58.89)	259 (59.68)	
— 8+	4,475 (9.06)	59 (13.59)	

Note:

* Chronic Opioid users refer to patients in trajectory 4 in Figure 2.

** Non-chronic users refer to patients in trajectory 1, 2, and 3 in Figure 2.

*** Other substance use includes anxiolytic, stimulant, hallucinogenic drugs, or abuse of unspecified drugs/medications.

**** Others including Indemnity and Preferred provider organization.

***** MME=Morphine Milligram Equivalent

Review Only

Table 2. Baseline and initial opioid prescription characteristics that predict chronic opioid use following hysterectomy among opioid naive patients, 2010-2015; N=49,844

Characteristics	Unadjusted OR (95% CI)	Adjusted OR* (95% CI)
Age group, years		
— 40-49	Ref.	Ref.
— < 40	1.45 (1.10, 1.90)	1.42 (1.08, 1.87)
— 50+	1.88 (1.52, 2.33)	1.54 (1.22, 1.95)
US census region		
— Northeast	Ref.	Ref.
— Midwest	1.19 (0.82, 1.73)	1.33 (0.91, 1.96)
— South	1.20 (0.84, 1.71)	1.31 (0.91, 1.88)
— West	1.59 (1.07, 2.37)	1.77 (1.18, 2.66)
Insurance type		
— Commercial	Ref.	Ref.
— Medicare	2.29 (1.79, 2.92)	1.58 (1.19, 2.11)
Type of hysterectomy		
— Vaginal hysterectomy	Ref.	Ref.
— Laparoscopic/Robotic	1.00 (0.75, 1.34)	1.03 (0.77, 1.39)
— Abdominal hysterectomy	1.54 (1.14, 2.09)	1.35 (0.98, 1.87)
— Year (2010-2013 vs 2014-2015)	1.13 (0.93, 1.37)	1.27 (1.05, 1.56)
— Length of stay (days)	1.13 (1.11, 1.16)	1.10 (1.07, 1.13)
— Charlson comorbidity index	1.34 (1.26, 1.42)	1.18 (1.10, 1.26)
Pain conditions		
— Fibromyalgia (Yes vs. No)	2.25 (1.56, 3.23)	1.58 (1.08, 2.31)
— Back pain (Yes vs. No)	2.14 (1.69, 2.72)	1.70 (1.32, 2.18)
Substance use		
— Alcohol	5.11 (2.50, 10.44)	2.28 (1.06, 4.87)
— Tobacco	2.19 (1.62, 2.96)	1.81 (1.33, 2.46)
Psychiatric medications		
— Antidepressants	2.19 (1.80, 2.66)	1.78 (1.44, 2.19)
— Benzodiazepines	2.56 (2.02, 3.25)	1.94 (1.51, 2.49)
— Psychostimulants	2.75 (1.71, 4.43)	2.07 (1.26, 3.41)
Initial Opioid Prescription Characteristics		
Type of opioid		
— Oxycodone	Ref.	Ref.
— Hydrocodone	1.15 (0.95, 1.41)	1.35 (1.07, 1.70)
— Other	1.33 (0.93, 1.89)	1.37 (0.95, 1.97)
Days' supply, days		
— ≤3	Ref.	Ref.
— 4-7	1.22 (0.96, 1.51)	1.41 (1.11, 1.79)
— 8+	1.80 (1.31, 2.47)	2.04 (1.43, 2.90)
Tertile MME, mg/day		
— ≤40	Ref.	Ref.
— 40.5-58.9	0.95 (0.75, 1.21)	1.24 (0.95, 1.62)
— 60.0+	1.24 (0.99, 1.55)	1.70 (1.28, 2.26)

Adjusted model: c-statistic = 0.73; Hosmer-Lemeshow Goodness-of-fit Test: $X^2=9.32$, $df=8$, $p=0.32$

*: Adjusted for baseline patient demographic and clinical characteristics listed in this table.

Table 3. Distribution of possible indications for chronic opioid use during the 6-month follow-up period after hysterectomy: shown as N (%).

Possible indications	All patients in cohort (N = 49,844)	Non-chronic Users (N = 49,410)	Chronic Opioid Users (N = 414)	P-value
Rheumatoid arthritis	6 (0.01)	6 (0.01)	0 (0.00)	NA
Chronic pancreatitis	73 (0.15)	67 (0.14)	6 (1.38)	<.0001
Sickle cell disease	7 (0.01)	7 (0.01)	0 (0.00)	NA
Headache syndromes	3,373 (6.77)	3,311 (6.70)	62 (14.29)	<0.0001
Back pain	4,652 (9.33)	4,511 (9.13)	141 (32.49)	<0.0001
Fibromyalgia	1,588 (3.19)	1,548 (3.13)	40 (9.22)	<0.0001
Additional anesthetic procedures*				<0.0001
0	45,840 (91.97)	45,621 (92.33)	219 (50.46)	
1	3,316 (6.65)	3,177 (6.43)	139 (32.03)	
2	524 (1.05)	479 (0.97)	45 (10.37)	
3+	164 (0.33)	133 (0.27)	31 (7.14)	

Note:

*: Number of additional procedures requiring anesthesia.

CONFLICT OF INTEREST DISCLOSURE

The Editors of *Pharmacoepidemiology and Drug Safety* recognize that most studies in pharmacoepidemiology cost money and thus pose a potential conflict of interest. As a conflict of interest may affect the assessment or judgment of an author, we ask that **all** authors (not just the Corresponding Author) complete the following form.

For Co-authors: Please complete questions 4-10. Completed forms should be saved, and emailed as an attachment to the Corresponding Author.

For Corresponding Authors: Please complete all questions. It is the responsibility of the Corresponding Author to submit completed forms on behalf of all co-authors via Manuscript Central at the point of manuscript submission.

Corresponding author only (Co-authors go to Question 4):

POTENTIAL STUDY INTERPRETATION CONFLICTS

1. Some or all of the data that were used in this study were provided by a company with a vested interest in the product being studied. No
2. The sponsor of this project had the right of commenting but the authors retained the right to accept or reject comments or suggestions. No
3. The sponsor of this project had the right of final editing and/or approval of the manuscript submitted. No

Corresponding author and Co-authors:

POTENTIAL FINANCIAL CONFLICTS

4. I, my spouse, or one of my dependent children is an employee of a company whose product is being studied. No
5. I, my spouse, or one of my dependent children has significant equity interest (>USD 10,000) in the company that owns the product being studied. No

1
2
3 6. In the past three years I have:
4

5 been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
6 product being studied, on issues related to the product being studied; No
7

8 been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
9 product being studied, on issues unrelated to the product being studied; No
10

11 received research or educational support from a company with a vested interest in the product(s)
12 being studied. Yes
13

14
15
16 7. A company whose product is being studied has provided funding to support the work on this
17 project. No
18

19
20
21 If you have answered YES to any of the above questions, or if you have additional personal, commercial or
22 academic conflicts of interest, please draft a statement to publish with the article. e.g., AB has been
23 reimbursed by Safe Drug Ltd. for international conference attendance.
24

25 N/A
26
27

28
29
30 8. Manuscript title (first six words are sufficient)
31

32 **Chronic Opioid Use in Women following Hysterectomy**
33
34

35 9. Author's full name (a separate form must be submitted for each author)
36

37 **Xuerong Wen**
38

39 10. In checking this box, I confirm I have completed this form to the best of my knowledge.
40
41

42
43
44
45
46 This form is available online by [clicking here](#)
47
48
49
50
51
52
53
54
55
56
57

CONFLICT OF INTEREST DISCLOSURE

The Editors of *Pharmacoepidemiology and Drug Safety* recognize that most studies in pharmacoepidemiology cost money and thus pose a potential conflict of interest. As a conflict of interest may affect the assessment or judgment of an author, we ask that **all** authors (not just the Corresponding Author) complete the following form.

For Co-authors: Please complete questions 4-10. Completed forms should be saved, and emailed as an attachment to the Corresponding Author.

For Corresponding Authors: Please complete all questions. It is the responsibility of the Corresponding Author to submit completed forms on behalf of all co-authors via Manuscript Central at the point of manuscript submission.

Corresponding author only (Co-authors go to Question 4):

POTENTIAL STUDY INTERPRETATION CONFLICTS

1. Some or all of the data that were used in this study were provided by a company with a vested interest in the product being studied. n/a
2. The sponsor of this project had the right of commenting but the authors retained the right to accept or reject comments or suggestions. n/a
3. The sponsor of this project had the right of final editing and/or approval of the manuscript submitted. n/a

Corresponding author and Co-authors:

POTENTIAL FINANCIAL CONFLICTS

4. I, my spouse, or one of my dependent children is an employee of a company whose product is being studied. No
5. I, my spouse, or one of my dependent children has significant equity interest (>USD 10,000) in the company that owns the product being studied. No

1
2
3 6. In the past three years I have:
4

5 been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
6 product being studied, on issues related to the product being studied; No
7

8 been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
9 product being studied, on issues unrelated to the product being studied; No
10

11 received research or educational support from a company with a vested interest in the product(s)
12 being studied. No
13

14
15
16 7. A company whose product is being studied has provided funding to support the work on this
17 project. No
18

19
20
21 If you have answered YES to any of the above questions, or if you have additional personal, commercial or
22 academic conflicts of interest, please draft a statement to publish with the article. e.g., AB has been
23 reimbursed by Safe Drug Ltd. for international conference attendance.
24
25
26
27

28
29
30 8. Manuscript title (first six words are sufficient)

31 **Chronic Opioid Use in Women following Hysterectomy**
32
33

34
35 9. Author's full name (a separate form must be submitted for each author)

36 **Kristen A. Matteson**
37

38
39 10. In checking this box, I confirm I have completed this form to the best of my knowledge.
40
41

42
43
44
45
46 This form is available online by [clicking here](#)
47
48
49
50
51
52
53
54
55
56
57

CONFLICT OF INTEREST DISCLOSURE

The Editors of *Pharmacoepidemiology and Drug Safety* recognize that most studies in pharmacoepidemiology cost money and thus pose a potential conflict of interest. As a conflict of interest may affect the assessment or judgment of an author, we ask that **all** authors (not just the Corresponding Author) complete the following form.

For Co-authors: Please complete questions 4-10. Completed forms should be saved, and emailed as an attachment to the Corresponding Author.

For Corresponding Authors: Please complete all questions. It is the responsibility of the Corresponding Author to submit completed forms on behalf of all co-authors via Manuscript Central at the point of manuscript submission.

Corresponding author only (Co-authors go to Question 4):

POTENTIAL STUDY INTERPRETATION CONFLICTS

1. Some or all of the data that were used in this study were provided by a company with a vested interest in the product being studied. n/a
2. The sponsor of this project had the right of commenting but the authors retained the right to accept or reject comments or suggestions. n/a
3. The sponsor of this project had the right of final editing and/or approval of the manuscript submitted. n/a

Corresponding author and Co-authors:

POTENTIAL FINANCIAL CONFLICTS

4. I, my spouse, or one of my dependent children is an employee of a company whose product is being studied. No
5. I, my spouse, or one of my dependent children has significant equity interest (>USD 10,000) in the company that owns the product being studied. No

1
2
3 6. In the past three years I have:
4

- 5 • been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
6 product being studied, on issues related to the product being studied; No
7
8 • been paid as a consultant (or in a similar capacity by a company with a vested interest in the
9 product being studies, on issues unrelated to the product being studied; No
10
11 • received research or educational support from a company with a vested interest in the product(s)
12 being studied. No
13

14
15
16 7. A company whose product is being studied has provided funding to support the work on this
17 project. No
18

19
20
21 If you have answered YES to any of the above questions, or if you have additional personal, commercial or
22 academic conflicts of interest, please draft a statement to publish with the article. e.g., AB has been
23 reimbursed by Safe Drug Ltd. for international conference attendance.
24
25
26
27

28
29
30 8. Manuscript title (first six words are sufficient)

31
32 **Chronic Opioid Use in Women following Hysterectomy**
33

34
35 9. Author's full name (a separate form must be submitted for each author)

36
37 **Hilary Aroke**
38

39 10. In checking this box, I confirm I have completed this form to the best of my knowledge.
40
41

42
43
44
45
46 This form is available online by [clicking here](#)
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CONFLICT OF INTEREST DISCLOSURE

The Editors of *Pharmacoepidemiology and Drug Safety* recognize that most studies in pharmacoepidemiology cost money and thus pose a potential conflict of interest. As a conflict of interest may affect the assessment or judgment of an author, we ask that **all** authors (not just the Corresponding Author) complete the following form.

For Co-authors: Please complete questions 4-10. Completed forms should be saved, and emailed as an attachment to the Corresponding Author.

For Corresponding Authors: Please complete all questions. It is the responsibility of the Corresponding Author to submit completed forms on behalf of all co-authors via Manuscript Central at the point of manuscript submission.

Corresponding author only (Co-authors go to Question 4):

POTENTIAL STUDY INTERPRETATION CONFLICTS

1. Some or all of the data that were used in this study were provided by a company with a vested interest in the product being studied. n/a
2. The sponsor of this project had the right of commenting but the authors retained the right to accept or reject comments or suggestions. n/a
3. The sponsor of this project had the right of final editing and/or approval of the manuscript submitted. n/a

Corresponding author and Co-authors:

POTENTIAL FINANCIAL CONFLICTS

4. I, my spouse, or one of my dependent children is an employee of a company whose product is being studied. No
5. I, my spouse, or one of my dependent children has significant equity interest (>USD 10,000) in the company that owns the product being studied. No

1
2
3 6. In the past three years I have:
4

- 5
- 6 • been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
7 product being studied, on issues related to the product being studied; No
 - 8 • been paid as a consultant (or in a similar capacity by a company with a vested interest in the
9 product being studies, on issues unrelated to the product being studied; No
 - 10 • received research or educational support from a company with a vested interest in the product(s)
11 being studied. No
- 12
13
14

15
16 7. A company whose product is being studied has provided funding to support the work on this
17 project. No
18

19
20
21 If you have answered YES to any of the above questions, or if you have additional personal, commercial or
22 academic conflicts of interest, please draft a statement to publish with the article. e.g., AB has been
23 reimbursed by Safe Drug Ltd. for international conference attendance.
24
25
26
27

28
29
30 8. Manuscript title (first six words are sufficient)

31
32 **Chronic Opioid Use of after Hysterectomy**
33
34

35 9. Author's full name (a separate form must be submitted for each author)

36
37 **Stephen J Kogut**
38

39 10. In checking this box, I confirm I have completed this form to the best of my knowledge.
40
41

42
43
44
45
46 This form is available online by [clicking here](#)
47
48
49
50
51
52
53
54
55
56
57

58 September 2016

CONFLICT OF INTEREST DISCLOSURE

The Editors of *Pharmacoepidemiology and Drug Safety* recognize that most studies in pharmacoepidemiology cost money and thus pose a potential conflict of interest. As a conflict of interest may affect the assessment or judgment of an author, we ask that **all** authors (not just the Corresponding Author) complete the following form.

For Co-authors: Please complete questions 4-10. Completed forms should be saved, and emailed as an attachment to the Corresponding Author.

For Corresponding Authors: Please complete all questions. It is the responsibility of the Corresponding Author to submit completed forms on behalf of all co-authors via Manuscript Central at the point of manuscript submission.

Corresponding author only (Co-authors go to Question 4):

POTENTIAL STUDY INTERPRETATION CONFLICTS

1. Some or all of the data that were used in this study were provided by a company with a vested interest in the product being studied. n/a
2. The sponsor of this project had the right of commenting but the authors retained the right to accept or reject comments or suggestions. n/a
3. The sponsor of this project had the right of final editing and/or approval of the manuscript submitted. n/a

Corresponding author and Co-authors:

POTENTIAL FINANCIAL CONFLICTS

4. I, my spouse, or one of my dependent children is an employee of a company whose product is being studied. No
5. I, my spouse, or one of my dependent children has significant equity interest (>USD 10,000) in the company that owns the product being studied. No

1
2
3 6. In the past three years I have:
4

- 5 • been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
6 product being studied, on issues related to the product being studied; No
7
8 • been paid as a consultant (or in a similar capacity) by a company with a vested interest in the
9 product being studied, on issues unrelated to the product being studied; No
10
11 • received research or educational support from a company with a vested interest in the product(s)
12 being studied. No
13

14
15
16 7. A company whose product is being studied has provided funding to support the work on this
17 project. No
18

19
20
21 If you have answered YES to any of the above questions, or if you have additional personal, commercial or
22 academic conflicts of interest, please draft a statement to publish with the article. e.g., AB has been
23 reimbursed by Safe Drug Ltd. for international conference attendance.
24
25
26
27

28
29
30 8. Manuscript title (first six words are sufficient)

31
32 **Chronic Opioid Use in Women following Hysterectomy**
33

34
35 9. Author's full name (a separate form must be submitted for each author)

36
37 **Lynn Taylor**
38

39 10. In checking this box, I confirm I have completed this form to the best of my knowledge.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60