1976

Percus-Yevick and Hypernetted-Chain Approximations

A. Choudry
University of Rhode Island

Pranab K. Banerjee
University of Rhode Island, pbanerjee@uri.edu

Narinder K. Allawadi

Follow this and additional works at: https://digitalcommons.uri.edu/phys_facpubs

Terms of Use

All rights reserved under copyright.

Citation/Publisher Attribution

Choudry, A., Banerjee, P. K., & Ailawadi, N. K.. (1976). Percus-Yevick and hypernetted-chain approximations. *Physical Review A*, 14(1), 434-437. doi: 10.1103/PhysRevA.14.434
Available at: <http://dx.doi.org/10.1103/PhysRevA.14.434>

This Article is brought to you for free and open access by the Physics at DigitalCommons@URI. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Percus-Yevick and hypernetted-chain approximations*

A. Choudry

Department of Physics, University of Rhode Island, Kingston, Rhode Island 02881

P. K. Banerjee

Department of Physics, Rhode Island Junior College, Warwick, Rhode Island 02886

Narinder K. Ailawadi

Institut für Theoretische Physik (Kondensierte Materie), Freie Universität Berlin, West Berlin, Germany

(Received 10 November 1975)

Percus-Yevick and hypernetted-chain models connecting the structure factor $S(k)$ with effective pair potential $\phi(r)$ have been studied in detail. Recent neutron-scattering data of Yarnell *et al.*, for liquid argon at 85°K, have been employed to generate these potentials. Using 0.01 uncertainty in the experimental data, the uncertainty in the pair potentials has been computed. Having developed a general method of error propagation analysis, the effect of experimental errors in $S(k)$ on Percus-Yevick and hypernetted-chain potentials is investigated and the resulting potentials are compared with Lennard-Jones and Barker, Fisher, and Watts potentials.

INTRODUCTION

Recent accurate measurements of structure factors $S(k)$ have led to a renewed interest in the approximate theories of liquids. Typically, the data for the measured structure factor are inserted in one or the other of integral equations used in the theory of liquids, and information about the pair potentials is obtained. This procedure has been used for example, for liquid metals, particularly liquid Na and K,¹⁻⁶ in order to calculate effective pair potentials.

More recently, Yarnell *et al.*⁷ have measured structure factor of liquid argon at 85°K with an accuracy of 0.01 by neutron-diffraction techniques. These experimental data are in agreement with the molecular-dynamics data on $S(k)$ of Verlet,⁸ based on the Lennard-Jones (L-J) potential (6-12),

$$\phi(r) = 4\epsilon[(\sigma/r)^{12} - (\sigma/r)^6] \quad (1)$$

with $\epsilon/k_B = 119.8^\circ\text{K}$ and $\sigma = 3.405 \text{ \AA}$ as parameters.

The data of Yarnell *et al.*⁷ are also in good agreement with the simulations based on a rather different pair potential, which includes three-body effects.⁹ Thus, one is led to the conclusion⁷ that liquid argon can be equally well described by the L-J potential, Eq. (1), or by the Barker, Fisher, and Watts (BFW) potential.⁹

In an effort to obtain some information about the effective pair interaction, the Percus-Yevick (PY) and hypernetted-chain (HNC) approximations have been studied for a long time.¹⁰ It has been pointed out^{8,11} that the PY equation (and perhaps HNC) leads to quite accurate pair potentials at low densities but breaks down at liquid densities. Also,

in order to obtain 10% accuracy in the calculated pair potentials, an accuracy of 1% is needed in the measured structure factor.^{8,11,12} More recently, it has been shown¹³ that the equilibrium structure of simple liquids is dominated by the repulsive part of the potential and that the attractive part of the potential can be treated as a perturbation.¹⁴

In this paper we reconsider the following problem: Given the equilibrium structure factor, can we obtain information about the pair potential? In particular, we restudy the PY and HNC equations for the reason that more accurate data on $S(k)$ are now available, and thus the limits of the reliability of the inversion process need to be thoroughly investigated; i.e., for a given accuracy in the structure-factor measurements, how much error is propagated in computing $\phi_{PY}(r)$ and $\phi_{HNC}(r)$, the potentials obtained from PY and HNC equations; and we examine if their differences from L-J and BFW are due to experimental errors. Such a comparison might indicate if either of the two (PY and HNC) is closer to L-J. On theoretical grounds, there seems to be no clear argument for preferring one approximation or the other; the same is true for the Born-Green equation as well, although for liquid metals, the potential derived from Born-Green theory is usually preferred because of long-range Friedel oscillations.

For this purpose, a detailed analysis of the effect of the uncertainties of the input structure-factor data on the derived pair potentials $\phi_{PY}(r)$ and $\phi_{HNC}(f)$ is carried out. In particular, we have attempted to develop a general method for treating the errors, which yields not just the magnitude

of the error in ϕ but the density distribution thereof in the r - ϕ space.

METHOD

Experimentally one counts the number of neutrons scattered at an angle θ to determine the differential cross section $\sigma(\theta)$:¹⁶

$$\frac{d\sigma}{d\Omega} = \sigma(\theta) = \langle f(\theta)^* f(\theta) \rangle. \quad (2)$$

For n scattering by a nucleus of mass M , atomic number A , $f(\theta)$ is given by¹⁵ [$m = M/(A+)$]

$$f(\theta) = -\frac{m}{2\pi\hbar^2} \int e^{i\vec{k}\cdot\vec{r}} V(\vec{r}) d\vec{r}. \quad (3)$$

The potential $V(r)$ is assumed to be of Fermi type and is obtained by introducing a coherent scattering length b and summing over all the N particles:

$$V(r) = \frac{2\pi\hbar^2}{m} \frac{b}{N} \sum_{i=1}^N \delta(\vec{r} - \vec{r}_i), \quad (4)$$

which yields

$$\sigma(\theta) = \frac{b^2}{N} \int e^{i\vec{k}\cdot\vec{r}} \left\langle \sum_{ij} \delta(\vec{r} + \vec{r}_i - \vec{r}_j) \right\rangle d\vec{r}. \quad (5)$$

The pair distribution function $g(r)$ is defined by

$$g(r) = \frac{1}{\rho N} \left\langle \sum_{i \neq j} \delta(\vec{r} + \vec{r}_i - \vec{r}_j) \right\rangle \quad (6)$$

so that

$$dn(r) = n 4\pi r^2 g(r) dr, \quad (7)$$

where $dn(r)$ is the number of particles in a spherical shell of radius r and thickness dr , n being the number of particles in a unit volume. Combining it with Eq. (5), we have

$$\sigma(\theta) = b^2 \left(\int \rho [g(r) - 1] e^{i\vec{k}\cdot\vec{r}} d\vec{r} + 1 + \rho \delta(k) \right) \quad (8)$$

from which one defines the structure factor $S(k)$:

$$S(\vec{k}) = 1 + \rho \int e^{i\vec{k}\cdot\vec{r}} [g(r) - 1] d\vec{r}. \quad (9)$$

By combining Eqs. (8) and (9), a simple relation between $\sigma(\theta)$ and $S(k)$ can be obtained. After integrating the angular part, $S(k)$ and $g(r)$ can be written as a pair of Fourier transforms:

$$S(k) = 1 + \frac{4\pi n}{k} \int [g(r) - 1] r \sin kr dr, \quad (10)$$

$$g(r) = 1 + A \int [S(k) - 1] k \sin kr dk, \quad (11)$$

with $A = 1/(2\pi n^2 r)$. This series of well-known steps Eqs. (2)–(11) is reviewed here to point out the sources and analysis of errors involved in this procedure and the subsequent calculation of the

potentials. To convert the experimental neutron counts into $\sigma(\theta)$, a number of experimental parameters, such as cell geometry, detector efficiency, etc. are needed and all of these have some errors in them. To calculate $S(k)$ from $\sigma(\theta)$, cross sections and limiting values of $S(k \rightarrow 0)$ and $g(r \rightarrow 0)$ are needed. Details of the error analysis in this procedure are discussed by Yarnell *et al.*⁷ Symbolically the entire above analysis can be thought of as a transformation $T_{s\sigma}$ such that

$$\rho_s(k) = T_{s\sigma} \rho_o(\theta), \quad (12)$$

where $\rho_o(\theta)$ is a density distribution function in the θ - σ plane and is identical with the scattergram of the experimental points in that plane; $\rho_s(k)$ is the corresponding density distribution function in k - s plane. Thus the fact that all experimental values of $S(k)$ are normally distributed about the mean $S_o(k)$ with a constant variance v^2 (e.g., $v = 0.01$) can be stated as

$$E\{S(k)\} = S_o(k) = \int_0^\infty S(k) N(S_o, v^2) dS|_k \quad (13)$$

and

$$E\{(S - S_o)^2\} = v^2, \quad (14)$$

where $N(S_o, v^2)$ is a normal distribution with mean S_o and variance v^2 , and $E\{x\}$ is the expected value of x . Knowing $S(k)$ and $g(r)$ [from Eq. (11)] we can calculate $g(r)$ by using Eq. (11) and then the PY and HNC potentials can be calculated by the well-known relations

$$\phi_{PY} = k_B T \ln[1 - c(r)/g(r)], \quad (15)$$

$$\phi_{HNC} = k_B T [g(r) - c(r) - 1 - \ln g(r)], \quad (16)$$

where

$$c(r) = A \int_0^\infty [S(k) - 1] \frac{k \sin kr}{S(k)} dk. \quad (17)$$

We shall now discuss how the errors in $S(k)$ are propagated through $g(r)$ into ϕ_{PY} and ϕ_{HNC} . The fact that the precision in calculating the pair potentials is strongly dependent on the measurement precision in $S(k)$, has been well established, e.g., Levesque and Verlet,¹¹ Gehlen and Enderby¹⁷ developed a model of linear propagation of errors and applied it to Na. However this model is rather restrictive and may lead to doubtful estimates. We shall now formulate the general problem and indicate a solution. Mathematically speaking, the experimental function $S(k)$ is a set of stochastic variables with arbitrary distributions in k - S space and thus all operations on $S(k)$, e.g., differentiation, integration, etc. should be done in conformity with the rules appropriate to stochastic variables. We introduce an experimental error func-

tion $\epsilon(k, S)$:

$$S(k) = S_0 + \epsilon(k, S) \quad (18)$$

so that

$$E\{\epsilon(k, S)\} = 0. \quad (19)$$

From Eq. (11) it can be shown that the error function $\Delta g(r)$ for $g(r)$ is given by the following stochastic integral:

$$\Delta g(r) = A \int_0^\infty \epsilon(k, S) k \sin kr dk. \quad (20)$$

It should be noted that $\Delta g(r)$ is also a set of stochastic variables and Eq. (20) may be interpreted as a transformation T_{gs} , in the spirit of Eq. (12), i.e.,

$$\Delta g(r) = T_{gs} \epsilon(S, k). \quad (21)$$

Similarly using Eqs. (15), (16), and (17) the following can be defined:

$$\Delta c(r) = T_{cs} \epsilon(S, k), \quad (22)$$

$$\Delta \phi_{PY} = T_{PS} \epsilon(S, k), \quad (23)$$

$$\Delta \phi_{HNC} = T_{HS} \epsilon(S, k). \quad (24)$$

All of the above involve stochastic integrals of the type in Eq. (20). Ballentine and Jones¹² have replaced the above stochastic operations by a linear relation in assuming that the T 's are matrices. As an illustration we shall indicate the evaluation of $\Delta g(r)$ from Eq. (20). For a finite number n of experimental points, in machine computation, Eq. (20) can be written as

$$\Delta g(r) = \sum_{i=1}^n \epsilon_i k_i \sin k_i r \Delta k_i, \quad (25)$$

which is a sum of n independent stochastic vari-

ables η_i where

$$\eta_i = \epsilon_i k_i \sin k_i r \Delta k_i. \quad (26)$$

The moment-generating function $\xi_i(t)$ of η_i is defined as

$$\xi_i(t) = \int_0^\infty \eta_i(s) e^{ts} ds \quad (27)$$

in terms of which (25) can be written as

$$\Delta g(r) = \int_0^\infty \left\{ \prod_{i=1}^n \xi_i(t) \right\} e^{-\epsilon t} dt, \quad (28)$$

which is a density distribution in r - g space. Expressions to be similarly evaluated corresponding to Eqs. (22), (23), and (24) are

$$\Delta c(r) = A \sum_{n=1}^\infty \left(\frac{-\epsilon(k)}{S_0} \right)^n \frac{k \sin kr}{S_0(k)} dk, \quad (29)$$

$$\Delta \phi_{HNC} = \Delta g(r) + \Delta c(r) - \ln(1 + \alpha), \quad (30)$$

$$\Delta \phi_{PY} = k_B T \ln \left[1 - \frac{c}{g} \left(\alpha + (1 + \alpha) \sum_{n=1}^\infty (-\beta)^n \right) e^{-\phi/kT} \right], \quad (31)$$

where $\alpha = \Delta g/g$ and $\beta = \Delta c/c$. Thus in principle it is possible to translate the density of experimental points in the k - s space into the corresponding density in r - ϕ space for arbitrary error functions.

CONCLUSION

Using the structure factor data of Ref. 7 and the error of 0.01, we have calculated HNC and PY potentials and the errors in them. These potentials are compared with L-J and BFW in Figs. 1 and 2, respectively (N.B. the errors in HNC and PY as displayed in these figures are magnified 10 times). The errors in PY (cf. Fig. 2) are much

FIG. 1. $\Delta \phi_{HNC}$ and ϕ_{HNC} compared with L-J and BFW.

FIG. 2. $\Delta \phi_{PY}$ and ϕ_{PY} compared with L-J and BFW.

more pronounced (up to 15%) than in HNC (cf. Fig. 1). In particular, the well depth of PY is rather severely affected by the errors in $S(k)$.^{8,11,12} It is interesting to note that the well depth of HNC agrees with L-J better than with BFW. Of course this agreement does not necessarily imply the superior validity of one or the other potential. However, one can conclude that by using the general analysis developed here, the differences from L-J and BFW are beyond experimental errors. A recent perturbation calculation of Madden and Fitts¹⁸ for liquid Ar based on HNC gives much

more accurate results, and our analysis supports this conclusion.

ACKNOWLEDGMENTS

We would like to thank the staff of the URI computer center for the generous use of their facilities. N. K. A. would like to thank Professor A. Rahman for discussions and his interest in this problem; A. C. would like to express his gratitude to Professor Vernier for extending kind hospitality at Faculté des Science, Université de Dijon.

*Supported in part by the Deutsche Forschungs Gemeinschaft.

¹Y. Waseda and K. Suzuki, in *Properties of Liquid Metals*, edited by S. Takeuchi (Taylor and Francis, London, 1973), p. 37.

²Y. Waseda and K. Suzuki, *Phys. Stat. Sol. B* **57**, 351 (1973).

³W. S. Howells and J. E. Enderby, *J. Phys. C* **5**, 1277 (1972).

⁴W. S. Howells, in Ref. 1, p. 43.

⁵N. K. Ailawadi, P. K. Banerjee, and A. Choudry, *J. Chem. Phys.* **60**, 2571 (1974); **60**, 5124 (1974).

⁶R. Kumaravadeivel, R. Evans, and D. A. Greenwood, *J. Phys. F* **4**, 1839 (1974).

⁷J. L. Yarnell, M. J. Katz, R. G. Wenzel, and S. K. Koenig, *Phys. Rev. A* **7**, 2130 (1973).

⁸L. Verlet, *Phys. Rev.* **165**, 201 (1967).

⁹J. A. Barker, R. A. Fisher, and R. O. Watts, *Mol. Phys.* **21**, 657 (1971).

¹⁰See, for example, P. G. Mikolaj and C. J. Pings, *J.*

Chem. Phys. **46**, 1401 (1967); H. Fredrikze, C. D. Andriesse, and E. Legrand, *Physica* **62**, 474 (1972).

¹¹D. Levesque and L. Verlet, *Phys. Rev. Lett.* **20**, 905 (1968).

¹²L. E. Ballentine and J. C. Jones, in Ref. 1, p. 51.

¹³D. Chandler and J. D. Weeks, *Phys. Rev. Lett.* **25**, 149 (1970).

¹⁴H. C. Andersen, J. D. Weeks, and D. Chandler, *Phys. Rev. A* **4**, 1597 (1971); H. C. Andersen, D. Chandler, and J. D. Weeks, *J. Chem. Phys.* **56**, 3812 (1972).

¹⁵See, for example, P. A. Egelstaff, *An Introduction to the Liquid State* (Academic, London, 1967).

¹⁶W. A. Schlup, Symposium on Inelastic Scattering of Neutrons by Condensed Systems, Brookhaven National Laboratory Report No. BNL 940, 1965 (unpublished).

¹⁷P. C. Gehlen and J. E. Enderby, *J. Chem. Phys.* **51**, 547 (1969).

¹⁸W. G. Madden and D. D. Fitts, *J. Chem. Phys.* **61**, 5475 (1974); *Mol. Phys.* **30**, 1809 (1975).