

2017

Advances in the application of amino acid nitrogen isotopic analysis in ecological and biogeochemical studies

Naohiko Ohkouchi

Yoshito Chikaraishi

Hilary Close

Brian Fry

Thomas Larsen

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

Citation/Publisher Attribution

Ohkouchi, N., Chikaraishi, Y., Close, H., Fry, B., Larsen, T., Madigan, D. J., McCarthy, M. D.,...Yokoyama, Y. (2017). Advances in the application of amino acid nitrogen isotopic analysis in ecological and biogeochemical studies. *Organic Chemistry*, 113, 150-174. doi: 10.1016/j.orggeochem.2017.07.009. Available at: <https://doi.org/10.1016/j.orggeochem.2017.07.009>

This Article is brought to you by the University of Rhode Island. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu. For permission to reuse copyrighted content, contact the author directly.

Advances in the application of amino acid nitrogen isotopic analysis in ecological and biogeochemical studies

Authors

Naohiko Ohkouchi, Yoshito Chikaraishi, Hilary Close, Brian Fry, Thomas Larsen, Daniel J. Madigan, Matthew D. McCarthy, Kelton McMahon, Toshi Nagata, Yuichi I. Naito, Nanako O. Ogawa, Brian N. Popp, Shawn Steffan, Yoshinori Takano, Ichiro Tayasu, Alex S. J. Wyatt, Yasuhiko T. Yamaguchi, and Yusuke Yokoyama

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

1 **Advances in the application of amino acid nitrogen isotopic analysis**
2 **in ecological and biogeochemical studies**

3
4 Naohiko Ohkouchi^{a,*}, Yoshito Chikaraishi^{a,m}, Hilary Close^b, Brian Fry^c, Thomas Larsen^d, Daniel J.
5 Madigan^e, Matthew D. McCarthy^f, Kelton W. McMahon^g, Toshi Nagata^h, Yuichi I. Naito^{a,n},
6 Nanako O. Ogawa^a, Brian N. Poppⁱ, Shawn Steffan^{j,k}, Yoshinori Takano^a, Ichiro Tayasu^l, Alex S.J.
7 Wyatt^h, Yasuhiko T. Yamaguchi^{h,o}, Yusuke Yokoyama^h

8
9 ^aDepartment of Biogeochemistry, Japan Agency for Marine-Earth Science and Technology, Japan

10 ^bRosenstiel School of Marine and Atmospheric Science, University of Miami, USA

11 ^cAustralian Rivers Institute, Griffith University, Australia

12 ^dLeibniz-Laboratory, University of Kiel, Germany

13 ^eHarvard University Center for the Environment, USA

14 ^fDepartment of Ocean Sciences, University of California, Santa Cruz, USA

15 ^gGraduate School of Oceanography, University of Rhode Island, USA

16 ^hAtmosphere and ocean Research Institute, The University of Tokyo, Japan

17 ⁱDepartment of Geology and Geophysics, University of Hawaii, USA

18 ^jUS Department of Agriculture, Agricultural Research Service, USA

19 ^kDepartment of Entomology, University of Wisconsin-Madison, USA

20 ^lResearch Institute of Humanity and Nature, Japan

21 ^mInstitute of Low Temperature Science, Hokkaido University, Japan

22 ⁿNagoya University Museum, Japan

23 ^oLake Biwa Environmental Research Institute, Japan

24
25 *Corresponding author

26 Naohiko Ohkouchi

27 Department of Biogeochemistry

28 Japan Agency for Marine-Earth Science and Technology

29 2-15 Natsushima-cho

30 Yokosuka 237-0061

31 Japan

32 Phone: +81-46-867-9790

33 Fax: +81-46-867-9775

34 Email: nohkouchi@jamstec.go.jp

35
36 *Organic Geochemistry*, 113, 150-174, DOI: 10.1016/j.orggeochem.2017.07.009

38 **Abstract**

39 Compound-specific isotopic analysis of amino acids (CSIA-AA) has emerged in the last decade as a
40 powerful approach for tracing the origins and fate of nitrogen in ecological and biogeochemical
41 studies. This approach is based on the empirical knowledge that source AAs (*i.e.*, phenylalanine),
42 fractionate ^{15}N very little ($<0.5\%$) during trophic transfer, whereas trophic AAs (*i.e.*, glutamic acid),
43 are greatly ($\sim 6\text{-}8\%$) enriched in ^{15}N during each trophic step. The differential fractionation of these
44 two AA groups can provide a valuable estimate of consumer trophic position that is internally
45 indexed to the baseline $\delta^{15}\text{N}$ value of the integrated food web. In this paper, we critically review the
46 analytical methods for determining the nitrogen isotopic composition of AAs by gas
47 chromatography/isotope-ratio mass spectrometry. We also discuss methodological considerations for
48 accurate trophic position assessment of organisms using CSIA-AA. We then discuss the advantages
49 and challenges of the CSIA-AA approach by examining published studies including trophic position
50 assessment in various ecosystems, reconstruction of ancient human diets, reconstruction of animal
51 migration and environmental variability, and assessment of marine organic matter dynamics. It is
52 clear that the CSIA-AA approach can provide unique insight into the sources, cycling, and trophic
53 modification of organic nitrogen as it flows through systems. However, some uncertainty still exists
54 in how biochemical, physiological, and ecological mechanisms affect isotopic fractionation of
55 trophic AAs. We end this review with a call for continued exploration of the mechanisms of AA
56 isotopic fractionation, through various studies to promote the evolution of the rapidly growing field
57 of CSIA-AA.

58

59 **Abbreviations**

60 AA: amino acid, EAA: essential amino acid, SAA: source amino acids, TAA: trophic amino acids,
61 Ala: alanine, Arg: arginine, Asn: asparagine, Asp: aspartic acid, Cys: cysteine, His: histidine, Glu:
62 glutamic acid, Gly: glycine, Ile: isoleucine, Leu: leucine, Lys: lysine, Met: methionine, Phe:
63 phenylalanine, Pro: proline, Ser: serine, Thr: threonine, Trp: tryptophan, Val: valine, CSIA:
64 compound-specific isotope analysis, TFA: trifluoroacetic acid, TFAA: trifluoroacetic acid
65 anhydride, Pv: pivaloyl, MOC: methoxycarbonyl, iPr: isopropyl, GC/IRMS: gas
66 chromatography/isotope-ratio mass spectrometry, HPLC: high-performance liquid chromatography,
67 TP: trophic position, TDF: trophic discrimination factor, OM: organic matter, POM: particulate
68 organic matter, DOM: dissolved organic matter, THAA: total hydrolysable amino acid

69

70 **Key words**

71 amino acid, nitrogen isotopic composition, trophic discrimination factor, trophic position, ecology,
72 biogeochemistry

73 **1. Introduction**

74

75 The stable nitrogen isotopic composition of organisms was first applied in the field of biogeoscience
76 more than half a century ago (*e.g.*, [Parwel et al., 1957](#); [Hoering and Ford, 1960](#); [Cheng et al., 1964](#)).
77 [Miyake and Wada \(1967\)](#) first reported that marine animals preferentially incorporate ^{15}N relative to
78 ^{14}N during metabolic processing of dietary nitrogen. These initial findings were later confirmed in
79 several seminal papers based on diet-controlled laboratory culture experiments and field studies that
80 provided further evidence of ^{15}N enrichment during heterotrophic processes (*e.g.*, [DeNiro and](#)
81 [Epstein 1981](#); [Minagawa and Wada 1984](#); [Fry, 2006 and references therein](#)). The stable nitrogen
82 isotopic composition provides unique insight into the dietary habits of animals, as well as
83 biogeochemical cycling of nitrogen because ^{15}N enrichment during trophic transfer integrates a
84 number of biochemical processes accompanying isotopic fractionation during nitrogen metabolism.
85 The nitrogen isotopic composition of organisms provides a unique approach for describing the
86 dietary habits of animals, a macroscale ecological phenomenon. Beyond ecological studies, this
87 approach has been widely applied to biogeochemical studies investigating the fate of nitrogen in
88 oceanographic, terrestrial and freshwater systems (*e.g.*, [Cline and Kaplan, 1975](#); [Wada et al., 1975](#);
89 [Wada, 1980](#); [Altabet and Francois, 1994](#)).

90 These early stable nitrogen isotope studies were based on bulk isotope analysis, which
91 integrates across all nitrogen containing entities in a sample. While certainly informative for many
92 applications, interpretation of bulk $\delta^{15}\text{N}$ data can be challenging as multiple independent factors
93 including baseline isotope values, trophic transfer, and microbial degradation, all can influence bulk
94 $\delta^{15}\text{N}$ values. Compound-specific isotopic analysis of amino acids (CSIA-AA) has emerged as a
95 powerful approach in many ecological and biogeochemical applications (*e.g.*, [Gaebler et al., 1963,](#)
96 [1966](#); [Macko and Estep, 1984](#); [Macko et al., 1986, 1987](#)), because the differential fractionation of
97 individual amino acids can disentangle the relative influences of baseline and trophic variability on
98 consumer $\delta^{15}\text{N}$ values. The nitrogen in an organism is predominantly contained in proteins, which
99 are long chains of amino acids (AAs) linked by peptide bonds. Consequently, the CSIA-AA
100 approach is based on the fact that the nitrogen isotopic composition of individual AAs in organic
101 matter reflects isotopic fractionation associated with various biochemical reactions of different
102 individual AA involved in nitrogen metabolism. An organism's $\delta^{15}\text{N}$ value also inherently reflects
103 the isotopic composition of inorganic nitrogen sources (*e.g.*, nitrate, nitrite, ammonia, and urea)
104 assimilated by primary producers at the base of the food web. With appropriate calibrations,
105 CSIA-AA can therefore provide uniquely specific information about multiple aspects of nitrogen
106 metabolism in organisms and ecosystem properties. CSIA-AA now has a broad range of applications,
107 including the trophic position assessment of a broad range of consumers in aquatic (*e.g.*, [McClelland](#)
108 [and Montoya, 2002](#); [Chikaraishi et al., 2009; 2014](#); [Hannides et al., 2009, 2013](#); [Bradley et al., 2014](#);
109 [Gutiérrez-Rodríguez et al. 2014](#)) and terrestrial ecosystems ([Chikaraishi et al., 2010, 2014](#); [Steffan et](#)

110 al. 2013), the identification of baseline isoscapes (the spatial pattern in isotopic signatures, Bowen,
111 2010) of nitrogen in marine systems, the assessment of the source and transformation of dissolved
112 and detrital organic matter in marine waters and sediments (e.g., Lorrain et al., 2009; McCarthy et al.,
113 2007; Calleja et al., 2013; Hannides et al., 2013; Sherwood et al. 2014; Batista et al., 2014;
114 Vokhshoori et al., 2014), tracing of animal migration (e.g., Dale et al., 2011; Madigan et al., 2014,
115 2016), and the reconstruction of food resource consumption by ancient humans (e.g., Hare et al.,
116 1991; Fogel et al., 1997; Naito et al., 2013a; Styling et al., 2010). While these studies clearly
117 demonstrated the potential of the CSIA-AA approach, they have also opened up many new questions
118 that suggest a wide range of potential future applications, as well as areas that need further research
119 to improve the interpretation of CSIA-AA data. Future work to address these case-specific problems
120 and the associated overarching challenges will push the evolution of this rapidly growing field and
121 improve CSIA-AA applications across a variety of scientific disciplines.

122 This paper reviews the most recent information about CSIA-AA analytical methods and
123 their applications to ecology, biogeochemistry, and related fields. It is an outcome of the workshop
124 “Technical Issues Integrating Advanced Isotope Analyses into Ecological Studies” organized in
125 association with the 10th International Conference on the Applications of Stable Isotope Techniques
126 to Ecological Studies (IsoEcol 10) held in Tokyo in April 2016. At the workshop, investigators with
127 widely different expertise discussed a broad range of issues related to the CSIA-AA methods and
128 reached the conclusion that it is now time to review both the analytical methods, as well as
129 underlying theoretical grounding of CSIA-AA applications, as a guide for future research. The
130 review covers many broad issues, but emphasis is placed on nitrogen isotopic composition of AAs
131 where greatest consensus has been reached. We also discuss how carbon isotopic composition of
132 AAs may also provide unique insights in ecological and biogeochemical studies and can be a
133 complementary approach to nitrogen CSIA-AA. The paper first explores analytical methodologies
134 and related issues (Sections 2 and 3), then follows with applications and case studies in various
135 fields (Section 4), before concluding with remarks addressing future perspectives and directions
136 (Section 5).

137

138 **2. Analytical Considerations**

139

140 **2.1. Amino acid extraction and separation**

141 AAs in sample material, such as an organism’s tissue (e.g., muscle), are extracted by a
142 simple hydrolysis procedure that breaks the peptide bonds of the constituent proteins. The hydrolysis
143 is generally conducted with 6 to 12 M HCl at 100° to 150°C for 1 h to 1 day. The AAs are
144 hydrophilic because of their short carbon skeletons and zwitterionic functional groups, including
145 -COOH, -NH₂, -SH, -OH, and imino groups (-NH-). Hydrophobic molecules produced by acid

146 hydrolysis (*e.g.*, lipids) should be eliminated, for example, with organic solvents by liquid/liquid
147 extraction prior to derivatization procedures.

148 In biological and most geochemical samples, AAs mostly exist as a “bound” form (*e.g.*,
149 protein and peptide), with “free” AAs being a minor fraction. Some biological samples, such as
150 calcareous and siliceous fossils, aggregated microbial samples, soils, sediments, and some biological
151 tissue, contain large amounts of interfering materials. In such samples, solid phase extraction is
152 required before derivatization. Cation-exchange chromatography is an effective method of removing
153 interfering materials from the extracts with sufficient recovery (*e.g.*, Dowex WX-8, 200-400 mesh,
154 [Metges and Petzke, 1997](#); Biorad AG50 W-X8, 200-400 mesh, [Hare et al., 1991](#); [Takano et al.,](#)
155 [2010](#)). Alternatively, target AAs can be separated by high-performance liquid chromatography
156 (HPLC) equipped with the fraction collector ([Broek et al., 2013](#); [Takano et al., 2015](#); [Bour et al.,](#)
157 [2016](#)). Significant nitrogen isotopic fractionation or exchange may occur with some types of column
158 resin ([Macko et al., 1987](#); [Hare et al., 1991](#); [Styring et al., 2012](#)) and therefore use of such a column
159 resin (*e.g.*, C18) should be avoided unless the isotopic fractionation is carefully evaluated. Finally,
160 for extremely complex geochemical sample matrixes, upstream HPLC isolation before derivatization
161 ([Broek et al., 2013](#)) can be required to purify AA sufficiently for accurate CSIA-AA.

162

163 **2.2. AA derivatization for precise determination of nitrogen isotopic composition**

164 AAs require derivatization to reduce polarity and increase their volatility in order to be
165 analyzed by GC/IRMS. The derivatization neutralizes polar carboxyl (-COOH), amino (-NH₂), and
166 hydroxyl (-OH) groups in AAs by replacing active hydrogen atoms with nonpolar moieties, resulting
167 in significant improvement in their chromatographic separation. Esterification of carboxyl groups
168 with an alcohol under acidic conditions and subsequent acylation of the amino group (and
169 simultaneous acetylation of hydroxyl group if AAs have a hydroxyl group) with an acid anhydride or
170 acid chloride, is a common chemical reaction for the derivatization (Fig. 1a).

171 Although a variety of reagents have been used over the last two decades, to our
172 knowledge, the following three derivatization reagents are most widely used in ecological and
173 geochemical studies: trifluoroacetyl-isopropyl ester (TFA/AA/iPr, Fig. 1b, *e.g.*, [McCarthy et al., 2007](#);
174 [Popp et al., 2007](#)), pivaloyl-isopropyl ester (Pv/AA/iPr, Fig. 1c, *e.g.*, [Metges et al., 1996](#); [Chikaraishi](#)
175 [et al., 2007](#)), and methoxycarbonyl (MOC) AA ester (Fig. 1d, *e.g.*, [Walsh et al., 2014](#); [Yarnes and](#)
176 [Herszage, 2017](#)). The first step of the derivatizations to TFA/AA/iPr and Pv/AA/iPr is the same
177 esterification with isopropanol to form the isopropyl esters of AAs. A major advantage of the use of
178 branched alcohol (*i.e.*, isopropanol) is that stable AA esters are obtained. The second step in the
179 TFA/AA/iPr and Pv/AA/iPr derivatizations is acylation with trifluoroacetic acid anhydride (TFAA)
180 or pivaloyl chloride (Pv-Cl), respectively. Because three atoms of fluorine, which is highly
181 electrophilic, increase the nucleophilicity of the carboxyl carbon of TFAA, acylation with TFAA is

182 much faster than that with Pv-Cl. MOC AA ester requires a rapid one-step derivatization, which
183 allows esterification of the carboxyl group and acylation of the amino group simultaneously at room
184 temperature within 5 min, although the hydroxyl group is not acetylated in this derivatization. The
185 TFA/AA/iPr and Pv/AA/iPr require strict hydrophobic conditions, whereas MOC AA ester works
186 well in both hydrophobic and hydrophilic conditions. Detailed derivatization procedures using each
187 reagent are described in the literature (*e.g.*, [Silfer et al., 1991](#); [Sacks and Brenna, 2005](#); [Chikaraishi
188 et al., 2007](#)).

189 For all derivatizations, great care should be taken with respect to the chemical properties
190 of the reagents and derivatives. First, because the ester groups in these derivatives are exchangeable
191 with water, no alcohols or other ester compounds, including many polar solvents, can be used. For
192 example, ethyl acetate, a convenient polar organic solvent, can exchange the isopropyl or methyl
193 ester group in the AA derivatives with its ethyl ester group (Fig. 2a). In general, suitable solvents for
194 the derivatives include ethers (*e.g.*, diethyl ether and tetrahydrofuran, although these solvents are
195 highly flammable) or chlorinated methanes (*e.g.*, dichloromethane and chloroform, although these
196 solvents are toxic). Second, most derivative reagents should be used in strict accordance with
197 exposure controls. In particular, Pv-Cl is acutely toxic. Third, because esterified AAs are unstable in
198 O₂ and water, even at 0°C, the derivatives must be stored at -20°C or lower (without O₂ and water, if
199 possible) until isotope analysis. Although TFA/AA/iPr and Pv/AA/iPr esters (*i.e.*, branched alcohol
200 esters) are relatively stable at low temperature (Fig. S1), they only survive for a few days to weeks at
201 room temperature. Finally, these derivatizations are not equally applicable to the isotopic
202 measurements of all 20 protein AAs. Arg, Asn, Cys, His, and Trp cannot be measured as
203 TFA/AA/iPr and Pv/AA/iPr derivatives because of degradation (including conversion to other
204 compounds) or less-quantitative reaction during derivatization. Although MOC AA esters can be
205 useful for the isotope measurement of most of these AAs (Asn, Cys, His, and Trp, except for Arg),
206 this derivatization is not appropriate for determining the isotope values of Glu, because two types of
207 Glu derivatives are produced with distinct isotopic compositions (Fig. S2).

208

209 **2.3. Nitrogen isotopic measurements of AAs**

210 In GC/IRMS, the nitrogen isotopic compositions of AAs are determined by analyzing the
211 ¹⁵N:¹⁴N ratios of N₂ molecules generated by combustion-reduction of the derivatives. The instrument
212 consists of a conventional gas chromatograph (GC) connected to a chemical reaction interface
213 including combustion and reduction furnaces ([Merritt and Hayes, 1994](#)). Individual AA derivatives
214 are separately eluted by GC, and combusted mainly into N₂, NO_x, CO₂, and H₂O in a combustion
215 furnace with CuO and NiO with Pt at 950°-1050°C. The NO_x generated by the combustion is
216 subsequently reduced to N₂ in a reduction furnace with Cu at 550°-650°C, and the H₂O and CO₂
217 generated during the combustion are eliminated using a liquid nitrogen trap. A countercurrent drier

218 can be used for H₂O elimination prior to the liquid nitrogen trap in some cases. To avoid isotopic
219 fractionation, a nucleophilic stationary phase (*e.g.*, HP-5: phenyl-methyl polysiloxane;
220 HP-INNOWAX: polyethylene glycols) is required for the GC separation of AA derivatives
221 ([Chikaraishi et al., 2010](#)).

222 The nitrogen isotopic composition of AAs is expressed in the standard δ notation relative
223 to atmospheric N₂ ($\delta^{15}\text{N}$, ‰ vs. AIR), which is calibrated to the internationally recognized scale
224 through comparison of the $\delta^{15}\text{N}$ values of multiple reference AAs. In a typical sequence, derivatives
225 of reference mixtures of 5-14 AAs with known $\delta^{15}\text{N}$ values, which should cover the $\delta^{15}\text{N}$ range of
226 the samples, are analyzed every 4-8 sample runs. At the beginning and end of each chromatography
227 run, 2-3 pulses of reference N₂ gas are discharged for all reference mixtures and samples (Fig. 3).
228 The regression line between the known (‰, vs. AIR) and mean measured values (‰, vs. reference
229 N₂ gas) represents the reproducibility of the isotope measurement (Fig. S3) and can be used to
230 normalize the measured values (‰, vs. reference N₂ gas) to the internationally recognized scale (‰,
231 vs. AIR) for both the reference mixtures and samples. In some laboratories norleucine and
232 aminoadipic acid with known $\delta^{15}\text{N}$ values are co-injected with each sample and additional internal
233 reference compounds that can be used for normalization (*e.g.*, [Hannides et al., 2009](#); [McCarthy et al.,](#)
234 [2013](#)). The average and standard deviation for the normalized values (1σ) and the difference in the
235 normalized and known values ($\Delta_{\text{normalized-known}}$) for the reference AAs are frequently used as evidence
236 of the precision and accuracy of the isotope measurement. Detection limits to achieve this level of
237 precision and accuracy depend on various factors, but they are highly correlated with the
238 signal/noise ratio of the GC/IRMS chromatogram (Fig. S4, [Chan et al., 2016](#)).

239 Baseline separation between the AA peaks on the GC/IRMS chromatogram is required to
240 obtain accurate $\delta^{15}\text{N}$ values of the AAs. When an AA peak is co-eluted with other AAs or impurities,
241 the isotopically heavy tail of the first peak underlies the isotopically light front of the second peak
242 ([Hayes et al., 1990](#)). For example, in case of Pv/AA/iPr derivatives, Glu and Phe generally show
243 good baseline separation, whereas Asp, Thr, Ser, and Met on the same chromatogram are
244 sequentially eluted without baseline separation (Fig. 3).

245 We should note that, in addition to the analysis by GC/IRMS, off-line process ([Broek et al.,](#)
246 [2013](#)) and HPLC-IRMS coupling may be useful in the future to determine nitrogen isotope ratios of
247 AAs ([Federherr et al., 2016](#)).

248

249 **3. Methodological considerations for trophic position assessment**

250

251 **3.1. Bulk versus CSIA-AA approach**

252 As noted above, stable nitrogen isotope analysis of bulk organisms and their tissues has
253 been used extensively for conventional estimation of the trophic positions of organisms in food

254 webs (e.g., Post, 2002; Fry 2006; Ohkouchi et al., 2015). The trophic position (TP_{bulk}) is generally
 255 calculated using equation 1, based on the empirical observation that the ^{15}N content of bulk
 256 organisms tends to increase with each trophic transfer in food webs (e.g., DeNiro and Epstein,
 257 1981; Minagawa and Wada, 1984).

258

$$259 \quad TP_{\text{bulk}} = (\delta^{15}\text{N}_{\text{sample}} - \delta^{15}\text{N}_{\text{pp}}) / TDF_{\text{bulk}} + 1 \quad (1)$$

260

261 where $\delta^{15}\text{N}_{\text{sample}}$ and $\delta^{15}\text{N}_{\text{pp}}$ are the $\delta^{15}\text{N}$ values of a target organism and the primary producers at the
 262 base of the food web, respectively. TDF_{bulk} is the trophic discrimination factor of $\delta^{15}\text{N}_{\text{bulk}}$ between
 263 prey and predator (DeNiro and Epstein, 1981; Minagawa and Wada, 1984). Many studies use a
 264 canonical TDF_{bulk} value of 3-4‰, however, a variety of TDF_{bulk} values are frequently used in studies
 265 focusing on specific tissues, such as collagen, or specific localized environments (e.g., Vander
 266 Zanden and Rasmussen, 2001; McCutchan et al., 2003; Martinez del Rio et al., 2009). The ‘bulk
 267 method’ has been successfully applied to various ecological studies and has thus helped expand our
 268 knowledge of feeding ecology greatly over the last four decades (Fry, 2006). However, the method
 269 suffers from several problems that can cause large uncertainty in the estimated TP_{bulk} values. The
 270 most important problem is that the $\delta^{15}\text{N}$ values of bulk tissues intrinsically reflect i) the trophic
 271 changes in the $\delta^{15}\text{N}$ value in the food web and ii) temporal or spatial changes in the $\delta^{15}\text{N}$ value at the
 272 base of the food web (Fig. 4a). The former (~3-4‰) is often much smaller than the latter (in some
 273 cases >10‰) (e.g., Hannides et al., 2009; Rolff, 2000; Dore et al., 2002; O’Reilly et al., 2002).

274 In contrast, trophic position ($TP_{\text{TAA/SAA}}$) estimated from CSIA-AA using equation 2 can
 275 constrain both trophic changes in the $\delta^{15}\text{N}$ value and baseline variation within a single organism (e.g.,
 276 McClelland and Montoya, 2002; Chikaraishi et al., 2007; McCarthy et al., 2007; Popp et al., 2007).

277

$$278 \quad TP_{\text{TAA/SAA}} = [(\delta^{15}\text{N}_{\text{TAA}} - \delta^{15}\text{N}_{\text{SAA}} + \beta_{\text{TAA/SAA}}) / \Delta_{\text{TAA/SAA}}] + 1$$

279 (2)

280

281 where $\delta^{15}\text{N}_{\text{TAA}}$ and $\delta^{15}\text{N}_{\text{SAA}}$ are the $\delta^{15}\text{N}$ values of the trophic and source AAs, respectively, from a
 282 single organism; $\beta_{\text{TAA/SAA}}$ is the isotopic difference between these AAs in primary producers at the
 283 base of the food web; and $\Delta_{\text{TAA/SAA}}$ is the difference in the TDF of the TAAs and SAAs during each
 284 trophic transfer ($\Delta_{\text{TAA/SAA}} = TDF_{\text{TAA}} - TDF_{\text{SAA}}$). Trophic amino acids (TAAs) (e.g., Ala, Asp, Glu,
 285 Ile, Leu, Pro, and Val) tend to show large ^{15}N enrichment (by ~3-8‰) relative to diet during trophic
 286 transfer, which likely reflects isotopic fractionation associated with deamination (a first step in
 287 transamination, Macko et al., 1986; Miura and Goto, 2012) as a dominant metabolic pathway for
 288 these AAs in consumers (Fig. 5a). Source amino acids (SAAs) (e.g., Met, Lys, and Phe) show little
 289 ^{15}N enrichment (~0-1‰) relative to diet during trophic transfer, which probably reflects the fact that

290 the initial steps in their metabolism are generally dominated by reactions that neither form nor cleave
 291 C-N bonds (Fig. 5a) and thus directly provide an estimate of the $\delta^{15}\text{N}_{\text{SAA}}$ value of the base of the
 292 food web. Therefore, *CSIA-AA derived TP values are independent of temporal or spatial changes in*
 293 *the $\delta^{15}\text{N}$ value at the base of the food web* (Fig. 4b).

294 [Chikaraishi et al. \(2009, 2010\)](#) first suggested the utility of Glu and Phe as a TAA and a
 295 SAA, respectively, with $\beta_{\text{Glu/Phe}}$ values of -3.4% for aquatic and $+8.4\%$ for terrestrial C3
 296 plant-based food webs, and with $\Delta_{\text{Glu/Phe}}$ values of 7.6% for both ecosystems. Later, it was found that
 297 the β value in vascular plants is increased by the deamination of Phe for lignin biosynthesis, a
 298 process specific to vascular plants (Fig. 5b; [Ohkouchi and Takano, 2014](#); [Naito et al., 2016a](#)).
 299 Therefore, algal vs. vascular grouping is a better classification than aquatic vs. terrestrial
 300 ([Chikaraishi et al., 2009, 2010](#)). Indeed, the observed β values in seagrasses (vascular plants from
 301 coastal marine environments) are similar to those of terrestrial vascular plants (e.g., [Vander Zanden](#)
 302 [et al., 2013](#); [Choi et al., 2017](#)). However, for simplified nomenclature we use the terms aquatic and
 303 terrestrial throughout this paper.

304

$$305 \quad [TP_{\text{Glu/Phe}}]_{\text{aqua}} = [(\delta^{15}\text{N}_{\text{Glu}} - \delta^{15}\text{N}_{\text{Phe}} - 3.4)/7.6] + 1 \quad (3)$$

$$306 \quad [TP_{\text{Glu/Phe}}]_{\text{terr}} = [(\delta^{15}\text{N}_{\text{Glu}} - \delta^{15}\text{N}_{\text{Phe}} + 8.4)/7.6] + 1 \quad (4)$$

307

308 Because of the large differences in $\beta_{\text{TAA/SAA}}$ values between aquatic and terrestrial producers, mixing
 309 models must be constructed so as to consider two potential food webs where both aquatic and
 310 terrestrial primary producers may serve as basal food resources. These environments include rivers
 311 ([Ishikawa et al., 2014](#)) and coastal marine ecosystems ([Vander Zanden et al., 2013](#); [Choi et al., 2017](#)).
 312 In this paper, and many others, there has been a focus on glutamic acid and phenylalanine as the
 313 canonical trophic and source amino acids, However, in principle, any combination of trophic and
 314 source amino acids can be used in equation 2 (e.g., [Decima et al., 2013](#); [Nielsen et al., 2015](#); [Bradley](#)
 315 [et al., 2015](#)) as long as $\beta_{\text{TAA/SAA}}$ and $\Delta_{\text{TAA/SAA}}$ values appropriate for the combination of trophic and
 316 source amino acids are used.

317

318 **3.2. Uncertainties and errors in the TP assessment**

319

320 **3.2.1. Variability in trophic discrimination factors**

321 Constant $\Delta_{\text{Glu/Phe}}$ (= $TDF_{\text{Glu}} - TDF_{\text{Phe}}$) or $\Delta_{\text{TAA/SAA}}$ values throughout the food web is
 322 prerequisite for estimating TP precisely. However, the stability of $\Delta_{\text{Glu/Phe}}$ has recently come under
 323 increasing scrutiny based on new laboratory and field studies (e.g., [Dale et al., 2011](#); [Matthews and](#)
 324 [Ferguson, 2014](#); [Chikaraishi et al., 2015](#); [McMahon et al., 2015a](#)). Comprehensive meta-analyses
 325 of CSIA-AA from wild animals with known TP values ([Nielsen et al., 2015](#); [Bradley et al., 2015](#))

326 and controlled feeding experiments (McMahon and McCarthy, 2016) that examine individual
327 TDF_{AA} values have addressed the following primary questions: i) what are the magnitude and
328 variability in $\Delta_{TAA/SAA}$ values across a wide range of consumer-resource relationships, and ii) are
329 there systematic underlying mechanisms driving this variability in predictable ways that could be
330 used to improve CSIA-AA-based estimates of consumer trophic dynamics.

331 These meta-analyses found large variability in $\Delta_{TAA/SAA}$ values. For example, McMahon
332 and McCarthy (2016) found the overall mean $\Delta_{Glu/Phe}$ value was $6.2 \pm 2.5\%$ across a wide range of
333 taxa, diet types, and modes of nitrogen excretion, consistent with other recent large scale analyses
334 of field-collected data for wild-caught marine consumers ($6.6 \pm 1.7\%$; Nielsen et al., 2015; $5.7 \pm$
335 0.3% ; Bradley et al. 2015). However, within this distribution there were also some very significant
336 excursions, with $\Delta_{Glu/Phe}$ values from 0 to $>10\%$ across 70 species (317 individuals) and 88 distinct
337 species-diet combinations. Some of the reported $\Delta_{Glu/Phe}$ values, particularly for animals with TP
338 values of less than 3, were within a small range (6 to 8%) that overlapped with the original $\Delta_{Glu/Phe}$
339 values of 7.0% (McClelland and Montoya, 2002) and 7.6% (Chikaraishi et al., 2007). However,
340 simply focusing on the mean can inherently obscure large variation underlying that mean. The
341 meta-analysis of controlled feeding studies by McMahon and McCarthy (2016) is also consistent
342 with large scale studies of wild consumers by Nielsen et al. (2015) and Bradley et al. (2015), which
343 together strongly suggest that the observed variability in $\Delta_{Glu/Phe}$ and $\Delta_{TAA/SAA}$ values is not simply
344 noise, but rather is predictably linked to consumer biochemistry. Below, we discuss two possible
345 underlying biochemical and physiological processes that influence $\Delta_{TAA/SAA}$: diet quality and
346 metabolic flux (e.g., mode of nitrogen excretion).

347

348 Diet quality: In the aquatic environment, there is a trend between TP and $\Delta_{Glu/Phe}$ across a wide
349 range of (although not all) consumers (Bradley et al., 2015; Nielsen et al. 2015; McMahon and
350 McCarthy, 2016). However, this trend was not observed in insects kept in ecologically realistic
351 pure cultures, representing three distinct communities from the terrestrial environment (Steffan et
352 al., 2013). Further, low variability in the TDF was observed among 15 consumer species,
353 representing a phylogenetically diverse group of consumers, from freshwater crustaceans and fish,
354 to terrestrial mammals, fungi, and bacteria (Steffan et al., 2015) Most primary consumers
355 examined in the marine environment (e.g., grazing teleost fishes, zooplankton, etc.) had $\Delta_{Glu/Phe}$
356 values between 6% and 8% , often not substantially different from the value of $\sim 7-8\%$ originally
357 reported by McClelland and Montoya (2002), and substantiated by Chikaraishi et al. (2007). In
358 contrast, most marine consumers with TP higher than 3 showed lower $\Delta_{Glu/Phe}$ values (Bradley et al.,
359 2015; Nielsen et al. 2015; McMahon and McCarthy, 2016). One hypothesis for the pattern of
360 decreasing $\Delta_{Glu/Phe}$ value with increasing TP is the effect of diet quality (defined here as the relative
361 AA composition of a food source relative to the needs of a consumer) on consumer $\delta^{15}N_{AA}$ values,

362 and thus on $\Delta_{\text{Glu/Phe}}$ values.

363 The diet quality hypothesis suggests that nitrogen isotope discrimination decreases as
364 dietary protein quality (degree of AA similarity between diet and consumer) increases (Hobson and
365 Clark, 1992; Roth and Hobson, 2000; Robbins et al., 2005, 2010; Mill et al., 2007; Florin et al.,
366 2011). McMahon et al. (2015a) showed that diet quality had a large and systematic effect on the
367 isotopic fractionation of individual AAs in an estuarine fish (*Fundulus heteroclitus*) fed
368 compositionally distinct diets. This study found a strong relationship between the TDF value of
369 most TAAs and protein quality between diet and consumer, and no change in TDF_{Phe} across diet
370 types. Furthermore, Chikaraishi et al. (2015) recently showed that with extreme manipulation of
371 dietary composition (*i.e.*, the relative composition of protein/fat/carbohydrates), vastly different
372 $\Delta_{\text{Glu/Phe}}$ values can be obtained in a single consumer. However, these two studies found opposite
373 trends in the $\Delta_{\text{Glu/Phe}}$ vs. diet quality relationship, defined as relative AA composition of a food
374 source relative to the needs of a consumer. McMahon et al. (2015a), showed that as the diet AA
375 composition converged on that of the consumers, the $\Delta_{\text{Glu/Phe}}$ values tended to decrease. In contrast,
376 Chikaraishi et al. (2015) indicated that the $\Delta_{\text{Glu/Phe}}$ values decreased as diet quality declined. While
377 both of these studies indicate that diet composition strongly affects individual AA isotopic
378 fractionation, more work is necessary to resolve the full relationship between diet quality and
379 $\Delta_{\text{Glu/Phe}}$ value.

380 The reason why $\Delta_{\text{Glu/Phe}}$ often varies with TP might reflect differences in diet quality
381 across different consumer-resource relationships within a food web. Generally, lower TP
382 consumers often feed on diets that are more compositionally distinct relative to their own tissues
383 (*e.g.*, zooplankton feeding on phytoplankton) than higher TP consumers (*e.g.*, fish feeding on other
384 fish). When feeding on low-quality diets, defined as having highly imbalanced AA composition
385 compared with consumer requirements, the consumer synthesizes scarce AAs *de novo* from surplus
386 AAs. Because TAAs enriched in ^{15}N relative to SAAs tend to be abundant in the organisms,
387 synthesis leads to the apparent increase in $\delta^{15}\text{N}_{\text{AA}}$ (Krueger and Sullivan, 1984; Roth and Hobson,
388 2000; Clements et al., 2009). Conversely, carnivores feeding on high-quality diets can meet more
389 of their AA requirements *via* direct isotopic routing of dietary AAs, which should reduce ^{15}N
390 enrichment of heavily transaminating AAs (*e.g.*, Glu) compared with consumers feeding on
391 low-quality diets (Schwarcz, 1991; Ambrose and Norr, 1993). It should be noted that Ishikawa et al.
392 (2017) recently showed that satiated and starved dobsonfly (*Protohermes grandis*) larvae had
393 similar $\Delta_{\text{Glu/Phe}}$ values (7.1‰ and 7.3‰, respectively), suggesting that the $\Delta_{\text{Glu/Phe}}$ value was
394 independent from starvation.

395

396 Mode of nitrogen excretion: There is also a clear pattern of lower $\Delta_{\text{Glu/Phe}}$ values for some urea/uric
397 acid-producing organisms relative to ammonia-producing organisms, largely driven by differences

398 in TDF_{Glu} but not TDF_{Phe} (Dale et al., 2011; Germain et al., 2013; Nielsen et al., 2015; McMahon
399 and McCarthy, 2016). The typically low $\Delta_{\text{Glu/Phe}}$ values for urea/uric acid producers may be
400 explained by the nitrogen storage and cycling capabilities of animals (Wilkie, 2002), or by the way
401 urea is produced in the liver (Dale et al., 2011). Key nitrogen-transferring enzymes preferentially
402 remove ^{14}N -amines during metabolism, resulting in the subsequent ^{15}N enrichment of residual
403 animal tissue and the excretion of ^{15}N -depleted nitrogenous waste (DeNiro and Epstein, 1981). The
404 final isotope value of a biochemical reaction depends not only on the number of steps and
405 associated ϵ values (*i.e.*, the maximal potential isotopic fractionation), but also on the relative
406 nitrogen fluxes through branch points in the reaction chain (*e.g.*, reviewed by Hayes et al., 2001;
407 Koch et al., 2007). Germain et al. (2013) proposed that this concept of variable nitrogen flux
408 through additional branch points in the ornithine-to-urea pathway probably underlies the offset in
409 $\Delta_{\text{Glu/Phe}}$ values for urea vs. ammonia-excreting organisms. In elasmobranchs, which have reduced
410 hepatic glutamate catabolism relative to ureotelic organisms, a lower ϵ value may be related to
411 their unique glutamate-glutamine-urea pathway (Dale et al., 2011). In addition, the recycling of
412 ^{15}N -depleted urea by gut microbes for subsequent AA synthesis is another possible explanation for
413 low $\Delta_{\text{Glu/Phe}}$ values in urea/uric acid-producing consumers (Davidson et al., 2003; Fouillet et al.,
414 2008).

415
416 Summary: Independent meta-analyses of controlled feeding studies (McMahon and McCarthy,
417 2016) and wild consumers (Bradley et al. 2015; Nielsen et al. (2015) have shown that both diet
418 quality and metabolic flux (*e.g.*, mode of nitrogen excretion) affect $\Delta_{\text{Glu/Phe}}$ values considerably.
419 These processes are not mutually exclusive, and both appear to impact TDF_{Glu} by affecting the flux
420 of nitrogen through transamination and deamination isotopic branch points. There are many
421 systems that appear to be well characterized by a single $\Delta_{\text{Glu/Phe}}$ value, where there are minimal
422 changes in diet quality and/or mode of nitrogen excretion within a food web (*e.g.*, Chikaraishi et al.,
423 2009, 2011; Ishikawa et al., 2014; Kruse et al., 2015; Miyachi et al., 2015). However, the accuracy
424 of $TP_{\text{Glu/Phe}}$ estimates may be improved by directly incorporating $\Delta_{\text{Glu/Phe}}$ variability into $TP_{\text{Glu/Phe}}$
425 estimates in systems where such changes do occur (*e.g.*, Lorrain et al., 2009; Dale et al., 2011;
426 Choy et al., 2012; Germain et al., 2013; Ruiz-Cooley et al., 2013, 2014; Matthews and Ferguson,
427 2014; McMahon et al., 2015b). This probably requires moving toward multi- Δ equations (*e.g.*,
428 Hoen et al., 2014), potentially averaging across multiple AAs (*e.g.*, Decima et al., 2013; Nielsen et
429 al., 2015; Bradley et al., 2014), although averaging across multiple AAs has been shown to
430 profoundly increase variability surrounding the TDF in terrestrial and freshwater systems (*e.g.*,
431 Table 1 in Steffan et al., 2015). While accounting for key transitions in diet quality and mode of
432 nitrogen excretion with multi- Δ equations improves TP estimates in many cases (*e.g.*, McMahon et
433 al., 2015a,b), diet quality and metabolic flux are likely not the only drivers of variability in

434 $\Delta_{TAA/SAA}$ values. Continued exploration of the underlying mechanisms controlling AA $\delta^{15}\text{N}$
 435 fractionation is critical to improve our ability to accurately estimate consumer TP with the
 436 CSIA-AA approach.

437

438 3.2.2. Propagation of error calculations for trophic position determination

439 For both ecological and geochemical / paleoceanographic applications, interpreting
 440 CSIA-AA based TP data requires a rigorous estimation of uncertainty in values being compared.
 441 However, uncertainty in TP based on nitrogen isotopic composition of AAs is more complex than
 442 standard uncertainties in measured isotopic values, because it must take into account analytical
 443 uncertainty in source and trophic AA isotopic measurements, as well as environmental uncertainty
 444 in β and Δ values. The combination of these uncertainties can be calculated using propagation of
 445 errors. The variability in the parameters used for TP determination can be modeled using Monte
 446 Carlo simulations, however it is also straightforward to propagate errors using a first-order Taylor
 447 series expansion (Ku, 1966), resulting in a formula easily solved in a spreadsheet or programmed
 448 into an algorithmic language (e.g., Matlab, R).

449 In general, for any result w that is a function of two or more experimentally determined
 450 independent variables, variance in w can be calculated by Taylor series expansion if the variance in
 451 the variables is known (e.g., Gelwicks and Hayes, 1990; Phillips and Gregg, 2001). In the case
 452 where $w = f(x, y, z)$, variance in w can be determined using the analytical solution of

453

$$454 \sigma_w^2 = (\partial w / \partial x)^2 \sigma_x^2 + (\partial w / \partial y)^2 \sigma_y^2 + (\partial w / \partial z)^2 \sigma_z^2 \quad (5)$$

455

456 The measured values of $\delta^{15}\text{N}_{TAA}$ and $\delta^{15}\text{N}_{SAA}$ have inherent analytical uncertainty and
 457 there is uncertainty in the values of β and Δ compiled in the literature. If we assume a general
 458 formulation of the equation used for calculation of TP as equation 2, uncertainty in TP can be
 459 determined by propagation of errors (e.g., Blum et al., 2013; Bradley et al., 2015) using the
 460 analytical solution of

461

$$462 \sigma_{TP}^2 = (\partial TP / \partial \delta^{15}\text{N}_{TAA})^2 \sigma_{\delta^{15}\text{N}(TAA)}^2 + (\partial TP / \partial \delta^{15}\text{N}_{SAA})^2 \sigma_{\delta^{15}\text{N}(SAA)}^2 \\ 463 + (\partial TP / \partial \beta_{TAA/SAA})^2 \sigma_{\beta(TAA/SAA)}^2 + (\partial TP / \partial \Delta_{TAA/SAA})^2 \sigma_{\Delta(TAA/SAA)}^2 \quad (6)$$

464

465 The exact solution to equation 6 has been published elsewhere (Bradley et al., 2015) and an
 466 equation for calculating the propagated variance in TP is summarized in equation 7.

467

$$468 \sigma_{TP}^2 = (1/\Delta_{TAA/SAA})^2 \sigma_{\delta^{15}\text{N}(TAA)}^2 + (-1/\Delta_{TAA/SAA})^2 \sigma_{\delta^{15}\text{N}(SAA)}^2 + (1/\Delta_{TAA/SAA})^2 \sigma_{\beta(TAA/SAA)}^2 \\ 469 + \{-1/\Delta_{TAA/SAA}^2 (\delta^{15}\text{N}_{TAA} - \delta^{15}\text{N}_{SAA} + \beta)\}^2 \sigma_{\Delta(TAA/SAA)}^2 \quad (7)$$

470

471 The analytical uncertainty in isotopic measurements of trophic and source AAs in samples must be
472 determined. Because the AA distribution in samples is more complex than that of artificial
473 mixtures of AAs, we suggest replicate analysis of each sample following the recommendations of
474 [Hayes et al. \(1990\)](#).

475 It has been suggested ([Hoen et al., 2014](#)) that the TP of a carnivore might best be
476 determined using separate Δ -values for herbivores and carnivores:

477

$$478 \quad TP = \{(\delta^{15}\text{N}_{\text{TAA}} - \delta^{15}\text{N}_{\text{SAA}} + \beta_{\text{TAA/SAA}} - \Delta_{\text{herbivore}}) / \Delta_{\text{carnivore}}\} + 2 \quad (8)$$

479

480 where $\Delta_{\text{herbivore}}$ is the ^{15}N enrichment in a TAAs relative to a SAA of a grazing herbivore and
481 $\Delta_{\text{carnivore}}$ is the ^{15}N enrichment in a TAAs relative to a SAA associated with each trophic transfer
482 for an omnivore or carnivore ([Hoen et al., 2014](#)). An expression for the variance in TP based on
483 two different Δ values is

484

$$485 \quad \sigma_{TP}^2 = (1/\Delta_{\text{carnivore}})^2 \sigma_{\delta^{15}\text{N}(\text{TAA})}^2 + (-1/\Delta_{\text{carnivore}})^2 \sigma_{\delta^{15}\text{N}(\text{SAA})}^2 + (1/\Delta_{\text{carnivore}})^2 \sigma_{\beta(\text{TAA/SAA})}^2 \\ 486 \quad + (-1/\Delta_{\text{carnivore}})^2 \sigma_{\Delta_{\text{carnivore}}}^2 + \{-1/\Delta_{\text{carnivore}} (\delta^{15}\text{N}_{\text{TAA}} - \delta^{15}\text{N}_{\text{SAA}} + \beta - \Delta_{\text{herbivore}})\}^2 \sigma_{\text{herbivore}}^2 \\ 487 \quad (9)$$

488

489 TP for animals feeding in aquatic and terrestrial environments can be calculated using the
490 nitrogen isotopic composition of AAs if the fraction of one of the binary components is
491 independently determined (*e.g.*, [Jarman et al., 2017](#)):

492

$$493 \quad TP = \{(\delta^{15}\text{N}_{\text{TAA}} - \delta^{15}\text{N}_{\text{SAA}} + f_1\beta_1 + (1-f_1)\beta_2) / \Delta_{\text{TAA/SAA}}\} + 1 \quad (10)$$

494

495 where β_1 and β_2 are the C_3 , C_4 , or aquatic plant ^{15}N enrichment in the same trophic and source AAs
496 measured in the sample, and f_1 is the fractional contribution of one of those plant types. The
497 propagated variance in TP when there is a binary mixture of feeding is given by

498

$$499 \quad \sigma_{TP}^2 = (1/\Delta_{\text{TAA/SAA}})^2 \sigma_{\delta^{15}\text{N}(\text{TAA})}^2 + (-1/\Delta_{\text{TAA/SAA}})^2 \sigma_{\delta^{15}\text{N}(\text{SAA})}^2 + \{(\beta_1 - \beta_2) / \Delta_{\text{TAA/SAA}}\}^2 \sigma_{f_1}^2 \\ 500 \quad + \{(1-f_1) / \Delta_{\text{TAA/SAA}}\}^2 \sigma_{\beta_2}^2 + \{f_1 / \Delta_{\text{TAA/SAA}}\}^2 \sigma_{\beta_1}^2 \\ 501 \quad + \{-1/\Delta_{\text{TAA/SAA}} (\delta^{15}\text{N}_{\text{TAA}} - \delta^{15}\text{N}_{\text{SAA}} + (1-f_1)\beta_2 + f_1\beta_1)\}^2 \sigma_{\Delta(\text{TAA/SAA})}^2 \\ 502 \quad (11)$$

503

504 Propagated uncertainty in f_1 and f_2 must be input into equation 11 and can be determined using
505 [Phillips and Gregg \(2001\)](#) or a similar approach.

506

507 **4. Applications of the CSIA-AA approach to ecological and biogeochemical studies**

508

509 **4.1. Food web analyses in aquatic ecosystems**

510 Several recent studies have used $\delta^{15}\text{N}_{\text{SAA}}$ variation to understand the baseline of food
511 webs in the North Pacific Subtropical Gyre ecosystem. [Hannides et al. \(2013\)](#) used differences in
512 $\delta^{15}\text{N}_{\text{SAA}}$ between zooplankton and suspended particles to demonstrate that deep water zooplankton
513 in the subtropical gyre probably depend on surface rather than *in situ* particulate food, either
514 through sinking of surface particles or vertical migrations. [Choy et al. \(2015\)](#) further showed that
515 surface productivity also fuels higher-order consumers in the North Pacific Subtropical Gyre food
516 web. A large range of $\delta^{15}\text{N}_{\text{Phe}}$ values in the tissues of both large and small pelagic micronekton
517 suggested that some components of the food web instead are fueled by slowly settling particles that
518 are highly modified by microbes. In contrast, there was no relationship between depth and $\delta^{15}\text{N}_{\text{SAA}}$
519 for large predatory fish ([Choy et al., 2015](#)), demonstrating how CSIA-AA may also be used to infer
520 species movement and foraging across large depth gradients in oceanic ecosystems, and thus the
521 dependence on a range of nutrient sources.

522 The *TP* of consumers can be estimated based only on consumer tissue $\delta^{15}\text{N}_{\text{AA}}$ (dashed
523 trophoclines in Fig. 6), with food chain length inferred from the *TP* of apex predators. [Chikaraishi
524 et al. \(2014\)](#) used these trophoclines to describe the food web of a coastal rocky shoreline
525 community in Japan. Based on 39 species ($n = 100$) covering macroalgae, gastropods, echinoderms,
526 bivalves, crustaceans, fish, and a cephalopod to document the food web structure, the study
527 documented a food web covering 4.5 trophic levels (Fig. 7a). Probably supported by macroalgae at
528 *TP* 1, the top predator in the system was the Kidako moray eel (*Gymnothorax kidako*) with an
529 average *TP* of 4.6. Despite a large variation in baseline $\delta^{15}\text{N}$ values, demonstrated by $\delta^{15}\text{N}_{\text{Phe}}$
530 values varying between 3.5‰ and 8.7‰, all algal samples had *TP* close to 1 with known
531 herbivores all close to *TP* 2, demonstrating the importance in knowing the baseline $\delta^{15}\text{N}$ value
532 across appropriate time and space scales. In Lake Baikal, [Ohkouchi et al. \(2015\)](#) reported
533 analytical results for seven species ($n = 53$) covering diatoms, amphipods, sculpins, and seals (Fig.
534 7b). The *TP* of seals, a top predator in the lake, was as high as 5.1, suggesting that the trophic
535 length of the lake was one unit longer than that calculated based on the $\delta^{15}\text{N}_{\text{bulk}}$ record.
536 Furthermore, the potential for baseline variation to confound analysis of spatial changes in *TP*
537 based on bulk $\delta^{15}\text{N}$ values has recently been highlighted by a study on Lake Superior food webs by
538 [Kruger et al. \(2016\)](#). TP_{bulk} suggested that the top predator (lake trout) spatially varied by up to *TP*
539 of 1; however, the $\delta^{15}\text{N}_{\text{AA}}$ values confirmed a common *TP* and the likelihood that baseline $\delta^{15}\text{N}$
540 variation confounded TP_{bulk} estimates. A recent study by [Papastamatiou et al. \(2015\)](#) is one of the
541 few to demonstrate variation in bulk isotopic composition due to trophic differences rather than

542 baseline differences. The authors combined acoustic tracking with CSIA-AA to demonstrate
543 trophic flexibility in giant trevally from deep water reefs on a Pacific atoll. Individuals showed
544 variability in their diel migration and feeding behavior that was mirrored in the wide range of *TP*
545 determined by CSIA-AA (*TP* 3.5 to 4.6).

546 In addition to demonstrating the utility of the $\delta^{15}\text{N}_{\text{AA}}$ approach, these studies highlight
547 the potential variability in food webs, such that higher-order consumers do not occupy a single *TP*.
548 Indeed, the capacity for intraspecific variation in *TP* has long been known based on theory and
549 empirical work (Polis, 1991; Polis and Strong, 1996). Conceptual examples of trophic omnivory
550 associated with ontogeny are provided in Fig. 6. Adults and juveniles of an anchovy prey species
551 are shown with varying *TP* of around 2, whereas the increase in the size classes of an apex
552 consumer (tuna) leads to increased *TP*, likely reaching a *TP* close to 4.5-5.0 at their maximum size
553 (Choy et al., 2015; Estrada et al., 2005). The degree of trophic omnivory within a food web could
554 be quantified based on the deviation of consumers from integer values (e.g., 2 for strict herbivores,
555 3 and 4 for strict carnivores) for *TP* derived from $\delta^{15}\text{N}_{\text{AA}}$. Recognition of intra-individual
556 variability in *TP*, as observed by Papastamatiou et al. (2015) in deep reef giant trevally, will be an
557 important outcome of future $\delta^{15}\text{N}_{\text{AA}}$ studies.

558 The trophic role of apex consumers will probably ultimately be elucidated by applying
559 CSIA-AA to *TP* estimates in aquatic systems. Several recent studies have highlighted the
560 importance of accurate *TP* estimates for apex consumers, and raised questions about the trophic
561 role of sharks as apex predators in coral reef ecosystems. Hussey et al. (2014) highlighted the
562 dependence of TP_{bulk} estimates on TDF_{bulk} values and suggested that the *TP* of apex shark species
563 may have been underestimated using $\delta^{15}\text{N}_{\text{bulk}}$. Hussey et al. (2015) later demonstrated an expanded
564 trophic complexity among large sharks using CSIA-AA. Hilting et al. (2013) used bulk stable
565 isotope analysis to suggest that apex predators in central Pacific reefs might be predominately
566 supported by benthic productivity and N_2 -fixation. However, strong conclusions could not be
567 drawn because of spatial variability in primary producer bulk isotopes (see also the CSIA-AA
568 findings of O'Malley et al. (2012) on two species of lobster in the Northwest Hawaiian Islands).
569 Conversely, based on limited bulk isotope analyses, Frisch et al. (2016) assigned Great Barrier
570 Reef tiger sharks (*Galeocerdo cuvier*) to pelagic productivity, whereas whitetip and blacktip reef
571 sharks were assigned to predominately benthic sources (65% and 72%, respectively). Coupling
572 $\delta^{15}\text{N}_{\text{bulk}}$ values with stomach contents analyses suggested these reef sharks occupy similar trophic
573 positions to large predatory fish, such as snapper, rather than acting as apex predators (Frisch et al.,
574 2016).

575 In addition to comprehensive studies at ecosystem scales, several studies have used
576 CSIA-AA tools to understand the habitat of cryptic species. One example is Miller et al. (2012),
577 who measured the $\delta^{15}\text{N}_{\text{AA}}$ values from leptocephali, the larvae of the Japanese eel (*Anguilla*

578 *japonica*), whose food source is unknown. The estimated mean *TP* of the eel larvae was 2.4, which
579 in that ecosystem was most consistent with a diet based on particulate organic matter (POM)
580 composed of detritus from multiple sources. [Ohkouchi et al. \(2013\)](#) reported the *TP* of deep-water
581 ram's horn squid (*Spirula spirula*), one of the most enigmatic cephalopods found commonly all
582 over the world. Such information is useful for conserving endangered species through developing
583 artificial diets for aquafarming.

584 Finally, it is important to note that CSIA-AA is also applicable to laboratory and
585 museum specimens. A laboratory experiment conducted over the period of one year indicated that
586 formalin-fixation does not affect $\delta^{15}\text{N}_{\text{AA}}$ values derived from an aquatic consumer ([Ogawa et al.,](#)
587 [2013](#)). [Ogawa et al. \(2013\)](#) used formalin-fixed samples to reconstruct historical variation (1916 to
588 1992 CE) in *TP* of isaza (*Gymnogobius isaza*), a pelagic gobiid fish from the eutrophic Lake Biwa,
589 Japan. The $\delta^{15}\text{N}_{\text{bulk}}$ value of isaza has increased greatly during the 20th century ([Ogawa et al., 2001;](#)
590 [Nakazawa et al., 2010](#)), which can be explained either by an increase in *TP* with the reorganization
591 of bio-communities because of eutrophication, or by the increase of $\delta^{15}\text{N}$ in the nitrogen pool
592 owing to denitrification. The CSIA-AA results strongly suggested that eutrophication did not affect
593 the *TP* of the fish in the lake, and that the $\delta^{15}\text{N}_{\text{AA}}$ value of the formalin-fixed fish reflected the $\delta^{15}\text{N}$
594 of the nitrogen pool of the lake accurately (Fig. 8). A large global archive of formalin-fixed
595 samples would offer a tool for reconstructing paleo-limnological changes, and for constraining the
596 ecological consequences of environmental change with CSIA-AA.

597

598 **4.2. Food web analyses in terrestrial ecosystems**

599 CSIA-AA has provided new insights into the trophic roles of terrestrial organisms and, as
600 in aquatic ecosystems, distinguished itself from traditional bulk isotopic approaches ([Chikaraishi et](#)
601 [al., 2011;](#) [Steffan et al., 2013](#)). [Chikaraishi et al. \(2011\)](#) showed that equation 2 is equally valid for
602 terrestrial systems and [Steffan et al. \(2013\)](#) demonstrated that accurate and precise *TP* could be
603 derived for higher-order carnivores, using the CSIA-AA method to measure *TP* across four trophic
604 levels in terrestrial insect communities. Recent work in regards to carbon CSIA-AA shows great
605 promise in filling gaps currently left open by nitrogen CSIA-AA approaches because carbon
606 isotopic composition among essential AAs, also called $\delta^{13}\text{C}_{\text{EAA}}$ fingerprints, can provide
607 information about trophic pathways from plant sources and gut/soil microbes to consumers in
608 terrestrial ecosystems ([Larsen et al., 2016a, 2016b](#)). Carbon and nitrogen CSIA-AA has also
609 revealed novel aspects of animal and microbial biology, proving CSIA-AA to be a powerful new
610 tool for examining modern and ancient biological communities ([O'Brien et al., 2002, 2004;](#)
611 [Chikaraishi et al., 2014;](#) [Steffan et al., 2015a](#)).

612

613 4.2.1. Trophic position estimation

614 The first evidence that CSIA-AA is a feasible method for *TP* analysis among terrestrial
615 organisms was obtained by using the $\delta^{15}\text{N}$ values of Glu and Phe of primarily herbivorous
616 organisms and their plant host material collected from a farm in Japan (Chikaraishi et al., 2011).
617 Because aphids are strict herbivores, they were ideal subjects for testing the accuracy of this tool,
618 and the estimated *TP* of the aphids was shown to be the expected value of ~ 2.0 . Carnivorous insect
619 specimens (e.g., lady beetles, wasps, and hornets) were also analyzed *via* CSIA to estimate their *TP*
620 values. The data provided interesting insights into the trophic ecology of these animals; however,
621 because most carnivores are free-roaming generalists, their actual *TP* values were not known and
622 they were not suitable for testing the accuracy of this tool. These early studies revealed that insect
623 *TP* remained constant through major ontogenetic shifts, including insect pupation (Chikaraishi et
624 al., 2011). During such metamorphoses, there is much synthesis of new tissues and organs, so it
625 was expected that there would be significant fractionation or routing of ^{15}N within the pupating
626 insect. The finding that arthropod metamorphosis left the $\delta^{15}\text{N}$ values largely unchanged was
627 critical to further applications of $\delta^{15}\text{N}_{\text{AA}}$ to insect food web ecology (Chikaraishi et al., 2011).

628 Steffan et al. (2013) used terrestrial insect populations in axenic culture to test whether top
629 carnivore *TP* could be reliably determined using $\delta^{15}\text{N}_{\text{AA}}$. In this study, two different insect
630 communities were maintained, each spanning four trophic levels, and each consuming an
631 ecologically realistic component of their diet. Steffan et al. (2013) showed that the $TP_{\text{Glu/Phe}}$ of
632 higher-order consumers (carnivorous insects) could be measured with high accuracy, and that the
633 $\Delta_{\text{Glu/Phe}}$ value was consistent between herbivores and tertiary carnivores. The $\Delta_{\text{Glu/Phe}}$ value for
634 herbivorous and carnivorous arthropods was similar to that found by Chikaraishi et al. (2009) for
635 marine fish and gastropods, showing that the *TP* formula using a $\Delta_{\text{Glu/Phe}}$ value $\sim 7.6\text{‰}$ and a $\beta_{\text{Glu/Phe}}$
636 value appropriate for each environment in equation 2 was applicable to a wide variety of
637 ecosystems. The consistency in the CSIA-AA findings across animal taxa and ecosystem types
638 observed on land (Chikaraishi et al., 2010; Steffan et al., 2013) provided a foundation to begin
639 investigating consumer *TP* in the field, at larger spatial scales and in more diverse communities.
640 Further CSIA-AA of directly sampled terrestrial organisms in the wild revealed a high degree of
641 trophic omnivory among 38 consumer species, providing some of the strongest empirical evidence
642 of the predominance of omnivory in food webs (Chikaraishi et al., 2014).

643 Novel contributions of CSIA-AA to terrestrial ecology have centered around the
644 microbiome and mainly the inclusion of microbes in trophic hierarchies. Studies involving multiple
645 phyla of fungi and bacteria, plus vertebrate and invertebrate animals, showed that the CSIA-AA
646 approach provides a new way to probe trophic ecology of the three domains of life (Steffan et al.,
647 2015a). Fungi are particularly important consumers and symbionts in many terrestrial systems
648 (Bardgett and Cook, 1998; Moore and de Ruiter, 2012). Showing that these organisms can be
649 integrated into food-chains has allowed more refined interpretations of animal trophic identity.

650 However, this also raises questions of how to interpret the *TP* values of detritivores. Recent work
651 has shown that microbes increase the *TP* values of detrital complexes, and when animals eat such
652 microbe-colonized complexes, the consumer *TP* values increase to the same degree (Steffan et al.,
653 2017). Given that detritivory is the dominant trophic paradigm on land (Coleman, 1996; Hagen et
654 al., 2012) and that microbes are the dominant consumers among the detritivores (Peterson and
655 Luxton, 1982; van der Heijden et al., 2008; Moore and de Ruiter, 2012), the ability to explicitly
656 integrate microbes into trophic hierarchies represents a major advance in trophic ecology. Common
657 detritivorous animals, such as earthworms, fruit flies, and springtails, exhibit *TP* values (2.4-2.8),
658 providing evidence of the degree to which they mix microbivory with herbivory (Steffan et al.,
659 2017). Detritivores form an immense prey base for predators in terrestrial systems (Haraguchi et al.,
660 2013; Hyodo et al., 2015), and this prey base tends to shape the trophic identity of most carnivores
661 (Coleman, 1996; Bardgett and Cook, 1998).

662

663 4.3.2. Recent discoveries in terrestrial biology and ecology

664 $\delta^{15}\text{N}_{\text{AA}}$ was used to reveal that leafcutter ants (*Acromyrmex*) in Neotropical rainforests are
665 trophically carnivorous (Steffan et al., 2015a). The ants feed almost exclusively on the fruiting
666 bodies of their fungal symbiont, *Leucoagaricus*. Since this fungus feeds solely on plant material,
667 the fungus is a strict herbivore, and the ants are strict carnivores. This finding implies that fungi,
668 not ants, are the dominant herbivores of the Neotropics. Interestingly, there is a third symbiont, a
669 bacterium, in the leafcutter ant fungus gardens that gathers in powdery white masses on the ant
670 exoskeleton (Currie et al., 2006). It was unclear whether this bacterium fed on the ants or some
671 other resource. CSIA-AA showed that the bacteria were feeding on ant tissues; thus, a bacterium
672 was the apex carnivore within the fungus-garden community (Fig. 9, Steffan et al., 2015a). Fungi
673 can also be predators, and $\delta^{15}\text{N}_{\text{AA}}$ values were used to demonstrate that an entomopathogenic
674 fungus, *Beauveria bassiana*, registered a *TP* of 3.0 after killing and consuming its prey, a
675 herbivorous caterpillar. At the other end of the trophic spectrum, Asiatic black bears (*Ursus*
676 *thibetanus*) were shown to feed remarkably low in the food-chain, registering near *TP* 2.0. Thus,
677 there are now multiple examples in the literature where the trophic tendencies of terrestrial
678 mammals (e.g., mice, bears) have been measured using $\delta^{15}\text{N}_{\text{AA}}$ (Nakashita et al., 2011; Steffan et
679 al., 2015a).

680 In agricultural contexts, CSIA-AA has been used to characterize the trophic roles of
681 organisms thought to be beneficial to crop protection (Steffan et al., 2015b). Carnivorous
682 arthropods are generally assumed to be helpful in suppressing herbivorous pest species, but
683 CSIA-AA has shown that only certain predator species contribute substantially to pest control.
684 Some carnivores are beneficial for crop protection, and some are neutral, and other species may
685 undermine crop protection efforts by feeding on the beneficial carnivores (Fig. 10). Knowledge of

686 which predator communities are likely to help or harm crop protection is useful for the ecological
687 management of crop fields.

688

689 **4.3. Applications to ancient humans and extinct mammals**

690 CSIA-AA has been used to study tissues, like bone collagen and scalp hair, from
691 archaeological and contemporary humans, and ancient soils in archaeological and anthropological
692 studies. These studies span various fields, including paleodiet, nutrition, paleopathology, and
693 ancient land use (e.g., [Hare et al., 1991](#); [Fogel, 1997](#); [Petzke et al., 2005](#)). Studies of paleodiet
694 mainly revolve around investigating i) marine protein consumption ([Naito et al., 2010a, 2010b](#);
695 [Styring et al., 2010](#)), ii) the importance of animal proteins relative to plant proteins in terrestrial
696 ecosystems ([Naito et al., 2013b, 2016b](#)), and iii) the importance of proteins from freshwater
697 resources relative to proteins from terrestrial resources ([Naito et al., 2013a](#)). Goal iii) is challenging
698 because distinguishing terrestrial and freshwater food consumption is difficult since these two
699 environments may share the same nitrogen sources (e.g., contributions of terrestrial primary
700 production to a stream ecosystem, [Naito et al., 2016a](#)).

701 $\delta^{15}\text{N}_{\text{Phe}}$ values in some archaeological contexts in animals mirrors their nitrogen source
702 owing to little trophic ^{15}N enrichment. For example, [Naito et al. \(2010, 2013b\)](#) examined coastal
703 and inland archaeological sites from the Jomon period in Japan (ca. 15,000 to 2,300 years BP). The
704 $\delta^{15}\text{N}_{\text{Phe}}$ values of animals in these contrasting ecosystems, including humans, were consistent
705 within each ecosystem, although there were differences between ecosystems (Fig. 11). The coastal
706 population showed $\delta^{15}\text{N}_{\text{Phe}}$ values between those of marine and terrestrial ecosystems, with values
707 closer to marine ecosystems, indicating heavier reliance of the humans on marine food resources.
708 However, the inland population had $\delta^{15}\text{N}_{\text{Phe}}$ values in the terrestrial ecosystem indicating purely
709 terrestrial food habits. In both cases, tracing the nitrogen source for humans was facile because
710 each ecosystem showed marked differences in $\delta^{15}\text{N}_{\text{Phe}}$ values. However, this is not the case in other
711 archaeological contexts where $\delta^{15}\text{N}_{\text{Phe}}$ values vary substantially within each ecosystem. $\delta^{15}\text{N}_{\text{Phe}}$
712 values in terrestrial prey animals can vary widely (>6‰), even for a single species from a single
713 site, which makes it difficult to trace the nitrogen source (Fig. 12). Nevertheless, this technique is
714 still useful for examining *TP* values of animals. Neanderthals from this site exhibited *TP* values of
715 2.7 to 2.8, similar to those of wolves (*TP* 2.9), suggesting that the Neanderthals ate meat-based
716 diets, with the possible addition of plant foods (Fig. 11).

717 CSIA-AA can also be used to investigate diets of extinct mammals, including woolly
718 mammoths (*Mammuthus primigenius*) ([Naito et al., 2016b](#); [Schwartz-Narbonne et al., 2015](#)), cave
719 bears (*Ursus spelaeus*) ([Naito et al., 2016c](#)), scimitar-toothed cats (*Homotherium serum*), and
720 short-faced bears (*Arctodus* spp.) ([Schwartz-Narbonne et al., 2015](#)). Based on the high $\delta^{15}\text{N}_{\text{Phe}}$
721 value of mammoths, it has been hypothesized that the mammoth occupied a distinct foraging niche

722 or habitat compared with other coeval herbivores, owing to the high $\delta^{15}\text{N}$ values of bulk collagen
723 arising from ^{15}N -enriched food sources (Naito et al., 2016b). This finding demonstrates the
724 separation of mixtures of environmental signals (e.g., aridity may elevate the $\delta^{15}\text{N}$ values of animal
725 body tissues: Heaton et al., 1986; Schwarcz et al., 1999) and dietary signals in $\delta^{15}\text{N}$ values of
726 collagen. However, the $\delta^{15}\text{N}$ values of body tissues, and probably the $\delta^{15}\text{N}_{\text{AA}}$ values, may also
727 encode physiological states, illness, and quality of diet (Fogel et al., 1989; Fuller et al., 2004, 2006;
728 Reitsema and Muir, 2015; Reitsema, 2013; Chikaraishi et al., 2015). In combination with studies
729 on contemporary humans and archaeological human remains, the number of study fields for
730 CSIA-AA, such as paleopathology, may expand (Fogel, 1997; Metges and Petzke, 1997; Petzke et
731 al., 2006, 2010; Romek et al., 2013).

732 Lastly, CSIA-AA has also been used to investigate past land use by humans. Preliminary
733 results suggest that the $\delta^{15}\text{N}$ values of Phe and Thr in the soil may be useful for distinguishing the
734 soil under grassland from that under cereal (Bol et al., 1998; Simpson et al., 1997, 1999). Although
735 the underlying mechanisms controlling the $\delta^{15}\text{N}$ dynamics of soil AAs are not well understood,
736 some AAs may provide clues for understanding past human activities like cultivation (Styring et al.,
737 2013), which is important because cultivars are rarely preserved in the archaeological record.

738

739 **4.4. CSIA-AA and isoscapes: application to ecogeochemistry and the detection of animal** 740 **migration**

741 Ecogeochemistry is the application of geochemical techniques to fundamental questions in
742 population and community ecology, and is inherently spatial (e.g., Bowen, 2010; Graham et al.,
743 2009; Ramos and Gonzalez-Solis, 2012; McMahan et al., 2013a). Consequently, accurate
744 interpretation of stable isotopic compositions in ecological or environmental studies requires
745 knowledge of the geospatial and temporal variability in isotope values at the base of the food web,
746 often referred to as isotope baselines (Post, 2002; McMahan et al., 2013b). Spatiotemporally explicit
747 maps of isotopic variability, termed isoscapes, have emerged as important tools for addressing
748 interrelated ecological questions about animal movement, habitat use, biogeochemical cycling, and
749 forensic science (e.g., West et al., 2010).

750 Effective application of isoscapes to ecological questions requires specific information
751 (Hobson et al., 2010). First, an isoscape must be established that characterizes systematic geospatial
752 variability in isotopic compositions across environmental gradients. Second, tissue turnover rates
753 that determine the period of spatial integration of isotopic composition for a particular animal tissue
754 must be constrained. Finally, the isotope fractionation factors between the consumer and diet, or
755 between animals and the ambient environment that offset geochemical values in animal tissues from
756 baseline isoscape values, must be estimated or quantified.

757 Bulk tissue or whole animal isotope analyses have been the primary tools in applications

758 of terrestrial and marine ecosystem isoscapes (Bowen et al., 2009; Jaeger et al., 2010; MacKenzie et
759 al., 2011; Hobson et al., 2012; Trueman et al., 2012; Clementz et al., 2014). However, in addition to
760 characterizing the geospatial structure of isotope data within a system, we also must account
761 accurately for how baseline isotope values are modified as they propagate through food webs to
762 upper trophic level consumers (Hobson et al., 2010). Thus, a major obstacle to interpreting bulk
763 tissue isotope values of consumers accurately is separating the relative effects of variability at the
764 base of the food web from trophic dynamics within the food web that links consumers to those
765 baselines (Post, 2002).

766 CSIA-AA can disentangle the relative effects of geographic and trophic dynamics on
767 consumer isotopic compositions (Chikaraishi et al., 2007, 2009; Popp et al., 2007; Lorrain et al.,
768 2009, 2015; Olson et al., 2010). The differential isotopic fractionation of individual AAs provides
769 direct access to information about integrated ecosystem isotopic baselines without the confounding
770 issue of trophic fractionation, and without the need to analyze and characterize all the trophic
771 linkages between the baseline isoscapes and upper trophic level consumers *a priori*. Below we
772 highlight several unique but complementary examples of how CSIA-AA, in the context of geospatial
773 isotopic variations, provides unprecedented links between animal ecology and biogeochemistry in
774 complex ecosystems.

775

776 4.4.1. Case study 1: Mussel isoscapes of the California coast

777 One promising CSIA-AA isoscape application for monitoring coastal biogeochemical
778 change is the creation of detailed maps of coastal isotopic baselines, based on CSIA-AA
779 measurements in filter feeding mollusks. Coastal system isoscapes are inherently challenging owing
780 to high temporal and spatial variability in primary production and biogeochemical cycles. Many
781 coastal regions are characterized by large seasonal swings in temperature, salinity, nutrient
782 availability, and terrestrial inputs, while high spatial variability in oceanographic conditions is driven
783 by coupled local winds, upwelling, and current patterns (*e.g.*, Walker and McCarthy, 2012; Walker
784 et al., 2014). The isotopic compositions of consumers can often integrate this environmental
785 variation. However, on the spatial scales of coastal processes, assigning mobile consumers to
786 specific locations can be difficult. Tissues of sessile filter-feeders, such as mussels, offer a solution
787 to this problem: they do not move, and specific tissues/organism sizes can be chosen to provide
788 additional control over the integrated time scales represented by the samples. Early work coupling
789 CSIA-AA proxies for isotopic baseline (*i.e.*, $\delta^{15}\text{N}_{\text{Phe}}$), coupled with high resolution sampling of filter
790 feeding consumers, has allowed the creation of isoscapes of baseline coastal primary production,
791 based on precisely known and replicable sampling locations (Vokhshoori et al., 2014; Vokhshoori
792 and McCarthy, 2014).

793 However, the reconstruction of baseline isoscapes based on $\delta^{15}\text{N}_{\text{AA}}$ also poses a number of

794 challenges, primarily with the interpretation of $\delta^{15}\text{N}_{\text{Phe}}$ values in mollusk bioarchives. The challenges
795 include clarifying the mix of littoral food sources using CSIA-AA proxy records, understanding
796 temporal/seasonal effects in this signal, and the requirement to understand the influence of bulk algal
797 $\delta^{15}\text{N}$ isoscapes on mollusk AA isotopic values. In two recent papers, Vokhshoori and coauthors
798 explored these problems using littoral *Mytilus californianus* collected from 28 sites on the California
799 coast, spanning ~10 degrees of latitude (32° to 42°N) within the California current system
800 (Vokhshoori et al., 2014; Vokhshoori and McCarthy, 2014). CSIA-AA values of adductor muscle
801 tissue from individuals of a similar size class were selected to represent approximately annual
802 integration timescales.

803 $\delta^{15}\text{N}_{\text{bulk}}$ values in mussels showed a strong linear trend with latitude (Fig. 13). Although
804 there were clear site-specific and region-specific offsets in $\delta^{15}\text{N}$ values, the overall data indicated a
805 strongly linear progressive change in $\delta^{15}\text{N}$ values, averaging 0.4‰ per degree of latitude, across the
806 coastal California current system. This change reflects the relative geographical influence of water
807 upwelled from the California undercurrent, which transports ^{15}N -rich nitrate poleward (e.g., Altabet
808 et al., 1999). The $\delta^{15}\text{N}_{\text{Phe}}$ values also tracked the $\delta^{15}\text{N}_{\text{bulk}}$ values, confirming the nutrient baseline as
809 the underlying driver for changes in bulk mussel tissue. Prior studies had indicated generally lower
810 $\delta^{15}\text{N}_{\text{nitrate}}$ values in more northern regions of this system (Altabet et al., 1999; Kienast et al., 2002;
811 Sigman et al., 2009), however the strength of the linear trend revealed by high resolution mussel
812 sampling was surprising. This result suggests that such mollusk-derived isoscapes can be used to
813 precisely define changes in the effects of coastal oceanography on baseline Isoscapes, as well as to
814 identify local regions of variation linked for example to upwelling patterns (Walker and McCarthy,
815 2012).

816 However, $\delta^{15}\text{N}_{\text{bulk}}$ records are inherently unable to reconstruct baseline $\delta^{15}\text{N}$ values
817 directly. $\delta^{15}\text{N}_{\text{bulk}}$ values were 2 to 4‰ higher than the expected range of $\delta^{15}\text{N}_{\text{nitrate}}$ values in this
818 system, probably because of a combination of trophic transfer and tissue-specific offsets.
819 Vokhshoori and McCarthy (2014) found that $\delta^{15}\text{N}_{\text{Phe}}$ values corresponded most closely to the range
820 of previously measured $\delta^{15}\text{N}_{\text{nitrate}}$, suggesting that $\delta^{15}\text{N}_{\text{Phe}}$ in mussels is a direct proxy for annual
821 average $\delta^{15}\text{N}_{\text{nitrate}}$. Finally, a constant $\delta^{15}\text{N}_{\text{bulk}}$ vs. $\delta^{15}\text{N}_{\text{Phe}}$ offset observed for all samples allowed the
822 construction of predicted “coastal nitrate” $\delta^{15}\text{N}$ values. The resulting isoscape was grounded in
823 high-resolution bulk sampling, but then calibrated to baseline $\delta^{15}\text{N}$ values based on CSIA-AA data
824 (Fig. 12). It is unclear how far isoscapes based on littoral species can be extrapolated. However, for
825 Monterey Bay, a direct comparison of mussel $\delta^{15}\text{N}$ data with a greater variety of more offshore
826 sample types (e.g., sinking POM, plankton tows, and surface sediments) suggested that, at least on
827 the timescales sampled, mussel $\delta^{15}\text{N}_{\text{Phe}}$ values reflect baseline $\delta^{15}\text{N}$ values in local coastal waters.

828 These results demonstrate the potential of CSIA-AA in sessile filter feeders to create the
829 first true baseline isoscapes of coastal production, with the potential for an extraordinary degree of

830 geographic and temporal resolution. Although bulk tissue analysis can indicate geographic trends,
831 coupling $\delta^{13}\text{C}_{\text{EAA}}$ fingerprinting with $\delta^{15}\text{N}_{\text{AA}}$ allows the fundamental ambiguity of organic matter
832 sources to be addressed, and can quantify the relative balance of sources underlying CSIA-AA
833 signals. $\delta^{15}\text{N}_{\text{Phe}}$ values can track baseline $\delta^{15}\text{N}$ values, and in systems with full nitrate utilization this
834 should also allow direct assembly of $\delta^{15}\text{N}_{\text{nitrate}}$ isoscapes. Therefore, it may be possible to monitor
835 fine-scale shifts in coastal nitrogen biogeochemical cycles linked to short- or longer-term
836 fluctuations in climate and physical forcing. However, several challenges remain, including
837 understanding in more detail the calibrations required to link measured $\delta^{15}\text{N}_{\text{AA}}$ values to average
838 primary production (or nitrate) isotope values, and investigating integration timescales, such as
839 potential seasonal bias and the effects of tissue type, organism size, and growth stage.

840

841 4.4.2. Case study 2: Detecting animal migration

842 Systematic variations in nitrogen isotopic compositions in the ocean, such as the mussel
843 isoscapes in Section 4.4.1 or those in the eastern tropical North Pacific (Olson et al., 2010), create
844 ecoregions with distinctive isotope ratios in baseline organisms (e.g., phytoplankton). These regional
845 differences allow the results of CSIA-AA to be used to recognize animal migrations. This approach
846 principally relies on certain AA isotopic compositions in animals having reached a steady state with
847 the $\delta^{15}\text{N}$ value at the base of the food web.

848 Marine animal migrations can be identified with CSIA-AA by two approaches. The first is
849 a chronological reconstruction of isotopic compositions of AAs in an archival tissue (e.g., otoliths,
850 fin spines) that represent the animal's environment at different stages of ontogeny. Older tissues can
851 represent an isotopic steady state with an environment different from an animal's current location,
852 which can then be compared with recently synthesized tissue that has AA isotopic compositions in a
853 steady state with the current location. The second approach is to compare isotopic compositions of
854 AAs in a non-archival tissue (e.g., muscle) across individuals that are a suspected mix of residents
855 and recent migrants to a particular environment. With this approach, the timeframe for
856 distinguishing residents from migrants is defined by the turnover time of nitrogen in the tissue
857 analyzed. The $\delta^{15}\text{N}_{\text{SAA}}$ values in animals record the isotopic composition at the base of the food web.
858 In addition, the difference in $\delta^{15}\text{N}_{\text{SAA}}$ and $\delta^{15}\text{N}_{\text{TAA}}$ values constrains potential *TP* variations between
859 residents and suspected migrants (e.g., Madigan et al., 2012b; Seminoff et al., 2012).

860 For example, $\delta^{15}\text{N}_{\text{AA}}$ values were used to study the foraging ecology and habitat use of the
861 brown stingray (*Dasyatis lata*) near Kaneohe Bay, Oahu, Hawaii (Dale et al., 2011). Although
862 quantitative stomach content analysis of *D. lata* indicated an ontogenetic shift to a higher *TP* in
863 larger, older specimens, the largest stingrays had the lowest $\delta^{15}\text{N}_{\text{bulk}}$ values. Lower $\delta^{15}\text{N}_{\text{bulk}}$ values
864 would indicate a decreased *TP* in the largest stingrays, contradicting stomach content analyses, if all
865 analyzed individuals were feeding in environments with similar baseline $\delta^{15}\text{N}$ values. However, Dale

866 [et al. \(2011\)](#) used differences in $\delta^{15}\text{N}_{\text{Glu}}$ and $\delta^{15}\text{N}_{\text{Phe}}$ to show that *TP* of *D. lata* increased with size
867 and that $\delta^{15}\text{N}_{\text{bulk}}$ values were independent of *TP*. These findings clearly indicated that the largest *D.*
868 *lata* were feeding in habitats that had distinctly lower $\delta^{15}\text{N}$ values at the base of the food web than
869 the environments where smaller stingrays foraged. One implication of this finding was that stingray
870 $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{bulk}}$ values reflected migration patterns better than *TP*. Both $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values,
871 examined as a function of size and stingray sex, revealed that changes in bulk isotopic compositions
872 closely coincided with the onset of sexual maturity, confirming Kaneohe Bay as a nursery habitat for
873 *D. lata* ([Dale et al. 2011](#)).

874 $\delta^{15}\text{N}_{\text{AA}}$ values have been used to recognize marine fish undergoing trans-Pacific
875 migrations ([Madigan et al., 2014, 2016](#)). Pacific bluefin tuna (*Thunnus orientalis*) inhabit the
876 western and eastern Pacific Ocean. All bluefin tuna spawn in the western Pacific and an unknown
877 proportion of these tuna migrate to the eastern Pacific early in their life. Once in the eastern Pacific,
878 these bluefin tuna migrants reside in the California Current ecosystem for several years and then
879 return to the western Pacific to spawn. Tracking these transoceanic migrations has been challenging;
880 however, large differences in baseline $\delta^{15}\text{N}$ values between the eastern and western Pacific Ocean
881 (e.g., [Navarro et al., 2013](#)) can be used to understand the timing and numbers of individuals
882 undergoing trans-Pacific migration better.

883 Recently, [Madigan et al. \(2012a, 2013\)](#) showed that the short-lived Fukushima-derived
884 radiocesium (^{134}Cs) content of bluefin tuna caught in the eastern Pacific unequivocally identified
885 bluefin tuna that had fed off the coast of Japan and migrated from the western Pacific. [Madigan et al.](#)
886 [\(2014\)](#) combined nitrogen isotope analyses of AAs with bluefin tuna containing Fukushima-derived
887 ^{134}Cs to evaluate the migration history of different year class bluefin tuna caught in the eastern
888 Pacific. Bluefin tuna in the eastern Pacific had a bimodal distribution of $\delta^{15}\text{N}_{\text{bulk}}$, with lower values
889 consistently found in bluefin tuna specimens with Fukushima-derived ^{134}Cs . Bluefin tuna with
890 Fukushima-derived ^{134}Cs had $\delta^{15}\text{N}$ values even lower than baseline organisms (krill, copepods)
891 found in the eastern Pacific Ocean. [Madigan et al. \(2014\)](#) also found that the $\delta^{15}\text{N}_{\text{SAA}}$ in eastern
892 Pacific bluefin tuna with Fukushima-derived ^{134}Cs were 7.7 to 8.7‰ lower than in fish that lacked
893 ^{134}Cs , including resident bluefin tuna, yellowfin tuna, and prey (Pacific saury and jack mackerel).
894 This indicated that $\delta^{15}\text{N}_{\text{SAA}}$ values were robust markers for distinguishing resident bluefin tuna from
895 recent migrants. In addition, the results of CSIA-AA indicated that differences in $\delta^{15}\text{N}_{\text{bulk}}$ values
896 were not due to trophic variability among bluefin tuna. Recently, [Madigan et al. \(2016\)](#) used the
897 CSIA-AA results for giant bluefin tuna caught in the western Pacific Ocean to validate the westward
898 trans-Pacific migration of sexually mature individuals from the eastern Pacific Ocean to spawning
899 grounds off the coast of Taiwan.

900 The findings of [Madigan et al. \(2014\)](#) have important implications for the sustainable
901 management of the bluefin tuna fishery in the eastern Pacific Ocean. The results of their study

902 indicated that the eastern Pacific bluefin tuna population was subsidized by a substantial number of
903 older individuals (*i.e.*, year class 2 to 3) from the western Pacific, which was not previously
904 recognized. In addition, knowledge of muscle turnover time in bluefin tuna (Madigan et al., 2012b)
905 sets limits on how quickly a migrant bluefin tuna would reach a nitrogen isotope steady state with
906 the new environment (and thus be classified as a resident based on CSIA-AA), and allows the date of
907 migrant arrival to be estimated. Madigan et al. (2014) found that the proportion of recent migrants to
908 residents decreased with increasing age, which is critical information for effectively managing this
909 heavily fished species. Unlike a radiogenic isotopic tracer that has finite utility for studying animal
910 migration in the ocean, CSIA-AA can be used *ad infinitum* and in other species. For example, the
911 same isotopic differences in the North Pacific Ocean were used to distinguish apparent eastern and
912 western Pacific migratory groups of endangered leatherback sea turtles (*Dermochelys coriacea*),
913 which provided unique evidence for foraging area philopatry among turtles nesting in Indonesia
914 (Seminoff et al., 2012). These CSIA-AA results clarify the interpretation of bulk tissue isotopic
915 variability in populations, and can be used to recognize and trace movements of many highly
916 migratory pelagic species.

917

918 4.4.3. Case study 3: Deep-sea coral

919 As Earth's climate changes, there is a growing need to put these changes and their
920 subsequent effects on ecosystem structure and function into a greater historical context (Corno et al.,
921 2007; Hoegh-Guldberg and Bruno, 2010). One of the most exciting new applications for CSIA-AA
922 is in paleoceanography, where parameters originally developed for ecology are being adapted as new
923 paleo-proxies in novel protein-rich archives. Biogenic skeletons of proteinaceous deep-sea corals
924 provide a remarkable geochemical archive of information about the structure and function of past
925 ocean ecosystems (Druffel, 1997; Robinson et al., 2014). These globally distributed corals represent
926 "living sediment traps", recording geochemical information about recently exported organic
927 materials in their exquisitely preserved accretionary protein skeletons (Roark et al., 2009; Guilderson
928 et al., 2013). Much of the recent work with proteinaceous deep-sea corals has focused on isotope
929 analysis of total skeletal material as a proxy for changes in surface ocean conditions (*e.g.*, Sherwood
930 et al., 2005; Williams et al., 2007; Hill et al., 2014). However, CSIA-AA results can provide
931 unprecedented reconstruction of past ocean conditions (Sherwood et al., 2011, 2014; Schiff et al.,
932 2014; Strzepek et al., 2014; McMahon et al., 2015c; Williams et al., 2017). The $\delta^{15}\text{N}_{\text{SAA}}$ values of
933 these consumers provide particularly faithful records of baseline nitrogen sources and cycling that
934 are otherwise seldom preserved in paleorecords.

935 Sherwood et al. (2011) first applied CSIA-AA to deep-sea corals to distinguish between
936 temporal (decadal to centennial) changes in nitrogen sources, while constraining changes in the
937 trophic structure of proteinaceous deep-sea corals in the Scotia-Maine region of the Northwest

938 Atlantic Ocean. They used the $\delta^{15}\text{N}_{\text{SAA}}$ values of *Primnoa resedaeformis* coral as a proxy for
939 increasing nitrate levels in the region, associated with externally driven shifts in slope water source
940 partitioning over the last 100 years. Given that slope water circulation in the Scotia-Maine region is
941 linked with broader scale climate variability associated with the North Atlantic Oscillation (Loder et
942 al., 2001; Pershing et al., 2001), these authors concluded that changes in nitrate source partitioning
943 may be tied to recent, human-caused changes in global climate.

944 More recently, Sherwood et al. (2015) determined $\delta^{15}\text{N}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{AA}}$ values recorded in
945 the skeletons of the very long-lived (>1,000 years) deep-sea proteinaceous corals *Kulamanamana*
946 *haumea* collected from the Hawaiian Archipelago. After nearly a millennium of minor oscillations,
947 coral $\delta^{15}\text{N}_{\text{bulk}}$ values decreased dramatically in the last 150 years. Using $\delta^{15}\text{N}_{\text{phe}}$ as a proxy for
948 baseline isotopic composition, Sherwood et al. (2015) calculated the relative contribution of
949 N_2 -fixation to export production in the North Pacific Subtropical Gyre. They found that increasing
950 N_2 -fixation in the subtropical gyre recently observed in the modern instrumental record (Karl et al.,
951 1997, 2001) is a continuation of a much longer centennial-scale trend, resulting in a 17% to 27%
952 increase in N_2 -fixation since the end of the Little Ice Age and the onset of the Industrial Era. These
953 authors suggested that this increase in N_2 -fixation might be attributed to Northern Hemisphere
954 climate change since the end of the Little Ice Age (Wilson et al., 2006; Mann et al., 2008).

955 In a complementary study of *K. haumea* in the North Pacific Subtropical Gyre,
956 McMahon et al. (2015c) reconstructed the first high-resolution records of changing plankton
957 community composition over the past millennium, using the AA carbon isotope fingerprinting
958 approach of Larsen et al. (2009, 2013). This study revealed three major plankton regimes
959 corresponding to Northern Hemisphere climatic periods over the past 1,000 years. The most recent
960 regime, which began during the warming and stratification period following the end of the Little Ice
961 Age (1850 CE; Corno et al., 2007; Dore et al., 2008), was characterized by an increase of
962 approximately 47% in the contribution of exported POM from N_2 -fixing cyanobacteria. These data
963 support the growing body of evidence that the last 150 years in the North Pacific Subtropical Gyre
964 have seen a major, and likely unique shift in plankton community dynamics and nitrogen cycling
965 associated with the end of the Little Ice Age. These studies illustrate the power of CSIA-AA
966 approaches to reconstructing past ocean ecosystem dynamics and biogeochemical cycling.

967

968 **4.5. CSIA-AA as an indicator of organic matter source and degradation state**

969

970 4.5.1. Patterns in microbial $\delta^{15}\text{N}_{\text{AA}}$ variability

971 The majority of OM in natural environments is not in living organisms, but exists as
972 detrital OM (e.g., Hedges, 1992; Eglinton and Repeta, 2004). Thus, production, alteration, and
973 degradation of detrital OM are key components in biogeochemical cycles, especially for carbon

974 and nitrogen, and they also play important roles in ecosystems. AAs represent a major fraction of
975 nitrogenous detritus, and are vital in biogeochemical cycles of OM in various environments such as
976 ocean water columns (Cowie and Hedges, 1994; McCarthy et al., 1996), marine sediments (Keil et
977 al., 2000), and soils (Schulten and Schnitzer, 1997). Therefore, $\delta^{15}\text{N}_{\text{AA}}$ values and patterns also
978 represent novel indicators for the sources and degradation state of detrital OM, especially for
979 organic nitrogen. In contrast to CSIA-AA in animal ecology (Section 4), however, CSIA-AA
980 studies of detrital OM must consider not only food chain processes, but also the subsequent effects
981 of metabolism of chemotrophic microbes (both heterotrophs and chemoautotrophs) on $\delta^{15}\text{N}_{\text{AA}}$
982 values and patterns. This remains a frontier area of CSIA-AA applications, and exactly how $\delta^{15}\text{N}_{\text{AA}}$
983 patterns are altered by microbial processes remains an area of active research. Importantly, in
984 contrast to metazoans, the metabolic plasticity of microbes allows for multiple means of AA
985 acquisition, including *de novo* synthesis, salvage incorporation (*i.e.*, uptake and incorporation of
986 existing AAs into bacterial biomass), as well as selected resynthesis (*i.e.*, heterotrophic synthesis).
987 This metabolic diversity is likely the reason that observed microbial $\delta^{15}\text{N}_{\text{AA}}$ fractionation patterns
988 are substantially more complex than metazoans. Based on the literature results, we propose that
989 $\delta^{15}\text{N}_{\text{AA}}$ patterns resulting from microbial heterotrophy can be classified into four main categories,
990 and that these patterns can be used as a conceptual framework for interpreting $\delta^{15}\text{N}_{\text{AA}}$ values in
991 detrital OM. Patterns indicating different microbial metabolisms may include changes in TP ,
992 $\delta^{15}\text{N}_{\text{SAA}}$ values, and an additional parameter, ΣV . Here, ΣV is a proxy for total heterotrophic
993 resynthesis, and is defined as $\Sigma V = 1/n \sum \text{Abs}(\chi_{\text{AA}})$, where deviation of each TAA is $\chi = \delta^{15}\text{N}_{\text{AA}} -$
994 $\delta^{15}\text{N}$ of average Ala, Asp, Glu, Ile, Leu, and Pro, and n is the total number of TAAs used in the
995 calculation (McCarthy et al., 2007, Fig. 14).

996

997 Pattern 1: Algae-like $\delta^{15}\text{N}_{\text{AA}}$ patterns from *de novo* AA synthesis: Pure culture experiments with
998 microbes have shown that when chemotrophic microbes (*i.e.*, both heterotrophs and
999 chemoautotrophs, including Eukarya, Bacteria, and Archaea) synthesize AAs *de novo* from
1000 inorganic nitrogen, the relative $\delta^{15}\text{N}_{\text{AA}}$ pattern normalized to $\delta^{15}\text{N}_{\text{Glu}}$ is very similar to that of algae
1001 (Fig. 14a, Yamaguchi, 2013). Applying standard formulas discussed above on such material
1002 indicates low TP values and low ΣV values, just as in fresh algal biosynthetic (Yamaguchi, 2013).
1003 Just as for algal production, the absolute $\delta^{15}\text{N}$ values depend on that of the nitrogen source and
1004 isotopic fractionation during uptake and synthesis of Glu (*e.g.*, Hoch et al., 1992; Fogel and
1005 Cifuentes, 1993; Chikaraishi et al., 2007; Ohkouchi and Takano, 2014), which is the sole source of
1006 most nitrogen in the other AAs (Bender, 2012).

1007 *De novo* synthesis of AAs from inorganic nitrogen by chemotrophic microbes might
1008 contribute greatly to detrital OM in some environments. For example, in environments with
1009 carbon-rich OM and abundant inorganic nitrogen, such as forest litter, some heterotrophic microbes

1010 use inorganic nitrogen as the main nitrogen source for AA synthesis. Another example is
1011 environments where chemoautotrophic microbes are the dominant primary producers, such as
1012 submarine hydrothermal vents. The algae-like *de novo* $\delta^{15}\text{N}_{\text{AA}}$ pattern of chemotrophic microbes
1013 could be useful for explaining $\delta^{15}\text{N}_{\text{AA}}$ values and patterns of detrital OM in such environments,
1014 although the effect of microbial heterotrophy must also be considered (see pattern 2-4).

1015 These results show that most algae and chemotrophic microbes covering the three domains
1016 of life generally show similar $\delta^{15}\text{N}_{\text{AA}}$ patterns for *de novo* AA synthesis. However, some differences
1017 in specific AAs may exist between domains or between microbial species (McCarthy et al., 2013;
1018 Yamaguchi, 2013; Maki et al., 2014). To use $\delta^{15}\text{N}_{\text{AA}}$ patterns as indicators for specific microbial
1019 groups, further microbial culture experiments are needed to verify interspecies differences and to
1020 understand the variation of microbial $\delta^{15}\text{N}_{\text{AA}}$ values mechanistically in terms of AA metabolic
1021 pathways.

1022
1023 Pattern 2: Animal-like changes in $\delta^{15}\text{N}_{\text{AA}}$ values (increases in *TP* value): Heterotrophic microbes can
1024 use existing AAs in environments (sometimes specific AAs) by metabolizing AAs as carbon and
1025 nitrogen sources for resynthesis, or by salvage incorporation. The enzymatic degradation processes
1026 of AAs, such as deamination or transamination, cause nitrogen isotopic fractionation (e.g., Macko
1027 and Estep, 1984; Macko et al. 1986). Experiments using axenic cultures of heterotrophic microbes
1028 across the three domains (Eukarya, Bacteria, and Archaea) have shown that the pattern of TDF_{AA}
1029 between microbial biomass and substrates (free AAs or complex media containing proteins) can be
1030 similar to that of animals, as evidenced by large positive TDF_{TAA} (e.g., +6 to +8‰ in Glu) and the
1031 small TDF_{Phe} (~0‰) (Steffan et al., 2015a; Yamaguchi, 2013, Fig. 14b). These results suggest that
1032 when microbes incorporate AAs from the environment, the AAs in the microbial biomass and the
1033 microbially produced OM show higher *TP* values, which would be distinct from the algae-like *de*
1034 *novo* $\delta^{15}\text{N}_{\text{AA}}$ pattern (pattern 1).

1035 However, the mechanisms behind the apparently similar *TDF* patterns may differ between
1036 animals and heterotrophic microbes, because these organisms often use different metabolic AA
1037 pathways. For example, the proposed mechanism for the small, stable TDF_{Phe} in animals *via* the
1038 phenylalanine hydroxylase pathway (Chikaraishi et al., 2007) would not apply to many microbes,
1039 which can synthesize Phe, and do not have this pathway (Yamaguchi, 2013). Alternatively, the small
1040 TDF_{Phe} in heterotrophic microbes may arise from the high energetic cost of Phe biosynthesis, which
1041 would strongly suppress Phe synthesis and degradation and result in the salvage incorporation of Phe
1042 from the culture media (Yamaguchi, 2013; Akashi and Gojobori, 2002). To better understand the
1043 mechanisms of the heterotrophic changes in microbial $\delta^{15}\text{N}_{\text{AA}}$ values, we propose that it is important
1044 to examine the AAs that were not analyzed in these first culture experiments (e.g., Met, Thr, Tyr,
1045 etc.), and to directly compare $\delta^{15}\text{N}_{\text{AA}}$ patterns in microbes that have different metabolic AA

1046 pathways, as has been done for $\delta^{13}\text{C}_{\text{AA}}$ (Scott et al., 2006).

1047

1048 Pattern 3: Scattered changes in $\delta^{15}\text{N}_{\text{AA}}$ values (large increase in ΣV value): Although pure culture
1049 experiments have demonstrated that the heterotrophic microbes can show $\delta^{15}\text{N}_{\text{AA}}$ changes similar to
1050 those of animals, the microbial $\delta^{15}\text{N}_{\text{AA}}$ changes in natural environments may also show patterns that
1051 are more scattered (Fig. 14c). For example, incubation experiments of natural marine microbes with
1052 algal DOM showed that microbial DOM reworking caused $\delta^{15}\text{N}_{\text{AA}}$ changes that were more scattered
1053 than those observed in pure culture experiments and in animals (Calleja et al., 2013). Large ^{15}N
1054 enrichment was observed for some AAs, such as Gly (>10‰), and small ^{15}N enrichment was
1055 observed for some TAAs such as Ile (~0‰). Similarly, incubation of plant materials in salt marsh
1056 sediments also showed highly scattered $\delta^{15}\text{N}_{\text{AA}}$ changes caused by microbial OM reworking and
1057 replacement, but little change in Phe (Fogel and Tuross, 1999). Microcosm experiments of an alga
1058 and a phagotrophic protist showed a scattered *TDF* pattern in the protist (e.g., +8‰ for Ala and ~0‰
1059 for Glu, Gutierrez-Rodriguez et al., 2014).

1060 These “scattered” $\delta^{15}\text{N}_{\text{AA}}$ changes caused by heterotrophic microbial resynthesis of only
1061 selected AAs can be quantified by relative ΣV values, (as defined above by the average deviation in
1062 the $\delta^{15}\text{N}$ values of the trophic AAs, Ala, Asp, Glu, Ile, Leu, and Pro; McCarthy et al., 2007).
1063 Changes in ΣV values caused by microbial OM reworking may also be decoupled from changes in
1064 $TP_{\text{Glu/Phe}}$ values, because the microbially-mediated changes in $\delta^{15}\text{N}_{\text{Glu}}$ values may be small in some
1065 settings, relative to changes in other source AA (e.g., Gutierrez-Rodriguez et al., 2014). Thus, large
1066 increase of ΣV values decoupled with $TP_{\text{Glu/Phe}}$ values has been hypothesized as a characteristic
1067 marker of microbial reworking (McCarthy et al., 2007). In contrast, while ΣV values also increase in
1068 animal trophic steps to some extent, the increase of ΣV values in animals are relatively small and
1069 usually coupled with increase of $TP_{\text{Glu/Phe}}$ values (McCarthy et al., 2007). The AAs used to calculate
1070 ΣV values may also vary, because some AAs often cannot be measured depending on analytical
1071 protocols and the status of samples. Therefore, relative inter-sample trends in ΣV values would be
1072 typically interpreted as diagnostic for relative degradation, whereas exact values are only generally
1073 comparable among studies.

1074 Mechanisms for selected AA $\delta^{15}\text{N}$ changes, leading to “scattered” $\delta^{15}\text{N}_{\text{AA}}$ changes linked
1075 to microbial reworking of OM in natural settings are still poorly understood, but we suggest several
1076 hypotheses. First, the quality of OM substrates, particularly AA content and AA imbalances between
1077 substrates and microbial biomass, may be an important factor controlling the $\delta^{15}\text{N}_{\text{AA}}$ changes by
1078 heterotrophic microbes, as has been suggested for animals (Chikaraishi et al., 2015; McMahon et al.,
1079 2015a, see Section 3.2.1). For example, substantial effects of the C:N ratio (i.e., AA content) of
1080 substrates on the microbial $\delta^{15}\text{N}_{\text{AA}}$ patterns were reported in microbial culture experiments using a
1081 single AA as the nitrogen source (Maki et al., 2014). Second, a mixture of *de novo* AA synthesis

1082 from inorganic nitrogen, coupled with direct AA incorporation from the environment (*i.e.*,
1083 combination of patterns 1 and 2) could also cause scattered $\delta^{15}\text{N}_{\text{AA}}$ values, due to selective microbial
1084 resynthesis of specific AAs. This mixed metabolism may be particularly important in settings with
1085 abundant available inorganic nitrogen. Third, the diversity of microbial AA metabolic pathways
1086 itself could also be a cause of the variation in $\delta^{15}\text{N}_{\text{AA}}$ patterns. Finally, while only internal processes
1087 within microbial cells are considered in the above three hypothesis, mixing between
1088 microbially-produced OM and residue of original substrate also needs to be considered for
1089 reworking of detrital OM. Because the patterns of $\delta^{15}\text{N}_{\text{AA}}$ fractionation may be different between
1090 intercellular and extracellular processes (see pattern 4), mixing of the two different OM pools could
1091 complicate $\delta^{15}\text{N}_{\text{AA}}$ patterns. To use the ΣV value properly as an indicator of heterotrophic microbial
1092 OM reworking, it is important to reproduce the scattered $\delta^{15}\text{N}_{\text{AA}}$ changes in highly controlled culture
1093 experiments with heterotrophic microbes whose AA metabolic pathways are well characterized.
1094 Such future controlled experiments should particular address if ΣV changes can be linked to specific
1095 AA, whose $\delta^{15}\text{N}$ values change under specific conditions. In addition, for assessing the factors
1096 controlling ΣV changes, it is important to culture microbes with substrates containing varying AA
1097 contents and compositions, or with substrates containing both inorganic nitrogen and AAs.

1098
1099 Pattern 4: Similar $\delta^{15}\text{N}_{\text{TAA}}$ and $\delta^{15}\text{N}_{\text{SAA}}$ increases, possibly by extracellular protein hydrolysis: The
1100 last $\delta^{15}\text{N}_{\text{AA}}$ pattern that has been observed is very different from any others discussed: linked
1101 increases in $\delta^{15}\text{N}$ values for both TAAs and SAAs (including Phe), with similar amplitudes for all
1102 AA, possibly due to isotopic fractionation associated with extracellular protein hydrolysis to
1103 oligomers (Fig. 14d) (Hannides et al., 2013). To assimilate AAs in natural environments,
1104 heterotrophic microbes usually need to conduct extracellular hydrolysis to degrade proteins into
1105 small molecules such as free AAs or small peptides (Hoppe et al., 2002). If preferential cleavage of
1106 ^{14}N -C peptide bonds in proteins occurs during microbial extracellular hydrolysis, the residual AAs in
1107 the proteins should show ^{15}N enrichment (Silfer et al., 1992). Furthermore, if nitrogen isotopic
1108 fractionation during peptide bond hydrolysis is similar among peptide bonds between various AAs,
1109 there should be similar increases in $\delta^{15}\text{N}$ values for TAAs and SAAs. Hannides et al. (2013)
1110 proposed this mechanism to explain the $\delta^{15}\text{N}_{\text{AA}}$ values of suspended POM observed in the
1111 mesopelagic ocean (Section 4.5.2), noting that $\delta^{15}\text{N}_{\text{AA}}$ changes across all AAs were consistent with a
1112 simple Rayleigh distillation mechanism, suggesting an external (as opposed to metabolic)
1113 fractionation process. It has been suggested that extracellular protein hydrolysis by heterotrophic
1114 microbes plays an important role in the biogeochemical cycles in many environments (*e.g.*, Arnosti,
1115 2011); thus, the effect of this mechanism on the $\delta^{15}\text{N}_{\text{AA}}$ values of detrital OM might be critical in
1116 various environments.

1117 However, nitrogen isotope fractionation of AAs during peptide bond hydrolysis has been

1118 experimentally investigated only for the abiotic hydrolysis of glycylglycine (Silfer et al., 1992), and
1119 there has been no experimental study of changes in $\delta^{15}\text{N}_{\text{AA}}$ during peptide bond hydrolysis by
1120 microbes. Future experimental studies using various microbes or enzymes are needed to verify this
1121 hypothesized $\delta^{15}\text{N}_{\text{AA}}$ pattern resulting from extracellular protein hydrolysis. Such studies must
1122 carefully separate measurement of microbial biomass from partially-hydrolyzed substrate in order to
1123 isolate the origins of the patterns described above.

1124

1125 4.5.2. Case studies: Suspended particles in the ocean

1126 As discussed above Section 4.5.1, $\delta^{15}\text{N}_{\text{AA}}$ analysis of detrital OM can provide a direct
1127 molecular-level view of $\delta^{15}\text{N}_{\text{bulk}}$ values of OM. In the ocean, early studies documented large
1128 increases in $\delta^{15}\text{N}_{\text{bulk}}$ values of POM from the mesopelagic surface ocean (e.g., Saino and Hattori,
1129 1980; Altabet et al., 1991). Hannides et al. (2013) evaluated the mechanisms of a $\delta^{15}\text{N}_{\text{bulk}}$ increase by
1130 applying CSIA-AA to POM in the North Pacific Subtropical Gyre. Their key observation was one of
1131 large similar increases in $\delta^{15}\text{N}$ values of SAAs and TAAs between the surface and mesopelagic
1132 POM. This resulted in constant TP values of POM with depth. The ΣV values also remained low and
1133 stable with depth. Thus, they concluded that the inclusion of high- TP material or heterotrophic
1134 microbial biomass in the POM pool (i.e., patterns 2 and 3) is unlikely to be the mechanism of ^{15}N
1135 enrichment for mesopelagic POM in the North Pacific Subtropical Gyre. They also suggested that
1136 microbial utilization of ^{15}N -enriched nitrate in the midwater as a nitrogen source for *de novo* AA
1137 synthesis (i.e., contribution of pattern 1) is not likely to be a major contributor to the $\delta^{15}\text{N}$ depth
1138 trends of POM.

1139 Hannides et al. (2013) also proposed that isotopic fractionation associated with
1140 heterotrophic degradation, probably driven by extracellular hydrolysis of protein (pattern 4), controls
1141 the $\delta^{15}\text{N}_{\text{AA}}$ values of midwater POM. The smaller magnitude of ^{15}N enrichment in Lys, which is
1142 around half that of most AAs, is consistent with the proposed hydrolytic mechanism, because Lys
1143 was the only measured AA with both an amide and an amino nitrogen (Hannides et al., 2013).
1144 However, the values for Thr do not appear consistent with the extracellular protein hydrolysis
1145 hypothesis. The depth changes in $\delta^{15}\text{N}_{\text{Thr}}$ values were very small in the POM measured by Hannides
1146 et al. (2013). In contrast to Lys, there is no obvious explanation for the $\delta^{15}\text{N}_{\text{Thr}}$ values. There is no
1147 experimental data on the nitrogen isotopic effect on Thr during microbial heterotrophic processes;
1148 thus, future studies on $\delta^{15}\text{N}_{\text{Thr}}$ during microbial degradation, including extracellular protein
1149 hydrolysis and heterotrophic resynthesis, will be important to explain the anomalous $\delta^{15}\text{N}_{\text{Thr}}$
1150 signature in POM and to clarify POM transformation processes in the ocean.

1151 Comparing $\delta^{15}\text{N}_{\text{AA}}$ and $\delta^{15}\text{N}_{\text{bulk}}$ would also provide useful new information about the
1152 biogeochemical cycling of organic nitrogen, including nitrogen fractions other than AAs.
1153 Specifically, $\delta^{15}\text{N}$ values of total hydrolysable AAs ($\delta^{15}\text{N}_{\text{THAA}}$) can be used as a proxy for total

1154 proteinaceous $\delta^{15}\text{N}$ values, estimated as the molar-weighted average of individual $\delta^{15}\text{N}_{\text{AA}}$ values
1155 (McCarthy et al., 2013; Calleja et al., 2013; Batista et al., 2014). When concentrations of AAs and
1156 bulk nitrogen are known, $\delta^{15}\text{N}$ values of the nitrogen fraction other than THAA (non-THAA) can be
1157 calculated by $\delta^{15}\text{N}$ mass balance. Accurate quantification of AAs and bulk nitrogen is, however,
1158 essential for these mass-balance calculations, but has been absent from many past CSIA-AA studies.
1159 We suggest that the concentration of AAs and bulk nitrogen should be routinely reported in future
1160 CSIA-AA studies, to better understand the relationship between $\delta^{15}\text{N}$ values of THAA and
1161 non-THAA in organisms and detrital OM (e.g., Cowie and Hedges, 1992; Amelung and Zhang,
1162 2001).

1163

1164 **5. Future work and challenges**

1165 We have reviewed the current “state of the art” of using nitrogen isotopic composition of
1166 AAs for estimating the *TP* of organisms, as well as broader applications to terrestrial and marine
1167 ecology and biogeochemical cycling. The CSIA-AA method provides information on diet sources
1168 that is more precise than classical bulk isotope methods and is now rapidly expanding into a
1169 number of fields, such as biomagnification of toxic chemicals (e.g., polychlorinated biphenyls)
1170 through the food web (Ohkouchi et al., 2016), and nitrogen exchange between symbionts and host
1171 organisms (Maeda et al., 2012). Although the advantages of CSIA-AA for studying a wide range of
1172 ecosystems are clear, at the same time the methods remain relatively new, and will benefit greatly
1173 from further improvement and development. We suggest the following as being among the main
1174 problems which need to be addressed in future studies.

1175 1) A prerequisite for the wider application of this tool for accurately estimating *TP* is a
1176 robust knowledge of the magnitude of TDF_{AA} , especially, but not only, for well documented source
1177 and trophic amino acid pairings such as Phe and Glu. As discussed in Section 3.2, the most
1178 appropriate $A_{\text{Glu/Phe}}$ values, for instance, for calculating *TP* in specific situations is still open to
1179 debate. In some cases, the CSIA-AA approach based on current understanding of TDF_{AA} has not
1180 produced ecologically realistic *TP* values (e.g., penguins in Lorrain et al. (2009), elasmobranchs in
1181 Dale et al. (2011), dragonfish in Choy et al. (2012), killer whales in Matthews and Ferguson (2014),
1182 sperm whales in Ruiz-Cooley et al. (2014)). The following questions thus need to be addressed
1183 regarding the trophic discrimination of amino acids. A) Is the TDF_{AA} of a given AA value constant
1184 or variable across a wide variety of food webs? B) Do TDF_{AA} values decrease with increasing *TP*
1185 (Hetherington et al., 2016)? C) Are TDF_{AA} value more constant in the terrestrial environment than
1186 in the aquatic environment, as suggested by the work of Steffan et al. (2013, 2015a)? And most
1187 broadly, it will be critical to determine to what extent TDF_{AA} variations depend on the specific
1188 biochemistry and physiology of organisms and their diet, as suggested may be the case by the
1189 feeding experiments of McMahon et al. (2015a) and Chikaraishi et al. (2015). To answer these

1190 questions, further work focused on understanding the biochemical, physiological, and ecological
1191 mechanisms underlying TDF_{AA} variability is required.

1192 2) In natural environments, microorganisms play critical roles in the food web. Although
1193 several studies have examined explicitly aspects of these roles (*e.g.*, Steffan et al. 2017), the effects
1194 of microbial activity on the isotopic compositions of AAs require further evaluation. Knowledge of
1195 these effects is extremely important, particularly in terms of understanding complex
1196 microbially-driven nitrogen cycling in ocean and soil environments using CSIA-AA.

1197 3) It is still difficult to estimate precisely the TP of multivorous feeders that integrate
1198 aquatic and terrestrial food webs feeders such as humans. In some cases, such as [Naito et al. \(2010\)](#),
1199 the $\delta^{15}N_{Phe}$ value can be used to distinguish between aquatic and terrestrial food sources, whereas
1200 in other cases it cannot. Development of techniques which will help expand the application of
1201 CSIA-AA tools across food webs could open broad new applications in both ecological and
1202 archaeological contexts.

1203 4) Although to date most CSIA-AA studies have relied heavily on the isotopic
1204 compositions of just two AAs, Glu and Phe, to determine TP , we need a more holistic application
1205 of the technique, such as by embracing the diversity in TDF_{AA} in 1) above, to fully exploit the
1206 utility of AA data for interpreting the diet and physiology of organisms (*e.g.*, [Bradley et al., 2015](#);
1207 [Nielsen et al. 2015](#)).

1208 5) Currently, we know very little about how D-AAs affect $\delta^{15}N_{AA}$ values. Because
1209 D-AAs are subject to different metabolic pathways, they should have distinct isotopic
1210 compositions from L-AAs ([Engel and Macko, 1986](#); [Takano et al., 2010](#); [Chan, 2016](#)), which may
1211 affect the overall $\delta^{15}N_{AA}$ value, even if they are minor components.

1212 Finally, we note that in addition to nitrogen isotopic composition, carbon isotopic
1213 composition of AAs can provide an independent measure of sources and metabolic processes, and
1214 has immense potential to help resolve some of the challenges outlined above. Furthermore, recent
1215 advances in measuring the radiocarbon of AAs may also provide detailed information on carbon
1216 transfer from the environment to consumers. This latter technique may be especially useful for soil
1217 ecosystems, where old carbon potentially makes significant contributions to microbial substrates,
1218 and should also be helpful for adding chronological information to the food web, as well as for
1219 identifying the source of AAs from various pools. While such applications are beyond the scope of
1220 the current review, development of appropriate methods is ongoing (*e.g.*, [Marom et al., 2014](#);
1221 [Takano et al., 2015](#); [Bour et al., 2016](#)). Ultimately, combining CSIA-AA with such new tools
1222 offers the promise of extraordinarily high-resolution delineation of food webs in space and time, as
1223 well as the potential to quantify food web linkages between and within aquatic and terrestrial
1224 systems at a new level of precision.

1225

1226 **Acknowledgments**

1227 We dedicate this paper to the memory of John Hayes, who pioneered CSIA through the
1228 establishment of GC/IRMS. We thank JST-CREST for financially supporting the workshop. NO,
1229 ASJW, and YTY were supported by the Japan Society for the Promotion of Science. TL was
1230 supported by the Cluster of Excellence 80 "The Future Ocean" under the Deutsche
1231 Forschungsgemeinschaft (DFG).

1232

1233 **References**

- 1234 Akashi, H., Gojobori, T., 2002. Metabolic efficiency and amino acid composition in the proteomes
1235 of *Escherichia coli* and *Bacillus subtilis*. Proceedings of the National Academy of Sciences of
1236 the United States of America 99, 3695–3700.
- 1237 Altabet, M.A., Francois, R., 1994. Sedimentary nitrogen isotopic ratio as a recorder for surface
1238 ocean nitrate utilization. Global Biogeochemical Cycles 8, 103-116.
- 1239 Altabet, M.A., Deuser, W.G., Honjo, S., Stienen, C., 1991. Seasonal and depth-related changes in
1240 the source of sinking particles in the North Atlantic. Nature 354, 136-139.
- 1241 Altabet, M.A., Pilskaln, C., Thunell, R., Pride, C., Sigman, D., Chavez, F., Francois, R., 1999. The
1242 nitrogen isotope biogeochemistry of sinking particles from the margin of the Eastern North
1243 Pacific. Deep-Sea Research Part I 46, 655-679.
- 1244 Ambrose, S.H., Norr, L., 1993. Carbon isotopic evidence for routing of dietary protein to bone
1245 collagen, and whole diet to bone apatite carbonate: purified diet growth experiments. In:
1246 Lamber, J., G. Grupe, G. (Eds.), Molecular Archaeology of Prehistoric Human Bone.
1247 Springer-Verlag, Berlin, Germany, pp. 1-37.
- 1248 Amelung, W., Zhang X, 2001. Determination of amino acid enantiomers in soils. Soil Biology and
1249 Biochemistry 33, 553–562.
- 1250 Arnosti, C., 2011. Microbial Extracellular Enzymes and the Marine Carbon Cycle. Annual Reviews
1251 of Marine Science 3, 401–425.
- 1252 Bardgett, R., Cook, R., 1998. Functional aspects of soil animal diversity in agricultural grasslands.
1253 Applied Soil Ecology 10, 263-276.
- 1254 Batista, F.C., Ravelo, A.C., Crusius, J., Casso, M.A., McCarthy, M.D., 2014. Compound specific
1255 amino acid $\delta^{15}\text{N}$ in marine sediments: A new approach for studies of the marine nitrogen cycle.
1256 Geochimica et Cosmochimica Acta 142, 553-569.
- 1257 Bender, D.A., 2012. Amino Acid Metabolism, 3rd Edition. Wiley-Blackwell.
- 1258 Bengtsson, J., Setälä, H., Zheng, D., 1996. Food webs and nutrient cycling in soils: Interactions and
1259 positive feedbacks. In: Polis, G.A., Winemiller, K.O. (Eds.), Food webs: Integration of Patterns
1260 and Dynamics. Chapman & Hall, New York, pp. 30-38.
- 1261 Blum, J.D., Popp, B.N., Drazen, J.C., Choy, A.C., Johnson, M.W., 2013. Mercury isotope evidence

1262 for methylation below the mixed layer in the central North Pacific Ocean. *Nature Geoscience* 6,
1263 879-884.

1264 Bol, R., Ostle, N.J., Petzke, K.J., Watson, A., Cockburn, J., 1998. Amino acid $^{15}\text{N}/^{14}\text{N}$ analysis at
1265 natural abundances: A new tool for soil organic matter studies in agricultural systems. *Isotopes*
1266 in *Environmental and Health Studies* 34, 87-93.

1267 Bour, A.L., Walker, B.D., Broek, T.A., McCarthy, M.D., 2016. Radiocarbon analysis of individual
1268 amino acids: Carbon blank quantification for a small-sample high-pressure liquid
1269 chromatography purification method. *Analytical Chemistry* 88, 3521-3528.

1270 Bowen, G.J., 2010. Isoscapes: spatial pattern in isotopic biogeochemistry. *Annual Review of Earth*
1271 *and Planetary Science* 38, 161-187.

1272 Bowen, G.J., West, J.B., Vaughn, B.H., Dawson, T.E., Ehleringer, J.R., Fogel, M.L., Hobson, K.,
1273 Hoogewerff, J., Kendall, C., Lai, C.T., Miller, C.C., 2009. Isoscapes to address large-scale earth
1274 science challenges. *EOS, Transactions American Geophysical Union* 90, 109-110.

1275 Bradley, C.J., Wallsgrove, N.J., Choy, C.A., Drazen, J.C., Hoen, D.K., Hetherington, E.D., Popp,
1276 B.N., 2015. Trophic position estimates of marine teleosts using amino acid compound specific
1277 isotopic analysis. *Limnology and Oceanography: Methods* 13, 476-493.

1278 Broek, T.A.B., McCarthy, M.D., 2014. A new approach to $\delta^{15}\text{N}$ compound-specific amino acid
1279 trophic position measurements: preparative high pressure liquid chromatography technique for
1280 purifying underivatized amino acids for stable isotope analysis. *Limnology and Oceanography*,
1281 *Methods* 12, 840-852.

1282 Broek, T.A., Walker, B.D., Andersen, D.H., McCarthy, M.D., 2013. High-precision measurement of
1283 phenylalanine $\delta^{15}\text{N}$ values for environmental samples: A new approach coupling high-pressure
1284 liquid chromatography purification and elemental analyzer isotope ratio mass spectrometry.
1285 *Rapid Communications in Mass Spectrometry* 27, 2327-2337.

1286 Calleja, M.L., Batista, F., Peacock, M., Kudela, R., McCarthy, M.D., 2013. Changes in compound
1287 specific $\delta^{15}\text{N}$ amino acid signatures and D/L ratios in marine dissolved organic matter induced
1288 by heterotrophic bacterial reworking. *Marine Chemistry* 149, 32-44.

1289 Chan, H.S., Ohkouchi, N., Takano, Y., Chikaraishi, Y., 2016. Nitrogen isotope analysis of amino
1290 acid contents in carbonaceous chondrites Yamato 980115 and Allan Hills A77003. *Earth*,
1291 *Planet, and Space* 68:7 DOI 10.1186/s-40623-016-0382-8

1292 Cheng, H.H., Bremner, J.M., 1964. Variations of nitrogen-15 abundance in soils. *Science* 146,
1293 1574-1575.

1294 Chikaraishi, Y., Kashiyama, Y., Ogawa, N.O., Kitazato, H., Ohkouchi, N., 2007. Metabolic control
1295 of nitrogen isotope composition of amino acids in macroalgae and gastropods: implications for
1296 aquatic food web studies. *Marine Ecology Progress Series* 342, 85-90.

1297 Chikaraishi, Y., Ogawa, N.O., Kashiyama, Y., Takano, Y., Suga, H., Tomitani, A., Miyashita, H.,

- 1298 Kitazato, H., Ohkouchi, N., 2009. Determination of aquatic food-web structure based on
1299 compound-specific nitrogen isotopic composition of amino acids. *Limnology and*
1300 *Oceanography Method* 7, 740-750.
- 1301 Chikaraishi, Y., Ogawa, N.O., Ohkouchi, N., 2010. Further evaluation of the trophic level estimation
1302 based on nitrogen isotopic composition of amino acids. In: Ohkouchi, N. Tayasu, I., Koba, K.
1303 (Eds.), *Earth, Life, and Isotopes*. Kyoto University Press, Kyoto, pp. 37-51.
- 1304 Chikaraishi, Y., Ogawa, N.O., Doi, H., Ohkouchi, N., 2011. $^{15}\text{N}/^{14}\text{N}$ ratios of amino acids as a tool
1305 for studying terrestrial food webs: a case study of terrestrial insects (bees, wasps, and hornets).
1306 *Ecological Research* 26, 835-844.
- 1307 Chikaraishi, Y., Steffan, S.A., Ogawa, N.O., Ishikawa, N.F., Sasaki, Y., Tsuchiya, M., Ohkouchi, N.,
1308 2014. High-resolution food webs based on nitrogen isotopic composition of amino acids.
1309 *Ecology and Evolution* 4, 2423-2449.
- 1310 Chikaraishi, Y., Steffan, S.A., Takano, Y., Ohkouchi, N., 2015. Diet quality influences isotopic
1311 discrimination among amino acids in an aquatic vertebrate. *Ecology and Evolution* 5,
1312 2048-2059.
- 1313 Choi, B., Ha, S., Lee, J., Chikaraishi, Y., Ohkouchi, N., Shin, K-H., 2017. Trophic interaction among
1314 organisms in a seagrass meadow ecosystem as revealed by bulk $\delta^{13}\text{C}$ and amino acid $\delta^{15}\text{N}$
1315 analyses. *Limnology and Oceanography Method*, in press.
- 1316 Choy, C.A., Davison, P.C., Drazen, J.C., Flynn, A., Gier, E.J., Hoffman, J.C., McClain-Counts, J.P.,
1317 Miller, T.W., Popp, B.N., Ross, S.W., 2012. Global trophic position comparison of two
1318 dominant mesopelagic fish families (Myctophidae, Stomiidae) using amino acid nitrogen
1319 isotopic analyses. *PLoS ONE* 7, e50133.
- 1320 Choy, C.A., Popp, B.N., Hannides, C.C.S., Drazen, J.C., 2015. Trophic structure and food resources
1321 of epipelagic and mesopelagic fishes in the North Pacific Subtropical Gyre ecosystem inferred
1322 from nitrogen isotopic compositions. *Limnology and Oceanography* 60, 1156-1171.
- 1323 Clements, K.D., Raubenheimer, D., Choat, J.H., 2009. Nutritional ecology of marine herbivorous
1324 fishes: ten years on. *Functional Ecology* 23, 79-92.
- 1325 Clementz, M.T., Fordyce, R.E., Peek, S.L., Fox, D.L., 2014. Ancient marine isoscapes and isotopic
1326 evidence of bulk-feeding by Oligocene cetaceans. *Palaeogeography Palaeoclimatology*
1327 *Palaeoecology* 400, 28-40.
- 1328 Cline, J.D., Kaplan, I.R., 1975. Isotopic fractionation of dissolved nitrate during denitrification in the
1329 eastern tropical North Pacific Ocean. *Marine Chemistry* 3, 271-299.
- 1330 Coleman, D.C., 1996. Energetics of Detritivory and Microbivory in Soil in Theory and Practice. In:
1331 Polis, G.A., Winemiller, K.O. (Eds.), *Food webs: Integration of Patterns and Dynamics*.
1332 Chapman & Hall, New York, pp. 39-50.
- 1333 Corno, G., Karl, D.M., Church, M.J., Letelier, R.M., Lukas, R., Bidigare, R.R., Abbott, M.R., 2007.

- 1334 Impact of climate forcing on ecosystem processes in the North Pacific Subtropical Gyre.
1335 Journal of Geophysical Research Oceans 112, C04021.
- 1336 Cowie, G.L., Hedges, J.I., 1994. Biochemical indicators of diagenetic alteration in natural organic
1337 matter mixtures. Nature 369, 304-307.
- 1338 Currie, C.R., Poulsen, M., Mendenhall, J., Boomsma, J.J., Billen, J., 2006. Coevolved crypts and
1339 exocrine glands support mutualistic bacteria in fungus-growing ants. Science 311, 81-83.
- 1340 Dale, J.J., Wallsgrave, N.J., Popp, B.N., Holland, K., 2011. Foraging ecology and nursery habitat
1341 use of a benthic stingray from stomach content, bulk and amino acid stable isotope analysis.
1342 Marine Ecology Progress Series 433, 221-236.
- 1343 Davidson, D.W., Cook, S.C., Snelling, R.R., Chua, T.H., 2003. Explaining the abundance of ants in
1344 lowland tropical rainforest canopies. Science 300, 969-972.
- 1345 Decima, M., Landry, M. R., Popp, B.N., 2013. Environmental perturbation effects on baseline $\delta^{15}\text{N}$
1346 values and zooplankton trophic flexibility in the southern California Current Ecosystem.
1347 Limnology and Oceanography 58, 624-634.
- 1348 DeNiro, M.J., Epstein, S., 1981. Influence of diet on the distribution of nitrogen isotopes in animals.
1349 Geochimica et Cosmochimica Acta 45, 341-351.
- 1350 Dore, J.E., Brum, J.R., Tupas, L.M., Karl, D.M., 2002. Seasonal and interannual variability in
1351 sources of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean.
1352 Limnology and Oceanography 47, 1595-1607.
- 1353 Dore, J.E., Letelier, R.M., Church, M.J., Lucas, R., Karl, D.M., 2008. Summer phytoplankton
1354 blooms in the oligotrophic North Pacific Subtropical Gyre: Historical perspective and recent
1355 observations. Progress in Oceanography 76, 2-38.
- 1356 Druffel, E.R., 1997. Geochemistry of corals: Proxies of past ocean chemistry, ocean circulation, and
1357 climate. Proceedings of the National Academy of Science USA 94, 8354-8361.
- 1358 Eglinton, T.I., Repeta, D.J., 2004. Organic matter in the contemporary ocean. In Treatise on
1359 Geochemistry, 2nd edition. Elsevier. pp. 151-189.
- 1360 Engel, M.H., Macko, S.A., 1986. Stable isotope evaluation of the origins of amino acids in fossils.
1361 Nature 323, 531-533.
- 1362 Estrada, J.A., Lutcavage, M., Thorrold, S.R., 2005. Diet and trophic position of Atlantic bluefin tuna
1363 (*Thunnus thynnus*) inferred from stable carbon and nitrogen isotope analysis. Marine Biology
1364 147, 37-45.
- 1365 Federherr, E., Kupka, H.J., Cerli, C., Kalbitz, K., Dunsbach, R., Loos, A., de Reus, M. Lange, L.,
1366 Panetta, R.J., Schmidt, T.C. (2016) A novel tool for stable nitrogen isotope analysis in aqueous
1367 samples. Rapid Communications in Mass Spectrometry 30, 2537-2544.
- 1368 Florin, S.T., Felicetti, L.A., Robbins, C.T., 2011. The biological basis for understanding and
1369 predicting dietary-induced variation in nitrogen and sulphur isotope ratio discrimination.

- 1370 Functional Ecology 25, 519-526.
- 1371 Fogel, M.L., 1997. Biogeochemical record of ancient humans. *Organic Geochemistry* 27, 275–287.
- 1372 Fogel, M.L., Cifuentes, L.A., 1993. Isotope fractionation during primary production. In: Engel, M.H.,
1373 Macko, M.A. (Eds.) *Organic Geochemistry: Principles and Applications*. Plenum, New York,
1374 pp. 73-98.
- 1375 Fogel, M.L., Tuross, N., 1999. Transformation of plant biochemicals to geological macromolecules
1376 during early diagenesis. *Oecologia* 120, 336–346.
- 1377 Fogel, M.L., Tuross, N., Owsley, D.W., 1989. Nitrogen isotope tracers of human lactation in modern
1378 and archeological populations. *Carnegie Institution of Washington Yearbook* 88, 111-117.
- 1379 Fouillet, H., Juillet, B., Bos, C., Mariotti, F., Gaudichon, C., Benamouzig, R., Tomé, D., 2008.
1380 Urea-nitrogen production and salvage are modulated by protein intake in fed humans: results of
1381 an oral stable-isotope-tracer protocol and compartmental modeling. *The American Journal of*
1382 *Clinical Nutrition* 87, 1702-1714.
- 1383 Frisch, A.J., Ireland, M., Rizzari, J.R., Lönnstedt, O.M., Magnenat, K.A., Mirbach, C.E., Hobbs,
1384 J-P.A., 2016. Reassessing the trophic role of reef sharks as apex predators on coral reefs. *Coral*
1385 *Reefs* 35, 459-472.
- 1386 Fry, B., 2006. *Stable Isotope Ecology*. Springer, New York, 308 pp.
- 1387 Fuller, B.T., Fuller, J.L., Sage, N.E., Harris, D.A., O'Connell, T.C., Hedges, R.E.M., 2004. Nitrogen
1388 balance and $\delta^{15}\text{N}$: why you're not what you eat during pregnancy. *Rapid Communications in*
1389 *Mass Spectrometry* 18, 2889-2896.
- 1390 Fuller, B.T., Fuller, J.L., Harris, D., Hedges, R.E.M., 2006. Detection of breastfeeding and weaning
1391 in modern human infants with carbon and nitrogen stable isotope ratios. *American Journal of*
1392 *Physical Anthropology* 129, 279–293.
- 1393 Gaebler, O.H., Choitz, H.C., Vitti, T.G., Vukmirovich, R., 1963. Significance of N^{15} excess in
1394 nitrogenous compounds of biological origin. *Canadian Journal of Biochemistry and Physiology*
1395 41, 1089-1097.
- 1396 Gaebler, O.H., Vitti, T.G., Vukmirovich, R., 1966. Isotope effects in metabolism of ^{14}N and ^{15}N
1397 from unlabeled dietary proteins. *Canadian Journal of Biochemistry* 44, 1249-1257.
- 1398 Gelwicks, J.T., Hayes, J.M., 1990. Carbon-isotopic analysis of dissolved acetate. *Analytical*
1399 *Chemistry* 62, 535-539.
- 1400 Germain, L.R., Koch, P.L., Harvey, J., McCarthy, M.D., 2013. Nitrogen isotope fractionation in
1401 amino acids from harbor seals: implications for compound-specific trophic position calculations.
1402 *Marine Ecology Progress Series* 482, 265-277.
- 1403 Graham, B.S., Koch, P.L., Newsome, S.D., McMahon, K.W., Aurioles, D., 2009. Using isoscapes to
1404 trace the movements and foraging behavior of top predators in oceanic ecosystems. In: West, J.,
1405 Bowen, G.J., Dawson, T.E., Tu, K.P. (Eds.), *Isoscapes: Understanding Movement, Pattern and*

- 1406 Process on Earth Through Isotope Mapping. Springer, New York, pp. 299-318.
- 1407 Guilderson, T.P., McCarthy, M.D., Dunbar, R.B., Englebrecht, A., Roark, E.B., 2013. Late Holocene
1408 variations in Pacific surface circulation and biogeochemistry inferred from proteinaceous
1409 deep-sea corals. *Biogeosciences* 10, 6019-6028.
- 1410 Gutiérrez-Rodríguez, A., Décima, M., Popp, B.N., Landry, M.R., 2014. Isotopic invisibility of
1411 protozoan trophic steps in marine food webs. *Limnology and Oceanography* 59, 1590–1598.
- 1412 Hagen, E.M., McCluney, K.E., Wyant, K.A., Soykan, C.U., Keller, A.C., Luttermoser, K.C., Holmes,
1413 E.J., Moore, J.C., Sabo, J.L., 2012. A meta-analysis of the effects of detritus on primary
1414 producers and consumers in marine , freshwater , and terrestrial ecosystems. *Oikos* 121,
1415 1507-1515.
- 1416 Hannides, C.C., Popp, B.N., Landry, M.R., Graham, B.S., 2009. Quantification of zooplankton
1417 trophic position in the North Pacific Subtropical Gyre using stable nitrogen
1418 isotopes. *Limnology and Oceanography* 54, 50-61.
- 1419 Hannides, C.C.S., Popp, B.N., Choy, C.A., Drazen, J.C., 2013. Midwater zooplankton and
1420 suspended particle dynamics in the North Pacific Subtropical Gyre: A stable isotope
1421 perspective. *Limnology and Oceanography* 58, 1931-1946.
- 1422 Haraguchi, T.F., Uchida, M., Shibata, Y., Tayasu, I., 2013. Contributions of detrital subsidies to
1423 aboveground spiders during secondary succession, revealed by radiocarbon and stable
1424 isotope signatures. *Oecologia* 171, 935-944.
- 1425 Hare, P.E., Fogel, M.L., Stafford, T.W.Jr., Mitchell, A.D., Hoering, T.C., 1991. The isotopic
1426 composition of carbon and nitrogen in individual amino acids isolated from modern and fossil
1427 proteins. *Journal of Archaeological Science* 18, 277-292.
- 1428 Hayes, J.M., Freeman, K.H., Hoham, C.H., Popp, B.N., 1990. Compound-specific isotopic analyses,
1429 a novel tool for reconstruction of ancient biogeochemical processes. *Organic Geochemistry* 16,
1430 1115-1128.
- 1431 Hayes, J.M., 2001. Fractionation of carbon and hydrogen isotopes in biosynthetic processes.
1432 *Reviews in Mineralogy and Geochemistry* 43, 225-277.
- 1433 Heaton, T.H.E., Vogel, J.C., Chevallerie, G. von la, Collett, G., 1986. Climatic influence on the
1434 isotopic composition of bone nitrogen. *Nature* 322, 822-823.
- 1435 Hedges, J.I., 1992. Global biogeochemical cycles: progress and problems. *Marine Chemistry* 39,
1436 67-93.
- 1437 Hetherington, E.D., Olson, R.J., Drazen, J.C., Lennert-Cody, C.E., Balance, L.T., Kaufmann, R.S.,
1438 Popp, B.N., 2017. Spatial food-web structure in the eastern tropical Pacific Ocean based on
1439 compound-specific nitrogen isotope analysis of amino acids. *Limnology and Oceanography*, in
1440 press.
- 1441 Hill, T.M., Myrvold, C.R., Spero, H.J., Guilderson, T.P., 2014. Evidence for benthic–pelagic food

1442 web coupling and carbon export from California margin bamboo coral archives.
1443 Biogeosciences 11, 3845-3854.

1444 Hilting, A.K., Currin, C.A., Kosaki, R.K., 2013. Evidence for benthic primary production support of
1445 an apex predator-dominated coral reef food web. *Marine Biology* 160, 1681-1695.

1446 Hobson, K.A., Clark, R.G., 1992. Assessing Avian Diets Using Stable Isotopes II: Factors
1447 Influencing Diet-Tissue Fractionation. *The Condor* 94, 189-197.

1448 Hobson, K.A., Barnett-Johnson, R., Cerling, T., 2010. Using isoscapes to track animal migration. In:
1449 West, J., Bowen, G.J., Dawson, T.E., Tu, K.P. (Eds.), *Isoscapes: Understanding Movement,
1450 Pattern and Process on Earth Through Isotope Mapping*. Springer, New York, pp. 273-298.

1451 Hobson, K.A., Van Wilgenburg, S.L., Wassenaar, L.I., Larson, K., 2012. Linking hydrogen ($\delta^2\text{H}$)
1452 isotopes in feathers and precipitation: sources of variance and consequences for assignment to
1453 isoscapes. *PLoS ONE* 7, e35137.

1454 Hoch, M.P., Fogel M.L., Kirchman D.L., 1992. Isotope fractionation associated with ammonium
1455 uptake by a marine bacterium. *Limnology and Oceanography* 37, 1447-1459.

1456 Hoegh-Guldberg, O., Bruno, J.F., 2010. The impact of climate change on the world's marine
1457 ecosystems. *Science* 328, 1523-1528.

1458 Hoen, D.K., Kim, S.L., Hussey, N.E., Wallsgrove, N.J., Drazen, J.C., Popp, B.N., 2014. Amino acid
1459 ^{15}N trophic enrichment factors of four large carnivorous fishes. *Journal of Experimental Marine
1460 Biology and Ecology* 453, 76-83.

1461 Hoering, T.C., Ford, H.T., 1960. The isotope effect in the fixation of nitrogen by *Azotobacter*.
1462 *Journal of American Chemical Society* 82, 376-378.

1463 Hoppe, H.-G., Armosti, C., Herndl, G.J., 2002. Ecological significance of bacterial enzymes in the marine
1464 environment. In: Burns, R.G., Dick, R.P. (Eds.), *Enzymes in the Environment: Activity, Ecology,
1465 and Applications*. Marcel Dekker, New York, pp. 73–108.

1466 Hussey, N.E., MacNeil, M.A., Siple, M.C., Popp, B.N., Dudley, S.F.J., Fisk, A.T., 2015. Expanded
1467 Trophic complexity among large sharks. *Food Webs* 4, 1-7.

1468 Hyodo, F., Matsumoto, T., Takematsu, Y., Itioka, T., 2015. Dependence of diverse consumers on
1469 detritus in a tropical rain forest food web as revealed by radiocarbon analysis. *Functional
1470 Ecology* 29, 423–429.

1471 Ishikawa, N.F., Kato, Y., Togashi, H., Yoshimura, M., Yoshimizu, C., Okuda, N., Tayasu, I., 2014.
1472 Stable nitrogen isotopic composition of amino acids reveals food web structure in stream
1473 ecosystems. *Oecologia* 175, 911-922.

1474 Ishikawa, N.F., Hayashi, F., Sasaki, Y., Chikaraishi, Y., Ohkouchi, N., 2017. Trophic discrimination
1475 factor of nitrogen isotopes within amino acids in the dobsonfly *Protohermes grandis*
1476 (Megaloptera: Corydalidae) larvae in a controlled feeding experiment. *Ecology and Evolution*,
1477 in press.

- 1478 Jaeger, A., Lecomte, V.J., Weimerskirch, H., Richard, P., Cherel, Y., 2010. Seabird satellite tracking
1479 validates the use of latitudinal isoscapes to depict predators' foraging areas in the Southern
1480 Ocean. *Rapid Communications in Mass Spectrometry* 24, 3456-3460.
- 1481 Karl, D.M., Letelier, R.M., Tupas, L., Dore, J., Christian, J., Hebel, D., 1997. The role of nitrogen
1482 fixation in biogeochemical cycling in the subtropical North Pacific Ocean. *Nature* 388,
1483 533-538.
- 1484 Karl, D.M., Bidigare, R.R., Letelier, R.M., 2001. Long-term changes in plankton community
1485 structure and productivity in the North Pacific Subtropical Gyre: the domain shift hypothesis.
1486 *Deep Sea Research Part II* 48, 1449-1470.
- 1487 Keil, R.G., Tsamakis, E., Hedges, J.I., 2000. Early diagenesis of particulate amino acids in marine
1488 systems. In: Goodfriend, G.A., Collins, M.J., Fogel, M.L., Macko, S.A., Wehmiller, J.F. (Eds.),
1489 *Perspectives in Amino Acid and Protein Geochemistry*. Oxford University Press, New York. pp.
1490 69-82.
- 1491 Kienast, S.S., Calvert, S.E., Pedersen, T.F., 2002. Nitrogen isotope and productivity variations along
1492 the northeast Pacific margin over the last 120 kyr: Surface and subsurface paleoceanography.
1493 *Paleoceanography* 17, 1055, doi:10.1029/2001PA000650
- 1494 Koch, P.L., 2007. Isotopic study of the biology of modern and fossil vertebrates. *Stable isotopes in*
1495 *Ecology and Environmental Science* 2, 99-154.
- 1496 Krueger, H.W., Sullivan, C.H., 1984. Models for carbon isotope fractionation between diet and bone.
1497 In: Turnland, J.F., Johnson, P.E. (Eds.), *Stable Isotopes in Nutrition ACS Symposium Series*,
1498 258. American Chemical Society, Washington D.C., pp. 205-222.
- 1499 Kruger, B.R., Werne, J.P., Branstrator, D.K., Hrabik, T.R., Chikaraishi, Y., Ohkouchi, N., Minor,
1500 E.C., 2016. Organic matter transfer in Lake Superior's food web: Insights from bulk and
1501 molecular stable isotope and radiocarbon analyses. *Limnology and Oceanography* 61, 149-164.
- 1502 Kruse, S., Pakhomov, E.A., Hunt, B., Chikaraishi, Y., Ogawa, N.O., Bathmann, U., 2015.
1503 Uncovering the trophic relationship between *Themisto gaudichaudii* and *Salpa thompsoni* in the
1504 Antarctic Polar Frontal Zone. *Marine Ecology Progress Series* 529, 63-74.
- 1505 Ku, H.H., 1966. Notes on the use of propagation of error formulas. *Journal of Research of the*
1506 *National Bureau of Standards* 70, 263-273.
- 1507 Larsen, T., Taylor, D.L., Leigh, M.B., O'Brien, D.M., 2009. Stable isotope fingerprinting: a novel
1508 method for identifying plant, fungal, or bacterial origins of amino acids. *Ecology* 90,
1509 3526-3535.
- 1510 Larsen, T., Ventura, M., Andersen, N., O'Brien, D.M., Piatkowski, U., McCarthy, M.D., 2013.
1511 Tracing carbon sources through aquatic and terrestrial food webs using amino acid stable
1512 isotope fingerprinting. *PloS ONE* 8, e73441.
- 1513 Larsen, T., Pollierer, M.M., Holmstrup, M., D'Annibale, A., Maraldo, K., Andersen, N., Eriksen, J.,

1514 2016a. Substantial nutritional contribution of bacterial amino acids to earthworms and
 1515 enchytraeids: A case study from organic grasslands. *Soil Biology and Biochemistry* 99, 21-27.
 1516 Larsen, T., Ventura, M., Maraldo, K., Triadó-Margarit, X., Casamayor, E.O., Wang, Y.V., Andersen,
 1517 N., O'Brien, D.M., 2016b. The dominant detritus-feeding invertebrate in Arctic peat soils
 1518 derives its essential amino acids from gut symbionts. *Journal of Animal Ecology* 85,
 1519 1275-1285.
 1520 Loder, J.W., Shore, J.A., Hannah, C.G., Petrie, B.D., 2001. Decadal-scale hydrographic and
 1521 circulation variability in the Scotia-Maine region. *Deep-Sea Research Part II* 48, 3-35.
 1522 Lorrain, A., Graham, B., Menard, F., Popp, B.N., Bouillon, S., Van Breugel, P., Cherel, Y., 2009.
 1523 Nitrogen and carbon isotope values of individual amino acids: a tool to study foraging ecology
 1524 of penguins in the Southern Ocean. *Marine Ecology Progress Series* 391, 293-306.
 1525 Lorrain, A., Graham, B., Popp, B.N., Allain, V., Olson, R.J., Hunt, B.P.V., Potier, M., Fry, B.,
 1526 Galvan-Magna, F., Menkes, C.E.R., Kaehler, S., Menard, F., 2015. Nitrogen isotopic baselines
 1527 and implications for estimating foraging habitat and trophic position of yellowfin tuna in the
 1528 Indian and Pacific Oceans. *Deep-Sea Research II* 113, 188-198.
 1529 MacKenzie, K.M., Palmer, M.R., Moore, A., Ibbotson, A.T., Beaumont, W.R., Poulter, D.J.,
 1530 Trueman, C.N., 2011. Locations of marine animals revealed by carbon isotopes. *Scientific*
 1531 *Reports* 1:21 doi:10.1038/srep00021.
 1532 Macko, S., Estep, M., 1984. Microbial alteration of stable nitrogen and carbon isotopic compositions
 1533 of organic matter. *Organic Geochemistry* 6, 787-790.
 1534 Macko, S.A., Fogel Estep, M.L., Engel, M.H., Hare, P.E., 1986. Kinetic fractionation of nitrogen
 1535 isotopes during amino acid transamination. *Geochimica et Cosmochimica Acta* 50, 2143-2146.
 1536 Macko S.A., Fogel, M.L., Hare, P.E., Hoering, T.C., 1987. Isotopic fractionation of nitrogen and
 1537 carbon in the synthesis of amino acids by microorganisms. *Chemical Geology* 65, 79-92.
 1538 Madigan, D.J., Baumann, Z., Fisher, N.S., 2012a. Pacific bluefin tuna transport Fukushima-derived
 1539 radionuclides from Japan to California. *Proceedings of the National Academy of Sciences USA*
 1540 109, 9483–9486.
 1541 Madigan, D.J., Litvin, S.Y., Popp, B.N., Carlisle, A.B., Farwell, C.J., Block, B.A., 2012b. Tissue
 1542 turnover rates and isotopic trophic discrimination factors in the endothermic teleost, Pacific
 1543 Bluefin tuna (*Thunnus orientalis*). *PLoS ONE* 7, e49220.
 1544 Madigan, D.J., Baumann, Z., Snodgrass, O.E., Ergul, H.A., Dewar, H., Fisher, N.S., 2013.
 1545 Radiocesium in Pacific bluefin tuna *Thunnus orientalis* in 2012 validates new tracer technique.
 1546 *Environmental Science and Technology* 2287-2294.
 1547 Madigan, D.J., Baumann, Z., Carlisle, A.B., Hoen, D.K., Popp, B.N., Dewar, H., Snodgrass, O.E.,
 1548 Block, B.A., Fisher, N.S., 2014. Reconstructing transoceanic migration patterns of Pacific
 1549 bluefin tuna using a chemical tracer toolbox. *Ecology* 95, 1674-1683.

- 1550 Madigan D.J., Chiang W-C., Wallsgrove N.J., Popp B.N., Kitagawa T., Choy C.A., Tallmon J.,
1551 Ahmed N., Fisher N.S., Sun C., 2016. Intrinsic tracers reveal recent foraging ecology of giant
1552 Pacific bluefin tuna at their primary spawning grounds. *Marine Ecology Progress Series* 553,
1553 253-266.
- 1554 Maeda, T., Hirose, E., Chikaraishi, Y., Kawato, M., Takishita, K., Yoshida, T., Verbruggen, H.,
1555 Tanaka, J., Shimamura, S., Takai, Y., Tsuchiya, M., Iwai, K., Maruyama, T., 2012. Algivore or
1556 phototroph? *Plakobranchnus ocellatus* (Gastropoda) continuously acquires kleptoplasts and
1557 nutrition from multiple algal species in nature. *PloS ONE* 7, e42024.
- 1558 Maki, K., Ohkouchi, N., Chikaraishi, Y., Fukuda, H., Miyajima, T., Nagata, T., 2014. Influences of
1559 nitrogen substrates and substrate C:N ratios on the nitrogen isotopic composition of amino acids
1560 from the marine bacterium *Vibrio harveyi*. *Geochimica et Cosmochimica Acta* 140, 521-530.
- 1561 Mann, M.E., Zhang, Z., Hughes, M.K., Bradley, R.S., Miller, S.K., Rutherford, S., Ni, F., 2008.
1562 Proxy-based reconstructions of hemispheric and global surface temperature variations over the
1563 past two millennia. *Proceedings of the National Academy of Sciences USA* 105, 13252-13257.
- 1564 Marom, A., McCullagh, S.O., Higham, T.F.G., Sinitsyn, A.A., Hedges, R.E.M., 2012. Single amino
1565 acid radiocarbon dating of Upper Paleolithic modern humans. *Proceedings of National*
1566 *Academy of Science USA* 109, 6878-6881.
- 1567 Martinez del Rios, C., Wolf, N., Carlton, S.A., Gunnes, L.Z., 2009. Isotopic ecology ten years after a
1568 call for more laboratory experiments. *Biological Reviews* 84, 91-111.
- 1569 Matthews, C.J.D., Ferguson, S.H., 2014. Spatial segregation and similar trophic-level diet among
1570 eastern Canadian Arctic/north-west Atlantic killer whales inferred from bulk and compound
1571 specific isotopic analysis. *Journal of Marine Biological Association of the United Kingdom* 94,
1572 1343-1355.
- 1573 McCarthy, M., Hedges, J., Benner R., 1996. Major biochemical composition of dissolved high
1574 molecular weight organic matter in seawater. *Marine Chemistry* 55, 281-297.
- 1575 McCarthy, M.D., Benner, R., Lee, C., Fogel, M.L., 2007. Amino acid nitrogen isotopic fractionation
1576 patterns as indicators of heterotrophy in plankton, particulate, and dissolved organic matter.
1577 *Geochimica et Cosmochimica Acta* 71, 4727-2744.
- 1578 McCarthy, M.D., Lehman, J., Kudela, R., 2013. Compound-specific amino acid $\delta^{15}\text{N}$ patterns in
1579 marine algae: Tracer potential for cyanobacterial vs. eukaryotic organic nitrogen sources in the
1580 ocean. *Geochimica et Cosmochimica Acta* 103, 104-120.
- 1581 McClelland, J.W., Montoya, J.P., 2002. Trophic relationships and the nitrogen isotopic composition
1582 of amino acids in plankton. *Ecology* 83, 2173-2180.
- 1583 McCutchan, J.H.Jr, Lewis, W.M., Kendall, C., McGrath, C.C., 2003. Variation in trophic shift for stable
1584 isotope ratios of carbon, nitrogen, and sulfur. *Oikos* 102, 378-390.
- 1585 McMahan, K.W., McCarthy, M.D., 2016. Embracing variability in amino acid $\delta^{15}\text{N}$ fractionation:

1586 Mechanisms, implications, and applications for trophic ecology. *Ecosphere* 7(12), e01511

1587 McMahon, K.W., Hamady, L., Thorrold, S.R., 2013a. Ocean ecogeochemistry – A review.

1588 *Oceanography and Marine Biology - an Annual Review* 51, 327-374.

1589 McMahon, K.W., Hamady, L., Thorrold, S.R., 2013b. A review of ecogeochemistry approaches to

1590 estimating movements of marine animals. *Limnology and Oceanography* 58, 697-714.

1591 McMahon, K.W., Elsdon, T., Thorrold, S.R., McCarthy, M., 2015a. Trophic discrimination of

1592 nitrogen stable isotopes in amino acids varies with diet quality in a marine fish. *Limnology and*

1593 *Oceanography* 60, 1076-1087.

1594 McMahon, K.W., Polito, M., Abel, S., McCarthy, M.D., Thorrold, S.R., 2015b. Carbon and nitrogen

1595 isotope fractionation of amino acids in an avian marine predator, the gentoo penguin

1596 (*Pygoscelis papua*). *Ecology and Evolution* 5, 1278-1290.

1597 McMahon, K.W., McCarthy, M.D., Sherwood, O.A., Larsen, T., Guilderson, T.P., 2015c.

1598 Millennial-scale plankton regime shifts in the subtropical North Pacific Ocean. *Science*, 350,

1599 1530-1533.

1600 Merritt, D.A., Hayes, J.M., 1994. Nitrogen isotopic analyses by isotope-ratio-monitoring gas

1601 chromatography/mass spectrometry. *Journal of the American Society for Mass Spectrometry* 5,

1602 387-397.

1603 Metges, C.C., Petzke, K.J., 1997. Measurement of $^{15}\text{N}/^{14}\text{N}$ isotopic composition in individual plasma

1604 free amino acids of human adults at natural abundance by gas chromatography-combustion

1605 isotope ratio mass spectrometry. *Analytical Biochemistry* 247, 158-164.

1606 Metges, C.C., Petzke, K.-J., Henning, U., 1996. Gas chromatography/combustion/isotope ratio mass

1607 spectrometric composition of *N*-acetyl- and *N*-pivaloyl amino acid esters to measure ^{15}N isotopic

1608 abundances in physiological samples: a pilot study on amino acid synthesis in the upper

1609 gastro-intestinal tract of minipigs. *Journal of Mass Spectrometry* 31, 367-376.

1610 Mill, A.C., Pinnegar, J.K., Polunin, N.V.C., 2007. Explaining isotope trophic-step fractionation: why

1611 herbivorous fish are different. *Functional Ecology* 21, 1137-1145.

1612 Miller, M.J., Chikaraishi, Y., Ogawa, N.O., Yamada, Y., Tsukamoto, K., Ohkouchi, N., 2012. A low

1613 trophic position of Japanese eel larvae indicates feeding on marine snow. *Biology Letters* 9,

1614 DOI: 10.1098/rsbl.2012.0826

1615 Minagawa, M., Wada, E., 1984. Stepwise enrichment of ^{15}N along food chains. Further evidence and

1616 the relation between ^{15}N and animal age. *Geochimica et Cosmochimica Acta* 48, 1135-1140.

1617 Miura, K., Goto, A.S., 2012. Stable nitrogen isotopic fractionation associated with transamination of

1618 glutamic acid to aspartic acid: implications for understanding ^{15}N trophic enrichment in

1619 ecological food webs. *Researches in Organic Geochemistry* 28, 13-17.

1620 Miyachi, S., Mayahara, T., Tsushima, K., Sasada, K., Kohno, E., Ogawa, N.O., Chikaraishi, Y.,

1621 Ohkouchi, N., 2015. Approach to determine individual trophic level and the difference in food

1622 sources of Japanese anchovy *Engraulis japonicus* in Sagami Bay, based on compound-specific
1623 nitrogen stable isotope analysis of amino acids. *Fisheries Science*, 81, 1053-1062.

1624 Miyake, Y., Wada, E., 1967. The abundance ratio of $^{15}\text{N}/^{14}\text{N}$ in marine environments. *Records in*
1625 *Oceanographic Works in Japan* 9, 32-53.

1626 Moore, J.C., de Ruiter, P.C., 2012. *Energetic Food Webs*. Oxford University Press, Oxford, 344 pp.

1627 Naito, Y.I., Chikaraishi, Y., Ohkouchi, N., Mukai, H., Shibata, Y., Honch, N. V., Dodo, Y., Ishida,
1628 H., Amano, T., Ono, H., Yoneda, M., 2010a. Dietary reconstruction of the Okhotsk Culture of
1629 Hokkaido, Japan, based on nitrogen composition of amino acids: implications for correction of
1630 ^{14}C marine reservoir effects on human bone. *Radiocarbon* 52, 671-681.

1631 Naito, Y.I., Honch, N.V., Chikaraishi, Y., Ohkouchi, N., Yoneda, M., 2010b. Quantitative evaluation
1632 of marine protein contribution in ancient diets based on nitrogen isotope ratios of individual
1633 amino acids in bone collagen: An investigation at the Kitakogane Jomon site. *American Journal*
1634 *of Physical Anthropology* 143, 31-40.

1635 Naito, Y.I., Chikaraishi, Y., Ohkouchi, N., Drucker, D.G., Bocherens, H., 2013a. Nitrogen isotopic
1636 composition of collagen amino acids as an indicator of aquatic resource consumption: insights
1637 from Mesolithic and Epipalaeolithic archaeological sites in France. *World Archaeology* 45,
1638 338-359.

1639 Naito, Y.I., Chikaraishi, Y., Ohkouchi, N., Yoneda, M., 2013b. Evaluation of carnivory in inland
1640 Jomon hunter-gatherers based on nitrogen isotopic compositions of individual amino acids in
1641 bone collagen. *Journal of Archaeological Science* 40, 2913-2923.

1642 Naito, Y.I., Bocherens, H., Chikaraishi, Y., Drucker, D.G., Wißing, C., Yoneda, M., Ohkouchi, N.,
1643 2016a. An overview of methods used for the detection of aquatic resource consumption by
1644 humans: compound-specific delta N-15 analysis of amino acids in archaeological materials.
1645 *Journal of Archaeological Science* 6, 720-732.

1646 Naito, Y.I., Chikaraishi, Y., Drucker, D.G., Ohkouchi, N., Semal, P., Wißing, C., Bocherens, H.,
1647 2016b. Ecological niche of Neanderthals from Spy Cave revealed by nitrogen isotopes of
1648 individual amino acids in collagen. *Journal of Human Evolution* 93, 82-90.

1649 Naito, Y.I., Germonpré, M., Chikaraishi, Y., Ohkouchi, N., Drucker, D.G., Hofreiter, M., Edwards,
1650 M.A., Hobson, K.A., Wißing, C., Bocherens, H., 2016. Evidence for herbivorous cave bears
1651 (*Ursus spelaeus*) in Goyet Cave, Belgium: implications for paleodietary reconstruction of fossil
1652 bears using amino acid $\delta^{15}\text{N}$ approaches. *Journal of Quaternary Science* 31, 598-606.

1653 Nakashita, R., Suzuki, Y., Akamatsu, F., Naito, Y.I., Sato-Hashimoto, M., Tsubota, T., 2011.
1654 Ecological application of compound-specific stable nitrogen isotope analysis of amino acids –
1655 A case study of captive and wild bears. *Researches in Organic Geochemistry* 27, 73-79.

1656 Nakazawa, T., Sakai, Y., Hsieh, C-H., Koitabashi, T., Tayasu, I., Yamamuro, N., Okuda, N., 2010.
1657 Is the relationship between body size and trophic niche position time-invariant in a predatory

1658 fish? first stable isotope evidence. PLoS ONE 5, e9120.

1659 Nielsen, J.M., Popp, B.N., Winder, M., 2015. Meta-analysis of amino acid stable nitrogen isotope
1660 ratios for estimating trophic position in marine organisms. *Oecologia* 178, 1-12.

1661 O'Brien, D.M., Fogel, M.L., Boggs, C.L., 2002. Renewable and nonrenewable resources: amino acid
1662 turnover and allocation to reproduction in Lepidoptera. *Proceedings of the National Academy
1663 of Sciences of the USA* 99, 4413–4418.

1664 O'Brien, D.M.O., Boggs, C.L., Fogel, M.L., 2004. Making eggs from nectar: The role of life history
1665 and dietary carbon turnover in butterfly reproductive resource allocation. *Oikos* 105, 279-291.

1666 Ogawa, N.O., Koitabashi, T., Oda, H., Nakamura, T., Ohkouchi, N., Wada, E., 2001. Fluctuations of
1667 nitrogen isotope ratio of gobiid fish specimens and sediments in Lake Biwa during the 20th
1668 century. *Limnology and Oceanography* 46, 1228-1236.

1669 Ogawa, N.O., Chikaraishi, Y., Ohkouchi, N., 2013. Trophic position estimates of formalin-fixed
1670 samples with nitrogen isotopic compositions of amino acids: an application to gobiid fish
1671 (Isaza) in Lake Biwa, Japan. *Ecological Research* 28, 697-702.

1672 Ohkouchi, N., Takano, Y., 2014. Organic nitrogen: Sources, fates, and chemistry. In: Falkowski,
1673 P.G., Freeman, K.H. (Eds.), *Treatise on Geochemistry*, 2nd edn. Elsevier, Amsterdam, pp.
1674 251-289.

1675 Ohkouchi, N., Tsuda, R., Chikaraishi, Y., Tanabe, K., 2013. A preliminary estimate of the trophic
1676 position of the deep-water ram's horn squid *Spirula spirula* based on the nitrogen isotopic
1677 composition of amino acids. *Marine Biology* 160, 773-779.

1678 Ohkouchi, N., Ogawa, N.O., Chikaraishi, Y., Tanaka, H., Wada, E., 2015. Biochemical and
1679 physiological bases for the application of carbon and nitrogen isotopes to the environmental and
1680 ecological studies. *Progress in Earth and Planetary Science* 2, doi:10.1186/s40645-015-0032-y.

1681 Ohkouchi, N., Shibata, H., Nomaki, H., Ogawa, N.O., Chikaraishi, Y., Goto, T., Fujikura, K.,
1682 Kitazato, H., 2016. A monitoring result of polychlorinated biphenyls (PCBs) in deep-sea
1683 organisms and sediments off Tohoku during 2012-2014: temporal variation and the relationship
1684 with the trophic position. *Journal of Oceanography* 72, 629-639.

1685 Olson, R.J., Popp, B.N., Graham, B.S., Lopez-Ibarra, G.A., Galvan-Magana, F., Lennert-Cody, C.E.,
1686 Bocanegra-Castillo, N., Wallsgrove, N.J., Gier, E., Alatorre-Ramirez, V., Balance, L.T., Fry, B.,
1687 2010. Food web inferences of stable isotope spatial patterns in copepods and yellowfin tuna in
1688 the pelagic eastern Pacific Ocean. *Progress in Oceanography* 86, 124-138.

1689 O'Malley, J.M., Drazen, J.C., Popp, B.N., Gier, E., Toonen, R.J., 2012. Spatial variability in growth and
1690 prey availability of lobsters in the northwestern Hawaiian Islands. *Marine Ecology Progress Series*
1691 449, 211-220.

1692 O'Reilly, C.M., Hecky, R.E., Cohen A.S., Plisnier, P.-D., 2002. Interpreting stable isotopes in food
1693 webs: recognizing the role of time averaging at different trophic levels, *Limnology and*

- 1694 Oceanography 47, 306-309.
- 1695 Papastamatiou, Y.P., Meyer, C.G., Kosaki, R.K., Wallsgrove, N.J., Popp, B.N., 2015. Movements
1696 and foraging of predators associated with mesophotic coral reefs and their potential for linking
1697 ecological habitats. Marine Ecology Progress Series 521, 155-170.
- 1698 Parwel, A., Ryhage, R., Wickman, F. E., 1957. Natural variations in the relative abundances of the
1699 nitrogen isotopes. Geochimica et Cosmochimica Acta 11, 165-170.
- 1700 Pershing, A.J., Greene, C.H., Hannah, C., Sameoto, D., Head, E., Mountain, D.G., Jossi, J.W.,
1701 Benfield, M.C., Reid, P.C., Durbin, T.G., 2001. Oceanographic responses to climate in the
1702 Northwest Atlantic Ocean. Oceanography 14, 76-82.
- 1703 Peterson, H., Luxton, M., 1982. A comparative analysis of soil fauna populations and their role in
1704 decomposition processes. Oikos 39, 287-388.
- 1705 Petzke, K.J., Boeing, H., Klaus, S., Metges, C.C., 2005. Carbon and nitrogen stable isotopic
1706 composition of hair protein and amino acids can be used as biomarkers for animal-derived
1707 dietary protein intake in humans. Nutrition Methodology 135, 1515-1520.
- 1708 Petzke, K.J., Feist, T., Fleig, W.E., Metges, C.C., 2006. Nitrogen isotopic composition in hair
1709 protein is different in liver cirrhotic patients. Rapid Communications in Mass Spectrometry 20,
1710 2973-2978.
- 1711 Petzke, K.J., Fuller, B.T., Metges, C.C., 2010. Advances in natural stable isotope ratio analysis of
1712 human hair to determine nutritional and metabolic status. Current Opinion Clinical Nutrition
1713 Metabolic Care 13, 532-540.
- 1714 Phillips, D.L., Gregg, J.W., 2001. Uncertainty in source partitioning using stable isotopes. Oecologia
1715 127, 171-179.
- 1716 Polis, G.A., 1991. Complex trophic interactions in deserts: an empirical critique of food-web theory.
1717 The American Naturalist 138, 123-155.
- 1718 Polis, G.A., Strong, D.R., 1996. Food web complexity and community dynamics. The American
1719 Naturalist 147, 813-846.
- 1720 Popp, B.N., Graham, B.S., Olson, R.J., Hannides, C.C.S., Lott, M.J., Lopez-Ibarra, G.A.,
1721 Galvan-Magana, F., Fry, B., 2007. Insight into the trophic ecology of yellowfin tuna, *Thunnus*
1722 *albacares*, from compound-specific nitrogen isotope analysis of proteinaceous amino acids. In:
1723 Dawson, T., Siegwolf, R. (Eds.), Stable Isotopes as Indicators of Ecological Change, Elsevier
1724 Academic Press, Terrestrial Ecology Series, pp. 173-190.
- 1725 Post, D.M., 2002. Using stable isotopes to estimate trophic position: models, methods, and
1726 assumptions. Ecology 83, 703-718.
- 1727 Ramos, R., González-Solís, J., 2012. Trace me if you can: the use of intrinsic biogeochemical
1728 markers in marine top predators. Frontiers in Ecology and the Environment 10, 258-266.
- 1729 Reitsemá, L.J., 2013. Beyond diet reconstruction: Stable isotope applications to human physiology,

1730 health, and nutrition. *American Journal of Human Biology* 25, 445–456.

1731 Reitsema, L.J., Muir, A.B., 2015. Growth velocity and weaning $\delta^{15}\text{N}$ “Dips” during ontogeny in
1732 *Macaca mulatta*. *American Journal of Physical Anthropology* 157, 347-357.

1733 Roark, E.B., Guilderson, T.P., Dunbar, R.B., Fallon, S.J., Mucciarone, D.A., 2009. Extreme
1734 longevity in proteinaceous deep-sea corals. *Proceedings of the National Academy of Sciences*
1735 of the USA 106, 5204-5208.

1736 Robbins, C.T., Felicetti, L.A., Sponheimer, M., 2005. The effect of dietary protein quality on
1737 nitrogen isotope discrimination in mammals and birds. *Oecologia* 144, 534-540.

1738 Robbins, C.T., Felicetti, L.A., Florin, S.T., 2010. The impact of protein quality on stable nitrogen
1739 isotope ratio discrimination and assimilated diet estimation. *Oecologia* 162, 571-579.

1740 Robinson, L.F., Adkins, J.F., Frank, N., Gagnon, A.C., Prouty, N.G., Roark, E.B., Van de Flierdt, T.,
1741 2014. The geochemistry of deep-sea coral skeletons: a review of vital effects and applications
1742 for palaeoceanography. *Deep-Sea Res Part II* 99, 184-198.

1743 Rolff, C., 2000. Seasonal variation in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of size-fractionated plankton at a coastal station
1744 in the northern Baltic Proper. *Marine Ecology Progress Series* 203, 47-65.

1745 Romek, K.M., Julien, M., Frasquet-Darrieux, M., Tea, I., Antheaume, I., Hankard, R., Robins, R.J.,
1746 2013. Human baby hair amino acid natural abundance ^{15}N -isotope values are not related to the
1747 ^{15}N -isotope values of amino acids in mother’s breast milk protein. *Amino Acids* 45, 1365-1372.

1748 Roth, J.D., Hobson, K.A., 2000. Stable carbon and nitrogen isotopic fractionation between diet and
1749 tissue of captive red fox: implications for dietary reconstruction. *Canadian Journal of Zoology*
1750 78, 848-852.

1751 Ruiz-Cooley, R.I., Balance, L.T., McCarthy, M.D., 2013. Range expansion of the jumbo squid in the
1752 NE Pacific: $\delta^{15}\text{N}$ decrypts multiple origins, migration, and habitat use. *PLoS ONE* 8, e59651.

1753 Ruiz-Cooley, R.I., Koch, P.L., Fiedler, P.C., McCarthy, M.D., 2014. Carbon and nitrogen isotopes
1754 from top predator amino acids reveal rapidly shifting ocean biochemistry in the Outer
1755 California Current. *PLoS ONE* 9, e110355.

1756 Sacks, G.L., Brenna, J. T., 2005. $^{15}\text{N}/^{14}\text{N}$ position-specific isotopic analyses of polynitrogenous
1757 amino acids. *Analytical Chemistry* 77, 1013-1019.

1758 Saino, T., Hattori A., 1980. ^{15}N Natural abundance in oceanic suspended particulate matter. *Nature*
1759 283, 752–754.

1760 Schiff, J.T., Batista, F., Sherwood, O.A., Guilderson, T.P., Hill, T.M., Ravelo, A.C., McMahon,
1761 K.W., McCarthy, M.D., 2014. Compound specific amino acid $\delta^{13}\text{C}$ patterns in a deep-sea
1762 proteinaceous coral: implications for reconstructing detailed $\delta^{13}\text{C}$ records of exported primary
1763 production. *Marine Chemistry* 166, 82-91.

1764 Schulten, H.R., Schnitzer M., 1997. The chemistry of soil organic nitrogen: a review. *Biology and*
1765 *Fertility of Soils* 26, 1-15.

- 1766 Schwarcz, H.P., 1991. Some theoretical aspects of isotope paleodiet studies. *Journal of*
1767 *Archaeological Science* 18, 261-275.
- 1768 Schwarcz, H.P., Dupras, T.L., Fairgrieve, S.I., 1999. ^{15}N Enrichment in the Sahara: in search of a
1769 global relationship. *Journal of Archaeological Science* 26, 629–636.
- 1770 Schwartz-Narbonne, R., Longstaffe, F.J., Metcalfe, J.Z., Zazula, G., 2015. Solving the woolly
1771 mammoth conundrum: amino acid ^{15}N -enrichment suggests a distinct forage or habitat.
1772 *Scientific Reports* 5, 9791. doi:10.1038/srep09791
- 1773 Scott, J. H., O'Brien, D.M., Emerson, D., Sun, H., McDonald, G.D., Salgado, A. Fogel, M.L., 2006.
1774 Examination of carbon isotopic effects associated with amino acid biosynthesis. *Astrobiology* 6,
1775 867-880.
- 1776 Seminoff, J., Benson, S.R., Arthur, K.E., Tomoharu, E., Dutton, P.H., Tapilatu, R.F., Popp, B.N.,
1777 2012. Stable isotope tracking of endangered sea turtles: Validation with satellite telemetry and
1778 $\delta^{15}\text{N}$ analysis of amino acids. *PLoS ONE* 7, e37403.
- 1779 Sherwood, O.A., Heikoop, J.M., Scott, D.B., Risk, M.J., Guilderson, T.P., McKinney, R.A., 2005.
1780 Stable isotopic composition of deep-sea gorgonian corals *Primnoa* spp.: a new archive of
1781 surface processes. *Marine Ecology Progress Series* 301, 135-148.
- 1782 Sherwood, O.A., Lehmann, M.F., Schubert, C.J., Scott, D.B., McCarthy, M.D., 2011. Nutrient
1783 regime shift in the western North Atlantic indicated by compound-specific $\delta^{15}\text{N}$ of deep-sea
1784 gorgonian corals. *Proceedings of the National Academy of Science USA* 108, 1011-1015.
- 1785 Sherwood, O.A., Guilderson, T.P., Batista, F.C., Schiff, J.T., McCarthy, M.D., 2014. Increasing
1786 subtropical North Pacific Ocean nitrogen fixation since the Little Ice Age. *Nature* 505, 78-81.
- 1787 Silfer, J.A., Engel, M.H., Macko, S.A., Jumeau, E.J., 1991. Stable carbon isotope analysis of amino
1788 acid enantiomers by conventional isotope ratio mass spectrometry and combined gas
1789 chromatography/isotope ratio mass spectrometry. *Analytical Chemistry* 63, 370-374.
- 1790 Silfer, J.A., Engel, M.H., Macko, S.A., 1992. Kinetic fractionation of stable carbon and nitrogen
1791 isotopes during peptide bond hydrolysis: Experimental evidence and geochemical implications.
1792 *Chemical Geology* 101, 211–221.
- 1793 Simpson, I.A., Bol, R., Dockrill, S.J., Petzke, K.J., Evershed, R.P., 1997. Compound-specific $\delta^{15}\text{N}$
1794 amino acid signals in palaeosols as indicators of early land use : a preliminary study.
1795 *Archaeological Prospection* 4, 147-152.
- 1796 Simpson, I., Bol, R., Bull, I., Evershed, R., Petzke, K., Dockrill, S., 1999. Interpreting early land
1797 management through compound specific stable isotope analyses of archaeological soils. *Rapid*
1798 *Communications in Mass Spectrometry* 13, 1315–1319.
- 1799 Steffan, S., Chikaraishi, Y., Horton, D.R., Ohkouchi, N., Singleton, M.E., Miliczky, E., Hogg, D.B.,
1800 Jones, V.P., 2013. Trophic hierarchies illuminated via amino acid isotopic analysis. *PLoS ONE*
1801 8, e76152.

1802 Steffan, S.A., Chikaraishi, Y., Currie, C.R., Horn, H., Gaines-Day, H.R., Pauli, J.N., Zalapa, J.E.,
1803 Ohkouchi, N., 2015a. Microbes are trophic analogs of animals. *Proceedings of the National*
1804 *Academy of Sciences USA* 112, 15119-15124.

1805 Steffan, S.A., Chikaraishi, Y., Dharampal, P.S., Pauli, J.N., Horn, H., Currie, C.R., Guédot, C.,
1806 Ohkouchi, N., 2017. Unpacking brown foodwebs: Animal trophic identity reflects rampant
1807 microbivory. *Ecology and Evolution*. in revision.

1808 Steffan, S.A., Chikaraishi, Y., Horton, D.R., Miliczky, E., Zalapa, J.E., Jones, V.P., Ohkouchi, N.
1809 2015b. Beneficial or not? Decoding carnivore roles in plant protection. *Biological Control* 91,
1810 34-41.

1811 Strzepek, K.M., Thresher, R.E., Revill, A.T., Smith, C.I., Komugabe, A.F., Fallon, S.F., 2014
1812 Preservation effects on the isotopic and elemental composition of skeletal structures in the
1813 deep-sea bamboo coral *Lepidisis* spp. (Isididae). *Deep Sea Res Part II* 99, 199-206.

1814 Styring, A.K., Sealy, J.C., Evershed, R.P., 2010. Resolving the bulk $\delta^{15}\text{N}$ values of ancient human
1815 and animal bone collagen via compound-specific nitrogen isotope analysis of constituent amino
1816 acids. *Geochimica et Cosmochimica Acta* 74, 241-251.

1817 Styring, A.K., Kuhl, A., Knowles, T.D.J., Fraser, R.A., Bogaard, A., Evershed, R.P., 2012. Practical
1818 considerations in the determination of compound-specific amino acid $\delta^{15}\text{N}$ values in animal and
1819 plant tissues by gas chromatography-combustion-isotope ratio mass spectrometry, following
1820 derivatisation to their N-acetylisopropyl esters. *Rapid Communications in Mass Spectrometry*
1821 26, 2328-2334.

1822 Styring, A.K., Manning, H., Fraser, R.A., Wallace, M., Jones, G., Charles, M., Heaton, T.H.E.,
1823 Bogaard, A., Evershed, R.P., 2013. The effect of charring and burial on the biochemical
1824 composition of cereal grains: Investigating the integrity of archaeological plant material.
1825 *Journal of Archaeological Science* 40, 4767-4779.

1826 Takano, Y., Kashiyama, Y., Ogawa, N.O., Chikaraishi, Y., Ohkouchi, N., 2010. Isolation and
1827 desalting with cation-exchange chromatography for compound-specific nitrogen isotope
1828 analysis of amino acids: application to biogeochemical samples. *Rapid Communications in*
1829 *Mass Spectrometry* 24, 2317-2323.

1830 Takano, Y., Chikaraishi, Y., Ohkouchi, N., 2015. Isolation of underivatized amino acids by ion-pair
1831 high performance liquid chromatography for precise measurement of nitrogen isotopic
1832 composition of amino acids: development of comprehensive LC x GC/C/IRMS method.
1833 *International Journal of Mass Spectrometry* 379, 16-25.

1834 Trueman, C.N., MacKenzie, K.M., Palmer, M.R., 2012. Identifying migrations in marine fishes
1835 through stable-isotope analysis. *Journal of Fish Biology* 81, 826-847.

1836 Vander Zanden, M.J., Rasmussen, J.B., 2001. Variations in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ trophic fractionation:
1837 Implications for aquatic food web studies. *Limnology and Oceanography* 46, 2061-2066.

- 1838 Vander Zanden, H.B., Arthur, K.E., Bolten, A.B., Popp, B.N., Lagueux, C.J., Harrison, E., Campbell,
1839 C.L., Bjorndal, K.A., 2013. Trophic ecology of a green turtle breeding population. *Marine*
1840 *Ecology Progress Series* 476, 237-249.
- 1841 van der Heijden, M.G.A., Bardgett, R.D., van Straalen, N.M., 2008. The unseen majority: Soil
1842 microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology Letters*
1843 11, 296–310.
- 1844 Vokhshoori, N.L., McCarthy, M.D., 2014. Compound-specific $\delta^{15}\text{N}$ amino acid measurements in
1845 littoral mussels in the California upwelling ecosystem: A new approach to generating baseline
1846 $\delta^{15}\text{N}$ isoscapes for coastal ecosystems. *PLoS ONE* 9, e98087.
- 1847 Vokhshoori, N.L., Larsen, T., McCarthy, M.D., 2014. $\delta^{13}\text{C}$ of amino acids in littoral mussels: A new
1848 approach to constructing isoscapes of primary production in a coastal upwelling system. *Marine*
1849 *Ecology Progress Series* 504, 59-72.
- 1850 Wada, E., 1980. Nitrogen isotope fractionation and its significance in biogeochemical processes
1851 occurring in marine environments. In, Goldberg, E.D., Horibe, Y., Saruhashi, K. (Eds.) *Isotope*
1852 *Marine Chemistry*, Uchida Rokkakudo, Tokyo, pp. 375-398.
- 1853 Wada, E., Kadonaga, T., Matsuo, S., 1975. ^{15}N abundance in nitrogen of naturally occurring
1854 substances and global assessment of denitrification from isotopic viewpoint. *Geochemical*
1855 *Journal* 9, 139-148.
- 1856 Walker, B.D., McCarthy, M.D., 2012. Elemental and isotopic characterization of dissolved and
1857 particulate organic matter in a unique California upwelling system: Importance of size and
1858 composition in the export of labile material. *Limnology and Oceanography* 57, 1757-1774.
- 1859 Walker, B.D., Guilderson, T., Okimura, K.M., Peacock, M., McCarthy, M.D., 2014. Radiocarbon
1860 signatures and size-age-composition relationships of major organic matter pools within a
1861 unique California upwelling system. *Geochimica et Cosmochimica Acta* 126, 1-17.
- 1862 Walsh, R.G., He, S., Yarnes, C., 2014, Compound-specific $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis of amino acids: a
1863 rapid, chloroformate-based method for ecological studies. *Rapid Communications in Mass*
1864 *Spectrometry* 28, 96-108.
- 1865 West, J.B., Bowen, G.J., Dawson, T.E., Tu K.P., (Eds.) 2010. *Isoscapes: Understanding Movement,*
1866 *Pattern and Process on Earth Through Isotope Mapping.* Springer, New York, 487 pp.
- 1867 Wilkie, M.P., 2002. Ammonia excretion and urea handling by fish gills: present understanding and
1868 future research challenges. *Journal of Experimental Zoology* 293, 284-301.
- 1869 Williams, B., Risk, M.J., Ross, S.W., Sulak, K.J., 2007. Stable isotope data from deep-water
1870 antipatharians: 400-year records from the southeastern coast of the United States of America.
1871 *Bulletin of Marine Science* 81, 437-447.
- 1872 Williams, B., Thibodeau, B., Chikaraishi, Y., Ohkouchi, N., Walnum, A., Grottoli, A., Colin, P.,
1873 2017. Consistency in coral skeletal amino acid composition offshore of Palau in the western

1874 Pacific warm pool indicates no impact of decadal variability in nitricline depth on primary
1875 productivity. *Limnology and Oceanography*, in press.

1876 Wilson, R., Tudhope, A., Brohan, P., Briffa, K., Osborn, T., Tett, S., 2006. Two-hundred-fifty years
1877 of reconstructed and modeled tropical temperatures. *Journal of Geophysical Research* 111,
1878 C10007, doi:10.1029/2005JC003188

1879 Wyatt, A.S.J., Lowe, R.J., Humphries, S., Waite, A.M., 2013. Particulate nutrient fluxes over a
1880 fringing coral reef: Source-sink dynamics inferred from carbon to nitrogen ratios and stable
1881 isotopes. *Limnology and Oceanography* 58, 409-427.

1882 Wyatt, A.S.J., Waite, A.M., Humphries, S., 2012. Stable isotope analysis reveals community-level
1883 variation in fish trophodynamics across a fringing coral reef. *Coral Reefs* 31, 1029-1044.

1884 Yamaguchi, Y.T., 2013. Biogeochemical dynamics of amino acids in marine sediments: constraints
1885 from compound-specific nitrogen isotopic composition and D/L ratio. PhD thesis, The
1886 University of Tokyo, p. 139.

1887 Yarnes, C., Herszage, J., 2017. The relative influence of derivatization and normalization procedures
1888 on the compound-specific stable isotope analysis of nitrogen in amino acids. *Rapid*
1889 *Communication in Mass Spectrometry*, in press.

1890

1891 **Table legends**

1892 Table 1. A summary of three types of derivatized amino acids used for the nitrogen isotopic analysis.

1893

1894 **Figure captions**

1895

1896 **Fig. 1.** Derivatization of amino acids for the nitrogen isotope analysis by GC/IRMS: Schemes of (a)
1897 basic chemical reaction, (b) TFA/AA/iPr ester, (c) Pv/AA/iPr ester, and (d) MOC AA ester.

1898

1899 **Fig. 2.** Ester exchange between amino acid derivatives and ethyl acetate.

1900

1901 **Fig. 3.** A representative chromatogram of GC/IRMS analysis of the nitrogen isotope analysis of
1902 amino acids as *N*-pivaloyl *O*-isopropyl esters.

1903

1904 **Fig. 4.** Schematic illustrations of the trophic position (*TP*) estimates by (a) bulk and (b) CSIA-AA
1905 methods. In the bulk method, the $\delta^{15}\text{N}$ values of consumers at the same *TP* frequently vary, due to
1906 temporal or spatial change in the $\delta^{15}\text{N}$ value at the basis of food web. In contrast, CSIA-AA method
1907 can estimate *TP* independent of change in the $\delta^{15}\text{N}$ value at the basis of food web (after [Naito et al.,](#)
1908 [2016](#)).

1909

1910 **Fig. 5.** (a) Initial steps of the dominant metabolism for glutamic acid and phenylalanine in animals,
1911 and (b) schematic illustrations of the relationship between $\delta^{15}\text{N}$ values of amino acids (Glu and Phe)
1912 and trophic position in aquatic and terrestrial food webs (after [Chikaraishi et al., 2009, 2010](#)).

1913

1914 **Fig. 6.** Schematic of two aquatic food webs differentiated based on the $\delta^{15}\text{N}$ values of Phe and Glu.
1915 Changes in baseline nitrogen sources cause each food web to be separated along the source amino
1916 acid axis; here an oceanic food web based on phytoplankton production (P) supported by *e.g.*
1917 N_2 -fixation (low $\delta^{15}\text{N}_{\text{Phe}}$) is separated from a benthic food web based on macroalgae production (M)
1918 supported by *e.g.* upwelling or terrestrial run-off (high $\delta^{15}\text{N}_{\text{Phe}}$). The potential for ‘trophic omnivory’
1919 can be evident as non-integer *TP*s; here the oceanic food web depicts potential ontogenetic changes
1920 in CSIA-AA-derived *TP* for fish across two *TP*s (anchovy and tuna). An example is also provided of
1921 a mobile apex predator (the tiger shark, *Galeocerdo cuvier*; *Ga*) potentially integrating across
1922 oceanic and benthic food webs at a given *TP*, leading to intermediate $\delta^{15}\text{N}_{\text{SAA}}$ value.

1923

1924 **Fig. 7.** Two examples of food web analysis by $\delta^{15}\text{N}_{\text{AA}}$: a) the coastal marine (a stony shore)
1925 ecosystem in Japan ([Chikaraishi et al., 2014](#)), and b) Lake Baikal ([Ohkouchi et al., 2015](#)).

1926

1927 **Fig. 8.** (a) Concentrations of nitrate (black) and phosphate (red) observed in the hypolimnetic water in
1928 the north basin of Lake Biwa, Japan. (b) Trophic position of gobiid fish Isaza (*Gymnogobius isaza*)
1929 estimated by the $\delta^{15}\text{N}_{\text{AA}}$. (c) $\delta^{15}\text{N}$ values of bulk muscular tissue, Glu, and Phe of formalin-fixed
1930 Isaza specimens. $\delta^{15}\text{N}$ values of bulk sediments were also shown (data from Ogawa et al. 2001). A
1931 grey band indicates the major eutrophication period in Lake Biwa (1960-1980, [Ogawa et al., 2013](#))
1932

1933 **Fig. 9.** Fungi can be carnivorous. Here, the fungus *Beauveria bassiana* has subdued and killed a
1934 caterpillar (larval *Spodoptera frugiperda*). The trophic position of the fungus is 3.0 (Steffan et al.,
1935 2015a), because functionally, this fungus is a strict carnivore.
1936

1937 **Fig. 10.** Isotopic approaches have been used to decode carnivore impacts on key ecosystem metrics,
1938 such as primary productivity (after [Steffan et al., 2015b](#)). Heterotrophic feeding induces trophic
1939 cascades, which directly and indirectly influence other trophic groups. The trophic tendency of any
1940 given species, coupled with its resource capture efficiency (% consumption of resource base),
1941 permits estimation of the consumers' impacts on plant protection.
1942

1943 **Fig. 11.** Nitrogen isotopic compositions of Phe and Glu of Holocene hunter-gatherers in Japanese
1944 archipelago. a) Kitakogane shell midden located near the coastal line of Hokkaido (Early Jomon
1945 period, *ca.* 6000-5300 cal BP) and b) Tochibara rockshelter site located at inland Nagano (Initial
1946 Jomon period, *ca.* 9100-9700 cal BP). Note that Kitakogane humans exhibit $\delta^{15}\text{N}_{\text{Phe}}$ closer to marine
1947 fauna than terrestrial fauna suggesting their strong reliance on marine foods while Tochibara humans
1948 exhibit $\delta^{15}\text{N}_{\text{Phe}}$ comparable to those of terrestrial fauna suggesting their reliance exclusively on
1949 terrestrial foods ([Naito et al., 2010b, 2013b](#)).
1950

1951 **Fig. 12.** Nitrogen isotopic compositions of Phe and Glu for Neanderthal and animal remains from
1952 Spy and Scladina caves in Pleistocene Belgium ([Naito et al., 2016b](#)).
1953

1954 **Fig. 13.** $\delta^{15}\text{N}_{\text{bulk}}$ trends (A) and $\delta^{15}\text{N}_{\text{Phe}}$ -calibrated baseline $\delta^{15}\text{N}$ isoscape (B) along the California
1955 coast, based on selected CSIA-AA within high-density bulk sampling of littoral mussels.
1956


1957 **Fig. 14.** Conceptual diagrams describing the proposed four patterns of $\delta^{15}\text{N}_{\text{AA}}$ fractionation of
1958 chemotrophic microbes (for details, see Section 4.5.1 in the main text). Eight AAs which have
1959 been commonly analyzed are selected for the diagrams. (a) Pattern 1. The $\delta^{15}\text{N}_{\text{AA}}$ pattern of *de*
1960 *novo* AA synthesis from inorganic nitrogen by chemotrophic microbes (closed circles: microbial
1961 biomass), which was observed in the pure culture experiments ([Yamaguchi, 2013](#)). The $\delta^{15}\text{N}_{\text{AA}}$
1962 values are normalized to the $\delta^{15}\text{N}_{\text{Glu}}$ value. (b) Pattern 2. The $\delta^{15}\text{N}_{\text{AA}}$ fractionation pattern of

1963 heterotrophic microbes relative to preformed AA in substrates, which was observed in the pure
1964 culture experiments (red squares: microbial biomass) (Stefan et al., 2015; Yamaguchi, 2013). The
1965 $\delta^{15}\text{N}_{\text{AA}}$ values of the substrates in b, c, and d (open circles) are set as the average pattern of algae
1966 (Chikaraishi et al., 2009; McCarthy et al., 2013), and are normalized to the $\delta^{15}\text{N}_{\text{Glu}}$ value. (c)
1967 Pattern 3. A possible example of the scattered $\delta^{15}\text{N}_{\text{AA}}$ fractionation by heterotrophic microbes
1968 relative to substrates in some settings (blue triangles: degraded materials), hypothesized from the
1969 results of incubation or microcosm experiments (Fogel and Tuross, 1999; Calleja et al., 2013;
1970 Gutierrez-Rodriguez et al., 2014). Note that the $\delta^{15}\text{N}$ fractionation value of each AA in this pattern
1971 is not well constrained and is likely variable. (d) Pattern 4. A hypothesized $\delta^{15}\text{N}_{\text{AA}}$ fractionation
1972 pattern during extracellular protein hydrolysis by heterotrophic microbes (green squares: residue of
1973 hydrolysis) (Hannides et al., 2013). Note that the magnitude of $\delta^{15}\text{N}$ fractionation would be
1974 variable, depending on the character of substrates and the degree of degradation.


| | TFA/AA/iPr | Pv/AA/iPr | MOC/AA ester |
|--------------------|------------|-----------|--------------|
| Available solvent | DCM | DCM | DCM or MeOH |
| Toxicity | High | Very high | High |
| Stability at -20°C | 1-2 years | 1-2 years | 1-2 weeks |
| Volatility | High | Low | Very high |

Table 1


(a) Basic chemical reaction



(b) TFA/AA/iPr



(c) Pv/AA/iPr



(d) MOC AA ester






Fig. 1

(a) Exchange of ester group



(b) Combustion of TFA/AA/iPr



(c) MOC derivatives of glutamic acid




Fig. 2





Fig. 3

(a) Bulk method



(b) CSIA-AA method





Fig. 4

(a) First step in metabolism



(b) Trophic enrichment in ^{15}N




Fig. 5




Fig. 6





Fig. 7




Fig. 8



Fig. 9




Fig. 10





Fig. 11




Fig. 12





Fig. 1: Bulk $\delta^{15}\text{N}$ trends (A) and a $\delta^{15}\text{N}_{\text{PDB}}$ calibrated baseline $\delta^{15}\text{N}$ isoscape (B) along the CA coast,





Fig. 14




Fig. S1 Comparison of stabilities of various pivaloyl esters of alanine. Triangle: methyl ester (Pv/Ala/Me), diamond: ethyl ester (Pv/Ala/Et), square: *n*-propyl ester (Pv/Ala/nPr), circle: isopropyl ester (Pv/Ala/iPr).




Fig. S2. Isotopic fractionation of MOC ester derivatization of Glu. The MOC ester (MOC Glu ester, filled circle) is depleted in ^{15}N whereas the cyclic ester (p-Glu ester, open circle) is enriched in ^{15}N . Molar ratio of these two derivatives (p-Glu/Glu) has a negative correlation with $p\text{H}$ during the derivatization, however, the mass balanced values (gray square) do not equal to the reference isotopic composition of glutamic acid (+45.7‰, broken line).




Fig. S3. A regression line between known (‰, vs. AIR) and measured values (‰, vs. reference N_2 gas).




Fig. S4. The accuracy and precision of the nitrogen isotope analysis of amino acids by GC/IRMS with respect to the signal/noise (S/N) ratio (Chan et al., 2016).