University of Rhode Island

DigitalCommons@URI

Open Access Master's Theses

2014

EARLY SPRING PHYTOPLANKTON DYNAMICS IN THE SUBPOLAR NORTH ATLANTIC: THE INFLUENCE OF HETEROTROPHIC-PROTIST HERBIVORY

Francoise Morison

University of Rhode Island, francoisemorison@gmail.com

Follow this and additional works at: <https://digitalcommons.uri.edu/theses>

Recommended Citation

Morison, Francoise, "EARLY SPRING PHYTOPLANKTON DYNAMICS IN THE SUBPOLAR NORTH ATLANTIC: THE INFLUENCE OF HETEROTROPHIC-PROTIST HERBIVORY" (2014). *Open Access Master's Theses*. Paper 423.

<https://digitalcommons.uri.edu/theses/423>

This Thesis is brought to you for free and open access by DigitalCommons@URI. It has been accepted for inclusion in Open Access Master's Theses by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

**EARLY SPRING PHYTOPLANKTON DYNAMICS
IN THE SUBPOLAR NORTH ATLANTIC:
THE INFLUENCE OF HETEROTROPHIC-PROTIST HERBIVORY**

**BY
FRANÇOISE MORISON**

**A THESIS SUBMITTED IN PARTIAL FULFILLMENT OF THE
REQUIREMENTS FOR THE DEGREE OF
MASTER OF SCIENCE
IN
OCEANOGRAPHY**

UNIVERSITY OF RHODE ISLAND

2014

MASTER OF SCIENCE THESIS

OF

FRANÇOISE MORISON

APPROVED:

Major Professor: Susanne Menden-Deuer

Thesis Committee: David Smith

Jason Kolbe

Nasser H. Zawia
DEAN of THE GRADUATE SCHOOL

UNIVERSITY OF RHODE ISLAND

2014

ABSTRACT

To assess the importance of herbivory by heterotrophic protists in relation to mixed-layer depth prior to the spring phytoplankton bloom, we measured phytoplankton growth and heterotrophic-protist grazing rates during the March/April 2012 EuroBasin Deep Convection cruise in the subpolar North Atlantic. We performed 15 dilution experiments during 2-4 visits at one shelf (160 m) and two deep (~1300 m) stations. Of the two deep stations, one had a mean mixed-layer depth of 476 m, whereas the other was stratified (46 m). Euphotic depth averaged ~70 m at both stations. Initial chlorophyll-*a* varied from 0.2 to 1.9 $\mu\text{g L}^{-1}$ at the deep mixed layer station and from 0.5 to 1.0 $\mu\text{g L}^{-1}$ at the stratified station. In 80 % of the experiments, growth rates exceeded grazing mortality rates, regardless of mixed layer depth. Large mixed layer depth coincided with phytoplankton growth and grazing mortality rates that varied over a similar range from ≤ 0 to 0.6 d^{-1} , and to an average grazing-impact representing 50% of primary production (PP). At the stratified station, phytoplankton growth rates varied from 0.18 to 0.41 d^{-1} , grazing mortality rates varied from 0.11 to 0.34 d^{-1} , and a temporal shift from a positive to a negative balance between growth and grazing rates caused the proportion of PP consumed to increase from 60% to 180%. Variations in *in situ* chlorophyll-*a* could not be explained where the mixed layer was deep, whereas at the stratified station the balance between rate estimates of phytoplankton growth and grazing mortality rates explained 98 % of measured changes in chlorophyll-*a*. These results suggest a difference in the dominant surface loss process at the two stations: grazing at the stratified station vs. potential sinking aided by vertical mixing where mixed layer was deep.

ACKNOWLEDGEMENTS

I thank my major professor and advisor Dr. Susanne Menden-Deuer for sending me to conduct work at sea, supporting me with advice and patience, and trusting my ability to complete the present work. Thanks go to the scientific coordinator of the Euro-Basin program, Mike St John, for letting me participate in the RV Meteor Deep Convection cruise, and along with the cruise's chief scientists Jan Backhaus and B. Christiansen, for providing cruise leadership. I am indebted to Maria Lund Paulsen (DTU Aqua/University of Bergen) for sharing with me some of the data she and Karen Riisgaard (DTU Aqua) collected on the cruise, and for their friendship. In fact, the collaboration and companionship of all the cruise participants from the University of Hamburg, DTU-Aqua, National Oceanography Center (Southampton, UK) not only made lugging incubation bottles to the "heli-deck" much easier, but just as importantly enhanced the 42 days spent at sea with their energy and enthusiasm. Recognition goes to Master of RV Meteor, M. Schneider and his crew for ensuring safe sailing in the late winter high seas of the North Atlantic, and for helping the scientific party achieve their mission. Thank you to past and present lab partners, in particular Sam DeCuello, Elizabeth Harvey, Hyewon Kim, Caitlyn Lawrence, and Amanda Montalbano, for sharing not just skills and ideas, but more importantly friendship. Last, but by no means least, thanks to my dear children John H. IV and Nathalie for cheering me along, and to my beloved husband and dedicated life partner, John H. III, for his patience, his encouragement, and his generous and unfailing love. I could not do it without their support.

PREFACE

This thesis was prepared using the manuscript format for submission to and according to publication guidelines of the journal *Deep Sea Research II*

TABLE OF CONTENTS

ABSTRACT.....	ii
ACKNOWLEDGEMENTS.....	iii
PREFACE.....	iv
TABLE OF CONTENTS.....	v
LIST OF TABLES.....	vi
LIST OF FIGURES.....	vii
MANUSCRIPT.....	1
INTRODUCTION.....	2
METHODS.....	6
RESULTS.....	16
DISCUSSION.....	24
CONCLUSIONS.....	37
APPENDIX.....	38
REFERENCES.....	41
FIGURE LEGENDS.....	52

LIST OF TABLES

TABLE	PAGE
Table 1. Summary of dilution experiments, with environmental conditions at sampling location.....	48
Table 2. Summary of results from dilution experiments, including initial chl-a concentration, and phytoplankton growth and heterotrophic protist grazing rates estimated by regression and 2-point analyses.....	49
Table 3. Biomass estimates of major groups of plankton contained in undiluted samples collected at the beginning of each experiment	50
Table 4. Results of dilution experiments previously conducted in the subpolar North Atlantic during or post bloom conditions, including chlorophyll concentrations and HP numerical abundance and biomass for comparison with this study's findings.	51

LIST OF FIGURES

FIGURE	PAGE
Figure 1. Map showing locations of sampling stations.....	54
Figure 2. Principal Component Analysis ordination of <i>in situ</i> environmental conditions	55
Figure 3. Multi-Dimensional Scaling of species composition of (a) diatom and (b) heterotrophic protist assemblages.....	56
Figure 4. Size distribution of dominant groups of heterotrophic protists in undiluted samples collected at the beginning of each experiment at the two oceanic stations	57
Figure 5. Phytoplankton growth rates vs. grazing mortality rates	58
Figure 6. Heterotrophic protist herbivory rates in relation to heterotrophic protist biomass.....	59
Figure 7. Calculated vs. observed net phytoplankton population growth.....	60
Figure 8. Selected CTD profiles of temperature and fluorescence-based chlorophyll- <i>a</i> from the four visits at the Iceland basin station (S1).....	61

Manuscript

(To be submitted to the journal *Deep Sea Research II*)

Early spring phytoplankton dynamics in the subpolar North Atlantic:

The influence of Heterotrophic-Protist Herbivory

Françoise Morison

Susanne Menden-Deuer*

Graduate School of Oceanography, University of Rhode Island, Narragansett, RI,

USA

*Corresponding author: Tel: (401) 874-6608

E-mail address: smenden@gso.uri.edu

Key Words: Heterotrophic protist, Grazing, Dilution, Mixed-layer depth, North Atlantic, Spring Bloom

Running head: Pre-bloom grazing in the subpolar North Atlantic

1. Introduction

In the subpolar North Atlantic, the yearly cycle of primary production (PP) is dominated by the annual recurrence of the spring phytoplankton bloom. The seasonal increase in phytoplankton biomass is of large ecological significance: not only does the bloom fuel marine food webs, it also influences earth's climate, since the associated drawdown of atmospheric CO₂ and the consequent sinking of some of the bloom biomass (Buesseler et al. 2007, Sarthou 2005, Turner 2002) contribute substantially to the strength of the global biological pump (Takahashi et al. 2009).

For a bloom (i.e. an accumulation of biomass) to occur, net phytoplankton population growth rate (i.e. accumulation rate) needs to be positive, that is phytoplankton intrinsic growth rate has to exceed the rate at which production is lost (Banse 1994). The process can be described by the equation $r = \mu - l$, where r is the phytoplankton biomass accumulation rate, μ is the phytoplankton growth rate, and l represents the rate of phytoplankton losses (Behrenfeld 2010).

From the earliest days of the extensive research devoted to identifying what triggers the North Atlantic spring bloom, of the two terms involved in the equation describing a bloom, μ has received the most attention (Behrenfeld & Boss 2014). In particular a large focus has been placed on the influence on μ of one physical variable: mixed layer depth (MLD), a proxy for, yet not always representative of, the actively mixing layer (Ferrari et al. 2014, Taylor & Ferrari 2011). Starting with Sverdrup (1953), the idea that the North Atlantic spring bloom begins when the mixed layer shoals above a “critical depth”, i.e. the depth of a mixed layer within which integrated phytoplankton production and losses are equal, has served as a paradigm in the

understanding of bloom formation (Henson et al. 2006, Siegel et al. 2002, Sverdrup 1953). Numerous observations, however, have been reported of early spring surface increases in phytoplankton biomass preceding stratification (e.g. Dale et al. 1999, Townsend et al. 1992,1994), challenging Sverdrup's classical bloom model. Yet consequent new hypotheses have continued to focus on potential factors driving μ , all involving the extent of vertical mixing: for example, rates of turbulent mixing (Huisman et al. 1999, 2002), heat-flux induced weakening of turbulent mixing (Ferrari et al. 2014, Taylor & Ferrari 2011), and eddy-driven stratification (Mahadevan et al. 2012). Thus traditionally, a disproportionate emphasis has been given to μ with the loss term being less studied.

Of all losses affecting PP, the largest is due to grazing (Banse 1994). In particular, herbivory by ubiquitous $<200\ \mu\text{m}$ heterotrophic protists (HP), such as ciliates and dinoflagellates, has been identified as the major fate of ocean PP (Calbet & Landry 2004, Sherr & Sherr 2009, Strom 2002). Thanks to their diverse feeding strategies, protist grazers can access a broad range of prey sizes, from bacteria to prey larger than they are (Aberle et al. 2007, Sherr & Sherr 2002). HP grow at rates similar to the cells they eat, allowing their numbers to often increase quickly after an increase in available prey (Sherr et al. 2003). From a plethora of studies performed across oceans to measure HP grazing rates, HP grazing impact has been estimated to average $\sim 69\%$ of PP (Calbet & Landry 2004, Schmocker et al. 2013). Although temporal and spatial exceptions exist, in which other loss processes such as viral lysis (Brussaard 2004) or nutrient starvation (Taylor et al. 1993) control phytoplankton biomass, HP herbivory has been established as the most significant loss factor in PP.

From the research that has considered the role of grazing losses in phytoplankton blooms, a consensus has emerged that seasonal high-latitude blooms happen because grazing cannot keep pace with phytoplankton growth. Various mechanisms have been considered, including proposed phytoplankton predation-avoidance strategies (Irigoien et al. 2005), and constraints on HP growth rates at temperatures $<5^{\circ}\text{C}$ (Rose & Caron 2007), or low prey pre-bloom availability (Sherr & Sherr 2009, Sherr et al. 2013), all yielding μ in excess of grazing.

Recent work by Behrenfeld and colleagues (Behrenfeld 2010, Behrenfeld & Boss 2014, Behrenfeld et al. 2013) has re-examined the importance of the physics of MLD, by considering its effects not only on phytoplankton growth as has been traditionally done, but also on the magnitude of grazing pressure. Behrenfeld (2010) suggested that a key process influencing variations in the North Atlantic phytoplankton biomass is the alteration by vertical mixing of the balance between μ and grazing. According to Behrenfeld's "dilution-recoupling" hypothesis (Behrenfeld 2010), deepening of the mixed layer in winter reduces predator-prey encounters, decreasing phytoplankton grazing losses below the very low but positive rates of winter phytoplankton growth, thus allowing blooms to *initiate* during winter. Behrenfeld (2010) further postulated that the gradual seasonal shoaling of the mixed layer "re-couples" predators with their prey, resulting in increased grazing pressure, which curbs phytoplankton biomass accumulation rate.

Despite the potential importance of HP grazing, its role in bloom development remains theoretical, as our understanding of pre-bloom grazing dynamics suffers from a shortage of available *in situ* grazing rates measured before or during early bloom

development. In particular, for the open North Atlantic ocean at high latitudes above 50° N, where winter mixed layer is typically large due to convection (Backhaus et al. 2003), existing bloom-related *in situ* measurements of HP grazing rates come from studies conducted during or after the bloom (Burkill et al. 1993, Gaul & Antia 2001, Gifford et al. 1995, Stelfox-Widdicombe et al. 2000, Wolfe et al. 2000), and to our best knowledge, there are no empirical data of grazing rates for the critical period that precedes the bloom.

The present research was performed during the early spring 2012 EuroBasin program “Deep Convection” research cruise, which intended to evaluate the response of the subpolar North Atlantic ecosystem to physical forcing during the transitional period when winter convection gradually weakens (Backhaus et al. 2003). We sought to evaluate the importance of grazing mortality in the dynamics of phytoplankton biomass prior to the spring bloom. We repeatedly measured HP grazing and phytoplankton growth rates from March 26 to April 28 2012, at one shelf and at two open ocean sampling stations. Although this paper presents results for all three stations sampled, it focuses on results from the two open ocean sites, which contrasted in MLD, a variable of recognized importance in dynamics of phytoplankton biomass. We found that at the oceanic sites, rates of both phytoplankton growth and HP grazing could be substantial, even when the mixed layer was deep, yet in most experiments growth rates exceeded grazing rates, regardless of MLD. Further analysis suggested that grazing was a dominant factor controlling phytoplankton biomass only when the mixed layer was shallow, and not when it was deep.

2. Methods

2.1 *Sampling sites*

Heterotrophic-protist herbivory was quantified during two to four visits at two ~1300-m deep open-ocean sites located in the Iceland Basin and the Norwegian Sea (S1 & S2), and at a ~160-m deep site located on the Shetland shelf (S3) (Fig. 1).

2.2 *Dilution experiments*

We measured HP grazing rates in 15 separate experiments using the Landry & Hassett (1982) dilution method (Table 1). Water containing the plankton assemblage for the experiments was collected using Niskin bottles mounted on a rosette sampler with a SBE911Plus Seabird Electronics Inc. CTD equipped with WET Labs ECO-FLNTU(RT)D chlorophyll sensor. Depth of water collection corresponded to the CTD fluorescence maximum (F-max), except at S2 on March 31st (no F-max) and on April 14th when two depths were sampled (F-max and 5 m). Water was gently transferred from the Niskin bottles into 10-L carboys via a silicone tube, to which a 200- μ m mesh was affixed in order to screen out larger grazers (Sherr et al. 2009). We further refer to this < 200- μ m fraction as whole seawater (WSW).

For nine experiments (one per visit at each station), we prepared five target dilutions (9, 18, 37, 75, 100 % WSW), which were distributed so as to increase our ability to detect potential non-linearity related to feeding thresholds at low prey concentration, their specific values arbitrarily chosen to facilitate measuring the volumes to be combined. For the first and second experiments at S1, the actual number of dilution levels achieved was reduced from five to four, due to loss of the

most diluted sample initial chl-*a* measurement for the 1st experiment. For the 2nd experiment, the 9% and 18% dilutions deviated from the target and had equal initial chl-*a* concentration (equivalent to a 16 % fraction of WSW). We performed six additional experiments using only three target dilutions (10, 37, and 100% WSW), thus decreasing confidence in the linear regression estimates of the grazing coefficient, but increasing the number of experiments that could be performed during a visit at a station.

Each dilution level was prepared by combining appropriate proportions of WSW and filtered seawater (FSW), the latter obtained by gravity-filtration of some of the collected water through a 0.2- μ m capsule filter (Pall). To minimize variations among replicates, each dilution was prepared in a single carboy as a large volume stock. Duplicate 2.4-L polycarbonate bottles were filled with each dilution level. To ensure sufficient nutrients for phytoplankton growth (Landry & Hassett 1982) bottles were amended with final concentrations of 8.82 μ M nitrate, 0.48 μ M phosphate, and 10 μ M silicate. To check for effects of nutrients addition and for nutrient limitation, one or two additional undiluted replicates were prepared without added nutrients.

Bottles were incubated for 24 hours. All incubations took place in on-deck ~250-L tanks. Bottles were suspended mid-water by strapping them onto bungee cords loosely stretched across the length of the tanks, which together with ship motion provided gentle agitation. Incubations were maintained at *in situ* temperature by flow-through of ambient seawater. Incubation temperature was recorded at 30-min intervals using in-tank Hobo data loggers. Incubation temperature was on average 0.9 ($\pm$ 1.1) °C higher than the temperature at collection depth, however departure occurred mainly

during the first leg of the cruise, when differences were the largest at the 1st experiments at S2 and S3.

To minimize chlorophyll bleaching, which is known to occur in light-sensitive polar phytoplankton (Caron et al. 2000, Smith & Sakshaug 1990), bottles were incubated in black neutral-density mesh-bags that reduced the light to 30% of surface irradiance. Incubations carried at collection-depth irradiance fail to truly replicate the average light regime experienced by cells in a mixed layer (Ross et al. 2011), therefore, in general, the same mesh screen was used regardless of water collection-depth. However, to investigate the effect of light (31 March) and of collection depth (14 April) on rate magnitudes, a set of two experiments were incubated simultaneously, one with and one without a mesh-bag (See Appendix).

2.3 *Phytoplankton growth and grazing mortality estimates*

Phytoplankton growth and HP grazing mortality rates were estimated from changes in extracted chlorophyll-*a* (chl-*a*) over the incubation period (Landry & Hassett 1982). Initial and final chl-*a* concentrations were determined from triplicate subsamples of each dilution stock and of each replicate bottle respectively. Subsamples ranged in volume from 60 to 500 ml depending on the *in situ* chl-*a* concentration and the dilution level. Chl-*a* extraction and determination followed Graff & Ryneerson (2011), except that extraction took place at room temperature for 12-15 hours in 96% ethanol (Jespersen & Christoffersen 1987).

Apparent phytoplankton growth (k , d⁻¹) in each bottle was estimated using the equation $k = 1/t \ln (P_t - P_0)$, where t = incubation time in days, and P_t and P_0 are

respectively the final and the initial chl-*a* concentrations. We used these estimates of k to determine the instantaneous phytoplankton growth rate (μ , d⁻¹) and the instantaneous grazing rate (g , d⁻¹) using two methods.

First, as is customary to the dilution method (Landry & Hassett 1982), the rates were determined from the linear regression analysis of k as a function of the dilution factor, where μ is the y-intercept and g the negative slope of the line. We tested the hypothesis that $g = 0$ for each regression. We applied dilution factors as determined from measured initial chl-*a* concentrations in the dilutions, which was on average 1.8 % lower than the target (± 3.9 %). For the first experiment at S3, malfunctioning of the fluorometer yielded inaccurate measurements of initial chl-*a* concentration in the diluted treatments, thus initial chl-*a* was assumed to equal WSW chl-*a* multiplied by the target dilution factor. We found no significant difference between k in undiluted treatments with and without nutrients (two-tailed paired t-test; $p = 0.63, 0.21$, and 0.15 for station 1, 2, and 3 respectively), and consequently combined all undiluted replicates in our analysis.

One of the dilution method's major assumptions is that k is linearly related to the dilution factor. In case of deviations from linearity, a linear regression is thus an inadequate method to determine g and μ (Worden & Binder 2003). We therefore tested whether the linearity assumption held for all dilution experiments or if significant deviations existed (ANOVA, $\alpha = 0.05$; Zar 2010). We found significant deviations from linearity in three regular (Table 2) and one "light" experiments (Appendix). Therefore we subsequently estimated μ and g using Worden & Binder's (2003) two-point method. In this method, the grazing rate is calculated as the

difference between apparent phytoplankton growth rates k in the lowest and highest fractions of WSW; k in the most dilute treatment ($8 \pm 2\%$ WSW in our study) serves as an estimate of phytoplankton instantaneous growth rate (without grazing). Rate estimates obtained using the two-point approach are considered conservative (Lawrence & Menden-Deuer 2012, Worden & Binder 2003), and as in previous studies (DeCuollo & Menden-Deuer unpublished data, Strom & Fredrickson 2008), we did not find the two-point estimated rates to significantly differ from rates obtained from regression analysis (two-tailed paired t-test, $p = 0.10$ for μ , $p = 0.84$ for g). To insure internal consistency, the two-point method was used for all experiments, and rates reported herein are those thus derived (Table 2).

The grazing impact of HP in terms of the proportion of primary production (PP) consumed was calculated as $\% \text{ PP} = g : \mu \times 100$ following Calbet & Landry (2004). For all calculations, negative growth rate and negative grazing rate estimates were corrected to $+0.01 \text{ d}^{-1}$ and zero respectively. No $\% \text{ PP}$ was calculated for experiments in which no significant phytoplankton growth was measured. For each experiment, we also calculated biomass-specific grazing rates on phytoplankton (G_{HP}) using the equation $G_{\text{HP}} = [(g)(\text{Chl})(\text{C:chl})] / \text{HP}$, in which g is the estimated specific HP grazing rate, Chl is experiment initial chl- a concentration, and HP is HP biomass (Strom et al. 2007, Strom & Frederickson 2008). As we did not measure C:chl ratio during this study, we used a ratio of 21, a value found to be a good estimate for phytoplankton communities of Norwegian coastal waters (Bratbak et al. 2011), although we recognize that this ratio is highly variable and poorly constrained (Geider 1987, Sathyendranath et al. 2009).

As a mean to assess the importance of the grazing loss term in determining phytoplankton biomass dynamics, using chl-*a* as a proxy for biomass, we compared observed (i.e. *in situ*) chl-*a* accumulation rates (r_{obs}) to the accumulation rates inferred from the balance between experimentally determined phytoplankton growth and mortality rates ($r_{calc} = \mu - g$). The observed accumulation rate was determined using the equation $r_{obs} = 1/t * \ln(P_t - P_0)$ where P_t and P_0 are chl-*a* concentrations (from initial experiment samples) at the end and the beginning of the time interval t separating two consecutive experiments at the same station.

2.4 *Biomass and composition of the plankton community*

For microplankton biomass estimates and composition analysis, well-mixed sub-samples of the initial undiluted treatments of each experiment were preserved with acidified Lugol's iodine at a final concentration of 2% (Menden-Deuer et al. 2001). Diatoms, dinoflagellates, and ciliates were enumerated by settling 50 ml for a minimum of 24 h following the Ütermohl (1958) method. Diatoms were identified to genus following Throndsen et al. (2007) and Kraberg et al. (2010). Dinoflagellates were divided into thecate and athecate groups, and when possible further identified to genus following Dodge (1982), or assigned to a morphotype (based on similarity of shape). Preservation of samples in acid Lugol's fixative does not allow differentiating between auto- and heterotrophic dinoflagellates. Furthermore, many autotrophic dinoflagellates are also phagotrophic (Stoecker 1999), thus all dinoflagellates were assumed heterotrophic. Ciliates were divided into loricate (tintinnids) and aloricate groups. Higher taxonomic identification of aloricate ciliates relying on shape was

attempted following Strüder-Kypke et al. (2002), to provide a qualitative description of the ciliate community, but due to its lack of reliability (Montagnes & Lynn 1991), it was not used for quantitative analysis. An exception was however made for the obligate mixotroph species *Mesodinium rubrum* (= *Myrionecta rubra*; Hansen et al. 2012), which could easily be distinguished from and were not included with other aloricate ciliates.

Linear dimensions were measured using ImageJ software (National Institute of Health) from images taken of all dinoflagellates and ciliates contained in each sample and, depending on abundance, of all or a subset of diatom cells (30-300 cells per genus). Cell volumes were calculated from linear dimensions using appropriate geometric shape algorithms.

To refine the analysis of heterotrophs, aloricate ciliates and dominant dinoflagellate types (*Gymno-Gyrodinium* morphotypes and *Protoperidinium* spp.) were further divided into three size categories (small: <20 µm, medium: 20-50 µm, and large: >50 µm) based on equivalent spherical diameter (ESD), which was calculated as the diameter of a sphere with volume $V = \frac{4}{3} \pi r^3$ equal to the cell's biovolume, and thus was obtained using the equation: $ESD = (\text{Biovolume} \div 0.524)^{1/3}$.

Biomass estimates were calculated by converting biovolumes into carbon content (µg C L⁻¹) applying the following conversion factors: tintinnid ciliates, Verity & Langdon (1984); aloricate ciliates, Putt & Stoecker (1989); all other plankton groups, Menden-Deuer & Lessard (2000).

2.5 *Ancillary data*

We used hydrological data collected by CTD to characterize *in situ* conditions at the depth of sample collection, as well as general environmental conditions encountered during the study. A total of 20, 14, and 9 full-depth CTD casts were available for S1, S2, and S3 respectively, which we used to generate estimates of mixed-layer depth (MLD), average mixed-layer temperature (T) and salinity, and MLD integrated chl-*a* concentration. In estimating MLD, we adopted a T threshold criterion of $-0.2\text{ }^{\circ}\text{C}$ from a reference depth of 10 m (de Boyer Montégut et al. 2004).

We also used CTD data to estimate surface photosynthetically available radiation (PAR), and depth of the euphotic zone (Z_{eu}). Due to lack of PAR data from <5 m depth, we estimated surface PAR as the y-intercept (depth = 0 m) of a linear regression of the natural log of PAR profiles, the slope of which yields the coefficient of vertical light extinction (k_i). We then used these estimates of k_i to determine Z_{eu} . We adopted the commonly used definition of Z_{eu} as the depth receiving 1% of surface irradiance (Margalef 1978, Reynolds 2006).

2.6 *Statistical analyses*

Principal component analysis (PCA) was used to summarize environmental variability. Included in the analysis were data from CTD casts used to collect water for the experiments of *in situ* temperature and salinity, and estimates of MLD, and Z_{eu} . Before analysis, non-normally distributed data were log-transformed, and to place all

variables on a comparable dimensionless scale, data were normalized to a mean of zero and a standard deviation of 1.

Patterns in the composition of the diatom and of the HP assemblages were investigated using the non-parametric multivariate statistics package Primer-E (Plymouth Routines in Multivariate Ecological Research, version 6; Clarke & Gorley 2006). To visualize multivariate patterns, multi-dimensional scaling (MDS) and cluster analyses (Clarke 1993) were performed on Bray-Curtis index-based similarity matrices. Those were obtained using biomass data that were 4th root transformed to even out contribution of all groups. Points close together represent samples that are similar in species composition. Stress values indicate how well the 2-D plot summarizes the rank-order relationships between samples. Values of stress <0.1 are considered to correspond to a good ordination and values <0.2 provide a less satisfactory but still useful picture. Statistical routines were performed to explore correlations between biotic and environmental patterns using Spearman rank correlation (RELATE).

Plankton biomass-based similarity matrices were also used to further compare plankton assemblages based on location (3 levels: S1, S2, S3) and on grazing magnitude. The latter factor was partitioned into three different levels of grazing activity relative to the overall average (zero, below average, and above average). To examine if species composition influenced whether grazing occurred at all, the analysis was repeated using only two levels (0= no grazing, 1= grazing).

To assess the nature and strength of relationships between species composition samples, analysis of similarities (ANOSIM) was performed on biomass-based

resemblance matrices. ANOSIM is a non-parametric permutation procedure that computes the global R statistic, which can range from -1 to 1, although negative values are unlikely (Clarke & Warwick 2001). Values approaching 1 indicate greater similarities within a group than among groups, whereas values approaching zero indicate equal similarities within and among groups (e.g. no group associations/clustering).

Additionally, a series of univariate analyses (linear regression and Pearson correlation) were performed using SigmaPlot[®] software to examine relationships between grazing rates and a series of potential driving factors. All statistical analyses were performed at an alpha level of 0.05. All rates and other estimates are expressed $\pm$ one standard deviation from the mean.

3. Results

3.1 *Spatial patterns of in situ conditions*

Contrasting environmental conditions over the sampling period distinguished the three stations. Data from all CTD casts performed over the entire duration of the cruise provided evidence that temporal variation in physical parameters were greater among than within stations.

Stations significantly differed in MLD (ANOVA, $p < 0.001$). S1 had a deep mixed layer, which averaged 476 ± 149 m and showed no shoaling progression. MLD at S2 was one order of magnitude shallower than at S1, averaging 46 ± 16 m. At the shallow (160 m) shelf station S3, MLD always reached the bottom. Consequently, MLD at S1 was repeatedly deeper and MLD at S2 was generally shallower than the euphotic depth, which was estimated to average 70 m (± 18 and 10 m at S1 and S2 respectively). On the shelf at S3, MLD was always deeper than the mean euphotic depth of 50 ± 10 m.

Stations also differed in temperature (T) and salinity. Mixed-layer average T was warmest at S1, where over the sampling period, it averaged $8.61 (\pm 0.23)$ °C, and T was coldest at S2 where it averaged $6.90 (\pm 0.24)$ °C. At S3, T averaged $7.77 (\pm 0.15)$ °C. Differences in mixed-layer average salinity among stations were small yet distinctive, averaging $35.28 (\pm 0.04)$, $35.18 (\pm 0.02)$, and $35.36 (\pm 0.01)$ PSU at S1, S2, and S3 respectively.

Accordingly, conditions recorded *in situ* at the time of sampling (Table 1) were characterized by significant differences among stations (ANOSIM global $R = 0.796$, p

= 0.002) driven by differences in *in situ* T and MLD (Fig. 2). Together the first two axes of the PCA explained 89.6% of the variance of the *in situ* data.

3.2 *Species composition of the plankton assemblage*

There were clear spatial differences in the species composition of the plankton assemblage of S1 and S2, as station-specific samples taken over a period of 33 days resembled each other most (Fig. 3). Both the diatom (Fig. 3a) and the HP assemblages (Fig. 3b) from each sample were strongly associated with location (ANOSIM $p \leq 0.002$), and S1 and S2 differed the most ($p = 0.002$). Temporal variability of HP assemblage among station-specific samples was greater at S1 than at S2, whereas the reverse was true for diatoms, which at S2 were scarce (see below). At S3, a shift in phytoplankton species composition and biomass (see below) caused the diatom assemblage in the two experiments to be <40% similar. Corresponding HP assemblage samples were <50% similar, with the sample from the 1st experiment at S3 resembling those of S2 the most. Both the diatom and the HP assemblages correlated with the multivariate pattern of environmental data characterized by the PCA (RELATE Spearman correlation= 0.518 and 0.47 respectively, $p = 0.002$).

3.2.1 *Phytoplankton community*

Initial chl-*a* levels during our experiments ranged from 0.17 $\mu\text{g L}^{-1}$ at S1 to 2.65 $\mu\text{g L}^{-1}$ at S3, and averaged 1.02 (± 0.54), 0.71 (± 0.22), and 1.60 (± 1.49) $\mu\text{g L}^{-1}$ at S1, S2, and S3 respectively (Table 2). Stations differed in which size fraction dominated the autotrophic community. At S1, except for the 1st visit when autotrophic

biomass was low and dominated by picoplankton, the $>50\ \mu\text{m}$ chl-*a* fraction made up of diatoms came to make up $\sim 50\%$ of total chl-*a* (M. Paulsen pers. comm.), yet diatom biomass in our samples remained low (Table 3), fluctuating between $0.3\text{--}16\ \mu\text{g C L}^{-1}$. Based on a C:chl-*a* ratio of 21 (Bratbak et al. 2011), diatoms would have comprised 9–40% of total chl-*a*, increasing to their maximum on April 10 and decreasing thereafter. The genera *Chaetoceros* and *Pseudonitzschia* dominated the diatom community, respectively representing up to 86 % (April 9) and 62% (April 28) of total diatom biomass. In contrast at S2, diatoms were quasi-absent (average biomass $<0.1\ \mu\text{g C L}^{-1}$), and the $>10\text{-}\mu\text{m}$ chl-*a* fraction never exceeded 8 % of total chl-*a* (M. Paulsen pers. comm.). Although not included in estimates of autotrophic biomass, a large number of cryptophytes identified as *Teleaulax* spp. were present in the S2 samples. At S3, diatoms increased in biomass from 0.56 to $87.7\ \mu\text{g C L}^{-1}$ between the 1st and the 2nd visits (Table 3). At the 2nd visit, the species *Detonula pumila* and *Ditylum brightwellii* together constituted most of the diatom biomass (64 % and 25 % respectively).

3.2.2 Heterotrophic-protist assemblage

Total HP biomass varied from $1.2\ \mu\text{g C L}^{-1}$ (S1) to $10.4\ \mu\text{g C L}^{-1}$ (S2), and averaged $2.8 \pm 1.3\ \mu\text{g C L}^{-1}$, $6.4 \pm 2.9\ \mu\text{g C L}^{-1}$, and $4 \pm 3.1\ \mu\text{g C L}^{-1}$ at S1, S2, and S3 respectively (Table 3). Aloricate ciliate biomass represented an average of $54 \pm 20\ %$, $88 \pm 6\ %$, $82 \pm 7\ %$ of the corresponding total heterotrophic protist biomass. Aloricate ciliates included strobilidiid species of the genera *Lohmanniella* and *Leegaardiella*, as well as species of the genus *Strombidium*. The majority (45–100% total biomass) of aloricate ciliates were $20\text{--}35\ \mu\text{m}$ (Fig. 4a & 4c), however, at S1 there was a temporal

increase in the proportion of the $>50\ \mu\text{m}$ ciliate size fraction from 0-47 % (Fig. 4a), whereas such large ciliates were absent from S2 (Fig. 4b). At the 2nd visit at S3, 71% of aloricate ciliate biomass was made up of organisms $>50\ \mu\text{m}$. Only four tintinnid ciliate genera (*Acanthostomella* sp., *Parafavella* sp., *Salpingella* sp., and *Stenosomella* sp.) were observed across stations, always in low numbers.

Dinoflagellate types differed with station. At S1, on all dates except for the 1st visit, 50-100 % of dinoflagellates were athecate gymnodinoid species. When thecate dinoflagellates were present, *Protoperidinium* spp. made up an average of $43 (\pm 39)\%$ of their biomass. At S2, an average of $52 (\pm 16)\%$ of dinoflagellate biomass was made up of small unidentifiable thecate forms. Some of these cells may have been autotrophs, and thus may have erroneously contributed to our estimates of total heterotrophic biomass, although their contribution only amounted to $0.1 - 1.5\ \mu\text{g C L}^{-1}$. These small forms also dominated among dinoflagellates at the first visit at S3, when the $<10\text{-}\mu\text{m}$ size dominated total chl-*a*. Size distribution of dinoflagellates varied among experiments (Fig. 4b & 4d), but at S1, dinoflagellates $>50\ \mu\text{m}$ represented ~50% of all dinoflagellates on three dates coinciding with experiments that yielded the three highest grazing rates (Fig. 4b). Such large dinoflagellates were never observed at S2 (Fig. 4d).

There was no within-station correlation between heterotrophic biomass and chl-*a* concentration (Pearson correlation, S1 & S2 $p=0.83$). One concern was that collection depth, which differed among experiments, might have affected concentration of protistan grazers and by extension grazing rates, but it did not

significantly influence either their numerical abundance ($p>0.45$) or their biomass ($p>0.43$).

3.3 *Results of dilution experiments*

Over the entire sampling period and across all stations, phytoplankton growth rates ranged from -0.06 to 0.63 d^{-1} and mortality rates due to HP grazing ranged from 0 to 0.56 d^{-1} (Table 2). In all but three measurements, growth rates exceeded grazing mortality rates (Fig. 5). The magnitude and variability of growth and grazing rates at S1 and S2 differed, with S1 exhibiting both higher rates and higher variability. At S1 growth and grazing rates varied over the same range (0 to 0.6 d^{-1}), although average growth rate $0.35 (\pm 0.03)$ exceeded average grazing rate $0.25 (\pm 0.04)\text{ d}^{-1}$ (Table 2). There was one exception to the general decoupling between growth and grazing rates at S1: on April 10, rates were highly coupled (0.60 and 0.56 d^{-1} respectively), and corresponded to the highest initial concentration of chl-*a* ($1.9\text{ }\mu\text{g L}^{-1}$) of all experiments (Table 2).

At S2, growth rates ranged from 0.18 to 0.41 d^{-1} and grazing rates ranged from 0.11 to 0.34 d^{-1} . Growth and grazing rates had similar averages ($0.24 \pm 0.02\text{ d}^{-1}$ and $0.22 \pm 0.03\text{ d}^{-1}$ respectively) (Table 2). On the last two sampling dates, the balance between phytoplankton growth and grazing rates changed from positive to negative.

On the Norwegian shelf (S3), only two experiments were performed at a two-week interval. The first experiment yielded no detectable grazing, and a very low grazing rate (0.04 d^{-1}) was measured the second time (Table 2), whereas phytoplankton growth rates were similar on both dates (0.23 and 0.27 d^{-1}).

Translated into carbon, HP grazed 0.44-21.95 and 1.4-7.29 $\mu\text{g C L}^{-1}$ per day at S1 and S2 respectively, which represented 25-400 % and 25-100 % of their body carbon at S1 and S2 respectively (Fig. 6). Biomass-specific ingestion rates varied between 0.0 and 9.0 d^{-1} at S1, where they averaged $2.43 \pm 3.44 \text{ d}^{-1}$, and varied between 0.29 and 0.71 d^{-1} at S2, where they averaged $0.53 \pm 0.14 \text{ d}^{-1}$.

3.4 *Influence of grazing on dynamics of phytoplankton biomass*

The two oceanic stations differed in the level to which *in situ* chl-*a* variations followed the dynamics inferred from the rates. Based on the average balance between growth and grazing rates and assuming no other losses than grazing, phytoplankton population at S1 would (on average) have doubled approx. every week, whereas at S2, it would have doubled approx. every month. At S1, measured variations in chl-*a* did not match those inferred by the rate estimates ($R^2 = 0.10$, $p = 0.61$) (Fig. 7a). We measured a 10-fold increase from 0.2 to 1.9 $\mu\text{g L}^{-1}$ between March 26 and April 10, which clearly exceeded the ~zero growth rates measured in the first two experiments. Initial experimental chl-*a* remained $\sim 1 \mu\text{g L}^{-1}$ for the rest of the sampling period despite growth rates exceeding grazing rates. CTD profiles, however, show various sub-surface chl-*a* increases at S1 beyond the maxima measured in our experiments (e.g. Fig. 8), which corresponded to small temperature increases, and after which (from April 18) the vertical extent of chl-*a* was well below the euphotic zone and closely coincided with MLD (Fig. 8). Based on CTD data, MLD-integrated chl-*a* concentration increased from $\sim 40 \text{ mg m}^{-2}$ at the 1st visit to 230-250 mg m^{-2} during visits 3 and 4.

In contrast with S1, at S2 the observed variation in chl-*a* closely matched the balance between experimentally estimated rates ($R^2 = 0.98$, $p = 0.009$) (Fig. 7b). During the 1st half of April, phytoplankton growth rates exceeded grazing rates and surface chl-*a* doubled from a mean of ~ 0.5 to $\sim 1.0 \mu\text{g L}^{-1}$. For the last two experiments at the end of April, grazing rates exceeded phytoplankton growth rates. Coinciding with the decoupling of growth and grazing rates on April 23 (balance = -0.2 d^{-1}), an overnight 20 % decrease in chl-*a* was observed. This decrease also corresponded to an overnight change in MLD from 29 m to 68 m.

At S3, based on estimated rates and assuming no other losses than grazing, phytoplankton population would have doubled every ~ 3 days, twice more often than indicated by the two-week increase in chl-*a* from 0.5 to $2.7 \mu\text{g L}^{-1}$. This 5-fold increase coincided with a tripling of HP biomass. The increase in HP biomass corresponded to a 31 % decrease in grazer abundance and thus to a shift to a four times larger average size of grazers.

Overall impact of grazing by HP on primary production (% PP) averaged $66 (\pm 66)$ %. Despite positive grazing rates being generally higher at S1 than at S2, the grazing impact was on average highest at S2 (Table 2). At S1 it averaged $50 (\pm 37)$ %, varying from 0-94 %. At S2 PP consumed averaged $106 (\pm 80)$ %, varying from 45-242 % (Table 2). At S3, the average PP consumed was 8%. (Table 2.)

3.4 *Few specific drivers of protistan herbivory*

Several factors potentially governing the magnitude of grazing rates were examined. Experiments that yielded no detectable grazing were not included in

univariate analyses. No correlation existed between grazing magnitude and either *in situ* or incubation temperature ($p \geq 0.85$). One of the parameter of interest in this study was MLD, but we found no within-station significant correlation between MLD and grazing rates (Pearson, $p \geq 0.23$). At S2 however, the lowest grazing rate (0.11 d^{-1}) was measured on April 13, when unlike all other times, the sample collection depth (35 m) was greater than our estimate of MLD (30 m), and when ciliate biomass was the lowest ($2.3 \mu\text{g C L}^{-1}$). Furthermore, maximum S2 grazing rate (0.34 d^{-1}) was measured when MLD was the shallowest (29 m). For S1 and S2 combined, grazing rates decreased with increasing collection depth ($R^2 = 0.54$, $p = 0.016$), but within-station relationship was not significant ($R^2 = 0.72$ and 0.34 and $p = 0.069$ and 0.304 for S1 and S2 respectively). When combined, S1 and S2 grazing rates significantly correlated with chl-*a* (Pearson coefficient= 0.721 , $p = 0.019$), but no significant correlation existed within each station ($p \geq 0.18$). No significant correlation existed between grazing rates and either HP biomass ($p = 0.94$ and 0.08 for S1 and S2 respectively), or HP numerical abundance ($p > 0.58$). At S2, daily grazing rates transformed into carbon ingested per day using a C:Chl ratio of 21 tended to covary with HP biomass (Fig. 6) but the correlation was not statistically significant ($\rho = 0.837$, $p = 0.077$). We further investigated the effect of species composition of each the autotrophic and the heterotrophic assemblage on grazing patterns/rates, using indices of grazing rate magnitude as a factor in analyses (see methods) and found no significant correlation.

4. Discussion

Our study is, to our best knowledge, the first among a plethora of published field measurements (e.g. Calbet & Landry 2004, Schmoker et al. 2013) to provide estimates of *pre-bloom* heterotrophic protist grazing rates in the subpolar North Atlantic. Such rare estimates are much needed to complement proposed hypotheses (Sherr & Sherr 2009, Rose & Caron 2007, Irigoien et al. 2005) about the role of HP feeding in the development of phytoplankton blooms. We also examined how MLD may have modulated the balance between μ and g , a process that has been proposed to be a major factor controlling variations in phytoplankton biomass, including when the spring bloom initiates (Behrenfeld 2010).

In this study, significant, positive rates of protistan herbivory were often measured, representing a potentially substantial loss of phytoplankton biomass, yet for the most part grazing could not keep pace with phytoplankton growth, regardless of MLD. Our findings further suggest that, at the two open ocean sites with contrasting MLD, different processes were driving phytoplankton losses from the surface layers.

4.1 *HP grazing rates and grazing impact*

Major assumptions of the dilution method and deviations thereof have been discussed at length (Dolan et al. 2000, Moigis 2006, Agis et al. 2007, and others summarized in Schmoker et al. 2013) and are not addressed here. More rarely mentioned is that the dilution method assumes that mortality rates are entirely due to grazing, when they may include phytoplankton mortality due to physiological senescence (Franklin et al. 2006), and perhaps more importantly viral lysis. Although

the magnitude of virus-induced mortality is poorly constrained due to the limitations of available methods, it may be significant (Brussaard 2004), potentially varying with the type of trophic interactions regulating carbon flow within the plankton (Ory et al. 2010). Due to their influence on carbon and energy fluxes (Moore et al. 2004, Suttle 2007), these two processes would deserve to be more routinely evaluated.

This being acknowledged, our results indicate that HP collected from F-max were active grazers of chl-*a* at the two oceanic sites sampled during this study, consuming 26-94 % and 45-242 % of daily PP at S1 and S2 respectively, i.e. >60% in 9 out of 11 experiments that yielded >0 grazing. With few exceptions, these values are similar or greater than the average estimate for other oceanic regions (70%) or polar and temperate regions (60%) (Calbet & Landry 2004). Grazing impact was however variable, as is characteristic of most studies, including previous studies conducted in the region at different times spanning May to July (Table 4).

In contrast to the % PP consumed, our estimates of grazing rates, which ranged from 0 to 0.56 d⁻¹, were at the lower end of the range of rates (0-1.48 d⁻¹) measured in previous studies (Table 4). Average grazing rates at the two oceanic stations differed slightly (by 0.03 d⁻¹) with a maximum difference of 0.2 d⁻¹. When comparing the two sites, temperature (T) difference has to be considered, since T influences ingestion rates (Hansen et al. 1997, Verity et al. 2002). Based on published values of Q₁₀ for ingestion rates (Verity et al. 2002), the largest difference in T between the two oceanic stations (2°C) could have produced a ~0.1 d⁻¹ difference in grazing rates, thus T may have marginally impacted the magnitude of the rates measured.

Grazing rates were particularly variable at S1. HP distribution has been found to be patchy at very fine scales (Montagnes et al. 1999), which may inherently confer variability among grazing rates resulting from different experiments due to differences in the species and size composition of HP assemblages. The relationship between variability of HP assemblage and grazing rates seem to be corroborated by our observations that HP assemblages were more variable at S1 than at S2. Furthermore, advection of adjacent water masses within the same study site is an unavoidable part of Eulerian studies of plankton in the open ocean (Aksnes et al. 1997) and may also explain some of the variability. Pelagic phytoplankton is often distributed in concentrated patches or layers, which can promote predators' aggregation (Menden-Deuer & Grünbaum 2006) and increase feeding rates (Menden-Deuer & Fredrickson 2010). For example, it is possible that for the experiment on April 10, when a sudden and rapid doubling of chl-*a* coincided with maximum estimates of μ and g , which were unusually coupled, results may have been affected by horizontal advection of a plankton patch. Treating our results from April 10 as an “outlier” would bring a temporal pattern to the rates we measured, with the first phase of the sampling period being associated with $\leq 0.1 \text{ d}^{-1}$ growth and grazing rates, followed by a steady increase to their highest values on the last sampling date. We were not, however, able to firmly establish if sampling occurred in a different water mass.

4.2 *MLD and mechanisms of uncoupling between μ and g*

In this study, we wanted to address the question of how much MLD may modulate the balance between μ and g . The “dilution-recoupling” hypothesis proposes

that winter mixed layer deepening dilutes predators and prey, reducing grazing and causing accumulation rates to become positive, whereas “recoupling” occurs once the mixed layer deepening stops, and eventually biomass accumulation rates become negative when vernal stratification sets in (Behrenfeld 2010). Most of the experiments we conducted yielded growth rates that exceeded grazing rates, providing a potential mechanism for phytoplankton biomass to accumulate and potentially form a bloom, regardless of mixed layer depth. MLD did seem to influence the magnitude of the balance between μ and g , which was larger at S1 than at S2. Nevertheless, although the magnitude of the balance is important in setting the accumulation rate, it is its sign (>0 or <0) that ultimately controls the potential for biomass to accumulate. The majority of our results indicate that MLD was not a main determinant of whether the μ - g balance was positive or negative. In particular, although there were exceptions, the fact that growth rates exceeded rates of grazing losses, including at S2, where the mixed layer depth was approx. half as deep than the euphotic depth, suggests that stratification by itself may not always be sufficient for grazing to become large enough to decrease phytoplankton biomass. At S2, early stratification had occurred. Although its mode of formation has not been firmly established, it was unlikely related to vernal warming, as air temperature ranged between 2-5 °C. Despite of stratification, food and/or temperature conditions at the end of winter may not have been conducive to active growth of HP, at least in the first part of the sampling season, when growth rates exceeded grazing rates.

Sherr & Sherr (2009) presented evidence that HP cannot prevent blooms due to generally low food availability, leading to low growth rates and low biomass of HP

under non-bloom conditions. We may have underestimated HP biomass by not including heterotrophic nanoflagellates (HNAN), which can contribute significantly to HP biomass (Stelfox-Widdicombe et al. 2000). According to our estimates, HP biomass remained low for the entire sampling period at all stations, particularly at S1 where it corresponded to the lower limit of the range of values previously published for the region at other times of the year (Table 4). Our statistical analyses did not provide evidence of a direct relationship between HP biomass and grazing rates. To further examine this relationship, we calculated biomass-specific grazing rates (G_{HP}) and compared them to maximum laboratory-determined rates. Published estimates of G_{HP} generated by laboratory experiments reported in Hansen et al. (1997) are almost always $>2\text{ d}^{-1}$ and many are $>6\text{ d}^{-1}$. These estimates come from experiments conducted for the most part at 18-20 °C. In comparison, our estimates exceeded 6 d^{-1} only once. At all other times, G_{HP} values were $<3\text{ d}^{-1}$ at S1 and $<1\text{ d}^{-1}$ S2, and even when adjusted for temperature, these values remain low, especially at S2. Overall low values of G_{HP} could have resulted from the C:Chl ratio of 21 used in their estimates. Although low, in our study this ratio may have been representative of the physiological state of phytoplankton having to acclimate to early spring low levels of light and active vertical mixing in abundant nutrients, all of which would contribute to substantial cell⁻¹ chl-*a* (Geider1987). Using this ratio to determine the relative contribution of diatoms to total chl-*a* yielded estimates that adequately compared to estimates made by other investigators on the cruise (M. Paulsen, pers. comm.). Furthermore, higher estimates of HP biomass (by including HNAN) would further reduce G_{HP} values. Low values indicate that HP were feeding below their potential

rates. Although the smallest grazers may have been feeding exclusively or alternatively on bacteria, generally low G_{HP} suggest HP may have been food limited. For the majority of the experiments, HP were found to consume between 25 and 100% of their body carbon daily. At S1, however, these values varied, and herbivory represented up to 400% of HP carbon biomass, but values <500% are considered low rations (Burkill et al. 1993). These values would again indicate that HP were food limited, although feeding on alternative prey such as bacteria or other protists cannot be ruled out.

The uncoupling between μ and losses needed for phytoplankton biomass to accumulate can be achieved if g is kept low or if μ is large/increases. At the time our study took place, mixed layer deepening at S1 had stopped. Instead we had entered the period when hypothesized increases in grazing pressure are compensated by increases in μ in response to improved light conditions (Behrenfeld 2010). Contributing to the uncoupling at S1, were high values of μ : in two thirds of the experiments at S1 growth rates were equivalent to doubling times of 1-2 days. Our estimates of μ were based on changes in chl- a , and thus could have been overestimated if differences in the light regime experienced by the cells in incubation vs. *in situ* caused the cells to increase their pigment concentration during the incubation period, however such photo-acclimation may be too slow to have significantly affected growth rate estimates measured over a 24-hour period (Landry et al. 1995). Increases in surface chl- a recorded in CTD profiles, whether they indicate *in situ* growth or horizontal advection, do provide evidence of the capacity of phytoplankton to sustain substantial growth rates at the latitudes and time of year we sampled. In fact, such increases in surface

phytoplankton biomass prior to stratification are characteristic of the process leading to the bloom climax (Townsend et al. 1994, Backhaus et al. 2003).

Furthermore, at S1, a difference in which size fraction of the phytoplankton had highest growth rates and which size was consumed may have maintained the uncoupling between μ and g . In particular, diatoms likely enhanced μ , but may not have been readily consumed. This remains speculative, as we did not measure either μ or g for different size fractions separately. Nonetheless diatoms can grow at high rates (Smayda 1997 cited in Tillmann 2004) and are physiologically adapted to the highly variable and low light regime induced by frequent mixing (Weeks et al. 1993). Thus diatoms likely enhanced total phytoplankton growth rates. Interestingly, μ values at S2, where very few diatoms were observed, were in general lower than at S1.

Furthermore, measures of HP grazing on mixed phytoplankton assemblages often show higher grazing rates on small cells (Gifford et al. 1995, Strom et al. 2007). In their investigation of taxon-specific grazing, Gaul & Antia (2001) reported grazing avoidance of diatoms and selective preference for small cells, although in their study, selective grazing may have been driven more by active growth of prey than by prey size. Although HP as a group can graze on a broad range of prey sizes, and individual grazers can adapt their own morphology to the size of the available prey, not all grazers can feed on all sizes (Strom 2002). Heterotrophic dinoflagellates are often the major consumers of diatoms (Lawrence & Menden-Deuer 2012, Levinsen & Nielsen 2002, Sherr et al. 2013, 2009, Strom & Frederickson 2008), but they were not abundant in our study. Interestingly, the highest grazing rates measured at S1 were associated with few, but at other time absent, $>50\ \mu\text{m}$ dinoflagellates. Dinoflagellates'

growth rates, low in comparison with other HP (Tillmann 2004), may have prevented them to keep pace with diatom growth. Ciliates, which typically feed on smaller particles (Strom 2002), may not have been actively consuming diatoms. Although they have been observed to feed on diatoms (Sherr et al. 2013), this type of feeding is likely restricted to larger ($>50\ \mu\text{m}$) types (Aberle et al. 2007), a few of which only appeared at S1 after 4/18, possibly responding to an increase in larger prey. Similarly at S3, dinoflagellate biomass was among the lowest we observed during our study ($<0.4\ \mu\text{g L}^{-1}$) and did not increase during the two weeks between sampling dates.

Simultaneously, ciliate size substantially increased concurrently with the change in the phytoplankton size distribution. Yet even the larger ciliates may only have been able to feed on diatoms at a slow rate due to a possible increase in the time needed to handle the prey (Irigoien et al. 2005). Clearly more has to be learned about the relation between the size structure of the phytoplankton community and the prey preferences and feeding interactions of the various predators. The size-related loophole hypothesis proposed by Irigoien et al. (2005) may well apply to early spring phytoplankton dynamics at high latitudes, when diatoms first start to grow.

High variability of the physical environment at S1 may have produced a patchy grazing response. Small but distinct surface increases in T recorded in CTD profiles support the idea that oceanic heat uptake was at times sufficient to confer stability to the water column and stall convective mixing, which even if transient, was sufficient to provide a window of opportunity for growth (Townsend et al. 1994, Taylor & Ferrari 2011). Such periods of increased water column stability were intermittent with periods of deep mixing, as evidenced by the presence of substantial chl-*a* at large

depths. This variability in the convective mixing regime, which is believed to be diurnal (Taylor & Stephens 1993), may have influenced the variability of S1 grazing rates, HP biomass, and HP species composition among experiments.

Thus at S1, a slow response of HP to increases in prey due to the episodic nature of the physical environment, and a seasonal shift in the size structure of the prey field, may have favored the uncoupling between μ and g .

4.4 *Mixed layer depth and grazing control of phytoplankton biomass*

At S1, the balance between μ and g could not account for ambient (i.e. depth of collection) chl-*a* variations. We are mindful of the caveats and assumptions associated with comparing observed rates of change in chl-*a* with the balance between experimental estimates of μ and g . Differences between the two may inevitably result from the difficulty to duplicate in incubation experiments all field conditions that can affect chl-*a* concentrations. This may be particularly true in regions of active deep mixing. In incubation bottles, plankton assemblages are artificially kept at one depth and isolated from the ambient turbulence, which vertically re-distribute plankton cells, thus changing light availability (Ross et al. 2011), and possibly altering encounter rates between predators and prey. Such differences can lead to experimental estimates that vary from true *in situ* values. Furthermore, our sampling frequency imposed long time intervals between experiments, obviously producing gaps in our data.

While the described caveats may have contributed at S1 to the lack of agreement between ambient changes in chl-*a* and the balance between μ and g , most previous studies have shown that the balance between phytoplankton growth and HP

grazing rates can rarely account for the variability of chl-*a*, possibly due to a general lack of equilibrium between growth and loss processes (Schmoker 2013). Such observations have been made even when sampling frequency was high (Lawrence & Menden-Deuer 2012). Furthermore, ambient departures from the μ - g balance may be inevitable considering that the dilution method provides estimates of *potential* grazing rates, which are obtained from a truncated plankton assemblage, from which mesozooplankton grazers, known to feed both on phytoplankton and heterotrophic protists (Calbet & Saiz 2005, Saiz & Calbet 2011), have been removed. The importance of such artifact in dilution experiments remains to be determined (Schmoker et al. 2013).

Nevertheless a poor match between ambient chl-*a* variability and the balance between our rate estimates would suggest a minimal control of grazing relative to other processes on the dynamics of chl-*a*. In particular, our data do not support the idea that decreases of surface chl-*a*, observed at S1 both in our experiments and in chl-*a* vertical profiles, were due to grazing, when phytoplankton growth rates exceeded grazing rates. In contrast to surface layers, vertically integrated phytoplankton accumulation rate was overall positive. The presence of chl-*a* below the euphotic zone, where phytoplankton growth cannot be sustained, indicates that down-mixing was a major loss process of primary production from the surface layers. As Backhaus et al. (2003) justly remarked, if concentrated within a shallow mixed layer, the observed vertically integrated biomass would be similar or even surpass spring bloom concentrations. Additionally, as the mixed layer shoals, some of the phytoplankton will inevitably become trapped below the thermocline (Backhaus et al. 2003,

Behrenfeld et al. 2013). Thus intermittent deep mixing in early spring may be an important mechanism for export of carbon, which may exceed sinking losses associated with surface blooms, much of which may be respired through grazing (Behrenfeld 2010).

In contrast with S1, at S2 the balance between phytoplankton growth and grazing-mortality rates was a reasonably good predictor of *in situ* phytoplankton population dynamics. This suggests that losses due to sinking may have been limited, and that the majority of the losses incurred were due to grazing, making grazing an important determinant of variations in phytoplankton biomass. Among phytoplankton species, diatoms are believed to drive carbon export because their heavy silicate frustules cause them to sink (Sarhou et al. 2005, Smayda 1970). At S2 the phytoplankton community was dominated by pico- and nanophytoplankton, and diatoms were rare. Small particles are less likely to sink, and their vertical retention may increase grazing opportunities. Thus the importance of grazing in determining variations in ambient phytoplankton biomass at S2 was likely influenced by species composition of the phytoplankton community.

We could not firmly establish the source of stratification at S2. Mesoscale variabilities are frequent in the Norwegian Sea (Hansen et al. 2010). S2 sat near the Iceland Faroe front, and thus stratification could have resulted from the mix of North Atlantic water transported in the Faroe Current with East Iceland Current water (Hansen & Østerhus 2000), or an eddy could have developed along the front. Early stratification other than through surface warming can also be driven by the formation of eddies induced by the slumping of the North-South density gradient associated with

the latitudinal differences in temperature (Mahadevan et al. 2012). Such stratification is believed to trigger early patchy phytoplankton blooms (Mahadevan 2012) dominated by diatoms (Alkire et al. 2012). At S2, however, no bloom and few diatoms were observed during the 30 days of the study, despite availability of ample macronutrients. Low contribution of diatoms to total phytoplankton has previously been observed in a nutrient-rich mesoscale eddy with shallow mixed layer (Stelfox-Widdicombe et al. 2000). Thus eddy-stratification may not always result in PP being dominated by diatoms.

4.5 Drivers of grazing magnitude

In this and previous studies (e.g. Lawrence & Menden-Deuer 2012, Strom et al. 2007), drivers of grazing magnitude remain elusive. As in previous studies (Lawrence & Menden-Deuer 2012, Menden-Deuer & Frederickson 2010, Sherr et al. 2009), there was no direct relationship between chl-*a* and grazing rates, confirming that many factors other than a coarse metric for phytoplankton quantity can affect grazing rates (Sherr & Sherr 2007). Failing to identify specific drivers of grazing magnitude may be due to the fact that the dilution method provides bulk estimates of grazing, which result from a poorly constrained multitude of complex feeding interactions. Planktonic trophic links include mixotrophy and omnivory (Caron et al. 2012, Flynn et al. 2012) and trophic cascades (Calbet & Saiz 2013). They involve taxonomically diverse organisms, which span a large size range and exhibit a variety of prey preferences and prey selection (Caron et al. 2012, Montagnes et al. 2008), and whose feeding behaviors respond in specific ways to the surrounding physical,

chemical, and biological conditions (Caron & Hutchins 2012). Teasing apart planktonic food webs both through laboratory experiments and in the field remains challenging but necessary to increase our ability to predict grazing losses.

Concluding remarks

Our study is one of the first to document HP grazing for the subpolar North Atlantic during the early spring period that precedes the seasonal increase in surface phytoplankton biomass. Although models can resolve the effect of physical and biological forcing on PP much more comprehensively than logistically-intensive field experiments ever will, such models can only be accurately parameterized if field measurements of key rate processes such as grazing are available. Our ability to predict how North Atlantic PP will respond to warming ultimately depends on a better understanding of what controls variations in phytoplankton biomass. Results of several modeling studies (Boyd & Doney 2002, Le Quere et al. 2003, Sarmiento et al. 2004) predict that, at high latitudes, increased thermal stratification will result in greater light and longer growing season afforded to photosynthetic organisms, which should increase present-day light-limited PP (Doney 2006, Riebesell et al. 2009). Although more studies comparing losses incurred by phytoplankton under different conditions of mixing depths are needed to generalize our findings, the different dominant loss factor – sinking in deep mixed layer, and grazing in shallow mixed layer - suggested by our results may imply that a longer period of stratification could reduce the export of organic carbon that occurs due to deep mixing before stratification and the spring bloom climax, whereas more PP could potentially be lost to respiration associated with HP grazing. In the field, higher geographical and temporal sampling resolution will be needed to capture the dynamics of the biophysical factors driving coupling/decoupling between phytoplankton growth and grazing-mortality rates.

APPENDIX

Light and collection depth experiments

This appendix reports and discusses results of two experiments conducted at S2 to investigate the effect of light (3/31) and of collection depth (4/14) on rate magnitudes. These two experiments were conducted in parallel to a regular experiment and similar methods were applied, except that they were incubated at surface irradiance, without the use a mesh-bag.

On March 31, the same bulk assemblage collected from a 20-m depth (night cast) was used in two parallel experiments incubated at two light regimes, in order to assess the effect of light on phytoplankton growth and grazing rates. Phytoplankton growth rate was 35% higher when replicates were incubated at mixed-layer adjusted light ($0.34 \pm 0.04 \text{ d}^{-1}$) than when unprotected by light screening mesh ($0.25 \pm 0.04 \text{ d}^{-1}$). Grazing rates remained unaffected by light (0.26 ± 0.06 and $0.25 \pm 0.1 \text{ d}^{-1}$ respectively).

On April 14, water collected from a 5-m depth was incubated at surface irradiance simultaneously with the experiment using water from F-max on that day. Both depths had similar *in situ* chl-*a* concentration ($0.6 \mu\text{g L}^{-1}$). Heterotrophic biomass was similar in the two samples ($7.1 \mu\text{g C L}^{-1}$ at 5m and $7.9 \mu\text{g C L}^{-1}$ at 30 m). Heterotrophic cells, however, were twice as abundant at 5 m ($9600 \text{ cells L}^{-1}$) than at 30 m ($4800 \text{ cells L}^{-1}$), reflecting a 55% smaller average size of heterotrophic protists at the surface. Phytoplankton growth rates at 5 m were approx. one third lower than at 30 m ($0.12 \pm 0.03 \text{ d}^{-1}$ vs. $0.41 \pm 0.05 \text{ d}^{-1}$), whereas grazing rates were approx. twice higher at 5 m ($0.32 \pm 0.06 \text{ d}^{-1}$) than at 30 m ($0.19 \pm 0.07 \text{ d}^{-1}$).

The decrease in phytoplankton growth rates in response to increased incubation irradiance may underline one of the caveats associated with estimating growth rates using the dilution method based on measured changes of chl-*a* (Landry et al. 1995). Exposing cells to higher irradiance relative to *in situ* likely caused phytoplankton cells to photo-acclimate, i.e. adjust their cellular pigment concentration (Geider 1987), yielding artificially lower estimates of growth. We did not estimate total phytoplankton numerical abundance before and after the experiment, so we cannot be certain whether growth rates were lower due to changes in cellular pigment concentration or an actual difference in the doubling time. In the second experiment, in which plankton assemblage were collected at two different depths, differences in growth rates could have resulted from differences in phytoplankton species composition, which we did not verify. Ross et al. (2011) showed, however, that growth rates based on chl-*a* or on carbon differed considerably from each other, particularly near the surface, due to cells photo-acclimating to higher light intensity by reducing their chl-*a* synthesis while uptaking carbon and thus increasing their C:Chl ratio. In experiments in which plankton were incubated in mesocosms at two different depths (and thus at two light regimes), chl-*a* increase at low light was found to be mostly due to photo-acclimation, and phytoplankton growth rates were higher when plankton were exposed to higher light (Calbet et al. 2012). Photo-acclimation, however, should not alter grazing estimates, as long as it affects all dilutions used to compute grazing rate equally (Landry et al. 1995).

In contrast to growth rates, which responded similarly to light in the two experiments, grazing rates responded differently. In the March 31st experiment, light

showed no effect on grazing. Calbet et al. (2012) measured similar grazing rates in differently illuminated mesocosms. On the other hand, Strom (2001) found that light can enhance ingestion, digestion, and growth rates of herbivorous protists, however the enhancement was relative to rates obtained in total darkness. Thus the difference in light between our two treatments may not have been large enough to produce an effect. Further experiments would be necessary to determine a light threshold above which grazing rates significantly increase. The April 14th experiments contained two variables (depth and incubation light) that confound the interpretation of the results. Nevertheless, if we take clues from the March 31st experiment, in which light only affected growth rates and not grazing rates, then the observed difference in grazing rates between the April 14th treatments were likely due to a difference in collection depth, or to factors associated with it. In the 5-m sample, which yielded the higher grazing rate, heterotrophic protists were twice as abundant but on average half the size of those collected at 30 m, which may have played a role, potentially through an increase in encounter rates. Additionally, since the autotroph community was dominated by small cells, a large proportion of it being composed of picoplankton, grazing rates may have been enhanced because of a better match between the smaller grazers found at 5 m and the size spectrum of the prey.

References

- Aberle, N., Lengfellner, K., Sommer, U., 2007. Spring bloom succession, grazing impact and herbivore selectivity of ciliate communities in response to winter warming. *Oecologia* 150, 668-681.
- Agis, M., Granda, A., Dolan, J.R., 2007. A cautionary note: examples of possible microbial community dynamics in dilution grazing experiments. *Journal of Experimental Marine Biology and Ecology* 341, 176-183.
- Alkire, M.B., D'Asaro, E., Lee, C., Perry, M.J., Gray, A., et al., 2012. Estimates of net community production and export using high-resolution Lagrangian measurements of O_2 , NO_3^- , and POC through the evolution of a spring diatom bloom in the North Atlantic. *Deep-Sea Research I* 64, 157-174.
- Aksnes, D.L., Miller, C.B., Ohman, M.D., Wood, S.N., 1997. Estimation techniques used in studies of copepod population dynamics – a review of underlying assumptions. *Sarsia* 82, 279-296.
- Backhaus, J.O., Nøst Hegseth, E., Wehde, H., Irigoien, X., Hattern, K., Logemann, K., 2003. Convection and primary production in winter. *Marine Ecology Progress Series* 251, 1-14.
- Banse, K. 1994. Grazing and zooplankton production as key controls of phytoplankton production in the open ocean. *Oceanography* 7, 13-20.
- Behrenfeld, M.J. 2010. Abandoning Sverdrup's critical depth hypothesis on phytoplankton blooms. *Ecology* 91, 977-89.
- Behrenfeld, M.J., Boss, E.S., 2014. Resurrecting the ecological underpinnings of ocean plankton blooms. *Annual Review of Marine Science* 4.6, 167-194.
- Behrenfeld M.J., Doney, S.C., Lima, I., Boss, E.S., Siegel, D.A., 2013. Annual cycles of ecological disturbance and recovery underlying the subarctic Atlantic spring plankton bloom. *Global Biogeochemical Cycles* 27, 526-540.
- Boyd, P., Doney, S.C., 2002. Modelling regional responses by marine pelagic ecosystems to global climate change. *Geophysical Research Letters* 29, 53-1-53-4.
- Bratbak, G., Jacquet, S., Larsen, A., Petterson, L.H., 2011. The plankton community in Norwegian coastal waters- abundance, composition, spatial distribution and diel variation. *Continental Shelf Research* 31, 1500-1514.
- Brussaard, C., 2004. Viral control of phytoplankton populations- a review. *Journal of Eukaryotic Microbiology* 51, 125-138.
- Burkill, P.H., Edwards, E.S., John, A.W.G., Sleigh, M.A., 1993. Microzooplankton and their herbivorous activity in the northeastern Atlantic Ocean. *Deep-Sea Research II* 40, 479-493.
- Buesseler, K.O., Lamborg, C.L., Boyd, P.W., Lam, P.J., Trull, T.W., et al., 2007. Revisiting carbon flux through the ocean's twilight zone. *Science* 316, 567-569.
- Calbet, A., Landry, M.R., 2004. Phytoplankton growth, microzooplankton grazing, and carbon cycling in marine systems. *Limnology and Oceanography* 49, 51-57.
- Calbet, A., Saiz, E., 2013. Effects of trophic cascades in dilution grazing experiments: from artificial saturated feeding responses to positive slopes. *Journal of Plankton Research* 35, 1183-1191.
- Calbet, A., Saiz, E., 2005. The ciliate-copepod link in marine ecosystems. *Aquatic Microbial Ecology* 38, 157-167.

- Calbet, A., Martinez, R.A., Isari, S., Zervoudaki, S., Nejstgaard, J.C., et al., 2012. Effects of light availability on mixotrophy and microzooplankton grazing in an oligotrophic plankton food web: Evidences from a mesocosm study in Eastern Mediterranean waters. *Journal of Experimental Marine Biology and Ecology* 424-425, 66-77.
- Caron, D.A., Hutchins, D.A., 2012. The effects of changing climate on microzooplankton grazing and community structure: drivers, predictions and knowledge gaps. *Journal of Plankton Research* 0, 1-18.
- Caron, D.A., Countway, P.D., Jones, A.C., Kim, D.Y., Schnestzer, A., 2012. Marine protistan diversity. *Annual Review of Marine Science* 4, 467-493.
- Caron, D.A., Dennett, M.R., Lonsdale, D.J., Moran, D.M., Shalapyonok, L., 2000. Microzooplankton herbivory in the Ross Sea, Antarctica. *Deep-Sea Research II* 47, 3249–3272.
- Clarke, K. R. 1993. Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* 18, 117-143
- Clarke, K.R., Gorley, R.N., 2006. *PRIMER v6: User Manual/Tutorial*. PRIMER-E, Plymouth.
- Clarke, K.R., Warwick, R.M., 2001. A further biodiversity index applicable to species lists: variation in taxonomic distinctness. *Marine Ecology Progress Series* 216, 265-278.
- Dale, T., Rey, F., Heimdal, B.R., 1999. Seasonal development of phytoplankton at a high latitude oceanic site. *Sarsia* 84, 419-435.
- de Boyer Montégut, C., Madec, G., Fischer, A.S., Lazar, A., Iudicone, D., 2004. Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *Journal of Geophysical Research* 109, C12003.
- Dodge, J.D. 1982. *Marine dinoflagellates of the British Isles*. London : Her Majesty's Stationery Office.
- Dolan, J.R., Gallegos, C.L., Moigis, A., 2000. Dilution effects on microzooplankton in dilution grazing experiments. *Marine Ecology Progress Series* 200: 127-139.
- Doney, S.C., 2006. Plankton in a warmer world. *Nature* 444, 695-696.
- Ferrari, R., Merrifield, S.T., Taylor, J.R., 2014. Shutdown of convection triggers increase of surface chlorophyll. *J Mar Syst* (in press).
- Flynn, K.J., Stoecker, D.K., Mitra, A., Raven, J.A., Glibert, P.M., et al., 2012. Misuse of the phytoplankton dichotomy: the need to assign organisms as mixotrophs within plankton functional types. *Journal of Plankton Research* 0, 1-9.
- Franklin, D.J., Brussaard, P.D., Berges, J.A., 2006. What is the role and nature of programmed cell death in phytoplankton ecology? *European Journal of Phycology* 41, 1-41.
- Gaul, W., Antia, A.N., 2001. Taxon-specific growth and selective microzooplankton grazing of phytoplankton in the Northeast Atlantic. *Journal of Marine Systems* 30, 241-261.
- Geider, R.J. 1987. Light and temperature dependence of the carbon to chlorophyll a ratio in microalgae and cyanobacteria: Implications for physiology and growth of phytoplankton. *New Phytologist* 106, 1-34.
- Gifford, D.J., Fessenden, L.M., Garrahan, P.R., Martin, E., 1995. Grazing by microzooplankton and mesozooplankton in the high-latitude North Atlantic Ocean: Spring versus summer dynamics. *Journal of Geophysical Research* 100, 6665-75.

- Graff, J.R., Ryneerson, T.A., 2011. Extraction method influences the recovery of phytoplankton pigments from natural assemblages. *Limnology and Oceanography-Methods* 9, 129-139.
- Hansen, B., Østerhus, S., 2000. North Atlantic-Nordic Seas exchanges. *Progress in Oceanography* 45, 109-208.
- Hansen, C., Kvaleberg, E., Samuelsen, A., 2010. Anticyclonic eddies in the Norwegian Sea; their generation, evolution and impact on primary production. *Deep-Sea Research I* 57, 1079-1091.
- Hansen, P.J., Moldrup, M., Tarangkoon, W., Garcia-Cuetos, L., Moestrup, Ø., 2012. Direct evidence for symbiont sequestration in the marine red tide ciliate *Mesodinium rubrum*. *Aquatic Microbial Ecology* 66, 63-75.
- Hansen, P.J., Bjørnsen, P.K., Hansen, B.W., 1997. Zooplankton grazing and growth: Scaling within the 2-2,000- μ m body size range. *Limnology and Oceanography* 42, 687-704.
- Henson, S.A., Robinson, I., Allen, J.T., Waniek, J., 2006. Effect of meteorological conditions on interannual variability in timing and magnitude of the spring bloom in the Irminger Basin, North Atlantic. *Deep-Sea Research I* 53, 1601-1615.
- Huisman, J., van Oostveen, P., Weissing, F.J., 1999. Critical depth and critical turbulence: Two different mechanisms for the development of phytoplankton blooms. *Limnology and Oceanography* 44, 1781-87.
- Huisman, J., Arrayás, M., Ebert, U., Sommeijer, B., 2002. How do sinking phytoplankton species manage to persist? *American Naturalist* 159, 245-254.
- Irigoin, X., Flynn, K.J., Harris, R.P., 2005. Phytoplankton blooms: a 'loophole' in microzooplankton grazing impact? *Journal of Plankton Research* 27, 313-321.
- Jespersen, A.M., Christoffersen, K., 1987. Measurements of chlorophyll-a from phytoplankton using ethanol as extraction solvent. *Archiv für Hydrobiologie* 109, 445-454.
- Kraberg, A., Baumann, M., Dürselen, C-D., 2010. Coastal phytoplankton: photo guide for Northern European Seas. München: Verlag Dr. Friedrich Pfeil.
- Landry, M.R., Hassett, R.P., 1982. Estimating the grazing impact of marine microzooplankton. *Marine Biology* 67, 283-288.
- Landry, M.R., Constantinou, J., Kirshtein, J., 1995. Microzooplankton grazing in the central equatorial Pacific during February and August, 1992. *Deep-Sea Research II* 42, 657-671.
- Lawrence C., Menden-Deuer, S., 2012. Drivers of protistan grazing pressure: seasonal signals of plankton community composition and environmental conditions. *Marine Ecology Progress Series* 459, 39-52.
- Le Quéré C., Aumont, O., Monfray, P., Orr, J., 2003. Propagation of climatic events on ocean stratification, marine biology, and CO₂: Case studies over the 1979-1999 period. *Journal of Geophysical Research* 108, 5-1-5-7.
- Levinson, H., Turner, J.T., Nielsen, T.G., Hansen, B.W., 2000. On the trophic coupling between protists and copepods in arctic marine ecosystems. *Marine Ecology Progress Series* 204, 65-77.
- Mahadevan, A., D'Asaro, E., Lee, C., Perry, M.J., 2012. Eddy-driven stratification initiates North Atlantic spring phytoplankton blooms. *Science* 337, 54-58.

- Margalef, R., 1978. Life-forms of phytoplankton as survival alternatives in an unstable environment. *Oceanologica Acta* 1, 493-509.
- Menden-Deuer, S., Fredrickson, K., 2010. Structure-dependent protistan grazing and its implication for the formation, maintenance, and decline of plankton patches. *Marine Ecology Progress Series* 420, 57-71.
- Menden-Deuer, S., Grünbaum, D., 2006. Individual foraging behaviors and population distributions of planktonic predators aggregating to phytoplankton layers. *Limnology and Oceanography* 51, 109-116.
- Menden-Deuer, S., Lessard, E.J., Satterberg, J., 2001. Effect of preservation on dinoflagellate and diatom cell volume and consequences for carbon biomass predictions. *Marine Ecology Progress Series* 222, 41-50.
- Menden-Deuer, S., Lessard, E., 2000. Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton. *Limnology and Oceanography* 45, 569-579.
- Moigis, A.G., 2006. The clearance rate of microzooplankton as the key element for describing estimated non-linear dilution plots demonstrated by a model. *Marine Biology* 149, 743-762.
- Montagnes, D.J.S., Lynn, D.H., 1991. Taxonomy of Choreotrichs, the major marine planktonic ciliates, with emphasis on the aloricate forms. *Marine Microbial Food Webs* 5, 59-74.
- Montagnes, D.J.S., Poulton, A.J., Shammon, T.M., 1999. Mesoscale, finescale and microscale distribution of micro- and nanoplankton in the Irish Sea, with emphasis on ciliates and their prey. *Marine Biology* 134, 167-179.
- Montagnes, D.J.S., Barbosa, A.B., Boenigk, J., Davidson, K., Jürgens, K., et al., 2008. Selective feeding behaviour of key free-living protists: avenues for continued study. *Aquatic Microbial Ecology* 53, 83-98.
- Moore, J.C., Berlow, E.L., Coleman, D.C., de Rulter, P.C., Dong, Q., et al., 2004. Detritus, trophic dynamics and biodiversity. *Ecology Letters* 7, 584-600.
- Ory, P., Hartmann, H.J., Jude, F., Dupuy, C., Del Amo, Y., et al., 2010. Pelagic food web patterns: Do they modulate virus and nanoflagellate effects on picoplankton during the phytoplankton spring bloom? *Environmental Microbiology* 12, 2755-72.
- Putt, M., Stoecker, D.K., 1989. An experimentally determined carbon:volume ratio for marine "oligotrichous" ciliates from estuarine and coastal waters. *Limnology and Oceanography* 34, 1097-1103.
- Reynolds, C.S. 2006. The ecology of phytoplankton. New York: Cambridge University Press.
- Riebesell, U., Körtzinger, A., Oschlies, A., 2009. Sensitivities of marine carbon fluxes to ocean change. *PNAS* 106, 20602-609.
- Rose, J.M., Caron, D.A., 2007. Does low temperature constrain the growth rates of heterotrophic protists? Evidence and implications for algal blooms in cold waters. *Limnology and Oceanography* 53, 886-895.
- Ross, O.N., Geider, R.J., Berdalet, E., Artigas, M.L., Piera, J., 2011. Modeling the effect of vertical mixing on bottle incubations for determining *in situ* phytoplankton dynamics. I. Growth rates. *Marine Ecology Progress Series* 435, 13-31.
- Saiz, E., Calbet, A., 2011. Copepod feeding in the ocean: scaling patterns, composition of their diet and the bias of estimates due to microzooplankton grazing during incubations. *Hydrobiologia* 666, 181-196.

- Sarmiento, J.L., Slater, R., Barber, R., Bopp, L., Doney, S.C., Hirst, A.C., et al., 2004. Response of ocean ecosystems to climate warming. *Global Biogeochemical Cycles* 18, GB 3003.
- Sarthou, G., Timmermans, K.R., Blain, S., Treguer, P., 2005. Growth physiology and fate of diatoms in the ocean: a review. *Journal of Sea Research* 53, 25-42.
- Sathyendranath, S., Stuart, V., Nair, A., Oka, K., Nakane, T., et al., 2009. Carbon-to-chlorophyll ratio and growth rate of phytoplankton in the sea. *Marine Ecology Progress Series* 383, 73–84.
- Schmoker, C., Hernández-León, S., Calbet, A., 2013. Microzooplankton grazing in the oceans: impacts, data variability, knowledge gaps and future directions. *Journal of Plankton Research* 0, 1-16.
- Sherr, E.B., Sherr, B.F., 2009. Capacity of herbivorous protists to control initiation and development of mass phytoplankton blooms. *Aquatic Microbial Ecology* 57, 253-262.
- Sherr, E.B., Sherr, B.F., 2007. Heterotrophic dinoflagellates: a significant component of microzooplankton biomass and major grazers of diatoms in the sea. *Marine Ecology Progress Series* 352, 187-187.
- Sherr, E.B., Sherr, B.F., 2002. Significance of predation by protists in aquatic microbial food webs. *Antonie van Leeuwenhoek* 81, 293-308.
- Sherr, E.B., Sherr, B.F., Ross, C., 2013. Microzooplankton grazing impact in the Bering Sea during spring sea ice conditions. *Deep-Sea Research II* 94, 57-67.
- Sherr, E.B., Sherr, B.F., Hartz, A.J., 2009. Microzooplankton grazing impact in the Western Arctic Ocean. *Deep-Sea Research II* 56, 1264-1273.
- Sherr, E.B., Sherr, B.F., Wheeler, P.A., Thompson, K., 2003. Temporal and spatial variation in stocks of autotrophic and heterotrophic microbes in the upper water column of the central Arctic Ocean. *Deep Sea Research I* 50: 557-571.
- Siegel, D.A., Doney, S.C., Yoder, J.A., 2002. The North Atlantic spring phytoplankton bloom and Sverdrup's critical depth hypothesis. *Science* 296, 730:733.
- Smayda, T.J. 1970. The suspension and sinking of phytoplankton in the Sea. *Oceanography and Marine Biology - An Annual Review* 8, 353–414.
- Smith, W.O., Sakshaug, E., 1990. Polar phytoplankton. In: Smith W.O. (Ed.). *Polar Oceanography, Part B*. San Diego, CA: Academic Press pp. 477–525.
- Stelfox-Widdicombe, C.E., Edwards, E.S., Burkill, P.H., Sleight, M.A., 2000. Microzooplankton grazing activity in the temperate and sub-tropical NE Atlantic: summer 1996. *Marine Ecology Progress Series* 208, 1-12.
- Stoecker, D.S. 1999. Mixotrophy among dinoflagellates. *Journal of Eukaryotic Microbiology* 46, 397-401.
- Strom, S.L. 2002. Novel interactions between phytoplankton and microzooplankton: their influence on the coupling between growth and grazing rates in the sea. *Hydrobiologia* 480, 41-54.
- Strom, S.L. 2001. Light-aided digestion, grazing and growth in herbivorous protists. *Aquatic Microbial Ecology* 23, 253-261.
- Strom, S.L., Fredrickson, K.A., 2008. Intense stratification leads to phytoplankton nutrient limitation and reduced microzooplankton grazing in the southeastern Bering Sea. *Deep-Sea Research II* 55, 1761–1774.

- Strom, S.L., Macri, E.L., Olson, M.B., 2007. Microzooplankton grazing in the coastal Gulf of Alaska: variations in top-down control of phytoplankton. *Limnology and Oceanography* 52, 1480–1494.
- Strüder-Kypke, M.C., Kypke, E.R., Agatha, S., Warwick, J., Montagnes, D.J.S., 2002. The user-friendly guide to coastal planktonic ciliate. <http://www.zooplankton.cn/Default.aspx?tabid=604&language=zh-CN>
- Suttle, C.A., 2007. Marine viruses – major players in the global ecosystem. *Nature Reviews/Microbiology* 5, 801–812.
- Sverdrup, H.U., 1953. On conditions of the vernal blooming of phytoplankton. *Journal du Conseil* 18, 287–295.
- Takahashi, T., Sutherland, S.C., Wanninkhof, R., Sweeney, C., Feely, R.A., et al., 2009. Climatological mean and decadal change in surface ocean pCO_2 and net sea–air CO_2 flux over the global oceans. *Deep-Sea Research II* 56, 554–77.
- Taylor, A.H., Stephens, J.A., 1993. Diurnal variations of convective mixing and the spring bloom of phytoplankton. *Deep-Sea Research II* 40: 389–408.
- Taylor, A.H., Harbour, D.S., Harris, R.P., Burkill, P.H., Edwards, E.S., 1993. Seasonal succession in the pelagic ecosystem of the North Atlantic and the utilization of nitrogen. *Journal of Plankton Research* 15, 875–891.
- Taylor, J.R., Ferrari, R., 2011. Shutdown of turbulent convection as a new criterion for the onset of spring phytoplankton blooms. *Limnology and Oceanography* 56, 2293–2307.
- Thrandsen J., Hasle, G.R., Tangen, K., 2007. Phytoplankton of Norwegian coastal waters. Oslo: Almatier Forlag As.
- Tillmann, U. 2004. Interactions between planktonic microalgae and protozoan grazers. *Journal of Eukaryotic Microbiology* 51, 156–168.
- Townsend, D.W., Cammen, L.M., Holligan, P.M., Campbell, D.E., Pettigrew, N.R., 1994. Causes and consequences of variability in the timing of spring phytoplankton blooms. *Deep-Sea Research I* 4, 747–765.
- Townsend, D.W., Keller, M.D., Sieracki, M.E., Ackleson, S.G., 1992. Spring phytoplankton blooms in the absence of vertical water column stratification. *Nature* 360, 59–62.
- Turner, J.T., 2002. Zooplankton fecal pellets, marine snow and sinking phytoplankton blooms. *Aquatic Microbial Ecology* 27, 57–102.
- Utermöhl, H., 1958. Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. *Internationale Vereinigung für Theoretische und Angewandte Limnologie: Mitteilungen*, 9:1–38.
- Verity, P.G., Langdon, C., 1984. Relationships between lorica volume, carbon, nitrogen, and ATP content of tintinnides in Narragansett Bay. *Journal of Plankton Research* 6, 859–868.
- Verity, P.G., Wassmann, P., Frischer, M.E., Howard-Jones, M.H., Allen, A.E., 2002. Grazing of phytoplankton by microzooplankton in the Barents Sea during early summer. *Journal of Marine Systems* 38, 109–123.
- Weeks, A., Conte, M.H., Harris, R.P., Bedo, A., Bellan, I., et al., 1993. The physical and chemical environment and changes in community structure associated with bloom evolution: the Joint Global Flux Study North Atlantic Bloom Experiment. *Deep-Sea Research II* 40, 347–368.

- Wolfe, G.V., Levasseur, M., Cantin, G., Michaud, S., 2000. DMSP and DMS dynamics and microzooplankton grazing in the Labrador Sea: application of the dilution technique. *Deep-Sea Research I* 47, 2243-2264.
- Worden, A.Z., Binder, B.J., 2003. Application of dilution experiments for measuring growth and mortality rates among *Prochlorococcus* and *Synechococcus* populations in oligotrophic environments. *Aquatic Microbial Ecology* 30, 159-174.
- Zar, J. H., 2010. *Biostatistical Analysis*. 5th Ed. Upper Saddle River, NJ: Pearson Prentice-Hall.

TABLES

Table 1. Details of source water used in dilution experiments that were performed during the M87 cruise, including water collection depth (CTD fluorescence maximum, except when marked *), water temperature (T) and salinity, mixed layer depth (MLD, estimated using potential temperature threshold criteria of -0.2 °C from a reference depth of 10 m), and euphotic depth (Z_{eu} = euphotic depth, i.e. the depth that receives 1% of surface photosynthetically available radiation). Incubations took place at mixed-layer adjusted irradiance. Not included are two experiments conducted at S2 at surface irradiance (See Appendix).

Date	CTD cast #	Water Collection Depth (m)	Dilution levels	T in situ (°C)	Salinity (PSU)	MLD (m)	Z_{eu} (m)
Station 1 (61°30'N – 011°00'W - Iceland Basin – Total depth: 1,345 m)							
26-Mar	424-1	30	4	8.7	35.30	601	108
9-Apr	523-1	30	4	8.6	35.29	524	64
10-Apr	541-1	18	3	8.6	35.30	452	65
18-Apr	611-1	40	5	8.7	35.29	551	47
19-Apr	624-1	35	3	8.6	35.28	642	63
28-Apr	679-1	25	5	8.6	35.28	498	79
Station 2 (62°50'N - 001°00'E - Norwegian Basin – Total depth: 1298 m)							
31-Mar	460-1	20*	3	7.1	35.19	48	Dark
13-Apr	564-1	35	3	6.4	35.14	30	62
14-Apr	578-1	30	5	6.6	35.15	36	65
23-Apr	649-1	20	5	6.8	35.18	29	64
24-Apr	659-1	35	3	7.0	35.19	68	53
Station 3 (60°20'N – 001°00'E - Shetland shelf – Total depth: 163 m)							
2-Apr	487-1	30	5	7.8	35.36	Bottom	63
16-Apr	605-1	30	5	7.8	35.37	Bottom	52

Table 2. Initial chlorophyll-*a* concentration (Chl-*a*, $\mu\text{g L}^{-1}$), phytoplankton growth (μ) and grazing mortality (*g*) rates, and grazing impact as % of primary production (% PP) consumed ($=100 \times g/\mu$). Rates are given per day. Values in parentheses represent standard deviation. In the 2-point analysis (see text for details), rates were determined using the undiluted and the lowest dilution treatments only. Net apparent growth rate in the undiluted treatment (k_l) is also given. Significance of deviation from linearity (DL) for linear regression (alpha= 0.05): NS= not significant S= significant. For calculations, <0 phytoplankton growth and grazing rates (marked *) were set to 0.01 and 0 respectively.

Date	Chl- <i>a</i>	-----2-point analysis-----				-----Regression analysis-----				% PP
		μ	g	k ₍₁₎	DL	μ	g	p	R ²	
Station 1										
26-Mar	0.17 (0.01)	-0.06* (0.001)	0.12 (0.10)	-0.19 (0.10)	NS	-0.11 (0.06)	0	0.431	0.08	n/a
9-Apr	0.99 (0.01)	0.05 (0.13)	-0.10* (0.15)	0.15 (0.06)	NS	-0.05 (0.08)	0	0.168	0.18	0
10-Apr	1.87 (0.03)	0.60 (0.02)	0.56 (0.08)	0.04 (0.04)	NS	0.65 (0.01)	0.61 (0.017)	<0.0001	0.99	94
18-Apr	1.13 (0.04)	0.29 (0.10)	0.08 (0.14)	0.22 (0.09)	NS	0.29 (0.05)	0	0.279	0.12	26
19-Apr	0.96 (0.03)	0.49 (0.03)	0.31 (0.07)	0.18 (0.06)	NS	0.51 (0.02)	0.34 (0.028)	0.0004	0.93	63
28-Apr	1.02 (0.02)	0.63 (0.11)	0.44 (0.12)	0.20 (0.05)	NS	0.61 (0.05)	0.47 (0.086)	0.0005	0.75	69
Average	1.02	0.35	0.25							50
Station 2										
31-Mar	0.49 (0.02)	0.34 (0.04)	0.25 (0.10)	0.10 (0.10)	NS	0.34 (0.04)	0.25 (0.05)	0.037	0.70	72
13-Apr	0.60 (0.01)	0.18 (0.04)	0.11 (0.06)	0.07 (0.05)	S					61
14-Apr	0.59 (0.03)	0.41 (0.05)	0.19 (0.07)	0.23 (0.04)	S					45
23-Apr	1.03 (0.03)	0.14 (0.04)	0.34 (0.06)	-0.20 (0.04)	NS	0.13 (0.02)	0.32 (0.04)	<0.0001	0.89	242
24-Apr	0.85 (0.02)	0.18 (0.03)	0.19 (0.03)	-0.02 (0.01)	S					110
Average	0.71	0.25	0.22							106
Station 3										
2-Apr	0.54 (0.04)	0.23 (0.03)	-0.03* (0.06)	0.26 (0.05)	NS	0.21 (0.03)	0	0.199	0.16	0
16-Apr	2.65 (0.05)	0.27 (0.05)	0.04 (0.09)	0.23 (0.08)	NS	0.28 (0.03)	0	0.202	0.16	15
Average	1.60	0.25	0.02							8

Table 3. Biomass estimates ($\mu\text{g C L}^{-1}$) of heterotrophic protists (HP, dinoflagellates and ciliates only) and of diatoms contained in undiluted samples collected at the beginning of each dilution experiment. Biomass of aloricate ciliates, tintinnids, and dinoflagellates are given as a percentage of total heterotrophic protist biomass.

Date	Total HP	% Aloricate ciliates	% Tintinnid ciliates	% Dino- flagellates	Diatoms
Station 1					
26-Mar	1.2	82.9	0.8	16.3	0.32
9-Apr	2.0	44.7	21.1	34.2	7.54
10-Apr	2.4	42.6	0.0	57.4	15.89
18-Apr	3.2	45.7	13.3	41.0	8.27
19-Apr	4.9	33.6	15.8	50.6	5.73
28-Apr	2.8	74.7	0.0	25.3	6.05
Station 2					
31-Mar	4.6	92.1	0.0	7.9	0.06
13-Apr	2.6	88.0	0.9	11.1	0.04
14-Apr	7.9	94.3	0.0	5.7	0.07
23-Apr	10.4	79.8	1.9	18.2	0.20
24-Apr	6.1	82.6	2.1	15.2	0.04
Station 3					
2-Apr	1.8	76.3	0.0	23.7	0.56
16-Apr	6.1	86.6	7.0	6.4	87.69

Table 4. Results of studies previously conducted at high latitudes of the subpolar North Atlantic during or after the spring bloom to quantify heterotrophic-protist herbivory, including grazing rates (g), proportion of primary production consumed (% PP), chlorophyll-a concentrations, and numerical abundance (10^3 cells L^{-1}) and biomass (μg C L^{-1}) of heterotrophic protist grazers (HP). Results from the present study are summarized for comparison.

Study	Month	Lat-Long	Chl-a (μg L^{-1})	g (d^{-1})	% PP	HP abundance	HP biomass
Gifford et al. 1995	05	59N-21W	0.59-2.89	0-1.01	56-64	3.3-6.9	n/a
Wolfe et al. 2000	05	56N-45W	1.7	0.12	150	n/a	n/a
Burkill et al. 1993	06	60N-20W	0.97	0.324	39	n/a	~3.9
Stelfox-Widdicombe et al. 2000	06	59N-20W	0.61-1.26	0.89-1.48	74	12.5-18.5	4.5-12.5
Gaul & Antia 2001	07	62N-11W	0.79-1.15	0.37-0.52	70-75	n/a	5.2-6.5
This study	03-04	61N-11W	0.17-1.87	0-0.56	0-94	0.4-2.0	1.2-4.9
This study	03-04	63N-02W	0.49-1.03	0.11-0.34	45-242	3.4-12.0	2.6-10.3
This study	03-04	60N-01E	0.54-2.65	0-0.04	0-15	1.4-2	1.8-6.1

FIGURE LEGENDS

Figure 1. Map showing locations of cruise sampling sites.

Figure 2. Principal Component Analysis (PCA) ordination of *in-situ* conditions at M87 stations 1 (▲), 2 (□), and 3 (●). PC1 accounts for 61.9 % of the variability (eigenvalue = 2.49). PC1 represents an axis of decreasing *in situ* Temperature (T) and mixed layer depth (MLD), each variable having a similar eigenvector value (0.618 for T and 0.590 for MLD). PC1 and PC2 together account for 89.6 % of the variation. Euphotic depth (Zeu) was the major contributor to PC2 (eigenvalue = -0.948 of a total of 1.11 for PC2) and varied more within than across stations. Samples belonging to the same station tended to segregate along the MLD and T gradient. Grouping was significant (ANOSIM global R = 0.796 p = 0.0002, 999 permutations).

Figure 3. Multidimensional scaling (MDS) analysis of species composition of (a) the diatom fraction of the autotrophic community and (b) heterotrophic protists assemblage at stations 1 (▲), station 2 (□) and station 3 (●). Overlaid contours represent gradual levels of similarities of 40, 50, and 60% between samples (CLUSTER analysis). Note that very few diatoms were present in samples at S2.

Figure 4. Size distribution of dominant heterotrophic protists in samples collected in the initial undiluted treatments of dilution experiments, at station 1 (a and c) and station 2 (b and d).

Figure 5. Phytoplankton growth rates vs. heterotrophic protist grazing rates at S1 (▲), S2 (□), and S3 (●). Dashed line represents 1:1 ratio. Error bars represent standard deviation.

Figure 6. Heterotrophic protist (HP) herbivory in relation to HP biomass at the Iceland basin (S1, ▲) and the Norwegian basin (S2, □). Ingested carbon was estimated using a carbon:chlorophyll-*a* ratio of 21. Reference lines represent 400, 200, 100, 50, and 25 % of body carbon ingested.

Figure 7. Daily net calculated ($\mu - g$) and net *in situ* phytoplankton accumulation rates, the latter estimated from changes in *in situ* chlorophyll-*a* concentration measured at the beginning of each experiment, for the two oceanic stations: (a) S1 ($R^2 = 0.10$, $p = 0.60$), (b) S2 ($R^2 = 0.98$, $p = 0.009$). The dashed line represents the 1:1 ratio. Apparent lack of error bars means that due to their small values, error bars are obscured by corresponding symbol.

Figure 8. Selected CTD profiles of temperature (left panel) and fluorescence-based chlorophyll-*a* (right panel) from the four visits to the Iceland Basin Station (S1).

Morison & Menden-Deuer Fig. 1

Morison & Menden-Deuer Fig.2

Morison & Menden-Deuer Fig. 4

Morison & Menden-Deuer Fig. 5

Morison & Menden-Deuer Fig. 6

Morison & Menden-Deuer Fig. 7

Morison & Menden-Deuer Fig. 8