

2012

Vibrio parahaemolyticus in Rhode Island Coastal Ponds and the Estuarine Environment of Narragansett Bay

Annie M. Cox
University of Rhode Island

Marta Gomez-Chiarri
University of Rhode Island, gomezchi@uri.edu

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

Terms of Use
All rights reserved under copyright.

Citation/Publisher Attribution

Cox, A. M., & Gomez-Chiarri, M. (2012). *Vibrio parahaemolyticus* in Rhode Island Coastal Ponds and the Estuarine Environment of Narragansett Bay. *Applied and Environmental Microbiology*, 78(8), 2996-2999. doi: 10.1128/AEM.07519-11
Available at: <http://dx.doi.org/10.1128/AEM.07519-11>

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Vibrio parahaemolyticus in Rhode Island Coastal Ponds and the Estuarine Environment of Narragansett Bay

Annie M. Cox^{a*} and Marta Gomez-Chiari^b

University of Rhode Island, Graduate School of Oceanography, Narragansett, Rhode Island, USA,^a and University of Rhode Island, Department of Fisheries, Animal, and Veterinary Sciences, Kingston, Rhode Island, USA^b

Quantification of the abundance of *Vibrio parahaemolyticus* in water and oysters from Rhode Island showed the presence of environmental strains and low levels of potentially pathogenic strains when water temperatures were $\geq 18^{\circ}\text{C}$, with peak levels in late July to early August. A higher abundance of the *trh* gene than of the *tdh* gene was observed.

Vibrio parahaemolyticus is a Gram-negative, halophilic bacterium occurring naturally in estuarine and coastal marine waters that is able to cause illness in humans (12, 14, 18, 19). Filter feeders, such as oysters, are capable of acquiring *V. parahaemolyticus* from the water column and concentrating it in their tissues (10). Because of the potential for human illness through consumption of raw or undercooked shellfish with *V. parahaemolyticus*, these bacteria pose a threat to shellfish aquaculture and fisheries through loss of revenue due to concerns about seafood safety and negative publicity in cases of outbreaks. Illnesses due to *V. parahaemolyticus* from consuming shellfish in Rhode Island and the Northeast United States are rare (3). Because of a strong association between *V. parahaemolyticus* levels and temperature (5, 7, 18, 19), there is an increased awareness of the danger posed by *V. parahaemolyticus* due to a rise in marine and estuarine water temperatures. In Narragansett Bay (Rhode Island), spring-summer sea surface temperatures have warmed by 1.6°C from 1959 to 2005

and are generally over 15.0°C from late May through mid-October (1). Warmer water temperatures could potentially extend the season in which *V. parahaemolyticus* could be a concern.

The total and relative abundances of environmental and potentially pathogenic *V. parahaemolyticus* bacteria in Rhode Island coastal waters have not been determined. A 1985 study showed the presence of *V. parahaemolyticus* in Narragansett Bay, but no pathogenic strains

Received 10 November 2011 Accepted 23 January 2012

Published ahead of print 3 February 2012

Address correspondence to Marta Gomez-Chiari, gomezchi@uri.edu.

* Present address: Southwest Fisheries Science Center, La Jolla, California, USA.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.07519-11




FIG 1 Partial map of Rhode Island showing the locations of the sampling sites in Narragansett Bay (NB1, NB2, and Fox Island) and coastal salt ponds (CP1 and CP2). (Map courtesy of Christopher Damon, adapted from the Rhode Island Geographic Information System [RIGIS; copyright 2011].)




FIG 2 Temporal variation in temperature (right axis, squares, broken line) and levels of *V. parahaemolyticus* *thl* (circles, solid line), *trh* (triangles, solid line), and *tdh* (diamonds, solid line) (left axis) in oysters collected from four Rhode Island farms during the summers of 2009 and 2010.

were detected (19). The purpose of this study was to determine the seasonal abundance of environmental and pathogenic *V. parahaemolyticus* in Rhode Island. Water (1 liter) and oyster ($n = 10$ to 15) samples were collected from two farms (NB1, 41°33.678'N, 71°18.431'W; NB2, 41°32.583'N, 71°25.351'W) in Narragansett Bay and two farms (CP1, 41°22.883'N, 71°31.954'W; CP2, 41°22.804'N, 71°33.720'W) in coastal salt ponds (Fig. 1). Water samples were also collected from a site near Fox Island (41°34.2'N, 71°23.4'W) within Narragansett Bay. A 3-tube most-probable-number (MPN) method combined with quantitative PCR detection of the *thl*, *tdh*, and *trh* genes was used for detection of *V. parahaemolyticus* (16, 20). After incubation of the MPN tubes, a 1-ml portion from each enrichment tube in the serial dilution series was transferred to cryotubes and stored at -80°C . Bacterial DNA was extracted by boiling, and quantitative PCR (qPCR) was used to score the tubes as positive or negative for *thl*, *trh*, and *tdh*. *Vibrio parahaemolyticus* densities (+1 to account for 0 values) were \log_{10} transformed for statistical analysis using one-way analysis of variance (ANOVA) or Kruskal-Wallis one-way ANOVA on ranks and the appropriate *post hoc* tests (IBM SPSS Statistics 19; IBM, Somers, NY; Sigmasat 3.1; Systat, Chicago, IL).

***Vibrio parahaemolyticus* densities in oysters and water.** *V. parahaemolyticus* was detected in Rhode Island oyster and water samples for approximately 15 weeks during the summers of 2009

and 2010 (Fig. 2 and 3). For most locations, densities of *V. parahaemolyticus* increased rapidly in a period of 2 to 3 weeks after water temperatures reached approximately 18°C , starting in early July for 2009 or mid-June for 2010. This study confirmed July to August as the period of highest risk for *V. parahaemolyticus*. This is similar to the case with the Chesapeake Bay, where *V. parahaemolyticus* first peaked in June and July in the water column after water temperature rose to 19°C (13, 18). The maximum level of total *V. parahaemolyticus* seen in oysters in this study (9×10^3 MPN g^{-1} oyster tissue) corresponds to a very low risk of gastroenteritis from oysters (10). Levels of total *V. parahaemolyticus* bacteria detected in Rhode Island oysters are comparable to levels observed in shellfish from other U.S. and European coastal areas ($<10^4$ MPN g^{-1}) (2, 5, 6, 11, 15, 16, 17, 18, 20).

Densities of *V. parahaemolyticus* in Narragansett Bay water samples in 2009 and 2010 (≤ 15 MPN ml^{-1}) are comparable to the maximum densities found in a 1985 study (4.95 CFU ml^{-1}) (19) (Fig. 3). Levels of *V. parahaemolyticus* in water collected from oyster farms were in general slightly higher than levels at the non-farm reference site at Fox Island. Maximum densities of *thl* in water were significantly higher in 2009 than in 2010 (Kruskal-Wallis one-way ANOVA; $P = 0.049$). The summer of 2009 was a colder summer with more precipitation, possibly explaining some of the differences between these 2 years.




FIG 3 Temporal variation in temperature (right axis, squares, broken line) and levels of *V. parahaemolyticus* *tlh* (circles, solid line), *trh* (triangles, solid line), and *tdh* (diamonds, solid line) (left axis) in water collected from four Rhode Island farms and Fox Island station during the summers of 2009 and 2010.

Abundance of pathogenic *V. parahaemolyticus* relative to total *V. parahaemolyticus*. For 21 of 48 (43.8%) oyster samples, only the *trh* gene was detected (*tdh* was nondetectable [Table 1]), showing a predominance of *trh*⁺ strains in Rhode Island. The percentage of pathogenic *V. parahaemolyticus* relative to total *V. parahaemolyticus*, calculated by dividing *trh* and *tdh* values by that for *tlh*, varied greatly both spatially and temporally for both years, and changes occurred in a matter of weeks (Table 1 and Fig. 2 and 3). Studies of other U.S. coastal waters (Gulf Coast and Pacific) typically show that most samples contain either a majority of *tdh*⁺ strains lacking *trh* or of *tdh*⁺ *trh*⁺ strains (6, 16, 20). Only one other study of U.S. waters (Chesapeake Bay) has shown a higher

prevalence of *trh*⁺ over *tdh*⁺ (18). However, studies from other countries, such as Norway, India, and France, have also shown a higher relative prevalence of *trh*⁺ (4, 8, 9). The relative percentage of pathogenic *V. parahaemolyticus* in oysters in Rhode Island (2.5 to 31.9% on average) (Table 1) is higher than what has been reported for other coastal locations in the United States (0.3 to 3.2%) (10). However, recent studies have shown that the percentage of pathogenic *V. parahaemolyticus* can be high (16).

Conclusions. This study provides a range of *V. parahaemolyticus* densities found in Rhode Island oysters and water samples. We detected a higher abundance of the *trh* gene than of the *tdh* gene. It is unknown if the levels of pathogenic *V. parahaemolyticus* mea-

TABLE 1 Percentages of pathogenic *V. parahaemolyticus* relative to total *V. parahaemolyticus* in water and oyster samples from Fox Island, Narragansett Bay, and two farms in the coastal ponds for 2009 and 2010 combined

Farm ^a	% <i>trh</i> or <i>tdh</i> prevalence [avg ± SD (range)] in:			
	Oysters		Water	
	<i>trh</i>	<i>tdh</i>	<i>trh</i>	<i>tdh</i>
NB1	2.5 ± 5.0 (0–17.8)	0.2 ± 0.3 (0–0.8)	6.2 ± 11.8 (0–31.8)	0 (0)
NB2	7.2 ± 11.2 (0–39.8)	0.4 ± 0.8 (0–2.4)	10.5 ± 16.7 (0–45.8)	0 (0)
CP1	31.9 ± 39.2 (0–100)	0.4 ± 1.3 (0–4.6)	15.7 ± 29.0 (0–100)	0 (0)
CP2	6.3 ± 4.7 (0.1–24.8)	3.5 ± 7.6 (0–24.8)	2.4 ± 3.0 (0–4.6)	1 ± 1.5 (0–3.8)
Fox Island	NT ^b	NT	7 ± 14.4 (0–38.9)	2.7 ± 8.1 (0–24.3)
Total	12.1 ± 23.4 (0–100)	1.1 ± 3.9 (0–24.8)	8.2 ± 17.1 (0–100)	0.7 ± 3.4 (0–24.3)

^a NB1 and NB2 are farms in Narragansett Bay; CP1 and CP2 are farms in coastal ponds.

^b NT, not tested.

sured in this study by qPCR were due to the presence of one or a few *trh*⁺ strains lacking *tdh* in relatively higher abundance or to the accumulation of low levels of multiple *tdh*⁺ *trh*⁺ strains and of *trh*⁺ strains lacking *tdh*. This study provides a baseline for further studies of the ecology of *V. parahaemolyticus* in coastal waters in a temperate estuary in the Northwest Atlantic.

ACKNOWLEDGMENTS

Funding for this project was provided by USDA AFRI Special Research Grant NIFA 2008-03240 to D. A. Bengtson, M. Gómez-Chiarri, and M. A. Rice. Equipment (Stratagene MX3005 and Nanodrop 8000) used for this study was available at the Rhode Island Genomics and Sequencing Center, supported in part by the National Science Foundation under an EPSCoR Grant (no. 0554548).

We thank the East Coast Shellfish Growers Association for their support and for providing oyster samples. We thank L. Maranda, B. D. Jenkins, P. S. Cohen, and C. A. Oviatt from the University of Rhode Island and S. Jones and C. A. Whistler from the University of New Hampshire for their helpful comments on the manuscript. Also, we thank Katherine Markey, Tyler Ferrara, and Paul Torbett for help with summer field and lab work and Christopher Damon from URI’s Environmental Data Center for the map in Fig. 1.

REFERENCES

- Collie JS, Wood AD, Jeffries HP. 2008. Long-term shifts in the species composition of a coastal fish community. *Can. J. Fish. Aquat. Sci.* 65:1352–1365.
- Cook DW, Bowers JC, DePaola A. 2002. Density of total and pathogenic (*tdh*⁺) *Vibrio parahaemolyticus* in Atlantic and Gulf Coast molluscan shellfish at harvest. *J. Food Prot.* 65:1873–1880.
- Daniels NA, et al. 2000. *Vibrio parahaemolyticus* infections in the United States, 1973–1998. *J. Infect. Dis.* 181:1661–1666.
- Deepanjali A, Kumar S, Karunasagar I, Karunasagar I. 2005. Seasonal variation in abundance of total and pathogenic *Vibrio parahaemolyticus* bacteria in oysters along the southwest coast of India. *Appl. Environ. Microbiol.* 71:3575–3580.
- DePaola A, Hopkins LH, Peeler JT, Wentz B, McPhearson RM. 1990. Incidence of *Vibrio parahaemolyticus* in U.S. coastal waters and oysters. *Appl. Environ. Microbiol.* 56:2299–2302.
- DePaola A, Kaysner CA, Bowers J, Cook DW. 2000. Environmental investigations of *Vibrio parahaemolyticus* in oysters after outbreaks in

- Washington, Texas, and New York (1997 and 1998). *Appl. Environ. Microbiol.* 66:4649–4654.
- DePaola A, Nordstrom JL, Bowers JC, Wells JG, Cook DW. 2003. Seasonal abundance of total and pathogenic *Vibrio parahaemolyticus* in Alabama oysters. *Appl. Environ. Microbiol.* 69:1521–1526.
- Deter J, et al. 2010. Ecology of pathogenic and non-pathogenic *Vibrio parahaemolyticus* on the French Atlantic coast. Effects of temperature, salinity, turbidity and chlorophyll *a*. *Environ. Microbiol.* 12:929–937.
- Ellingsen AB, Jorgensen H, Wagley S, Monshaugen M, Rorvik LM. 2008. Genetic diversity among Norwegian *Vibrio parahaemolyticus*. *J. Appl. Microbiol.* 105:2195–2202.
- FDA. 2005. Quantitative risk assessment on the public health impact of pathogenic *Vibrio parahaemolyticus* in raw oysters. FDA, Washington, DC.
- Hervio-Heath D, et al. 2002. Occurrence of pathogenic vibrios in coastal areas of France. *J. Appl. Microbiol.* 92:1123–1135.
- Jones SH, Summer-Brason B. 1998. Incidence and detection of pathogenic *Vibrio* sp. in a northern New England estuary, USA. *J. Shellfish Res.* 17:1665–1669.
- Kaneko T, Colwell RR. 1978. The annual cycle of *Vibrio parahaemolyticus* in Chesapeake Bay. *Microb. Ecol.* 4:135–155.
- Levin RE. 2006. *Vibrio parahaemolyticus*, a notably lethal human pathogen derived from seafood: a review of its pathogenicity, characteristics, subspecies characterization, and molecular methods of detection. *Food Biotechnol.* 20:93–128.
- Martinez-Urtaza J, et al. 2008. Environmental determinants of the occurrence and distribution of *Vibrio parahaemolyticus* in the Rias of Galicia, Spain. *Appl. Environ. Microbiol.* 74:265–274.
- Nordstrom JL, Vickery MCL, Blackstone GM, Murray SL, DePaola A. 2007. Development of a multiplex real-time PCR assay with an internal amplification control for the detection of total and pathogenic *V. parahaemolyticus* bacteria in oysters. *Appl. Environ. Microbiol.* 73:5840–5847.
- Oberbeckmann S, Wichels A, Wiltshire KH, Gerdt G. 2011. Occurrence of *Vibrio parahaemolyticus* and *Vibrio alginolyticus* in the German Bight over a seasonal cycle. *Antonie Van Leeuwenhoek* 100:291–307.
- Parveen S, et al. 2008. Seasonal distribution of total and pathogenic *Vibrio parahaemolyticus* in Chesapeake Bay oysters and waters. *Int. J. Food Microbiol.* 128:354–361.
- Watkins WD, Cabelli VJ. 1985. Effect of fecal pollution on *Vibrio parahaemolyticus* densities in an estuarine environment. *Appl. Environ. Microbiol.* 49:1307–1313.
- Zimmerman AM, et al. 2007. Variability of total and pathogenic *Vibrio parahaemolyticus* densities in northern Gulf of Mexico water and oysters. *Appl. Environ. Microbiol.* 73:7589–7596.