

2019

Uptake of hydrophobic organic compounds, including OCPs and PBDEs, and perfluoroalkyl acids (PFAAs) in fish and blue crabs of the lower Passaic River (NJ, USA)

Mohammed A. Khairy
University of Rhode Island

Gregory O. Noonan

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

**The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.**

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Khairy, M. A., Noonan, G. O. and Lohmann, R. (2019), UPTAKE OF HYDROPHOBIC ORGANIC COMPOUNDS, INCLUDING OCPs AND PBDES, AND PERFLUOROALKYL ACIDS (PFAAS) IN FISH AND BLUE CRABS OF THE LOWER PASSAIC RIVER (NJ, USA). *Environ Toxicol Chem*. Accepted Author Manuscript. doi:10.1002/etc.4354
Available at: <https://doi.org/10.1002/etc.4354>

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors

Mohammed A. Khairy, Gregory O. Noonan, and Rainer Lohmann

1 **Running head: Uptake sources of HOCs in biota?**

2
3 **Uptake of hydrophobic organic compounds, including OCPs and PBDEs, and**
4 **perfluoroalkyl acids (PFAAs) in fish and blue crabs of the lower Passaic River (NJ, USA)**

5 **Mohammed A. Khairy ^{†,‡}, Gregory O. Noonan [§], and Rainer Lohmann ^{‡*}**

6
7 [†]Graduate School of Oceanography, University of Rhode Island, Narragansett, Rhode Island 02882 USA

8 [‡]Department of Environmental Sciences, Faculty of Science, Alexandria University, 21511 Moharam Bek,
9 Alexandria, Egypt

10 [§] US Food and Drug Administration, Center for Food Safety and Applied Nutrition, College Park, MD 20740, USA

11 *Corresponding author. E-mail address: rlohmann@uri.edu

12 Phone: 401-874-6612; Fax 401-874-6811

13
14
15 **Keywords: HOCs, bioaccumulation, porewater, PFASs, sediment**

16

17

18

19 **Abstract**

20

21 The bioavailability and bioaccumulation of sedimentary hydrophobic organic compounds (HOCs)
22 is of concern at contaminated sites. Passive samplers have emerged as a promising tool to measure
23 the bioavailability of sedimentary HOCs and possibly to estimate their bioaccumulation. We thus
24 analyzed HOCs including organochlorine pesticides (OCPs), polybrominated diphenyl ethers
25 (PBDEs), polychlorinated biphenyls (PCBs) and polychlorinated dibenzo-p-dioxins/furans
26 (PCDD/Fs) in sediment, porewater and riverwater using low density polyethylene (LDPE) passive
27 samplers, and in 11 different finfish species and blue crab from the lower Passaic River.
28 Additionally, perfluorinated alkyl acids (PFAAs) were measured in grab water samples, sediment
29 and fish. Best predictors of bioaccumulation in biota were either porewater concentrations (for
30 PCBs and OCPs), or sediment organic carbon (PBDEs and PFAAs), including black carbon
31 (OCPs, PCBs and some PCDD/F congeners) normalized concentrations. Measured lipid-based
32 concentrations of the majority of HOCs exceeded the chemicals' activities in porewater by at least
33 2-fold, suggesting dietary uptake. Trophic magnification factors were > 1 for moderately
34 hydrophobic analytes ($\log K_{ow} = 6.5 - 8.2$) with low metabolic transformation rates ($< 0.01 \text{ day}^{-1}$),
35 including longer alkyl chain PFAAs. For analytes with lower ($4.5 - 6.5$) and higher (>8.2) K_{ows} ,
36 metabolic transformation was more important in reducing trophic magnification.

37

38

39 **INTRODUCTION**

40
41 Rivers in highly urbanized/industrialized areas are impacted by various classes of chemicals, often
42 resulting in adverse ecological effects, such as habitat destruction, and various physiological and
43 reproductive disorders in the river fauna. Once in the ecosystem, pollutants can accumulate in biota
44 from the surrounding water, sediments and/or porewater depending on the physicochemical
45 properties of the pollutants, feeding habits, trophic position and metabolic rates. The significance
46 of bioaccumulation in fish and the accompanied ecological and human health risks have been
47 previously highlighted in studies performed by the U.S. EPA (US EPA 1992a; US EPA 1992b).
48 Typically, organic chemicals are prioritized with respect to their persistence, bioaccumulation
49 potential and toxicity (PBT chemicals) (Gobas et al. 2009). Recently, the use of chemical
50 properties to predict the bioaccumulation potential of a chemical has been questioned (Borgå et al.
51 2012).

52
53 In our recent work (Khairy et al. 2014), we assessed the biomagnification of polychlorinated
54 biphenyls (PCBs), and polychlorinated dibenzo-*p*-dioxin/dibenzofurans (PCDD/Fs) in fish and
55 crabs collected from the Passaic River estuary. In general, the dominant exposure pathways for
56 legacy hydrophobic organic contaminants (HOCs) in biota were porewater and sediments rather
57 than the riverwater. Here, we aimed to expand on these results by investigating a wider range of
58 contaminants with more diverse structures and properties. Targeted compounds include
59 organochlorine pesticides (OCPs), polybrominated diphenyl ethers (PBDEs) and perfluorinated
60 alkyl acids (PFAAs), including perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA).
61 As PFAAs and some OCPs display much greater aqueous solubility than the legacy HOCs, it is
62 unclear whether sediment-bound contamination remains a primary exposure pathway to biota in
63 the river. These compounds also cover a wide range of resistance to metabolism (US EPA 2016a),

64 which should, in conjunction with varying degrees of lipophilicity, affect their uptake and
65 magnification in the food web.

66
67 While recent work established the presence of PBDEs at relatively high concentrations in all the
68 environmental compartments of the lower Passaic River, and deduced diffusive fluxes of PBDEs
69 from sediment to overlying water (Khairy and Lohmann 2017), little is known regarding their
70 uptake sources and their bioaccumulation potential.

71
72 Unlike legacy HOCs, PFAAs are ionic, amphiphilic compounds (Key et al. 1997), and thus have
73 different physicochemical properties. They have been globally detected in all environmental
74 compartments (Labadie and Chevreuil 2011; Loi et al. 2011; Fang et al. 2014). PFAAs are
75 persistent and tend to bioaccumulate (PFAAs with longer fluorinated carbon chain) (Conder et al.
76 2008), undergo long range transport (LRT) and cause adverse ecological and health effects (Lau
77 et al. 2007). Anionic PFAAs are proteinophilic and generally found at highest concentrations in
78 blood and liver (Becker et al. 2010) in contrast to the lipophilic legacy pollutants.

79
80 Studying the bioaccumulation potential of HOCs requires the ability to measure the freely
81 dissolved fraction in sediments (porewater) and/or the water column, where bioavailability is
82 expected to increase with an increase in the dissolved concentrations (You et al. 2006). Passive
83 samplers, such as low density polyethylene (LDPE) sheets, have emerged as an inexpensive and
84 effective technique for measuring the freely dissolved concentrations of HOCs. Traditionally,
85 geochemical approaches have been used to predict porewater concentrations (sediment
86 concentrations and organic carbon content), due to the difficulties associated with the measurement
87 of the dissolved fraction in sediment's porewater directly. However, since the development of

88 passive sampling devices, estimation of porewater concentrations have become feasible (
89 Fernandez et al. 2009; Ghosh et al. 2014; Khairy and Lohmann 2017).

90
91 The original equilibrium partitioning theory (Di Toro et al. 1991) linked porewater concentrations
92 of HOCs to their accumulation in the lipid of organisms. As LDPE samplers similarly accumulate
93 HOCs from porewater, recent studies indicated that LDPE concentrations of HOCs (at
94 equilibrium) can be used to predict their bioaccumulation (Friedman et al. 2009). In fact, recent
95 work by Jahnke et al. (2014) suggested that passive sampling can be used to estimate the
96 bioaccumulation potential of HOCs, and indicated that the majority of studies resulted in tissue
97 concentrations below the equilibrium concentration with porewater (Jahnke et al. 2014). Being
98 able to rely on passive samplers to predict (maximum) bioaccumulation in aqueous foodwebs
99 would certainly be very useful for the assessment of contaminated sites.

100
101 In the current study, we investigated the bioaccumulation of OCPs and PBDEs in 11 fish species
102 and *Callinectes sapidus* (blue crab), collected from the fresh-brackish portion of the Passaic River
103 estuary. PFAAs data for only 8 fish species were included in the current study. To that end, we
104 calculated the bioaccumulation factors (BAFs) for OCPs, PBDEs and PFAAs, compared the
105 accumulated amounts of HOCs in LDPE and in the biota and investigated the best predictors for
106 HOCs and PFAAs in biota using sediment geochemistry and LDPE (active water samples for
107 PFAAs). Since sampling of biotic and abiotic compartments were done at the same time intervals
108 and all the samples were analyzed in the same lab, we present here a uniquely consistent set of
109 BAFs. Our objectives were to (i) assess the sources of contaminants accumulated by the biota
110 (sediments, porewater, diet, or overlying water), (ii) assess the suitability of using LDPE to predict

111 the bioaccumulation of HOCs in biota and (iii) derive bioaccumulation factors and assess the
112 trophic magnification (or dilution) of target analytes.

113

114 **MATERIALS AND METHODS**

115 *Samples collection, extraction, analysis for OCPs and PBDEs*

116 Detailed description of the sampling of biota and sediments, preparation and deployment of the
117 LDPE in the river water (for OCPs and PBDEs), the stable isotope analysis in the biota tissues, the
118 determination of porewater concentrations using LDPE tumbling experiment and the uncertainty
119 analysis can be found in in Figures S1-S3, Table S1, text in the Supplementary information (SI)
120 and in Khairy et al. (2014, 2015) and Khairy and Lohmann (2017). More information about the
121 extraction of all matrices, instrumental analysis, quality assurance and the determination of
122 porewater concentrations is available in the SI (Tables S2-S6) and is briefly summarized below.

123 Finfish specimens (n = 350 representing 10 species) including benthic feeders (*Callinectes*
124 *sapidus*, *Anguilla rostrata*, *Fundulus heteroclitus*, *Hybognathus regius*, *Morone americana*),
125 benthic-pelagic taxa (*Fundulus diaphanus*, *Lepomis spp.*, *Esox americanus*, and juvenile *Morone*
126 *saxatilis*), and pelagic species (*Menidia menidia* and *Dorosoma cepedianum*) and 7 *Callinectes*
127 *sapidus* (blue crab) specimens were collected during August-November, 2011 at three locations in
128 the Passaic River estuary (Figure S1). For marine transients, fish collections were restricted to the
129 extent practical to young-of-year that had a recent history of exposure to contaminants in the
130 system. See Table (S1) for more details on the biota samples.

131 Sediment samples were collected from 18 different locations (Figure S2) along the lower Passaic
132 River during September to November, 2011, from the mudflats at low tide. Detailed description

133 of the sampling methodology and sampling locations can be found in Khairy et al. (2015). Total
134 organic carbon (TOC) and black carbon (BC) content in the sediments were determined as detailed
135 in Gustafsson et al. (1997).

136 LDPE passive samplers were fastened to an anchored rope and suspended in the water column ~
137 1-2 m below the surface at six different locations along the lower Passaic River (Figure S1) to
138 sample the freely dissolved OCPs and PBDEs.

139 Freely dissolved porewater concentrations were determined by shaking (equilibrating) LDPE
140 passive sampling sheets (25 μm thick) with sediment-water (containing sodium azide as a biocide)
141 slurry for 9 weeks according to the method detailed in Lohmann et al. (2005).

142 Biota and sediment samples were extracted and subjected to cleanup according to the methods
143 detailed elsewhere (Khairy et al. 2014, 2015). LDPE (both deployed in the river water and
144 porewater) were cold extracted twice (24 hours each) in dichloromethane followed by hexane,
145 concentrated and were not subjected to any further cleanup steps.

146 *Collection of water samples, extraction and analysis of all the samples for PFAAs*

147 We used the same sediment and biota samples that were collected for OCPs and PBDEs. Water
148 samples (~ 20 cm below the surface) were collected in pre-cleaned (inner walls were cleaned with
149 basic methanol followed by methanol and left overnight to dry) one-liter polypropylene bottles
150 with polypropylene lids at 8 different locations along the lower segment of the Passaic River
151 (Figure S3). Samples were kept in an ice bath until preserved in a freezer for analysis. Water
152 samples (400 mL each) were analyzed for PFAAs (PFHxA - PFDoDA, PFBS, PFHxS and PFOS)
153 in the U.S. Food and Drug Administration's (FDA) Laboratory in Maryland (USA) according to
154 the method detailed in Young et al. (2013). Sediments and fish samples were analyzed at

155 University of Rhode Island (URI), USA. Extraction and cleanup was done according to the U.S.
156 EPA method for the analysis of PFAAs (EPA 2011) (see SI, Tables S2 and S3 for more
157 information). PFAAs were measured in 8 fish species [*Morone saxatilis* (29 – 33 cm), *Anguilla*
158 *rostrate* (110 cm), *Morone Americana* (8 -16 cm), *Fundulus diaphanous*, *Hybognathus regius*,
159 *Menidia menidia*, *Dorosoma cepedianum*, and *Fundulus heteroclitus*], but not in the blue crabs
160 due to a lack of extra tissue after PBDE and OCP analysis.

161 *Quality assurance*

162 Procedural blanks, field blanks (LDPE), matrix spikes, and duplicate samples (20% of the total
163 samples) were included with each sample batch and were carried throughout the entire analytical
164 procedure in a manner identical to the samples. BDE-47 was detected in the blanks (10-20 % of
165 the least detected concentration in the different matrices). Similarly, detected PFAAs in the blank
166 samples were 12-25 % of the sample concentrations and accordingly, correction of samples for the
167 blank detects was performed.

168 Recoveries of the surrogate standards in the biota, LDPE and sediment samples generally ranged
169 from 65 - 102% and 72-104 % for OCPs and PBDEs respectively. Matrix spikes recoveries (Table
170 S4) ranged from 94 – 104 % for OCPs and PBDEs with RSD % < 10 %. Reproducibility of the
171 results ranged from 7.0 – 17 % and 5.0 – 18.5 % for OCPs and PBDEs respectively.

172 Recoveries of PFAAs in the matrix spikes (Table S5) were 87-133% for the majority of the target
173 PFAAs and < 55 % for C₁₃ - C₁₈- perfluorocarboxylic acid and C₁₀-perfluorosulfonate.
174 Accordingly, these PFAAs were excluded from the discussion. Recoveries of the surrogate
175 standards ranged from 57 – 120 % (Table S5). Limit of detection (LOD) corresponded to a signal-
176 to-noise ratio of 3 (Table S6). Variability between duplicates was <20%.

177 *Physicochemical properties*

178 Octanol-water partitioning coefficient (K_{ow}) values of OCPs were obtained from Schenker et al.
179 (2005). K_{ow} values for PBDEs were obtained from Yue and Li (2013). Polyethylene-water
180 partitioning coefficients (K_{PE-w}) for OCPs and PBDEs were obtained from Lohmann (2012).
181 Lipid-water partitioning coefficients (K_{lip-w}) were estimated from K_{ow} according to the method of
182 Endo et al. (2011). Organic carbon-water partitioning coefficient (K_{OC}) were calculated as shown
183 in Xia (1998) .

184 For PFAAs, K_{OC} were obtained from obtained from Higgins and Luthy (2007). Missing values
185 were obtained by correlating the available values against those obtained from Labadie and
186 Chevreuil (2011). Protein-water partitioning coefficients (K_{PW}) were obtained from Kelly et al.
187 (2009). Missing values were obtained from correlating the available values against K_{OCs} ($n = 8$;
188 $R^2 = 0.98$); K_{ow} values were obtained from Wang et al. (2011). K_{PW} rather than K_{lip-w} were used
189 with PFAAs as these compounds bind to proteins rather than to general lipids. All the
190 physicochemical properties used in the current study are given in Table S7.

191 Estimated compound specific metabolic biotransformation rates (K_{mtrS}) normalized to 10 g tissue
192 weight were obtained from the BCFBAF function in EPI Suite 4.11 (US EPA 2016b).

193 *Calculation of BAFs, TMFs and predicting biota concentrations*

194 Bioaccumulation factors (BAFs, L/kg lipid or ww) were calculated as the ratio of lipid normalized
195 tissue concentrations of PBDEs and OCPs (C_{lip} , ng/kg lipid) to the freely dissolved river water
196 concentrations from PE-samplers (C_w , ng/L), or fresh weight concentrations of PFAAs (; ng/kg
197 ww) to grab water sample concentrations (C_w , ng/L) as follows (equation 1) (Schwarzenbach et al.
198 2005):

199
$$\text{BAF} = \frac{C_{\text{lip}(ww)}}{C_w} \quad (1)$$

200 TMF values were calculated as follows (equation 2):

201
$$\text{TMF} = e^b \quad (2)$$

202 where b is the slope of the linear relationship between the natural log of lipid-normalized tissue
 203 concentrations and the trophic position of each species. TMFs >1 indicates that contaminants are
 204 biomagnifying whereas values < 1 indicate that contaminants are not taken up by the organism or
 205 that they are metabolized (trophic dilution).

206 Lipid-based concentrations (ng/g) of HOCs were estimated from each sediment's sorbent phase
 207 [organic carbon (OC) and black carbon (BC)], porewater LDPE, and riverwater LDPE and
 208 compared to those measured in tissues (C_{lip}) as follows: (equation 3-5):

209
$$C_{\text{lip(OC)}} = \frac{C_{\text{sed}}}{f_{\text{OC}} K_{\text{OC}}} K_{\text{lip-w}} \quad (3)$$

210 Where, C_{sed} is the sediment concentration (ng/g) and f_{OC} is the fraction of OC in sediments.
 211 Similarly, lipid concentrations based on OC + BC ($C_{\text{lip,OC+BC}}$) were estimated using a Freundlich
 212 coefficient of $n = 0.7$:

213
$$C_{\text{lip(OC+BC)}} = \frac{C_{\text{sed}}}{f_{\text{OC}} K_{\text{OC}} + f_{\text{BC}} K_{\text{BC}} C_{\text{PW}}^{n-1}} K_{\text{lip-w}} \quad (4)$$

214 where f_{BC} is the fraction of BC in the sediment and K_{BC} is the black carbon-water partitioning
 215 coefficient. We used site specific K_{BC} values in the current study.

216 Lipid concentrations from PE-derived porewater [$C_{lip(PW)}$] and riverwater [$C_{lip,(W)}$] were estimated
217 as follows:

$$218 \quad C_{lip[PW(W)]} = C_{[PW(W)]} * K_{lip-w} \quad (5)$$

219 For PFAAs, protein-based concentrations were predicted only from sediment's OC and from river
220 water using equations 3 and 5. In both equations, K_{lip-w} was replaced with K_{PW} . Concentrations of
221 PFAAs in the samples were normalized to a generic total protein content (25 %) assigned for fish
222 tissues as shown in Kelly et al. (2009) and references therein.

223 *Statistical Analysis*

224 Statistical tests (ANOVA, t-test and correlations) were performed with SigmaPlot (version 11,
225 Germany); regression analysis with IBM SPSS V25.

226

227 **RESULTS**

228 *Uncertainty analysis*

229 Uncertainties associated with the estimated porewater and river water concentrations (equation S4)
230 from LDPE ranged from 41 – 65 % and 40 – 64 % respectively. Uncertainties associated with
231 predicted tissue concentrations from sediment's OC were the lowest (22 – 58 %) followed by
232 predicted concentrations from porewater (41 – 65 %), river water (40 – 80 %) and sediment's OC
233 + BC (28 - 88 %).

234 *Concentration of organic pollutants*

235 *OCPs*: OCP concentrations in the biota samples are given in Figure 1A-C and Table S8. In all
236 species, the following descending order was observed: DDTs (900 – 21,000 ng/g lipid) >
237 chlordanes (840 – 4,700 μ g/g lipid) > hexachlorobenzene (HCB) (11 – 330 ng/g lipid) > α -

238 endosulfan (<LOD – 290 ng/g lipid) > HCHs (8.0 – 83 ng/g lipid). Concentrations of HCB, α -
239 endosulfan and DDTs were 2.0 – 10 times higher in the blue crab samples (Figure 1A) than the
240 other biota as was previously observed for PAHs, PCBs and PCDD/Fs (Khairy et al. 2014).
241 Average concentrations of HCB in the benthopelagic (including the top predator) and pelagic
242 species (164 and 200 ng/g lipid respectively) were 3.0 fold higher than in the benthic species (70
243 ng/g lipid). Similarly, average concentration of DDTs in the benthopelagic species (9,000 ng/g
244 lipid) was 3.0-4.0 fold higher than in benthic (3,000 ng/g lipid) and pelagic species (2,000 ng/g
245 lipid; Figure 1A).

246
247 DDTs (24-52 ng/g dw) were the dominant pesticide in the sediment samples (Table S9), followed
248 by chlordanes (12-35 ng/g dw) and α -endosulfan (1.0 – 11 ng/g dw). In the porewater (Table S10)
249 and surface water samples (Table S11) HCHs (2.6 – 9.0 and 1.4 – 3.1 ng/L respectively) dominated
250 followed by α -endosulfan (0.20 – 5.7 and <LOD – 0.90 ng/L) and chlordanes (0.30 – 1.1 and 0.16
251 – 0.48 ng/L).

252
253 **PBDEs:** Σ_{12} PBDEs concentrations ranged from 120 ng/g lipid (*Fundulus heteroclitus*) to 2,500
254 ng/g lipid (*Callinectes sapidus*) in the biota (Table S12), 1.0-11 ng/g dw in sediments (Table S13),
255 19 – 40 pg/L in porewater (Table S14) and 12 -31 pg/L in surface water (Table S15). All the biotic,
256 sediment, porewater and surface water samples were dominated by BDE-47 and BDE-99 (Figure
257 1D-F). Other congeners that were detected included BDE-183 in sediments, and BDE-100 and
258 BDE-49 in the biota, sediments and porewater (estimated by LDPE).

259 **PFAAs:** Concentrations of PFAAs ranged from 130 ng/g ww (*Fundulus heteroclitus*) to 350 ng/g
260 ww (*Anguilla rostrata*) (Table S16). All the samples were dominated by PFOS (52 – 76 % of the
261 total PFAA, Figure 1G) followed by PFDoA, PFUnDA, PFOA (4.0 – 10 % each) and PFHxA.

262 PFBA, PFPeA and PFHpS were below LOD in all the samples. In all samples, concentrations of
263 perfluorinated sulfonic acids (PFSAs) were significantly (t- test, $p < 0.05$) higher than
264 perfluorinated carboxylic acids (PFCAs).

265 PFAAs in the sediment (Figure 1H) samples ranged from 8.0 ng/g dw (km 3.7) to 15 ng/g dw (km
266 24) (Table S17). PFOS was the dominant compound (42 % of total concentrations). The longer
267 chain PFCAs (PFDoA: 35 %; PFUnDA:11 %), PFOA (5.0 %) and PFDA (4.0 %) were easily
268 detected in sediment, probably related to their greater affinity for particles and hence accumulation
269 in sediments (sorption to OC) due to increasing hydrophobicity with increasing number of CF_2
270 groups (Ahrens et al. 2010). PFBA, PFPeA and PFBS were below LOD in all the samples, whereas
271 PFHpA and PFHpS were below LOD in the majority of the samples.

272 Concentrations of PFAAs in the surface water ranged from 44 ng/L (km 3.7) to 120 ng/L (km 22.4)
273 (Table S17). Similar to results for sediments and biota, PFOS dominated in surface water samples,
274 followed by PFOA and PFHxA (Figure 1I).

275 *Tissue concentrations versus abiotic compartments*

276 LDPE concentrations of HOCs [in riverwater ($C_{PE(RW)}$) and porewater ($C_{PE(PW)}$)] at equilibrium
277 (ng/kg PE), OC-normalized sediment concentrations ($C_{sed(OC)}$; ng/kg OC), BC-normalized
278 sediment concentrations ($C_{sed(BC)}$; ng/kg BC) and OC+BC-normalized sediment concentrations
279 ($C_{sed(OC+BC)}$) were compared with lipid-normalized concentrations (C_{lip}) of the target analytes in
280 the food web of the lower Passaic River (Figure S4). For PFAAs, the comparison was only made
281 with $C_{sed(OC)}$. Significant linear relationships were observed between log measured C_{lip} and log
282 $C_{PE(RW)}$ (Figure S4A), log $C_{PE(PW)}$ (Figure S4B) and log sediment (Figure S4C-E) concentrations

283 (including PFAAs). Coefficient of determinations (R^2) ranged from 0.79 to 0.82 and slopes of the
284 best fits were insignificantly different from 1 (slopes = 1.2-1.3, $p < 0.001$).

285 C_{lip} for all HOCs were predicted from equations 3-5 using the OC ($C_{lip(OC)}$) and the OC + BC
286 models ($C_{lip(OC+BC)}$), porewater ($C_{lip(PW)}$) and riverwater ($C_{lip(RW)}$) LDPE (Figure 2) and compared
287 to measured concentrations. Similarly, protein-based tissue concentrations of PFAAs were
288 predicted from $C_{lip(OC)}$ and $C_{lip(RW)}$ based on measured riverwater concentrations. To facilitate the
289 comparison between HOCs and PFAAs, measured (in biota) and predicted protein-based
290 concentrations of PFAAs from sediment and riverwater were normalized to the lipid content. This
291 step was done by converting the predicted protein-based concentrations (from abiotic
292 compartments) to equivalent wet weight concentrations and then normalizing to the measured lipid
293 content in tissues. In addition to the regression lines, a factor difference of ± 10 (represented by
294 the dashed red lines) was used to evaluate the predictive ability of the abiotic compartments. This
295 factor was previously used to evaluate the ability of using passive samplers as a surrogate for
296 bioaccumulation of HOCs (Joyce et al. 2016). When points occur within the dashed lines, it
297 indicated similarities in the affinities of geochemical natural sorbents (OC and BC), lipid and
298 LDPE to the organic pollutants. Mono-through tri-chlorinated dioxins and furans and HCHs were
299 excluded from the comparison because of their high biotransformation potential and/or
300 uncertainties in the physicochemical properties.

301 Highly significant linear relationships (all regression parameters are shown in Figure 2) were
302 observed between log measured C_{lip} and predicted log $C_{lip(OC)}$ (Figure 2A), $C_{lip(OC+BC)}$ (Figure 2B)
303 and $C_{lip(PW)}$ for HOCs only (Figure 2C) and $C_{lip(RW)}$ (Figure 2D) for all species. In general,
304 PCDD/Fs were less well predicted than other classes of pollutants.

305 Jahnke et al. (Jahnke et al. 2014) used passive samplers to derive the chemical activity of several
306 PCBs and HCB in sediment and biota and to investigate the possibility of using passive samplers
307 as a metric for the thermodynamical potential for the bioaccumulation of HOCs. According to
308 their results and other cited studies, measured C_{lip} /predicted C_{lip} from porewater were either below
309 or near their equilibrium benchmark values (0.5 – 2.0). In the current study, 2,3,7,8-TCDD,
310 2,3,7,8-TCDF, PCB 52, 101, 105, 118, 128, 138, 153, 180 and 187, BDE-28, 49, 47, 100, 154,
311 HCB, chlordanes and DDTs in the majority of the fish and blue crab samples showed values greater
312 than the equilibrium benchmark based on porewater (> 2).

313 *Bioaccumulation and biomagnification of HOCs and PFAAs*

314 Calculated BAFs for OCPs, PBDEs (both in L/kg lipids) and PFAAs (in L/kg PW) are given in
315 Tables (S18 – S20). Average BAFs ranged from 4.1×10^3 (α -HCH) – 7.7×10^7 (o,p'-DDE), $7.1 \times$
316 10^5 (BDE-2) – 5.3×10^8 (BDE-154) and 4.8×10^2 (PFBS) – 1.1×10^5 (PFDoA) for OCPs, PBDEs
317 and PFAAs respectively. Calculated BAFs for PBDEs were the highest (Table S18) followed by
318 OCPs (Table S19) and PFAAs (Table S20). Additionally, PBDE BAFs were higher than those
319 previously calculated for PAHs in the same samples (Khairy et al. 2014) but lower than BAFs
320 previously calculated for PCDD/Fs (4.0×10^3 – 6.0×10^8) (Khairy et al. 2014) and PCBs ($2.0 \times$
321 10^3 – 2.0×10^9) (Khairy et al. 2014). Reported BAF values for OCPs and PBDEs in the current
322 study were higher than values previously reported for freshwater fish from China (2.0×10^3 – 8.0
323 $\times 10^5$ and 2.0×10^2 – 3.2×10^6 for OCPs and PBDEs respectively) (Wu et al. 2008; Wang et al.
324 2012) and for Lake Trout in Lake Michigan (Streets et al. 2006). Calculated BAFs for PFAAs
325 were within the range of values calculated for PFAAs in literature (1.2×10^1 - 1.0×10^5) (Kwadijk
326 et al. 2010; Labadie and Chevreuil 2011).

327

328 Trophic levels of the fish species were calculated here based on the levels of nitrogen isotopes in
329 the tissues (Khairy et al. 2014). Blue crabs were excluded because of their relatively high detected
330 concentrations of OCPs and PBDEs despite of their low trophic position, which is probably due to
331 accidental sediment ingestion. Calculated TMFs were significantly greater than 1 ($p < 0.05$) for
332 o,p'-DDD (4.1), HCB (3.9), o,p'-DDE (3.6), p,p'-DDE (3.1), p,p'-DDT (2.1), heptachlor (2.0),
333 oxychlordan (1.8), BDE-47 (3.1), BDE-154 (2.5), Σ_{12} PBDEs (2.4), and BDE-100 (2.1) (Table
334 S21).

335

336 **DISCUSSION**

337 *Comparing tissue concentrations to sediment, riverwater and porewater concentrations*

338 In this study, significant linear relationships were observed between $\log C_{PE(PW)}$, $C_{PE(RW)}$, $C_{sed(OC)}$
339 and $C_{sed(OC+BC)}$ and $\log C_{lip}$ in biota. Similar relationships were previously observed for PCBs
340 (Friedman et al. 2009; Joyce et al. 2015; Joyce et al. 2016) and PBDEs (Joyce et al. 2015) when
341 tissue concentrations were compared to LDPE concentrations adjusted for equilibrium.
342 Coefficients of determination (R^2) reported in the current study (for all classes of HOCs + PFAAs
343 in sediments) were within the range previously observed for PCBs (0.59-0.94) (Friedman et al.
344 2009; Joyce et al. 2015; Joyce et al. 2016) and DDTs (0.72) but higher than for PBDEs (0.59)
345 (Joyce et al. 2015).

346 In contrast, slopes of the regression lines in the current study (1.2 – 1.3, $p < 0.001$) were higher
347 than those observed in the previous studies for PCBs, PBDEs and DDTs (0.74 – 1.0) (Friedman et
348 al. 2009; Joyce et al. 2015; Joyce et al. 2016). The slopes we observed indicated that accumulated
349 amounts of HOCs in tissues exceeded those predicted from LDPE or sediment geochemistry. This
350 implies that biotransformation of most HOCs did not play a significant role in the fish samples

351 collected from the lower Passaic River. These results suggest that the linear relationships with
352 abiotic compartments could be used to predict the bioaccumulation potential in the fish. Best
353 predictions of C_{lip} for HOCs and PFAAs were obtained from either $C_{sed(OC+BC)}$ and/or $C_{PE(PW)}$
354 (Figure 2), as demonstrated by higher R^2 , lower SE and slopes closer to 1.

355 All the OCPs, PFAAs and the majority of PBDE, PCDD/F and PCB congeners occurred within a
356 factor of 10 of measurements, although lipid-based concentrations were overestimated for many
357 congeners. Importantly, PFAAs were not separated from the other HOCs and the regression
358 parameters with and without PFAAs did not differ (Figure 2). This possibly implies a similar
359 affinity of HOCs and PFAAs to OC. Moreover, tissue concentrations of PFOA through PFDoA,
360 PFOS and PBDEs were well predicted from $C_{sed(OC)}$ within a factor difference ranging from 1.0 to
361 2.8 (Figure 2A).

362 Lipid-normalized concentrations of OCPs and PCBs were similarly better predicted using the
363 $C_{PE(PW)}$ and $C_{sed(OC+BC)}$ models (Figure 2B, C) within a factor difference of 1.0 – 4.0. Higher factor
364 differences (measured /predicted > 2.0) was observed for some PCB congeners (PCB 28, 52, 101,
365 153, 138, 180, which are known to be strong bioaccumulative) and OCPs such as chlordanes and
366 DDTs, presumably due to their uptake from food and their biomagnification potential (see below).
367 As for OCs, regression parameters for the measured versus predicted tissue concentrations from
368 riverwater were similar with and without adding PFAAs (Figure 2D), which again indicates a
369 similar lipid-water partitioning behavior for HOCs and the more polar PFAAs. However, many
370 PFAAs were underestimated by riverwater, with factor differences (measured/predicted) > 10.
371 Only the more soluble PFHxA and PFHpA were better predicted from riverwater (factor
372 difference: 1.5 – 2.5) than sedimentary OC, $C_{sed(OC)}$. Unfortunately, concentrations of PFAAs were
373 not measured in porewater and the predictive ability of $C_{PE(PW)}$ and $C_{sed(OC+BC)}$ cannot be examined.

374 However, based on regression output of $C_{\text{sed(OC)}}$ and $C_{\text{PE(RW)}}$, we assume that the uptake of PFAAs
375 probably occurred from porewater and diet similar to HOCs.

376 When comparing measured to predicted concentrations of the legacy pollutants, we did not account
377 for the biotransformation rates in the biota. We assumed biotransformation was negligible for
378 several reasons: First, significant linear relationships were observed between calculated log
379 bioaccumulation factors (BAFs) and log K_{OWs} (Figure 3) indicating either equilibrium or steady
380 state for the HOCs. Second, as shown in Figure 3, log BAFs for all the analytes (except BDE-99)
381 were above the 1:1 line. In other words, calculated log BAFs were higher than their corresponding
382 log K_{OWs} , which is an indicative of low/non-biotransformation except for BDE-99. Third, lipid-
383 based concentrations of HOCs were higher than sediment concentrations (normalized to OC, BC
384 and OC + BC) and LDPE concentrations (ng/kg PE) (Figure S4). Finally, Wirgin et al. (2011)
385 indicated that fish in the Hudson River, which is located in the same region of the Passaic River,
386 have developed a mechanism to resist biotransformation due to evolutionary changes from the
387 exposure to legacy pollutants.

388 *Comparison of porewater and foodweb chemical activities*

389 In contrast to other studies [25 and references therein], measured C_{lip} of the majority of the HOCs
390 in the current study were higher than the porewater's equilibrium benchmark set by Jahnke et al
391 (2014). The benchmark value was derived as a factor of 2 of the chemicals' activities in porewater
392 as estimated by passive samplers in sediment and biota directly. Possible reasons for these
393 divergent results in our study include (i) the structure of the foodwebs, (ii) the relative importance
394 of pelagic vs trophic coupling, and (iii) relying on a different approach of comparing chemical
395 activities in biota relative to abiotic compartments.

396 We rule out metabolism having a significant influence on these persistent chemicals as we
397 mentioned in the previous section. Similarly, trophic levels in the Swedish lake's food-chain were
398 even greater (assumed to reach a TL of up to 4.4), than in the Passaic River, where the top predator
399 was at a TL of 4.0 (Khairy et al. 2014).

400 We argue that the exceedance of porewater chemical benchmark observed in biota of the Passaic
401 River is due to the strong benthic coupling, as we described previously (Khairy et al. 2014; Khairy
402 and Lohmann 2017). In other words, the fish in the Swedish lake were likely strongly affected by
403 water column concentrations (not measured), which we assume were much below porewater
404 concentrations. In the Passaic River, differences between porewater and water column were only
405 0.6 - 11. Lastly, we inferred chemical activities in biota by lipid-normalizing HOC concentrations,
406 while we used passive samplers in porewater and water column. This introduces more uncertainty,
407 but the good agreement observed by numerous studies between lipid-normalized and passive
408 sampler-derived HOC concentrations implies this is of minor importance (Friedman et al. 2009;
409 Friedman and Lohmann 2014; Joyce et al. 2016). Overall, our results suggest that predicting the
410 ecological risk to aquatic biota using measurements of porewater concentrations is not sufficient
411 unless adjusted by the calculated BAFs in benthically-coupled system.

412 *Bioaccumulation of the organic pollutants*

413 Significant log BAFs – log K_{OW} ($R^2 = 0.80 - 0.91$; $p < 0.001$) and log BAFs – log K_{lip-w} ($R^2 = 0.87$
414 $- 0.94$; $p < 0.001$) linear relationships were observed for PBDEs and OCPs in the biota using both
415 riverwater (BAF^{RW} ; Figures 3A, S5A) and porewater (BAF^{PW} ; Figures 3B, S5B) for calculating
416 BAFs, indicating that the observed partitioning between lipids (in biota) and water can be
417 approximated from the octanol – water equilibrium partitioning. In our previous work, we
418 indicated and quantified diffusive fluxes of PBDEs from the porewater to the overlying riverwater,

419 which could explain the tight coupling of HOC concentrations and profiles in porewater and
420 riverwater (Figure 1). Lower log BAFs^{RW} were generally observed for BDE-99 than for the tri-
421 through hepta-brominated congeners (Figure 3A) possibly due to its biotransformation
422 (metabolism) in fish species (Hu et al. 2010). Similar findings were previously observed in aquatic
423 species collected from China (Wu et al. 2008) and in Lake Michigan's trout (Streets et al. 2006).
424 BDE-183 also showed lower BAF values compared to penta- and hexa-congeners, probably due
425 to the limited uptake induced by its large molecular size. However, calculated log BAFs^{RW} and
426 BAFs^{PW} for the majority of the fish samples were higher than their corresponding log K_{OWs} (Figure
427 3) and log K_{lip-w} (Figure S5) indicating a possible dietary uptake. We argue that the exceedance of
428 K_{OW}-predicted BAFs^{RW} and BAFs^{PW} is the result of the influence of the porewater on riverwater
429 (via diffusive fluxes).

430
431 Calculated BAFs for PFAA increased significantly with increasing log K_{OW} (Figure 3C) and alkyl
432 chain length (Figure 4D; $R^2 = 0.60 - 0.65$; $SE = 0.42 - 0.45$; $p < 0.01$). Log BAF of the longer
433 chain PFUNDA, PFDODA and PFOS (average: 4.1, 4.4 and 3.7 respectively) were generally
434 higher than the shorter chain PFAAs. Additionally, values for PFOS were higher than all the
435 PFAAs other than the above-mentioned long chain ones indicating a possible higher
436 bioaccumulation potential for PFOS. Based on the TSCA bioaccumulation criteria (Hong et al.
437 2015), only the longer chain PFAAs and PFOS appear to be bioaccumulative to very
438 bioaccumulative in the fish species. ($3 \leq \log BAF < 5$), which agrees well with other studies (Hong
439 et al. 2015).

440

441 *Biomagnification of pollutants in the food web*

442 All the HOCs (DDTs, HCB, oxychlordane, BDE-47, 100, 154, some PCB and PCDD/F congeners)
443 that showed significant trophic magnification potential ($TMF > 1$, $p < 0.05$) were the analytes that
444 exceeded the equilibrium porewater benchmark value set by Jahnke et al. (2014) indicating the
445 importance of the dietary intake and strong benthic coupling in the Passaic River. All the other
446 investigated OCPs and PBDEs showed either insignificant trophic dilution (γ - and δ -HCH) or
447 trophic magnification. For PFAAs (Table S22), significant trophic magnification was only
448 observed for PFUnDA (1.9), PFOS (1.8), PFDA (1.6) and PFDoA (1.5), as was observed in
449 previous studies (Fang et al. 2014; Xu et al. 2014). However, calculated TMFs for PFAAs were
450 generally lower than values calculated for OCPs and PBDEs.

451
452 For PFCAs, TMF values were relatively stable for $C_5 - C_8$ perfluorinated acids, then increased for
453 $C_8 - C_{10}$, but decreased again for PFDoA, which is probably due its molecular size, limiting its
454 uptake. For PFSAs, PFOS showed the highest TMF value among all the PFAAs, which agrees
455 well with previous findings in other food webs worldwide (Conder et al. 2008).

456

457 *Role of metabolic transformation*

458 To investigate the relationships between hydrophobicity, metabolic transformation (k_{mtr}) and
459 TMFs, a conditional probability table was computed using TMFs and presented graphically in
460 Figure (4) based on cumulative frequencies. The figure indicated that the maximum probability of
461 obtaining a $TMF > 1$ were observed for HOCs having $\log K_{ow} > 7.5$ and characterized by low
462 metabolic transformation rates ($k_{mtr} \leq 0.01 \text{ day}^{-1}$). Starting from $\log K_{ow} > 6.5$, high probability
463 rates of $TMFs > 1$ were observed. This range included dioxins, PCBs, PBDEs, OCPs and the longer
464 alkyl chain PFAAs ($> C_8$). Exceptions were observed at lower $\log K_{ows}$ (4.5 – 6.5) for some OCPs
465 and higher molecular weight PAHs, but still at low metabolic transformation rates ($\log k_{mtr} \leq 1.25$

466 equivalent to $k_{\text{mtr}} \leq 0.06 \text{ day}^{-1}$) indicating that at these lower K_{OW} ranges, there is an increase in
467 the importance of k_{mtr} in controlling trophic magnification. Additionally, calculated TMFs for PCB
468 189, 195, 206 and 209 (1.4 – 1.97) were higher than TMF values calculated for OCDD, OCDF
469 and BDE-183 (0.70 – 1.72) although all have $\log K_{\text{OW}}$ values > 8.0 (8.02 – 8.79). This can be
470 explained knowing that the latter group has higher biotransformation rates ($\log k_{\text{mtr}}$ s: -1.96 to -
471 1.46) than the former group ($\log k_{\text{mtr}}$ s: -3.04 to -3.30). From all the investigated HOCs with TMFs
472 > 1 ($n = 59$), only four had $\log k_{\text{mtr}} > -1.5$ ($K_{\text{mtr}} > 0.032 \text{ day}^{-1}$). All these findings suggest that k_{mtr}
473 values are probably more important than hydrophobicity in explaining the biomagnification of
474 HOCs. The uncertainty related to the estimated k_{mtr} values could be substantial, however.

475 476 **CONCLUSIONS**

477 Although banned decades ago, the contamination by legacy pollutants is widespread in the aquatic
478 environment of the lower Passaic River. Our results indicate that $C_{\text{PE(PW)}}$ and $C_{\text{PE(RW)}}$ of HOCs and
479 - for some HOCs and PFAAs - $C_{\text{sed(OC)}}$ can be used to estimate C_{lip} in biota. In contrast to previous
480 results, though, measured C_{lip} in the current study were above the porewater equilibrium
481 benchmark concentration (0.5-2). Our results imply that in systems with strong benthic coupling,
482 bioaccumulation needs to be considered to predict HOC and PFAA concentrations in top predators.
483 Measured tissue concentrations of PFOA, PFNA, PFUnDA, PFDoA and PFOS were 1.4 – 2.7
484 times higher than predicted tissue concentrations from sediment OC. This agrees well with their
485 bioaccumulative nature as suggested by TSCA bioaccumulation criteria (Hong et al. 2015) and
486 suggests dietary uptake. Both hydrophobicity and metabolic transformations are needed to assess
487 the biomagnification potential for both the lipophilic contaminants and PFAAs. Future work
488 should confirm how useful chemical activities in sediment are for being able to predict the
489 chemicals' body burden in a given foodweb. This will always depend on the specifics of the

490 ecosystem under consideration, including variations in the environmental conditions, foodweb
491 characteristics and the physicochemical properties of sediments in the area under investigation.
492 Yet our study indicated that porewater concentrations from passive sampling for HOCs and/or
493 sediment geochemistry for PDBEs and PFAAs were good predictors of lipid-normalized
494 concentrations. There are additional factors, though, that will influence this predictive ability
495 between sites and methods: first, variabilities arising from using different passive samplers and/or
496 using unstandardized methods for the same passive sampler (Jonker et al. 2018); second,
497 uncertainties associated with the use of partitioning coefficients in particular for compounds such
498 as PFAAs that bind to proteins; and finally, expected differences from estimating porewater
499 concentrations using *in situ* vs *ex situ* samplers. Accordingly, these factors should be addressed in
500 future research.

501

502 **SUPPLEMENTAL DATA**

503 Supplemental Data on sampling, chemical analysis, physicochemical properties, spatial
504 distribution, BAFs and TMFs are available on the Wiley Online Library at DOI:
505 10.1002/etc.xxxx.

506

507 **ACKNOWLEDGMENT**

508 We acknowledge the Hudson River Foundation for funding this project (Hudson River Award #
509 HRF 2011-5), partial support by the National Institute of Environmental Health Sciences grants
510 P42ES027706 and SERDP grant ER 2538, and thank M. Weinstein and K. Barrett (Manhattan
511 College) for sampling support.

512

513 **Data availability statement:**

514 Data pertaining to this manuscript are available in the Supplemental Data.

515

516 REFERENCES

- 517 Becker AM, Gerstmann S, Frank H. 2010. Perfluorooctanoic Acid and Perfluorooctane
518 Sulfonate in Two Fish Species Collected from the Roter Main River, Bayreuth, Germany.
519 Bull Environ Contam Toxicol. 84(1):132–135. doi:10.1007/s00128-009-9896-0. [accessed
520 2016 Jun 14]. <http://link.springer.com/10.1007/s00128-009-9896-0>.
- 521 Borgå K, Kidd KA, Muir DCG, Berglund O, Conder JM, Gobas FAPC, Kucklick J, Malm O,
522 Powell DE. 2012. Trophic magnification factors: considerations of ecology, ecosystems, and
523 study design. Integr Environ Assess Manag. 8(1):64–84.
- 524 Chaoyang Yue, Loretta Y. Li. 2013. Filling the gap: Estimating physicochemical properties
525 of the full array of polybrominated diphenyl ethers (PBDEs). Environ Pollut. 180:312–323.
- 526 Conder JM, Hoke RA, Wolf W de, Russell MH, Buck RC. 2008. Are PFCAs
527 Bioaccumulative? A Critical Review and Comparison with Regulatory Criteria and Persistent
528 Lipophilic Compounds. Environ Sci Technol. 42(4):995–1003. doi:10.1021/es070895g.
529 [accessed 2016 Jun 14]. <http://pubs.acs.org/doi/abs/10.1021/es070895g>.
- 530 Endo S, Escher B, Goss K. 2011. Capacities of membrane lipids to accumulate neutral
531 organic chemicals. Environ Sci Technol. 45(14):5912–5921. [accessed 2016 May 3].
532 <http://pubs.acs.org/doi/abs/10.1021/es200855w>.
- 533 EPA US. 2011. Draft Procedure for Analysis of Perfluorinated Carboxylic Acids and
534 Sulfonic Acids in Sewage Sludge and Biosolids by HPLC/MS/MS.
- 535 Fang S, Chen X, Zhao S, Zhang Y, Jiang W, Yang L, Zhu L. 2014. Trophic magnification
536 and isomer fractionation of perfluoroalkyl substances in the food web of Taihu Lake, China.
537 Environ Sci Technol. 48(4):2173–2182.
- 538 Fernandez LA, MacFarlane JK, Tcaciuc AP, Gschwend PM. 2009. Measurement of Freely
539 Dissolved PAH Concentrations in Sediment Beds Using Passive Sampling with Low-Density
540 Polyethylene Strips. Environ Sci Technol. 43(5):1430–1436. doi:10.1021/es802288w.
- 541 Friedman CL, Lohmann R. 2014. Comparing sediment equilibrium partitioning and passive
542 sampling techniques to estimate benthic biota PCDD/F concentrations in Newark Bay, New
543 Jersey (U.S.A.). Environ Pollut. 186:172–9. doi:10.1016/j.envpol.2013.12.002. [accessed
544 2016 May 31]. <http://www.sciencedirect.com/science/article/pii/S0269749113006246>.
- 545 Ghosh U, Kane Driscoll S, Burgess RM, Jonker MTO, Reible D, Gobas F, Choi Y, Apitz SE,
546 Maruya KA, Gala WR, et al. 2014. Passive sampling methods for contaminated sediments:
547 Practical guidance for selection, calibration, and implementation. Integr Environ Assess
548 Manag. 10(2):210–223. doi:10.1002/ieam.1507.
- 549 Gobas FAPC, de Wolf W, Burkhard LP, Verbruggen E, Plotzke K. 2009. Revisiting
550 bioaccumulation criteria for POPs and PBT assessments. Integr Environ Assess Manag.
551 5(4):624–637.

552 Gschwend PM, MacFarlane JK, Reible DD, Lu X, Hawthorne SB, Nakles D V, Thompson T.
553 2011. Comparison of polymeric samplers for accurately assessing PCBs in pore waters.
554 *Environ Toxicol Chem.* 30(6):1288–1296. doi:10.1002/etc.510.

555 Gustafsson Ö, Haghseta F, Chan C, MacFarlane J, Gschwend PM. 1997. Quantification of
556 the Dilute Sedimentary Soot Phase: Implications for PAH Speciation and Bioavailability.
557 *Environ Sci Technol.* 31(1):203–209. doi:10.1021/es960317s. [accessed 2016 Mar 4].
558 <http://dx.doi.org/10.1021/es960317s>.

559 Higgins CP, Luthy RG. 2007. Modeling Sorption of Anionic Surfactants onto Sediment
560 Materials: An a priori Approach for Perfluoroalkyl Surfactants and Linear Alkylbenzene
561 Sulfonates. *Environ Sci Technol.* 41(9):3254–3261. doi:10.1021/es062449j.

562 Hong S, Khim JS, Wang T, Naile JE, Park J, Kwon B-O, Song SJ, Ryu J, Codling G, Jones
563 PD, et al. 2015. Bioaccumulation characteristics of perfluoroalkyl acids (PFAAs) in coastal
564 organisms from the west coast of South Korea. *Chemosphere.* 129:157–163.
565 doi:<https://doi.org/10.1016/j.chemosphere.2014.06.023>.

566 Hu G, Dai J, Xu Z, Luo X, Cao H, Wang J, Mai B, Xu M. 2010. Bioaccumulation behavior
567 of polybrominated diphenyl ethers (PBDEs) in the freshwater food chain of Baiyangdian
568 Lake, North China. *Environ Int.* 36(4):309–315. doi:10.1016/j.envint.2010.01.002.

569 Jahnke A, MacLeod M, Wickström H, Mayer P. 2014. Equilibrium Sampling to Determine
570 the Thermodynamic Potential for Bioaccumulation of Persistent Organic Pollutants from
571 Sediment. *Environ Sci Technol.* 48(19):11352–11359. doi:10.1021/es503336w.

572 Kelly BC, Ikonomidou MG, Blair JD, Surridge B, Hoover D, Grace R, Gobas FAPC. 2009.
573 Perfluoroalkyl contaminants in an Arctic marine food web: trophic magnification and
574 wildlife exposure. *Environ Sci Technol.* 43(11):4037–4043.

575 Key BD, Howell RD, Criddle CS. 1997. Fluorinated Organics in the Biosphere. *Environ Sci*
576 *Technol.* 31(9):2445–2454. doi:10.1021/es961007c. [accessed 2016 Jun 13].
577 <http://pubs.acs.org/doi/abs/10.1021/es961007c>.

578 Khairy MA, Barrett K, Lohmann R. 2015. The changing sources of polychlorinated dibenzo-
579 p-dioxins and furans in sediments from the lower Passaic River and Newark Bay, New
580 Jersey, USA. *Environ Toxicol Chem.* 35(3):in review. doi:10.1002/etc.3223.

581 Khairy MA, Lohmann R. 2017. Using Polyethylene Passive Samplers to Study the
582 Partitioning and Fluxes of Polybrominated Diphenyl Ethers in an Urban River. *Environ Sci*
583 *Technol.* 51(16). doi:10.1021/acs.est.7b02418.

584 Khairy MA, Weinstein MP, Lohmann R. 2014. Trophodynamic behavior of hydrophobic
585 organic contaminants in the aquatic food web of a tidal river. *Environ Sci Technol.* 48(21).
586 doi:10.1021/es502886n.

587 Kwadijk CJAF, Korytár P, Koelmans AA. 2010. Distribution of Perfluorinated Compounds
588 in Aquatic Systems in The Netherlands. *Environ Sci Technol.* 44(10):3746–3751.
589 doi:10.1021/es100485e.

590 Labadie P, Chevreuil M. 2011. Partitioning behaviour of perfluorinated alkyl contaminants
591 between water, sediment and fish in the Orge River (nearby Paris, France). *Environ Pollut.*

592 159(2):391–397. doi:10.1016/j.envpol.2010.10.039.

593 Lau C, Anitole K, Hodes C, Lai D, Pfahles-Hutchens A, Seed J. 2007. Perfluoroalkyl acids: a
594 review of monitoring and toxicological findings. *Toxicol Sci.* 99(2):366–394.

595 Lohmann R. 2012. Critical Review of Low-Density Polyethylene's Partitioning and
596 Diffusion Coefficients for Trace Organic Contaminants and Implications for Its Use As a
597 Passive Sampler. *Environ Sci Technol.* 46:606–618.

598 Lohmann R, MacFarlane JK, Gschwend PM. 2005. Importance of Black Carbon to Sorption
599 of Native PAHs, PCBs, and PCDDs in Boston and New York Harbor Sediments. *Environ Sci*
600 *Technol.* 39(1):141–148. doi:10.1021/es049424+.

601 Loi EIH, Yeung LWY, Taniyasu S, Lam PKS, Kannan K, Yamashita N. 2011. Trophic
602 magnification of poly-and perfluorinated compounds in a subtropical food web. *Environ Sci*
603 *Technol.* 45(13):5506–5513.

604 Schenker U, MacLeod M, Scheringer M, Hungerbühler K. 2005. Improving Data Quality for
605 Environmental Fate Models: A Least-Squares Adjustment Procedure for Harmonizing
606 Physicochemical Properties of Organic Compounds. *Environ Sci Technol.* 39(21):8434–
607 8441. doi:10.1021/es0502526. [accessed 2016 May 5]. <http://dx.doi.org/10.1021/es0502526>.

608 Schwarzenbach RP, Gschwend PM, Imboden DM. 2005. *Environmental Organic Chemistry*.
609 John Wiley & Sons. [accessed 2016 May 30].
610 <https://books.google.com/books?hl=en&lr=&id=77ShpUHTZCYC&pgis=1>.

611 Shoeib M, Vlahos P, Harner T, Peters A, Graustein M, Narayan J. 2010. Survey of
612 polyfluorinated chemicals (PFCs) in the atmosphere over the northeast Atlantic Ocean.
613 *Atmos Environ.* 44(24):2887–2893. doi:10.1016/j.atmosenv.2010.04.056.

614 Streets SS, Henderson SA, Stoner AD, Carlson DL, Simcik MF, Swackhamer DL. 2006.
615 Partitioning and Bioaccumulation of PBDEs and PCBs in Lake Michigan †. *Environ Sci*
616 *Technol.* 40(23):7263–7269. doi:10.1021/es061337p. [accessed 2016 Jun 13].
617 <http://pubs.acs.org/doi/abs/10.1021/es061337p>.

618 Di Toro DM, Zarba CS, Hansen DJ, Berry WJ, Swartz RC, Cowan CE, Pavlou SP, Allen HE,
619 Thomas NA, Paquin PR. 1991. Technical basis for establishing sediment quality criteria for
620 nonionic organic chemicals using equilibrium partitioning. *Environ Toxicol Chem.*
621 10(12):1541–1583. doi:10.1002/etc.5620101203.

622 US EPA. 1992a. National study of chemical residues in fish: volume I. doi:EPA 823-R-92-
623 008a.

624 US EPA. 1992b. National study of chemical residues in fish: volume II. doi:EPA 823-R-92-
625 008b.

626 US EPA. 2016a. Drinking water health advisory for PFOA and PFOS. :103. doi:EPA 822-R-
627 16-005.

628 US EPA. 2016b. Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.11.

629 Wang D-Q, Yu Y-X, Zhang X-Y, Zhang S-H, Pang Y-P, Zhang X-L, Yu Z-Q, Wu M-H, Fu

630 J-M. 2012. Polycyclic aromatic hydrocarbons and organochlorine pesticides in fish from
631 Taihu Lake: Their levels, sources, and biomagnification. *Ecotoxicol Environ Saf.* 82:63–70.
632 doi:10.1016/j.ecoenv.2012.05.010.

633 Wang Z, MacLeod M, Cousins IT, Scheringer M, Hungerbühler K. 2011. Using
634 COSMOtherm to predict physicochemical properties of poly- and perfluorinated alkyl
635 substances (PFASs). *Environ Chem.* 8(4):389–398.

636 Wirgin I, Roy NK, Loftus M, Chambers RC, Franks DG, Hahn ME. 2011. Mechanistic Basis
637 of Resistance to PCBs in Atlantic Tomcod from the Hudson River. *Science* (80-).
638 331(6022):1322 LP-1325.

639 Wu J-P, Luo X-J, Zhang Y, Luo Y, Chen S-J, Mai B-X, Yang Z-Y. 2008. Bioaccumulation
640 of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) in wild
641 aquatic species from an electronic waste (e-waste) recycling site in South China. *Environ Int.*
642 34(8):1109–13. doi:10.1016/j.envint.2008.04.001. [accessed 2016 Jun 2].
643 <http://www.sciencedirect.com/science/article/pii/S0160412008000640>.

644 Xia GS. 1998. Sorption behavior of nonpolar organic chemicals on natural sorbents (PhD
645 dissertation). John Hopkins University, Baltimore MD. John Hopkins University.

646 Xu J, Guo C-S, Zhang Y, Meng W. 2014. Bioaccumulation and trophic transfer of
647 perfluorinated compounds in a eutrophic freshwater food web. *Environ Pollut.* 184:254–61.
648 doi:10.1016/j.envpol.2013.09.011. [accessed 2016 Apr 25].
649 <http://www.sciencedirect.com/science/article/pii/S026974911300482X>.

650 You J, Landrum PF, Lydy MJ. 2006. Comparison of Chemical Approaches for Assessing
651 Bioavailability of Sediment-Associated Contaminants. *Environ Sci Technol.* 40(20):6348–
652 6353. doi:10.1021/es060830y.

653 Young WM, South P, Begley TH, Noonan GO. 2013. Determination of Perfluorochemicals
654 in Fish and Shellfish Using Liquid Chromatography–Tandem Mass Spectrometry. *J Agric*
655 *Food Chem.* 61(46):11166–11172.

656

657

658 **Figure captions**

659

660 **Figure 1: Concentrations of OCPs, PBDEs and PFAAs in fish/blue crabs (A, D, G), sediments**
661 **(B, E, H) and porewater/riverwater (C, F, I) of the lower Passaic River.**

662 **Figure 2: Measured versus model-predicted lipid-based concentrations of OCPs, PCBs,**
663 **PCDD/Fs, PFAAs and PBDEs in fish/blue crabs based on partitioning from (A)**
664 **sediment OC, (B) sediment OC + BC, (C) porewater, and (D) riverwater in the**
665 **lower Passaic River. Points represent the average values for all fish/blue crabs;**
666 **error bars represent the standard deviation; dashed red lines represent the 95 %**
667 **prediction intervals.**

668 **Figure 3: Linear regression relationships between log K_{ow} and log BAFs calculated for**
669 **OCPs and PBDEs based on riverwater (A), porewater (B), for PFAAs based on**
670 **riverwater (C), and log BAFs against number of fluorinated carbons for PFAAs**
671 **(D). TSCA: toxic substance control act.**

672 **Figure 4: 3D plot showing the probability of having a trophic magnification factor (TMF)**
673 **value > 1 based on the values of both metabolism (log k_{mtr}) and octanol-water**
674 **partitioning (log K_{ow}). Blue color represents the greatest probability (100 %) and**
675 **the dark orange color represent the least probability.**

676