

University of Rhode Island

DigitalCommons@URI

---

Senior Honors Projects

Honors Program at the University of Rhode  
Island

---

5-2021

## Dialectical Behavior Therapy for Dual Diagnosis: Treating Adolescent Substance Use Disorder and Co-Occurring Mood Disorders

Melody R. Gately

University of Rhode Island, melodygately3@gmail.com

Follow this and additional works at: <https://digitalcommons.uri.edu/srhonorsprog>



Part of the [Mental Disorders Commons](#), [Psychiatric and Mental Health Commons](#), and the [Substance Abuse and Addiction Commons](#)

---

### Recommended Citation

Gately, Melody R., "Dialectical Behavior Therapy for Dual Diagnosis: Treating Adolescent Substance Use Disorder and Co-Occurring Mood Disorders" (2021). *Senior Honors Projects*. Paper 897.

<https://digitalcommons.uri.edu/srhonorsprog/897><https://digitalcommons.uri.edu/srhonorsprog/897>

This Article is brought to you for free and open access by the Honors Program at the University of Rhode Island at DigitalCommons@URI. It has been accepted for inclusion in Senior Honors Projects by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

# MELODY GATELY (Psychology)

## Dialectical Behavior Therapy for Dual Diagnosis: Treating Adolescent Substance Use Disorder and Co-Occurring Mood Disorders

Sponsor: Rebecca Lebeau (Psychology)

Dialectical Behavior Therapy (DBT) is a comprehensive, evidence-based treatment method that was originally established to treat individuals with Borderline Personality Disorder (BPD; Bedics, 2020). Research has since provided evidence that DBT can be useful for treatment of mood disorders, suicidality, and behavioral health issues such as self-injury and Substance Use Disorder (SUD; Bedics, 2020). Based on these findings, DBT may be a promising treatment option for a variety of mental health conditions. This paper will review the literature and synthesize existing evidence in order to evaluate the potential of DBT as a successful treatment for adolescents diagnosed with SUD and a co-occurring Mood Disorder. It will also act as a foundation for the design of a therapeutic workbook that focuses on helping adolescents with this form of dual diagnosis to incorporate DBT into their lives. Research shows that there are notable similarities between individuals with BPD and typical adolescent behavior (Aguirre, 2014). Both populations appear to struggle with emotion regulation and impulsivity (Aguirre, 2014). Additionally, both groups have been associated with being at an increased risk for developing SUD when compared to the general population (Aguirre, 2014). These characteristic similarities among adolescence and BPD symptomatology indicate that DBT could be effective for adolescents as it has been for BPD. Further, research has shown DBT to have biological and psychological impacts on emotional regulation (Bedics, 2020). This, coupled with research showing that more than half of adolescents in community SUD treatment programs also meet diagnostic criteria for another mental health disorder, make DBT a strong choice for intervention in this dually diagnosed population (NIDA, 2021). Surprisingly, even with the research literature suggesting that many adolescents are dually diagnosed, less than one quarter of SUD programs, and even fewer mental health programs, actually meet criteria for services that are capable of treating those who are dually diagnosed (NIDA, 2021). This indicates a drastic need within this population for integrated treatment options that are adequate and effective.