

2021

Cryptic metabolisms in anoxic subseafloor sediment

Arkadiy I. Garber

Gustavo A. Ramírez
University of Rhode Island

Sean M. McAllister

William Orsi

Steven D'Hondt
University of Rhode Island, dhondt@uri.edu

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

Citation/Publisher Attribution

Garber, A. I., Ramírez, G. A., McAllister, S. M., Orsi, W., & D'Hondt, S. (2021). Cryptic metabolisms in anoxic subseafloor sediment. *Environmental Microbiology Reports*, 13(5), 696-701. <https://doi.org/10.1111/1758-2229.12983>

Available at: <https://doi.org/10.1111/1758-2229.12983>

This Report is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons-group@uri.edu.

Cryptic metabolisms in anoxic subseafloor sediment

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Brief Report

Cryptic metabolisms in anoxic subseafloor sediment

Arkadiy I. Garber, ^{1†} Gustavo A. Ramírez, ^{2,3,4†*}
Sean M. McAllister, ^{5,6} William Orsi ⁷ and
Steven D'Hondt ²

¹School of Life Sciences, Arizona State University, Tempe, AZ.

²Graduate School of Oceanography, University of Rhode Island, Narragansett, RI.

³Department of Marine Sciences, University of North Carolina, Chapel Hill, NC.

⁴College of Veterinary Medicine, Western University of Health Sciences, Pomona, CA.

⁵Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, WA.

⁶Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, Seattle, WA.

⁷Department of Earth & Environmental Sciences, Paleontology and Geobiology, Ludwig-Maximilians-Universität München, Munich, 80333, Germany.

Abstract

Microbial gene expression in anoxic subseafloor sediment was recently explored in the Baltic Sea and the Peru Margin. Our analysis of these data reveals diverse transcripts encoding proteins associated with neutralization of reactive oxygen species, including catalase, which may provide an *in situ* source of oxygen. We also detect transcripts associated with oxidation of iron and sulfur, and with reduction of arsenate, selenate and nitrate. Given limited input of electron acceptors from outside the system, these results suggest that the microbial communities use an unexpectedly diverse variety of electron acceptors. Products of water radiolysis and their interactions with sediment continuously provide diverse electron acceptors and hydrogen. Cryptic microbial utilization of these oxidized substrates and H₂ may be an important mechanism for multi-million-year survival under the extreme energy limitation in subseafloor sediment.

Received 8 March, 2021; accepted 6 June, 2021. *For correspondence. E-mail garamirez@uri.edu. †Both authors contributed equally and are considered co-first authors.

Introduction

Marine sediment is estimated to host a third of Earth's Bacteria and Archaea (Kallmeyer *et al.*, 2012; Bar-On *et al.*, 2018). Although cells are abundant, mean per-cell rates of metabolic activity are generally very low in subseafloor sediment [>1 meter below seafloor (mbsf)] (D'Hondt *et al.*, 2002; Jørgensen, 2011; Lomstein *et al.*, 2012). Despite their low metabolic rates, cells inhabiting subseafloor sediment are generally alive (Schippers *et al.*, 2005; Ramírez *et al.*, 2018) and capable of growth (Morono *et al.*, 2011).

In anoxic subseafloor sediment along continental margins, sulfate reduction, fermentation, and methanogenesis, fueled primarily by organic matter oxidation, are the dominant net metabolic activities (D'Hondt *et al.*, 2002). Along continental margins, where rates of anaerobic activity are relatively high, detectable dissolved sulfate and net sulfate reduction are typically limited to sediment within and above the sulfate–methane transition zone (SMTZ) (D'Hondt *et al.*, 2002; Orsi *et al.*, 2013). Below this zone, dissolved sulfate is usually below detection and net respiration is primarily hydrogenotrophic methanogenesis (D'Hondt *et al.*, 2002).

The energetic processes that enable long-term microbial survival in deeply buried marine sediment are not fully understood. Long-term survival in this habitat may be influenced by cryptic energy-yielding activities that are not represented by net activities traditionally recognized from dissolved chemical profiles. For example, water radiolysis is a ubiquitous process in marine sediment (Sauvage *et al.*, 2021). Water radiolysis, and the interaction of its products with sediment, generates diverse biologically accessible chemicals (Sauvage *et al.*, 2021), leading us to hypothesize that radiolytic products may be used by sedimentary microbes for energy. To explore this possibility, we analysed the only publicly available metatranscriptomes from deep (5–159 mbsf range) anoxic sediment, sampled from the Peru Margin (Orsi *et al.*, 2013) and the Baltic Sea (Zinke *et al.*, 2017; Zinke *et al.*, 2019). Using a custom set of hidden Markov models [HMMs (Johnson *et al.*, 2010)], we targeted genes associated with reactive oxygen

species neutralization and energy-yielding reactions in deeply buried, energy-starved, anoxic sedimentary environments.

Results

Geochemical profiles

Geochemical profiles show that all sites in this study are anoxic (Fig. 1). Detailed geochemical descriptions are found elsewhere (Shipboard Scientific Party, 2003; Expedition 347 Science Party, 2015a; Expedition 347 Science Party, 2015b). Briefly, at Baltic Sea Integrated Ocean Drilling Program Sites M0059 and M0063 (Fig. 1A and B), sulfate disappearance in the shallowest sediment (Expedition 347 Science Party, 2015a; Expedition 347 Science Party, 2015b) suggests that dissolved inorganic carbon (DIC) is the predominant net electron acceptor throughout most of the sediment column. At Peru Margin Ocean Drilling Program Site 1229 (Fig. 1C), sulfate penetrates deeper, and, at greater depths, diffuses upward from brine in deeper sediment (Shipboard Scientific Party, 2003). Sulfate is below detection and methane concentrations are high between 30 and

90 mbsf at Site 1229, indicating net methanogenesis (net DIC reduction) between those depths.

Energy-yielding reactions

We detected protein-encoding transcripts (PETs) linked to diverse energy-yielding metabolisms, derived largely from Chloroflexi and Firmicutes in the Baltic Sea subseafloor sediment and Actinobacteria and Proteobacteria in the Peru Margin subseafloor sediment (Fig. 2). Among these are PETs related to the reduction of sulfite, nitrate (as reported by Orsi *et al.*, 2013), arsenate and selenate. In the Peru Margin sediment, iron oxidation is predicted at depth from the presence of the *cyc2* gene fragments (McAllister *et al.*, 2020). Sulfur oxidation is predicted from the presence of the *sdo* [sulfur dioxygenase (Liu *et al.*, 2014)] and *sqr* [sulfide-quinone oxidoreductase (Griesbeck *et al.*, 2002)] gene transcripts in the Baltic Sea sediment (12–42 mbsf) and in all but one sample (5 mbsf) from the Peru Margin. Our detection of *sdo* transcripts is consistent with recent radiotracer-based recognition of a cryptic sulfur cycle in 'sulfate-depleted' marine sediment (Jørgensen *et al.*, 2019).




Fig 1. Sulfate (cyan), methane (red) and cell counts (black) profiles for sites (A) M0059E, (B) M0063E and (C) 1229D. Data from Expedition 347 Science Party (2015a), Expedition 347 Science Party (2015b) and Shipboard Scientific Party (2003).




Fig 2. Dot plots showing lineage-specific expression levels of PETs related to catabolic metabolisms and ROS neutralization. Transcripts from different taxonomic groups are colour-coded at the phylum level (Class level for Proteobacteria). Sites are depicted by depth in meters below the seafloor (mbsf) and colour coded by sulfate presence (orange) or depletion (purple). The dot sizes represent transcript copies, which represent the number of gene variants of each gene detected in the metatranscriptomes.

Reactive oxygen species neutralization

We detected PETs for neutralization of ROS in all surveyed metatranscriptomes (Fig. 2). PETs were identified for diverse ROS neutralization genes, including peroxidoreductin, hydroperoxide reductase and superoxide dismutase (Fig. 2). PETs homologous to the dye-decolorizing peroxidase (Dyp), which performs hydrogen peroxide-dependent oxidation of recalcitrant carbon polymers such as lignin (Chen *et al.*, 2015), and heme peroxidase (Hansel *et al.*, 2012), implicated in indirect extracellular manganese oxidation by *Roseobacter* spp. (Andeer *et al.*, 2015), were also detected. Predicted taxonomic affiliations of transcripts indicate lineage-specific expression of ROS-related PETs (Fig. 2). Firmicutes, Chloroflexi and Proteobacteria drive expression patterns. Cyanobacteria, Chlorobi, Bacteroidetes, Aquificae and various Archaeal PETs are present but rare. Baltic Sea ROS-related PETs belong to Firmicutes, Chloroflexi, and, to a smaller degree, Proteobacteria. In the Peru Margin, catalase transcripts are derived mostly from Proteobacteria.

Discussion

Catabolic activities

PETs related to the oxidation of iron and sulfur were detected in our survey (Fig. 2). In principle, sulfur can be oxidized with manganese or iron oxides as the electron

acceptor (Jørgensen *et al.*, 2019). In contrast, energy-yielding microbial oxidation of iron typically requires highly oxidized electron acceptors, such as oxygen, hydrogen peroxide, superoxide and nitrate (Jørgensen *et al.*, 2020). However, all of these samples are from much greater sediment depths than oxygen and nitrate appear to penetrate from the overlying ocean (D'Hondt *et al.*, 2004). And while oxidized manganese can also serve as an electron acceptor for oxidizing iron, there is no evidence, as far as we know, that Mn(IV) and Fe(II) can be metabolically coupled by microorganisms. We also found *nar* transcripts exclusively in the 91-mbsf Peru Margin sample, the approximate depth where downward-diffusing methane is intercepted by sulfate diffusing up from brine below (D'Hondt *et al.*, 2004), corroborating a previous report (Orsi *et al.*, 2013). Overall, our survey of catabolic metabolisms suggests the presence of a cryptic source of oxidizing power in deep anoxic sediment.

ROS-related activities

The presence of ROS-related PETs in deep sediment also indicates widespread neutralization and use of ROS in this system. The detection of transcripts encoding dye-decolorizing peroxidase suggests the presence and microbial use of H₂O₂ to oxidize substrates and degrade recalcitrant organic polymers (Chen *et al.*, 2015). Moreover, heme peroxidases, thought to be secreted from

cells, react with H_2O_2 and produce superoxide, which can then oxidize Mn(II) to Mn(IV). Interestingly, fungal peroxidases, including manganese and heme peroxidases, are also thought to result in extracellular manganese oxidation (Hansel *et al.*, 2012). Our observations, despite the known limitations of transcriptional surveys (see Notes on transcriptomic interpretation in the SI), support that H_2O_2 and perhaps other, more transient, ROS are present, at least ephemerally, in this ecosystem.

Potential source of oxidants

We report the presence of PETs associated with neutralization of ROS, as well as PETs associated with reduction of arsenate, selenate and nitrate in deep subseafloor sediment, where DIC and sulfate are typically assumed to be the primary terminal electron acceptors. PETs associated with oxidation of sulfur and iron are also expressed, indicating that these communities are primed to utilize these energy-generating pathways. We have considered whether these results could reflect signatures left over from the oxygenated world (i.e. constitutive functional gene expression rather than a true physiological response to the environment). Constitutive expression of functional genes from early in the sediment history might conceivably occur if microbes persist from that time without reproduction (D'Hondt *et al.*, 2014; Kirkpatrick *et al.*, 2019). However, recent studies have identified active cell division by dominant lineages of anaerobic subseafloor microbes (Atribacteria and Chloroflexi) in subseafloor sediment as old as 8 million years (Vuillemin *et al.*, 2020a; Vuillemin *et al.*, 2020b). In actively reproducing cells, the transcriptomic profiles change as net growth increases (Vuillemin *et al.*, 2020a, Vuillemin *et al.*, 2020b) and constitutive expression of functional genes should be a minor component of the transcriptome compared to the transcriptome that is activated as a part of the true physiological response to the environment. In accordance with this line of reasoning, residual transcription of oxygen utilization genes (e.g. terminal oxidases) is not observed in our analysis, indicating that transcripts not useful in the anoxic sediment are either not expressed or expressed at levels below detection limit.

Our observations collectively suggest the presence of an *in situ* source of biologically available oxidizing power in deep anoxic sediment. Although redox-active antimicrobials (RAAs) might explain PETs for ROS (Imlay, 2008), the independent detection of PETs related to diverse catabolic redox activities suggests that the ROS transcriptional patterns may not be exclusively driven by RAAs. Unexpected redox processes can be directly and indirectly supported by water radiolysis, which is ubiquitous in marine sediment and other wet

geological environments (Lin *et al.*, 2006; Blair *et al.*, 2007). Water radiolysis appears to be the predominant source of electron donors (H_2) and diverse electron acceptors in marine sediment older than a few million years (Sauvage *et al.*, 2021). Reaction of oxidized radiolytic products with sedimentary minerals, as well as various catalases and peroxidases, may generate redox species that are otherwise generally absent from sediment beneath the SMTZ [e.g. hydrogen peroxide, sulfite/sulfate, iron(III)].

Concluding remarks

Our results provide evidence for potential cryptic redox cycling in deeply buried marine sediment. Using a small, highly curated database of marker genes of biogeochemical significance, we predict the occurrence of unexpectedly diverse microbial activities, including various forms of catabolic metabolism and oxidative defence, that go undetected by standard geochemical measurements in this habitat. This study greatly expands the potential metabolic diversity associated with anoxic subseafloor sediment and highlights reactions that may provide long-term (multi-million-year scale) support for microbial communities buried in deep anoxic sediment.

Materials and methods

Sample collection

Information regarding collection, handling and processing of samples for the original transcriptomic studies can be found in the following publications: Peru Margin - Orsi *et al.*, 2013; Baltic Sea - Zinke *et al.*, 2017; Zinke *et al.*, 2019 (see the Extended Methods section of the SI for additional details).

Data acquisition and processing

Metatranscriptomic data from Peru Margin and Baltic Sea marine sediment were downloaded from the Sequence Read Archive (SRA) using the *SRA Toolkit v.2.8.2* (SRA Toolkit Development Team, NCBI). Site metadata, sequence trimming, assembly and read map details are provided in the Extended Methods section of the SI.

Annotation

To identify transcripts relevant to cryptic metabolisms and ROS neutralization, we developed two novel bioinformatics tools: *LithoGenie* and *RosGenie*. These software packages and the HMM libraries employed by each tool are public: <https://github.com/Arkadiy-Garber/LithoGenie> and <https://github.com/Arkadiy-Garber/RosGenie>. Software details,

including marker gene selection, HMM calibrations and contaminant screening are provided in the Annotation section of the SI.

Acknowledgements

We thank Andreas P. Teske for insightful commentary and suggestions on an early draft of this manuscript. We also thank two anonymous reviewers for their thoughtful comments. This project was funded in part by the US National Science Foundation, through the Center for Dark Energy Biosphere Investigations (C-DEBI; grant NSF-OCE-0939564). This is C-DEBI publication 569. This is JISAO contribution no. 2020-1070; PMEL contribution no. 5104.

Funding Information

G.A.R. was funded by an NSF C-DEBI post-doctoral fellowship. S.D. and G.A.R. were funded by the U.S. National Science Foundation through grants NSF-OCE-1130735 and NSF-OCE-0939564. S.M.M. was funded by the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) under the NOAA Cooperative Agreement NA15OAR4320063.

Author Contributions

G.A.R., S.D. and A.I.G. conceived the study; A.I.G. and G.A.R. performed bioinformatics analyses; A.I.G., G.A.R., S.M.M., W.O. and S.D. interpreted data and wrote the manuscript.

References

Andeer, P.F., Learman, D.R., McIlvin, M., Dunn, J.A., and Hansel, C.M. (2015) Extracellular haem peroxidases mediate Mn(II) oxidation in a marine *Roseobacter bacterium* via superoxide production. *Environ Microbiol* **17**: 3925–3936.

Bar-On, Y., Phillips, R., and Milo, R. (2018) The biomass distribution on Earth. *Proc Natl Acad Sci U S A* **115**: 6506–6511.

Blair, C.C., D'Hondt, S., Spivack, A.J., and Kingsley, R.H. (2007) Radiolytic hydrogen and microbial respiration in subsurface sediments. *Astrobiology* **7**: 951–970.

Chen, C., Shrestha, R., Jia, K., Gao, P.F., Geisbrecht, B.V., Bossmann, S.H., et al. (2015) Characterization of dye-decolorizing peroxidase (DyP) from *Thermomonospora curvata* reveals unique catalytic properties of A-type DyPs. *J Biol Chem* **290**: 23447–23463.

D'Hondt, S., Jørgensen, B.B., Miller, D.J., Batzke, A., Blake, R., Cragg, B.A., et al. (2004) Distributions of microbial activities in deep subseafloor sediments. *Science* **306**: 2216–2221.

D'Hondt, S., Rutherford, S., and Spivack, A.J. (2002) Metabolic activity of subsurface life in deep-sea sediments. *Science* **295**: 2067–2070.

D'Hondt, S., Wang, G., and Spivack, A.J. (2014) The under-ground economy (energetic constraints of subseafloor life). Chapter 2.3 in earth and life processes discovered from subseafloor environment – a decade of science achieved by the Integrated Ocean Drilling Program (IODP). In *Series Developments in Marine Geology*. Stein, R., Blackman, D., Inagaki, F., and Larsen, H.-C. (eds). Amsterdam/New York: Elsevier, pp. 127–148.

Expedition 347 Science Party. (2015a) Site M0063. In Andrén, T., Jørgensen, B.B., Cotterill, C., Green, S., The Expedition 347 Scientists, Proc. IODP, 347: College Station, TX (Integrated Ocean Drilling Program).

Expedition 347 Science Party. (2015b) Site M0059. In Andrén, T., Jørgensen, B.B., Cotterill, C., Green, S., The Expedition 347 Scientists, Proc. IODP, 347: College Station, TX (Integrated Ocean Drilling Program).

Griesbeck, C., Schutz, M., Schodl, T., Bathe, S., Nausch, L., Mederer, N., et al. (2002) Mechanisms of sulfide-quinone reductase investigated using site-directed mutagenesis and sulfur analysis. *Biochemistry* **41**: 11552–11565.

Hansel, C.M., Zeiner, C.A., Santelli, C.M., and Webb, S.M. (2012) Mn(II) oxidation by an ascomycete fungus is linked to superoxide production during asexual reproduction. *Proc Natl Acad Sci U S A* **109**: 12621–12625.

Imlay, J.A. (2008) Cellular defenses against superoxide and hydrogen peroxide. *Annu Rev Biochem* **77**: 755–776.

Johnson, L., Eddy, S., and Portugaly, E. (2010) Hidden Markov Model speed heuristic and iterative HMM search procedure. *BMC Bioinformatics* **11**: 431.

Jørgensen, B.B. (2011) Deep subseafloor microbial cells on physiological standby. *Proc Natl Acad Sci U S A* **108**: 18193–18194.

Jørgensen, B.B., Andrén, T., and Marshall, I.P.G. (2020) Sub-seafloor biogeochemical processes and microbial life in the Baltic Sea. *Environ Microbiol* **22**: 1688–1706.

Jørgensen, B.B., Findlay, A.J., and Pellerin, A. (2019) The biogeochemical sulfur cycle of marine sediments. *Front Microbiol* **10**: 849.

Kallmeyer, J., Pockalny, R., Adhikari, R.R., Smith, D.C., and D'Hondt, S. (2012) Global distribution of microbial abundance and biomass in subseafloor sediment. *Proc Natl Acad Sci U S A* **109**: 16213–16216.

Kirkpatrick, J.B., Walsh, E.A., and D'Hondt, S. (2019) Microbial selection and survival in subseafloor sediment. *Front Microbiol* **10**: 956.

Lin, L.-H., Wang, P.-L., Rumble, D., Lippmann-Pipke, J., Boice, E., Pratt, L.M., et al. (2006) Long-term sustainability of a high-energy, low-diversity crustal biome. *Science* **314**: 479–482.

Liu, H., Xin, Y., and Xun, L. (2014) Distribution, diversity, and activities of sulfur dioxygenases in heterotrophic bacteria. *Appl Environ Microbiol* **80**: 1799–1806.

Lomstein, B.A., Langerhuus, A.T., D'Hondt, S., Jørgensen, B.B., and Spivack, A.J. (2012) Endospore abundance, microbial growth and necromass turnover in deep sub-seafloor sediment. *Nature* **484**: 101–104.

McAllister, S.M., Polson, S.W., Butterfield, D.A., Glazer, B. T., Sylvan, J.B., Chan, C.S., et al. (2020) Validating the Cyc2 neutrophilic iron oxidation pathway using metatranscriptomics of Zetaproteobacteria iron mats at marine hydrothermal vents. *mSystems* **5**: e00553-19.

- Morono, Y., Terarda, T., Nishizawa, M., Ito, M., Hillion, F., Takahata, N., *et al.* (2011) Carbon and nitrogen assimilation in deep subseafloor microbial cells. *Proc Natl Acad Sci U S A* **108**: 18295–18300.
- Orsi, W.D., Edgcomb, V.P., Christman, G.D., and Biddle, J. F. (2013) Gene expression in the deep biosphere. *Nature* **499**: 205–208.
- Ramírez, G.A., Jørgensen, S.L., Zhao, R., and D'Hondt, S. (2018) Minimal influence of extracellular DNA on molecular surveys of marine sedimentary communities. *Front Microbiol* **9**: 2969.
- Sauvage, J., Flinders, A., Spivack, A.J., Pockalny, R., Dunlea, A. G., Anderson, C.H., *et al.* (2021) The contribution of water radiolysis to marine sedimentary life. *Nat Commun* **12**: 1297.
- Schippers, A., Neretin, L.N., Kallmeyer, J., Ferdelman, T., Cragg, B.A., Parkes, R.J., *et al.* (2005) Prokaryotic cells of the deep sub-seafloor biosphere identified as living bacteria. *Nature* **433**: 861–864.
- Shipboard Scientific Party. (2003) Site 1229. In D'Hondt, S.L., Jørgensen, B.B., Miller, D.J., *et al.*, Proc. ODP, Init. Repts., 201: College Station, TX (Ocean Drilling Program), pp. 1–78.
- Vuillemin, A., Kerrigan, Z., D'Hondt, S., and Orsi, W.D. (2020a) Exploring the abundance, metabolic potential and gene expression of subseafloor Chloroflexi in million-year-old oxic and anoxic abyssal clay. *FEMS Microbiol Ecol* **96**: fiae223.
- Vuillemin, A., Vargas, S., Coskun, O.K., Pockalny, R., Murray, R.W., Smith, D.C., *et al.* (2020b) Atribacteria reproducing over millions of years in the Atlantic abyssal subseafloor. *mBio* **11**: e01937-20.
- Zinke, L.A., Glombitza, C., Bird, J.T., Røy, H., Jørgensen, B. B., Lloyd, K.G., *et al.* (2019) Microbial organic matter degradation potential in Baltic Sea sediments is influenced by depositional conditions and in situ geochemistry. *Appl Environ Microbiol* **85**: e02164–e02118.
- Zinke, L.A., Mullis, M.M., Bird, J.T., Marshall, I.P.G., Jørgensen, B.B., Lloyd, K.G., *et al.* (2017) Thriving or surviving? Evaluating active microbial guilds in Baltic Sea sediment. *Environ Microbiol Rep* **9**: 528–536.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix S1. Supporting information.